

This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + Refrain from automated querying Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at http://books.google.com/



Soc. 1996 d. 322 GenPer. 49 Ps. 1996 - d. 590 · ·

•

-

.

•

-- -

		,	
		٠	
		1	
	-		

		٠	

SMITHSONIAN

MISCELLANEOUS COLLECTIONS.

VOL. XXVII.



AND EXPERIMENTS PROGUERS KNOWLEDGE FOR MEN."—SWITHSON.

WASHINGTON:
PUBLISHED BY THE SMITHSONIAN INSTITUTION.
1883.

			•	
		,		
	•			
				-

ADVERTISEMENT.

The present series, entitled "Smithsonian Miscellaneous Collections," is intended to embrace all the publications issued directly by the Smithsonian Institution in octavo form; those in quarto constituting the "Smithsonian Contributions to Knowledge." The quarto series includes memoirs, embracing the records of extended original investigations and researches, resulting in what are believed to be new truths, and constituting positive additions to the sum of human knowledge. The octavo series is designed to contain reports on the present state of our knowledge of particular branches of science; instructions for collecting and digesting facts and materials for research; lists and synopses of species of the organic and inorganic world; museum catalogues; reports of explorations; aids to bibliographical investigations, etc., generally prepared at the express request of the Institution, and at its expense.

The assignment of a work to one or the other of the two series will sometimes depend upon whether the required illustrations can be presented more conveniently in the quarto or the octavo form.

In the Smithsonian Contributions to Knowledge, as well as in the present series, each article is separately paged and indexed, and the actual date of its publication is that given on its special title page, and not that of the volume in which it is placed. In many cases works have been published, and largely distributed, years before their combination into volumes.


While due care is taken on the part of the Smithsonian Institution to insure a proper standard of excellence in its publications, it will be readily understood that it cannot hold itself responsible for the facts and conclusions of the authors, as it is impossible in most cases to verify their statements.

S. F. BAIRD, Secretary S. I.



TABLE OF CONTENTS.

- ARTICLE I. (358.) THE CONSTANTS OF NATURE. Part IV. ATOMIC WEIGHT DETERMINATIONS: A DIGEST OF THE INVESTIGATIONS PUBLISHED SINCE 1814. By GEORGE F. BECKER. 1880. Pp. 152.
 - ARTICLE II. (441.) THE CONSTANTS OF NATURE. Part V. A RECAL-CULATION OF THE ATOMIC WEIGHTS. By FRANK WIGGLES-WORTH CLARKE. 1882. Pp. 293.
 - ARTICLE III. (437.) CHECK LIST OF PUBLICATIONS OF THE SMITHSONIAN INSTITUTION, December, 1881. 1881. Pp. 22.
 - ARTICLE IV. (478.) CATALOGUE OF PUBLICATIONS OF THE SMITHSONIAN INSTITUTION (1846–1882) WITH AN ALPHABETICAL INDEX OF ARTICLES IN THE SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE, MISCELLANEOUS COLLECTIONS, ANNUAL REPORTS, BULLETINS AND PROCEEDINGS OF THE U.S. NATIONAL MUSEUM, AND REPORT OF THE BUREAU OF ETHNOLOGY. BY WILLIAM J. RHEES. 1882. Pp. 342.



SMITHSONIAN MISCELLANEOUS COLLECTIONS.

——— 358 ——— THE

CONSTANTS OF NATURE.

PART IV.

ATOMIC WEIGHT DETERMINATIONS:

A DIGEST

OF THE INVESTIGATIONS PUBLISHED SINCE 1814.

BY

GEORGE F. BECKER.



WASHINGTON: SMITHSONIAN INSTITUTION. AUGUST, 1880.



•

ADVERTISEMENT.

The following forms the fourth part of a general work on the "Constants of Nature," of which the first three are as follows:

- Part I and Supplement.—Specific Gravities, Boiling Points and Melting Points, by F. W. Clarke.
- Part II.—A Table of Specific Heats for Solids and Liquids, by F. W. Clarke.
- Part III.—Tables of Expansion by Heat for Solids and Liquids, by F. W. Clarke.

The manuscript of the present work has been presented to the Smithsonian Institution by Mr. G. F. Becker and is published at the expense of its fund.

S. F. BAIRD, Secretary Smithsonian Institution.

WASHINGTON, AUGUST, 1880.



PREFACE.

Of the fundamental importance of the most accurate attainable knowledge concerning the true atomic weights of the elements there can be no two opinions. If the enormous mass of known facts relating to the properties of matter is ever to be brought under wide generalizations, it is with the simple substances that a beginning must be made, and with the simplest property of these substances, the relative weights of their ultimate particles. Berzelius held this view and the labors of Mendelejeff, Meyer and others leave no question as to the fact of a relation between the atomic weights and the properties of simple and compound matter. Accurate information on the subject, however, is not easily attainable; different writers on chemistry follow different authorities, and some even take a mean . between the results arrived at by experimenters of different degrees of skill and accuracy, or assume some convenient number without experimental foundation. Nowhere, to my knowledge, is there even an approximately complete list of the determinations that have been made.

Forced back, myself, upon the original memoirs for information, I believed that I should do other chemists a service in presenting to them a short but systematic digest of each investigation on the subject, including the following points, so far as they could be ascertained: The nature of the material experimented upon, and the method of its preparation; the experimental method adopted to effect the determination, and the number of experiments; the mean result reached by the experiments, and the extreme difference between the results; such a record of the constants employed in the calculation as will enable any one to recalculate the results for different constants; and the place in literature where the original paper is to be found.

The following pages are the result. From the information

he will find in them, the experienced chemist will, in most cases, I think, be able to decide which determination offers the best guarantees for accuracy, or at least between which determinations his choice must lie, forming his judgment to a great extent independently of the comparative reputation of the observers—not always a safe guide where one is, in a general way, the unquestionable superior of the other—and no guide at all when the names carry on the whole an equal weight. As a record of the direction investigations have taken and of analytical methods of the most exact character also, I hope that this digest may not be without value.

As this compilation would serve rather to mislead than to assist investigators, unless it be accurate and practically exhaustive, it seems proper to explain the manner in which it has been prepared. Believing it best to work independently of any previous compilations, I selected as my base the three great German journals—Poggendorff's Annalen, Liebig's Annalen, and Erdmann's Journal für Praktische Chemie. My choice was determined not only by the position these journals take in chemico-physical science, but by the fact that their indices are admirable, and their tone cosmopolitan; all of them, until lately, having furnished their readers with the scientific news of the time, and with abstracts from and translations of the important papers published elsewhere and in whatever language, as well as with original investigations. The indices of these journals I read through from beginning to end, making an extract of every entry which bore on the subject of atomic weights, or which I suspected might do so. In studying the articles thus reached, every reference to other atomic weight determinations was preserved, and the originals, so far as possible, sought out; a task in which the Royal Society's Catalogue of Scientific Papers was of the greatest assistance. Having exhausted the supply of information in these journals, I turned to Berzelius' Jahresbericht, and to its continuation edited by Kopp, Liebig et al., and made a study of their contents by the same method. Later, I made a similar systematic study of the Annales de Chimie et de Physique, the Bericht der Deutschen Chemischen Gesellschaft, the Chemical PREFACE. 3

News, Fresenius' Zeitschrift für Analytische Chemie, the Journal of the Chemical Society, the Proceedings of the Royal Society, and the Philosophical Transactions, and of Silliman's American Journal of Science. I have also made some use of the Philosophical Magazine, and a great deal of use of the Paris Comptes Rendus. These publications are not so indexed as to make their contents readily available; but what appears in the Comptes Rendus is pretty sure to be noticed elsewhere, and I scarcely think that any determinations there published have escaped me. I have also made use of the Bibliothèque Universelle, Archives des Sciences of Geneva, (an incomplete set, unfortunately,) the Zeitschrift für Berg-Hütten-und Salinen-Wesen im Preussischen Staate, Thomson's Annals of Philosophy, Gilbert's Annalen der Physik und der Physikalischen Chemie, the British Association Reports, the Transactions of the Royal Society of Edinburgh, the Transactions of the Academies of Brussels and of St. Petersburg, and have consulted numerous works on chemistry, particularly Berzelius' Lehrbuch der Chemie and Gmelin-Kraut's Handbuch der Chemie.

I have not thought it necessary, or even desirable, to extend my search for atomic weight determinations further back than Wollaston's famous "Table of Equivalents," published in the Philosophical Transactions for 1814. true that numerous determinations had been made before that time, but, with the exception of those mentioned by Wollaston, few which can be of either interest or value to the chemist of the present day, except from a purely historical point of view. From Wollaston's table onwards, I have not felt that the purposes of this paper permitted of any selection between atomic weight determinations, however valueless many of them might appear to my own judgment. Indeed, it has cost me more labor to put many ill-made and ill-reported investigations into proper form for this digest than was required for a majority of those determinations upon which I set the highest value. In the attempt to make a complete collection of the determinations since the time indicated, a few may have escaped my search; but, if so, they must have fallen singularly dead upon the chemical world, and would be unlikely to repay further labor in seeking them. On the other hand, I have rigidly excluded atomic weights calculated from analyses never designed so to be used. Any chemist, upon whose experiments we could rely, would proceed in a very different manner in making an atomic weight determination, from that which he would select for an ordinary analysis, and to put his credit at stake by calculating atomic weights from analyses not designed for this use is alike unfair to him and to the scientific public, which is asked to receive as an atomic weight determination what really is not such.

The purpose of this paper is distinctly not critical, and the remarks I have added to, or inserted in, the digest are simply explanatory. I have, however, frequently mentioned criticisms which have appeared in literature when they seemed pertinent.

As for the accuracy with which the digests have been made, I may state that the preponderating importance of this point has been constantly before my mind. In the effort to crowd the maximum amount of information into the fewest words, I have had occasion to refer to most of the papers digested a number of times, and at long intervals. I have always taken advantage of such occasions, as well as those on which I have met with a reprint, translation or abstract of a determination, to verify the rough draughts of my digests. Only in a couple of instances have I thus discovered a trifling error. On the other hand, I have been able to detect and point out numerous misprints and miscalculations in the original sources. While, therefore, I cannot hope entirely to have escaped error in the thousands of values I have copied, and the almost equal number of calculations I have made, I have strong hopes that the accuracy of this digest will be found at least on a par with that of the original papers.

When, as is the case with provoking frequency, chemists have given their analytical data, but have omitted to state the atomic weights, or other constants, assumed in calculating their results, I have recalculated their data with accepted constants, which I have in each case stated. I have also, in many instances, recalculated determinations of importance,

in which constants varying considerably from those now received were assumed. I have further reduced the determinations originally given in terms of O = 100, or of O = 15.96, to O = 16. No confusion, however, will be found between the numbers for which the original investigators are responsible and my own. All values which I have calculated are in italics, or, with my explanations, enclosed in square brackets. The only arithmetical operation I have permitted myself to perform without these indications is a multiplication or division by two; and even in such cases it will usually appear from the digest itself that this operation has been performed.

The abbreviations of the literary references are essentially those adopted in the Royal Society's Catalogue of Scientific Papers. The first reference in each case is to the source upon which I have depended. When two references are necessary, they are connected by the word and. When my authority is not the original source, that to which it is accredited in my authority is also mentioned.

In conclusion, I shall be grateful to any one who, by drawing my attention to omissions or mistakes, will assist me in perfecting a labor which has occupied all my available time for twenty months.

BERKELEY, CAL., April, 1878.

POSTSCRIPT.

In preparing the following paper, I designed making it preliminary to a discussion of the various determinations and of the value to be assigned to each, and in this work I had already made some progress. After presenting this paper to the Institution, however, I learned that Prof. F. W. Clarke had been for some time engaged on a similar undertaking, and to him I gladly resigned the discussion of the data here compiled. The two papers will appear in the same form, and may be regarded as complementary.

G. F. B.

·				
,				
	•			
	•	•		
				r

ATOMIC WEIGHT DETERMINATIONS.

ALUMINIUM.

The specific heat of aluminium, as determined by Regnault and by Kopp, and the vapor density of volatile compounds, as determined by Deville and Troost and by Odling, indicate that the atomic weight of this element is about 27.5. (Gmelin-Kraut, Handbuch der Chemie, 1, 39; and L. Meyer, Moderne Theorien der Chemie, 50.)

J. J. Berzelius: 27.267 (O = 16).

100 parts of anhydrous aluminic sulphate decomposed by heat, gave 29.984 parts of oxide. Preparation not described. Number of experiments, probably 1. In Berzelius' Lehrbuch these data are calculated for S=200.75, and give Al=170.9~(O=100,) or 27.344 (O=16.) [If S=32, the data give Al=27.267.] (Poggend. Ann., 8, 1826, 187.*)

T. Thomson: 30 (O = 16).

Thomson found, probably from analysis of the sulphate, (see appendix.) that 125 Al = 100 O. Thomson supposed aluminic oxide to be a protoxide. [If it is a sesqui-oxide, the data give Al at 30.] (Thomson's System of Chemistry, 7th ed., 1, 1831, 454.)

W. W. MATHER: 20.55 (O = 16).

According to this chemist 0.646 grammes of chloride, prepared according to Woehler, gave 2.055 grammes argen-

^{*}This article by Berzelius, which contains the particulars of a large part of his earlier atomic weight determinations, will be referred to frequently in the course of this paper. It is unfortunately full of misprints, all of which are, by no means, corrected in the table of errata at the end of the volume. The correctly printed values of the atomic weights discussed in it are to be found in *Poggend. Ann.*, 10, 1827, 389.

[†] It must be remarked, in justice to Dr. Thomson, that his atomic weight determinations are, properly speaking, of a different nature from those of other chemists. So thoroughly persuaded was he of the truth of Prout's hypothesis, (that the atomic weights of the elements are all exact multiples of that of hydrogen,) that his experiments were directed merely towards ascertaining which multiple, in any case, was to be adopted.

tic chloride, and 0.2975 aluminic oxide. (Silliman's Amer. Journ., 27 1835, 138, 241.) Berzelius points out the inconsistency of these data. (Berzelius' Jahresbericht, 15, 1835, 138.)

C. Tissier: 27.12 (O = 16).

Determined by dissolving aluminium in chlorhydric acid, evaporating to dryness with excess of nitric acid and decomposing the nitrate by heat. The aluminium employed contained 0.135 per cent. sodium. 1.985 of this metal gave 3.645 oxide. [If Na = 23, these data give Al = 27.12.] The metal was prepared by heating aluminic fluoride with purified sodium in a graphite crucible. (Paris Comptes Rend., 46, 1858, 1105.)

J. Dumas: 27.446 (O = 16).

Determined by six experiments on the titration of aluminic chloride with argentic nitrate. The mean result was Al = 13.723 (0=8); extreme difference 0.09. The aluminic chloride, which had been prepared on a large scale, was purified by sublimation over iron-filings and over aluminium filings, and by a third sublimation in a current of hydrogen over aluminium filings, after which it was melted. Experiments on the oxidation of aluminium were found unsatisfactory on account of the difficulty of obtaining the metal pure. They gave Al at from 13.74 to 13.89. Dumas takes Ag = 108; Cl = 35.5. (Ann. de Chim. et de Phys., (3,) 65, 1859, 151.)

W. Odling: 27.5 (O = 16).

Determined from the vapor density of aluminium methide and ethide at 220° and upwards. (*Phil. Mag.*, (4,) 29, 1865, 316.)

—. Isnard: 27 (O = 16).

Pure aluminium dissolved in chlorhydric acid, evaporated and heated to redness, gives 15 of its weight in oxide. (Paris Comptes Rend., 66, 1868, 508.)

Pelouze and Fremy give 27.357 (O = 16); 170.98 (O = 100,) for the atomic weight of aluminium, and assert that this value is derived from the composition of potash-alum, but they give no authority for the value. The experiments were made by precipitation with barium chloride. (Traité de Chimie, 8d ed., 1, 50.)

ANTIMONY.

From the specific heat of antimony, as determined by Bunsen, Regnault, and others, and from the vapor density of volatile compounds, as determined by Mitscherlich, Loewig and Schweizer and others, it is certain that the atomic weight must be about 120. (Gmelin-Kraut, l. c.; and L. Meyer, l. c.)

J. J. Berzelius: 129.03 (O = 16); 806.452 (O = 100).

100 parts of pure antimony, oxidized with nitric acid, evaporated to dryness, and heated to redness, gave 124.8 antimonic antimoniate. The number of experiments and the preparation of the metal are not given. (Poggend. Ann., 8, 1826, 23.)

R. Schneider: 120.3 (O = 16); 751.9 (O = 100).

Determined by experiments on the reduction of native antimonic ter-sulphide in a current of hydrogen. The only foreign substance to be found in the mineral was silicic acid, which was determined in each case. The temperature was kept as low as possible, and the amount of sulphide volatilized, and of that undecomposed by the process, was determined. The mean composition, as ascertained by eight experiments, was 71.48 antimony—extreme difference, 0.078; and 28.52 per cent. sulphur. The atomic weight was calculated from the mean for S = 200. (Pogend. Ann., 98, 1856, 293.) Schneider published a preliminary note in Poggend. Ann., 97, 1856, 483, in which, from a portion of the above-mentioned experiments, he deduced the value 120.25.

H. Rose and Weber: 120.626 (O = 16).

Rose published this determination expressly as a confirmation of Schneider's value. Antimony ter-chloride was dissolved in water containing tartaric acid, and decomposed by hydrogen sulphide. Sulphur was removed from the filtrate by ferric sulphate, and the chlorine determined with argentic nitrate. 2.162 antimony chloride were found equivalent to 4.097 argentic chloride. [If Ag = 107.93 and Cl = 35.457, these data give Sb = 120.626; or, for O = 100, Sb = 753.92.] Rose, adopting some other values gets 1508.67 [twice 754.34.] He also recalculates some earlier

analyses of the ter-chloride, and the penta-chloride (*Poggend. Ann.*, 3, 1825, 443) made by himself by the same method, which give respectively 1512.91 and 1508.6. (*Poggend. Ann.*, 98, 1856, 455.)

W. P. Dexter: 122.836 (O = 16); 764.6 (O = 100).

Attempts were made to determine the atomic weight of antimony from its reducing action on the chloride of gold, but no constant result was obtained. Berzelius' method (vide supra) was, therefore, adopted. From the mean of ten irreproachable experiments Dexter deduces the value 1529.2; extreme difference, 3. The metal was prepared as follows: From antimony tartrate, sodium metantimonate was prepared, and antimonic acid separated out with nitric acid. The antimonic acid was reduced with carbon, and melted with another portion of antimonic acid to remove traces of sodium, etc. It was also heated in a current of hydrogen to remove traces of oxide. The investigation was carried out in Bunsen's laboratory, and with his assistance. (Poggend. Ann., 100, 1857, 563.)

J. Dumas: 122 (O = 16).

Neither the reduction of cervantite nor of the sulphide, nor the oxidation of metallic antimony gave accordant results. Dumas, therefore, resorted to the analysis of the ter-chloride with argentic nitrate. The chloride was prepared by three different methods, and was dissolved in water acidulated with tartaric acid. Seven experiments gave an average of 121.975; extreme difference, 0.69. Ag = 108; Cl = 35.5. (Ann. de Chim. et de Phys., (3,) 1859, 175.)

F. Kessler: 122.24 (O = 16).

In four experiments crystals of antimony ter-oxide were employed. This oxide had been sublimed in a current of pure, dry carbonic acid. A known weight of the compound was nearly oxidized in a chlorhydric acid solution by a known, slightly insufficient, weight of potassic chlorate. The remainder was titrated with a standard solution of potassic bi-chromate, and countertitrated with ferrous chloride. The mean result was Sb = 122.16. In three experiments metallic antimony was employed. It was prepared by reducing the precipitate formed when ammonic hydrate is added to stibium-ammonium tartrate. The metal was oxidized in chlorhydric acid solution by potassic chlorate, (not weighed,) and reduced to antimony ter-chloride by

stannous chloride. The excess of this reagent was chloridized by mercuric chloride, calomel being separated by filtration. The experiment was continued exactly as in the cases where the oxide was taken to start with. The mean of the experiments on metallic antimony was 122.84. The mean of the seven experiments above described is 122.24; extreme difference, 0.94. K = 39.12; Cl = 107.97. Kessler also made experiments by Rose's method, but got discordant results. (Poggend. Ann., 113, 1861, 145.)

B. Unger: 119.76 (O = 16).

Determined by analysis of sodium sulph-antimonate, (Schlippe's Salt.) (Kopp's Jarresbericht, 1871, 325; Arch. der Pharm., (2,) 147, 193; 148, 1.) A single determination by a method from which great accuracy could not be expected. S = 32; Na = 23. (J. P. Cooke, Jr., in Proc. Amer. Acad., 13, 6.)

J. P. Cooke Jr.: 120 (O = 16).

Cooke objects to the determinations of Dexter and Dumas, on the ground that there is no sufficient evidence of the absence of higher or lower compounds of the same elements in the salts employed.

In two experiments antimony was dissolved and precipitated as sulphide, which was heated to 240° before weigh-The formation of free S was prevented, occluded tartaric acid was determined, but occluded oxy-chloride was neglected. The experiments gave each Sb = 120.6 for S = 32. In thirteen experiments Sb was dissolved in a minimum of nitric acid, and the solution boiled over bullets of Sb to complete saturation. The sulphide was then precipitated in an atmosphere of carbon di-oxide. The precipitate contained no free S. The oxy-chloride was driven off at 180° and determined. The tartaric acid was decomposed at 210° and determined. The errors are opposed and min-The mean of the weighings of sulphide, dried at 180°, gave Sb = 119.994 for S = 32; extreme difference, 1.01. The mean of weighings of sulphide heated to 210° gave Sb = 120.295; extreme difference, 1.07. General mean Sb = 120.145. Fifteen analyses of antimonious bromide gave the Br contents at 66.6665 per cent. for Ag = 108, Br = 80, with an extreme difference of 0.195. This composition gives Sb 120. In seven experiments the iodide was analyzed. For I = 127 and Ag = 108, it gave a mean of 76.051 per cent. Sb, or Sb = 120. It was also shown that the chloride cannot be prepared free from exy-chloride, and that its Sb and Cl contents correspond to Sb = 120. Metallic Sb was prepared by reduction of sodic antimoniate, or of oxide, with potassic cyanide, or by Liebig's method. In all cases it was fused for several hours under its own oxide. The haloid salts were purified by fractional recrystallization and distillation, in part in a current of carbon di-oxide. (Proc. Am. Acad., 13, 1877, 1.)

ARSENIC.

The specific heat of metallic arsenic, as determined by Regnault, and the vapor density of a number of volatile compounds, as determined by Dumas, Mitscherlich, Bunsen, and others, prove that the atomic weight of this element must be in the neighborhood of 75. (Gmelin-Kraut, l. c.; and L. Meyer, l. c.)

J. J. Berzelius: 75.1 (O = 16); 469.4 (O = 100).

2.203 grammes of arsenious acid, heated with sulphur in a distilling apparatus in such a manner that sulphurous acid, but no sulphur, could escape, set free 1.069 grammes sulphurous acid. If S=200.75, the value follows. (Poggend. Ann., 8, 1826, 22; and Lehrbuch, 5 ed., 3, 1205.)

J. Dumas: 75 (O = 16).

Dumas found the vapor density of arsine 2.695. [This value multiplied by 28.94278 gives As = (sensibly) 75.] (Ann. de Chim. et de Phys., 33, 1826, 337.)

J. Pelouze: 75 (O = 16); 468.75 (O = 100).

A known weight of arsenic ter-chloride was introduced into a nitric acid solution of a known weight of perfectly pure silver, the chloride being in slight excess. The excess of chloride was then titrated with decimal silver solution.* As the mean of three experiments Pelouze found As = 937.50; extreme difference, 0.8. Ag = 1349.01; Cl = 443.2. The ter-chloride was repeatedly distilled to free it from excess of chlorine. It was colorless, dissolved com-

^{*}This method, which has been frequently employed in the determination of atomic weights, will be referred to as "Pelouze's method."

pletely in chlorine, and boiled between 184° and 185°. (Paris Comptes Rend., 20, 1845, 1047.)

J. Dumas: 74.94 (O = 16).

Determined by four experiments on the titration of arsenic ter-chloride with argentic nitrate, the ter-chloride being prepared in several lots. The number is the mean of the experiments; the extreme difference being 0.15. Dumas takes Ag = 108; Cl = 35.5. (Annal. de Chimie et de Physique, (3,) 55, 1859, 174.)

F. Kessler: 75.2 (O = 16).

In six experiments arsenious acid was titrated with potassic bichromate and counter-titrated with ferrous chloride. The number so obtained was 75.15. In twelve experiments a known weight of arsenious acid was oxidized in caustic potash solution by potassic chlorate, the arsenious acid being slightly in excess, acidified with chlorhydric acid and the excess of arsenious acid titrated with potassic bichromate and counter-titrated with ferrous chloride. The oxidizing action of the potassic bichromate was experimentally determined. The number obtained from these experiments was 75.24. Five experiments were made with acid instead of alkaline solutions of arsenious acid; they gave 75.15. The arsenious acid was colorless, transparent, volatilized without any residue, and was thoroughly dessicated. Kessler assumed K = 39.12; Cl = 107.97. (Poggend. Annal., 95, 1855, 210; 113, 1861, 140.)

BARIUM.

The specific heat of barium compounds, especially of the chloride, as determined by Regnault and by Kopp, shows that the atomic weight of this element lies in the neighborhood of 137. (*Gmelin-Kraut*, l. c.)

Wollaston and Klaproth. 139.2 (O = 16); 870 (O = 100).

Klaproth found that 100 parts of carbon di-oxide were equivalent to 352.57 parts barium oxide, and that 34 parts sulphuric anhydride were equivalent to 66 parts of barium

oxide. If C = 75.4, and S = 200, the value follows. (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius: 136.79 (O = 16).

100 parts of barium chloride gave 138.08 and 138.06 parts argentic chloride. [If Ag = 107.98, and Cl = 35.457, the above value follows.] Berzelius also determined barium from the sulphate; 100 parts barium chloride gave 112.17 and 112.18 parts sulphate. Calculated for S = 200.75 this determination is almost identical with the other; Berzelius, however, expressly adopts the former. [Calculated for S = 32.0742, it gives 135.74.] (Poggend. Annal., 8, 1826, 189, and Lehrbuch der Chemie, 5th ed., 3, 1229.)

E. TURNER: 137.4 (O = 16).

Turner determined the chlorine contents of barium chloride at 34.016 per cent. by precipitation with silver. This number was the mean of the best two experiments made, and the value follows from it on the assumption that Cl = 35.42. The barium chloride was prepared from native carbonate by solution in chlorhydric acid, precipitation of impurities with barium oxide, ignition of the chloride, treatment with alcohol, and recrystallization. (*Phil. Trans.*, 119, 1829, 291.*)

T. Thomson: 136 (O = 16); 850 (O = 100).

Thomson had formerly determined this atomic weight at 875 by mixing potassic sulphate with barium chloride in such proportions that the supernatant liquid contained no sensible amount of either sulphuric acid or barium. Turner having shown the fallacy of this method, Thomson substituted ammonium sulphate, and also sulphuric acid for the potassium salt, and found 9.5006 barium oxide equivalent to 5.00 sulphuric anhydride. He also analyzed the chloride with argentic nitrate, assuming silver = 1375, and chlorine = 450, and reached the same conclusion with reference to barium. (Thomson's System of Chemistry, 7th ed., 1, 1831, 426.)

^{*}Turner made the discovery in the course of this investigation that barium sulphate carries down other salts, such as potassic sulphate, which cannot be extracted from the precipitate by any degree of washing, and that determinations, with barium sulphate, are consequently unreliable. Although Berzelius drew attention to the importance of the observation, and Thomson was obliged to acknowledge errors in his work from this cause, the fact was for a long time nearly forgotten, as can readily be proved from the contents of this digest.

—. SALVETAT: 136 (O = 16); 850 (O = 100).

Determined from the loss of weight ensuing on the decomposition of barium carbonate by sulphuric acid. Details not given. (*Paris Comptes Rendus*, 17, 1843, 318.)

J. Pelouze: 137.28 (O = 16); 858.01 (O = 100).

Into a nitric acid solution of a known weight of perfectly pure silver, a known and slightly more than equivalent weight of barium chloride was introduced. The excess was titrated with decimal silver solution. The value is the mean result of three experiments, which give an extreme difference of 0.22 for O=100. The barium chloride was purified by recrystallizations continued till determinations gave a constant result, and was dessicated in part at 200° , and in part at a temperature just below redness. Pelouze took Ag=1349.01, and Cl=443.2. (Paris Comptes Rendus, 20, 1845, 1047.)

C. Marignac: 187.08 (O = 16); 856.77 (O = 100).

Determined by six experiments on the equivalence of silver and barium chloride performed by Pelouze's method, (vide supra.) 100 silver were found equivalent to 96.365 barium chloride; extreme difference, 0.038; hence the value taken. Marignac takes Ag = 1349.01, and Cl = 443.2. The barium chloride was purified as follows: Commercial chloride was crystallized from boiling aqueous solution; the crystals were heated to redness, dissolved in boiling water, treated with carbon di-oxide, filtered and crystallized, and these crystals were washed with alcohol and again recrystallized. Determinations were made at each stage and the purification was continued until constant results were obtained. (Liebig, Annal., 68, 1848, 214; Bibl. Univ., Arch. des Sciences, 8, 265.)

H. STRUVE: 136.26 (O = 16).

100 parts of barium chloride gave 112.0938 parts of sulphate as a mean of two experiments; extreme difference, 0.005. S = 32; Cl = 35.4624. (Liebig, Annal., 80, 1851, 204; Oefversigt af Kongl. vel. Acad. Foehr., 6, 165.)

T. Andrews: 137.578 (O = 16).

Andrews obtained this number from two nearly coincident experiments of which he gives no details. (Brit. Assoc. Rep., 1852, pt. 2, 33.)

C. Marignac: 137.16 (O = 16).

Three experiments were made on the titration of air-dried barium chloride in crystals by Pelouze's method, (vide supra.) Five grammes of the salt required for precipitation (1) 4.4205; (2) 4.4195; (3) 4.4210 grammes silver. Three experiments were made on the conversion of the same barium chloride into sulphate. Ten grammes of the salt gave (1) 9.543; (2) 9.544; (3) 9.542 grammes sulphate. In each of the latter experiments the water was determined, and was found to vary no more than 0.0005 grammes. Comparison of the two series gives for Ag = 108, S = 16, and O = 8; barium equal to (1) 68.57; (2) 68.61; (3) 68.55; in mean 68.58, or one-half of 137.16. This result is independent of the possible trace of water the chloride might have con-In another series of three experiments the water was driven off at a low red heat and determined, and the salt analyzed by Pelouze's method. It was proved that barium chloride is not decomposed at the temperature employed. (1) gave 68.61; (2) 58.59; and (3) 68.55, or a mean of 68.583. The salt for the experiments marked (1) was prepared by recrystallization and precipitation with alcohol; that for (2) by a repetition of the same process, and for (3) by resolution of (2) and precipitation with chlorhydric acid gas. Marignac proved that the precipitated argentic chloride contained entirely insignificant traces of barium salt. Cl = 35.5. (Bibl. Univ., Archives des Sciences, Nouv. Série., 1, 1858, 209.)

J. Dumas: 137 (O = 16).

Determined by fifteen experiments on the titration of barium chloride with argentic nitrate, which give a general average of 68.516 with an extreme difference of 0.11. The barium chloride was prepared from pure nitrate and pure carbonate, and from commercially pure chloride after it had been freed from lead by precipitation with barium sulphide. The chloride was precipitated from solution by chlorhydric acid gas and melted in a current of chlorine to prevent oxidation. Ag = 108; Cl = 35.5. (Annales de Chimie et de Physique, (3,) 55, 1859, 137.)

BERYLLIUM.

The atomic heat of beryllium has been determined by J. Emerson-Reynolds by direct comparison with that of silver.

In a calorimetric apparatus constructed for the purpose, the amount of heat given off during cooling by 108 parts of silver heated to 100° was found to be equal to that communicated by a little more than 9.2 parts of beryllium under the same conditions. Assuming the atomic weight of the metal to be 9.2, the atomic heat found would be 5.91. The smallness of this number the observer accounts for by supposing that there was a trace of platinum present introduced by the use of platinum vessels in the course of reduction. (Phil. Mag., (5,) 3, 1877, 38.)

J. J. Berzelius: 14.5 (O = 16).

Berzelius analysed the salt formed by saturating dilute sulphuric acid with beryllium oxide. From the amount of barium sulphate obtained he inferred that the atomic weight of beryllium was 331.261 on the supposition that the oxide was Be, + O₃ and that the salt was neutral. Berzelius took O = 100; S = 200.75, and Ba = 855.29. [Awdejew having discovered that this salt is basic, this value is reduced to 90.63; or, for O = 16, to 14.5.] Berzelius accepted Awdejew's determination in preference to his own. (Poggend. Annal., 8, 1826, 187; and Lehrbuch der Chemie, 5th ed., 3, 1225.)

T. Thomson: 36 (O = 16).

Experiments not given. The value is four times nine, and may have arisen from a mistake as to saturation. (System of Chem. 7 ed., 1, 1831, 459.)

—. Awdejew: 13.85 (O = 16); 86.58 (O = 100).

Beryllium sulphate, in chlorhydric acid solution, was decomposed with barium chloride. In the filtrate the excess of barium chloride was precipitated with sulphuric acid, and the beryllium oxide thrown down with ammonia, dried, heated, and weighed. The beryllium sulphate was prepared from pure carbonate by treatment with sulphuric acid and precipitation with alcohol. It was purified by recrystallization. Four experiments were made, the mean of which calculated for S = 201.165, gave Be = 58.084 with an extreme difference of 1.955. (Poggend. Annal., 56, 1842, 106.) Weeren recalculated these analyses for S = 200 and got 57.72, [or $\frac{2}{3}$ of 86.58.] (Poggend. Annal., 92, 1824, 124.)

J. Weeren: 13.83 (O = 16); 86.46 (O = 100).

Weeren followed the same method as Awdejew, except that he precipitated the beryllium with ammonium sulphide, the oxide being soluble in excess of ammonia. The mean of four experiments gave 57.64, the extreme difference being 1.52 for O = 100, [57.64 is $\frac{3}{2}$ of 86.46.] Weeren took S = 200. (Poggent. Annal., 92, 1854, 124.)

G. KLATZO: 13.89 (O = 16).

Klatzo made five analyses of the sulphates containing seven and four molecules of water, precipitating the sulphuric acid as barium sulphate, and the beryllium as oxide by means of ammonia. From a comparison of the sum of the oxide found in all the analyses with the total amount of barium sulphate found, Klatzo deduces Be = 9.227, for Ba = 137, and S = 32. [If Ba is taken equal to 137.16, and S = 32.07, and if each of the analyses is calculated for itself, Be = 13.89. The extreme difference is 0.45.] The sulphates were purified by recrystallization, and treatment with alcohol. (Erdmann's Journ. für Prak. Chemie, 106, 1868, 227; Klatzo, Ueber die Constitution der Beryllerde, Dorpat, 1868.)

L. F. Nilson and O. Pettersson have redetermined the specific heat of beryllium within a few weeks. They find the specific heat 0.4079, corresponding to a trivalent metal and a sesqui-oxide. The investigation seems to have been made with great care, while that of Emerson-Reynolds was merely preliminary. (Berlin, Bericht der chem. Ges., 11, 1878, 386.)

BISMUTH.

Dulong and Petit, Regnault, and Kopp, have determined the specific heat of Bismuth. It corresponds to an atomic weight of about 210. (*Gmelin-Kraut*, l. c.)

P. Lagerhjelm: 212.86 (O = 16); 1830.377 (O = 100).

Metallic Bismuth was oxidized in a weighed vessel by nitric acid, and the nitric acid expelled by heat. 10 grammes of bismuth gave 11.1275 oxide. (Berzelius' Lehrbuch der Chemie, 5th ed., 3, 1216; Stockholm, Akad. Handl., 34, 1813, 219.)

19

R. SCHNEIDER: 208 (O = 16); 1299.98 (O = 100).

Determined by eight experiments on the conversion of metallic bismuth into oxide by solution in nitric acid and decomposition of the nitrate in the same vessel. The escaping gases were led through nitric acid, and the bismuth caught in this way was separately converted into oxide and weighed. In four experiments the bismuth was prepared by the reduction of basic nitrate, and for the other four by the reduction in hydrogen of the oxide formed in those which preceded. 100 bismuth oxide were found to contain a mean of 89.655 metal; extreme difference, 0.048. (Poggend. Annal., 82, 1851, 303.)

J. Dumas: 210.44 (O = 16).

Determined by seven experiments on bismuth chloride, which was decomposed in solution by sodium carbonate, and the sodium chloride thus formed titrated with silver solution. The value taken is the mean result. The extreme difference is 1.12. Dumas takes Ag = 108, and Cl = 35.5. The bismuth chloride was prepared by the action of chlorine on bismuth, and was purified by fractional distillation over bismuth. That employed in the experiments was colorless. (Annal. de Chimie et de Physique, (3,) 55, 1859, 176.)

BORON.

The specific gravities of a number of volatile compounds of boron have been determined by Dumas, Woehler and Deville, and others, and correspond to an atomic weight of about 11. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

H. F. Weber has discovered that the specific heat of boron rises rapidly with the temperature, becoming nearly constant at 600°. Above this temperature its specific heat is 0.5, and its atomic heat 5.5. (*Poggend. Annal.*, 154, 1875, 575.)

J. J. Berzelius: 11.01 (O = 16).

Davy's investigations having shown that boracic acid contains about 68 per cent. oxygen, and having thus established the formula of borax, Berzelius determined the atomic weight from the water contents of that salt. He found in three experiments, without variation, 47.1 per cent. Gmelin-Kraut recalculates this composition with Stas' atomic

weights, and gets the value given. (Poggend. Annal., 8, 1826, 19.)

A. LAURENT: 10.86 (O = 16).

Laurent found that borax retains some water even when melted, which, however, can be expelled by the addition of iceland spar. By repeating Berzelius' experiments, and adding a known weight of spar, he found the water contents in two experiments 47.15 and 47.20. He did not regard the experiments as accurate. Gmelin-Kraut recalculates these data with Stas' atomic weights, and gets B = 10.91 and 10.81. (Paris Comptes Rendus, 29, 1849, 5.)

Woehler and Deville: 10.87 (O = 16).

These chemists titrated the bromide and the chloride of boron with argentic nitrate. They do not offer the analyses as atomic weight determinations, but Dumas applies the data to this object. Taking Ag = 108, and Cl = 35.5, Dumas calculates from the analysis of the chloride prepared by the action of H Cl on B, B = 11; from the analysis of the chloride prepared by the action of Cl on B, B = 10.6; from the analysis of the bromide prepared by the action of bromine on boron, B = 11. (Annal. de Chimie et de Physique, (3,) 62, 1858, 88; 65, 1859, 129.)

T. Thomson: 10.67 (O = 16).

Thomson supposed boracic acid to be composed of one atom of boron and two of oxygen, and concluded from Davy's and his own experiments that the atom of B was exactly equal to that of O. For the correct composition of the acid his value 'must be reduced one-third. (System of Chem., 7th ed., 1, 1831, 214.)

BROMINE.

Mitscherlich determined the vapor density of bromine, and Regnault the specific heat in a solid condition at very low temperatures. Both of these constants correspond to an atomic weight of 80. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

A. J. Balard: 75 (O = 16); 468.85 (O = 100). 1.27 potassium bromide decomposed with sulphuric acid gave a residue of 0.978 potassic sulphate. [If this analysis is calculated with Stas' atomic weights, it gives Br = 74.65.] In another experiment 100 parts of argentic bromide reduced with zinc, the excess of which was extracted with sulphuric acid, gave 58.9 parts silver. [Calculated with Stas' data this gives Br = 75.3.] Balard mentions no special precautions in the preparation of his salts for this determination. (Annal. de Chimie et de Physique, 32, 1826, 357, 362.)

J. Liebig: 75.29 (O = 16); 470.55 (O = 100).

2.521 potassic bromide precipitated with argentic nitrate gave 4.041 argentic bromide. The potassic bromide was obtained by adding potassic hydrate to an alcoholic solution of bromine until the solution began to lose color. (Annal. de Chimie et de Physique, 33, 1826, 331.)

J. J. Berzelius: 78.264 (O = 16); 489.15 (O = 100).

Berzelius suspected that insufficient precautions had been taken in the preceding determinations to get rid of chlorine. He washed bromine for a long time, and converted it into zinc bromide and ammonium bromide. These salts he partially precipitated with argentic nitrate to get rid of chlorine. From the filtrate he precipitated argentic bromide which he washed, dried, and melted. 7.202 of this bromide, decomposed in a current of chlorine, yielded 5.546 argentic chloride; 7.8805 bromide gave 6.069 chloride. If Ag = 1351.607, and Cl = 442.652, the mean value of Br is as above; difference, 0.09. (Poggend. Annal., 14, 1828, 565; Kongl. vet. Akad. Handl., 1828.)

C. Loewig: 75.76 (O = 16).

According to Gmelin-Kraut, Handbuch der Chemie, the determination was published in a treatise entitled Brom und Seine Chemische Verhältnisse, Heidelberg, 1829.

C. Marignac: 79.957 (O = 16).

In three experiments a known weight of silver was dissolved in nitric acid, precipitated with potassium bromide, and the argentum bromide dried at 200° and weighed. [For Ag = 107.93 these experiments give Br = 79.938, with an extreme difference of 0.018.] In vacuo this result is, according to Stas, 79.968. In seven experiments a known weight of silver was precipitated by a determinate amount of potassic bromide by titration. [If K = 39.187, and Ag = 107.93,

this gives bromine = 79.924 with an extreme difference of 0.046.] In vacuo this becomes, according to Stas, 79.945. In four experiments potassium bromate was decomposed by heat, and the potassic bromide weighed. [For K=39.137 these experiments give bromine at 80.11 with an extreme difference of 0.56. These latter are evidently much less accurate than the preceding, and I have therefore averaged the first and second series in vacuo.] The KBr was prepared by heating bromate purified by recrystallization. (Berzelius' Lehrbuch der Chemie, 5th ed., 3, 1194; Bibl. Univ., 46, 1848, 357.)

W. WALLACE: 79.74 (O = 16).

Determined by analysis of arsenic ter-bromide, by titration with argentic nitrate, according to the method of Pelouze, (see arsenic, Pelouze's determination.) Three experiments were made, giving a mean of 79.738; extreme difference, 0.051. As = 75; Ag = 107.97. The arsenic and bromine were directly combined, and the compound was purified by fractional distillation and recrystallization. (Phil. Mag., (4,) 18, 1859, 279.)

J. Dumas: 80 (O = 16).

Determined by three experiments on the conversion of argentum bromide into chloride in a current of dry chlorine. The mean is 80.03; the extreme difference is 0.18. Silver is taken at 108, and chlorine at 35.5. The argentum bromide was prepared with bromine free from iodine, and was purified from chlorine by digestion with argentum bromide. (Annal. de Chemie et de Physique, (3,) 55, 1859, 162.)

J. S. Stas: 79.952 (O = 16).

Four complete syntheses (the weight of each of the constituents, and of the compound being determined) were made of argentum bromide, a known weight of silver being converted into sulphate, and precipitated with a known weight of bromine which had been converted into hydrobromic acid. The mean result was that 100 Ag = 74.0805 Br; with an extreme difference of 0.004. Two analyses of argentic bromate, made by reducing the salt in suspension with sulphurous acid, gave for the molecular weight of the bromide 187.84, and 187.90, mean 187.87. A comparison of these data gives Br = 79.940. [This, I think, must be a misprint for 79.949.] Fourteen experiments were made on the equivalence of KBr and Ag by Pelouze's method, (see

As, Pelouze's determination.) The mean result was that 100 Ag = 110.345 KBr; extreme difference, 0.029. This gives Br = 79.958 for Ag = 107.93, and K = 89.137. The bromate of silver was prepared from potassic bromate and silver salts. For the preparation of Ag see Stas' determination of it. The potassic bromate was prepared by the action of chlorine on a mixture of KBr and KHO. The bromide was prepared by the action of heat on bromate, by treating bromine with KHO, and in other ways. No reagents were probably ever prepared with such care as those employed in this and the accompanying determinations. The weights are all in vacuo. (Stas, Untersuch. über Chem. Proport., Leipzig, 1867.)

CADMIUM.

Regnault, Kopp, and Bunsen have determined the specific heat of cadmium, which corresponds to an atomic weight of 112. Deville and Troost determined the density of cadmium vapor at above 1000°. It answers to an atomic weight of 114. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

F. STROMEYER: 111.48 (O = 16); 696.767 (O = 100).

Stromeyer found that 100 parts of cadmium combine with 14.352 parts of oxygen to form the oxide. (Berzelius' Lehrbuch der Chemie, 5th ed., 3, 1219; Schweigger's Journ., 22, 1818, 362.)

T. Thomson: 112 (O = 16); 700 (O = 100).

Thomson says that he has shown this to be the true value by analysis of the sulphate in two different states. (System of Chem., 7th ed., 1, 1831, 555.)

K. von Hauer: 112 (O = 16); 700 (O = 100).

Determined by nine experiments on the reduction of cadmium sulphate to sulphide in a current of hydrogen sulphide under pressure. The mean of the experiments gave Cd = 55.999; extreme difference, 0.16. Von Hauer took S = 16. The sulphate was purified by repeated recrystallizations and by conversion into oxide. It was dried at 200°. The sulphide was in each case carefully examined for undecomposed sulphate. (Erdmann's Journ. für Prak. Chem., 72, 1857, 346.)

J. Dumas: 112.24 (O = 16).

Determined by six experiments on the titration of cadmium chloride with argentic nitrate. The mean of all the experiments was Cd = 56.12; extreme difference, 0.49. The third experiment varies considerably from the rest, and Dumas seems inclined to omit it in the average. If it is left out, the mean becomes 56.06; extreme difference, 0.29. Dumas takes Cl = 35.5; Ag = 108. The cadmium chloride was prepared in two lots by solution of cadmium in chlorhydric acid, evaporation and melting for several hours in a current of chlorhydric acid gas. (Annal. de Chimie et de Physique, (3,) 55, 1859, 158.)

E. Lenssen: 112.06 (O = 16).

Three experiments were made on the decomposition of cadmium oxalate, the salt and the resulting oxide being weighed. The mean result was Cd = 56.03; extreme difference, 0.19. C = 6. The oxalate was prepared from pure chloride by precipitation with oxalic acid, washing and drying at 150° . It was carefully tested, and was found to be anhydrous. (Erdmann's Journ. für Prak. Chemie, 79, 1860, 281.

CÆSIUM.

The great similarity between cæsium and the other alkaline metals renders the deduction of its atomic weight from its equivalent sufficiently certain.

Kirchhoff and Bunsen: 123.35 (O = 16).

Determined by three experiments on the analysis of the chloride with argentic nitrate. The value is the mean; extreme difference, 0.13. The cossium was separated from the other alkalies by extracting a mixture of oxides and carbonates with alcohol. It was converted into chloride by precipitation with platinum chloride, reduction of the double chloride in hydrogen and solution. These operations were repeated until the cossium salt gave sensibly the same results after successive purifications. Its purity was also tested spectroscopically. Silver was taken at 107.94, and chlorine at 85.46. (Poggend. Annal., 113, 1861, 363.)

Johnson and Allen: 133 (O = 16).

Determined by four experiments on the precipitation of cæsium chloride with argentic nitrate. The mean result was Cs = 133.036; the extreme difference, 0.842. Ag = 107.94; Cl = 35.46. Cæsium and rubidium were separated by partial crystallization of their bitartrates. The cæsium bitartrate was converted into chloride by precipitation with platinum chloride, reduction and solution. The nitrate formed on the precipitation of the cæsium chloride with silver was reconverted into cæsium chloride and redetermined, and so on. The purity of the salt was tested spectroscopically. (Silliman's Amer. Journ., (2,) 35, 1863, 96.)

R. W. Bunsen: 183 (O = 16).

Determined by three experiments on the precipitation of cæsium chloride with argentic nitrate. The mean result was 132.99; extreme difference, 0.02. Ag = 107.94; Cl = 35.46. In order to prepare pure chloride, a mixture of cæsium and rubidium salts was converted into carbonates, a little more tartaric acid was added than was necessary to form acid tartrate with the rubidium and neutral tartrate with the cæsium, and the mixture was exposed on a filter to the action of a saturated atmosphere of aqueous vapor. The cæsium salt is deliquescent, and gradually passes through the filter, while the rubidium salt is unaffected. The exesium tartrate was turned into pure chloride by repeated precipitation with platinum chloride, reduction in hydrogen and solution. The determinations were made on the product of successive purifications, and only those were taken into consideration which were made after analysis showed a constant composition. The spectroscope was employed to test the purity of the salt. (Poggend. Annal., 119, 1863, 5.)

—. Mercer: 133 (O = 16).

The fact of this determination, without details, is mentioned by Frankland. (Chem. News, 8, 1863, 18.)

R. Godeffroy: 132.557 (O = 16).

Derived from the mean of four analyses of cossium chloride with argentic nitrate, the extreme difference being 0.185. Cl = 85.5; Ag = 108. The cossium was separated from the other alkalies by the fractional crystallization of their alums continued until the cossium compound was

spectroscopically pure. The aluminium was removed with ammonia, the sulphuric acid with barium chloride and traces of barium with ammonium carbonate. The cæsium chloride, which was not deliquescent, was dried at 150°. (Liebig's Annal., 181, 1876, 185.)

CALCIUM.

Bunsen has determined the specific heat of calcium. It corresponds to an atomic weight of 40. (Gmelin-Kraut, l. c.)

F. H. Wollaston: 40.736 (O = 16); 254.6 (O = 100).

Wollaston found that 43.7 parts of carbon di-oxide saturated 56.8 parts of lime. If C = 75.4, the value follows. (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius: 40.32 (O = 16); 252.075 (O = 100).

301 parts of anhydrous calcium chloride gave 775 parts argentic chloride. If Cl = 443.28 and Ag = 1349.66 the value follows. This analysis, made in 1818, was erroneously calculated from a mistake in setting down its results and the atomic weight of Ca was taken at 256.019. (Poggend. Annal., 8, 1826, 189; and Lehrbuch der Chemie, 5th ed., 3, 1227.)

J. Dumas: 40 (O = 16).

Three experiments were made on the calcination of calcium carbonate which contained 0.08 per cent. of ferric oxide and silicic acid. The weight of the residue was in mean 56.07, or, subtracting 0.03, 56.04, with an extreme difference of 0.08. These figures give almost exactly 40. The weighings are reduced to vacuum. (Paris Comptes Rendus, 14, 1842, 537.)

—. SALVETAT: 40 (O = 16); 250 (O = 100).

It is to be inferred from the context that this determination was made from the loss of weight ensuing on the decomposition of calcium carbonate by heat or sulphuric acid. (Paris Comptes Rendus, 17, 1843, 318.) C. Marignac: 40.208 (O = 16); 251.3 (O = 100).

Determined by precipitating calcium chloride with argentic nitrate; Ag = 1349.01; Cl = 443.2. Marignac laid no weight on this determination finding it impossible to prepare calcium chloride which did not show an alkaline reaction. The presence of caustic lime would make the result erroneously high; no doubt Berzelius' early analysis was defective from the same cause. (Berzelius' Jahresbericht, 24, 1844, 103; Bibl. Univ., 46, 1843, 367.)

Erdmann and Marchand: 40.007 (O = 100).

Four experiments were made on the calcination of calcium carbonate enclosed in a double platinum crucible in a wind-furnace, till the weight was constant. A mean of 56 per cent. calcium oxide was found with an extreme difference of 0.05. This gives Ca = 40 for C = 12. Two experiments were made by decomposing calcium carbonate by sulphuric acid. These gave a mean of 43.99 carbonic acid; difference, 0.02. The value taken is the mean of all experiments. The carbonate was prepared by precipitating calcic chloride with ammonium carbonate, and drying at 160° to 180°. Confirmatory experiments were made on iceland spar. The weighings are reduced to vacuum. (Erdmann's Journ, für Prak, Chem., 26, 1842, 472.)

mann's Journ. für Prak. Chem., 26, 1842, 472.)

Berzelius maintained that Erdmann and Marchand employed material containing water, chlorine and magnesium. Erdmann and Marchand answered that there could be no magnesium and was no chlorine but that they had convinced themselves that spar is the only compound of certain and constant composition. Berzelius replied that they then admitted that their carbonate contained water. Erdmann and Marchand appealed to their experiments on spar, upon which Berzelius made experiments showing that spar, too, retains water at 200°. This Erdmann and Marchand denied and finally assert that all the carbonic acid is not driven off at any attainable temperature, and that their results were therefore too high instead of being too low. The error they estimate to exactly cover the difference between their averages and 40. (Erdmann's Journ. für Prak. Chem., 31, 1844, 257; 37, 1846, 75; 50, 1850, 287.)

ERDMANN and MARCHAND: 40.062 (O = 16); 250.89 (O = 100).

The spar experiments referred to above. Six analyses were made as before, giving a mean of 56.028 oxide; extreme

difference 0.047. (Erdmann's Journ. für Prak. Chemie, 31, 1844, 268.) Another experiment, in which the absence of water was proved, gave 56.03 lime. The weighings are reduced to vacuum. (Erdmann's Journ. für Prak. Chem., 37, 1846, 77.)

J. J. Berzelius: 40.264 (O = 16); 251.651 (O = 100).

Five experiments were made on the conversion of caustic lime into sulphate. The value is the mean for 8 = 200.75; extreme difference 0.962 for O = 100. The lime was carefully purified and burnt, but Berzelius says nothing of testing it for carbonic acid, upon which Erdmann and Marchand found an objection. Berzelius expresses himself ill satisfied with the results. (Liebig's Annal., 46, 1843, 241; also Lehrbuch der Chemie, 5th ed., 3, 1228.)

J. Dumas: 40.02 (O = 16).

Five experiments were made on the titration of calcium chloride with argentic nitrate. They give a mean of 20.065, but Dumas considers only three of them as entitled to a voice. These give 20.01; extreme difference, 0.03. The calcium chloride was prepared by dissolving marble in chlorhydric acid, digestion with lime water, filtration, evaporation, treatment with chlorhydric acid and heating in a current of chlorine. For the three experiments averaged the chloride was kept melted in the current of gas for from 8 to 10 hours. Ag = 108; Cl = 85.5. (Annal. de Chimie et de Physique, (3,) 55, 1859, 129.)

CARBON.

The specific gravity of gaseous carbon compounds shows that the atomic weight must be nearly 12. (Gmelin-Kraut, l. c.) Weber has shown that the specific heat of carbon at high temperatures obeys Dulong and Petit's law.

F. H. Wollaston: 12.064 (O = 16); 75.4 (O = 100).

Biot and Arago found the specific gravity of carbon dioxide 1.5196, and that of oxygen, 1.1036. Calculation from these data gives the value. (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius: earlier determinations.

In 1817 Berzelius attempted to determine the atomic weight of carbon by two analyses of plumbic carbonate. [These analyses calculated for Pb = 206.926 (Stas,) give C = 11.998 and 11.984, or 74.99 and 74.90.] Considering the difference too great, he calculated the atomic weight from Biot and Arago's determination of the specific gravities of carbon di-oxide and oxygen, 1.10859 and 1.51961. Berzelius gives 75.33 as the result; [I make it 75.394.] Subsequently, (1819,) Berzelius and Dulong determined these specific gravities more accurately at 1.524 and 1.1026 whence he calculated C = 76.487. This number was accepted until Dumas showed it to be false, although in the mean time carbon di-oxide had been shown to be a condensible gas. According to Dumas, Berzelius at one time accepted a value 76.52 of which I have found no account. In Berzelius' Lehrbuch, 3, 1174, 76.48 is a misprint for 76.437. (Berzelius' Lehrbuch der Chemie, 5th ed., 3, 1197, et passim.)

T. Thomson: 12 (O = 16); 75 (O = 100).

Thomson found the specific gravity of carbon di-oxide 1.52678. Assuming the specific gravity of oxygen at 1.1111, chiefly to accord with the supposition that air is a compound containing 20 per cent. of oxygen, he calculates the atomic weight of carbon at 75. (Erdmann's Journ. für Prak. Chem., 8, 1836, 872; Records of General Science, by R. D. Thomson, 1836, 179.)

J. Dumas: about 12.16 (O = 16); 76 (O = 100).

From analysis of well crystallized naphthaline, Dumas infers that the atomic weight of carbon cannot be so high as 76.44, and must be nearly as above. (*Poggend. Annal.*, 44, 1838, 110.)

J. J. Berzelius: 12.23 (O = 16); 76.458 (O = 100).

One experiment was made on the decomposition of plumbic carbonate by heat, which gave C = 76.405. [If Pb = 206.926, the data give C = 12.185, or 76.157.] Another experiment was made on the oxalate, which gave C = 76.511. Berzelius regards these results as confirmatory of the value 76.488. The plumbic carbonate was prepared by precipitating the nitrate with ammonium carbonate. The oxalate was obtained by decomposing the acetate with oxalic acid. (Liebig's Annal., 30, 1839, 241.)

G. Fownes: 12.12 (O = 16).

Determined by three analyses of naphthaline with cupric oxide, the usual precautions being observed. The value is the mean; extreme difference, 0.14. The naphthaline was purified by slow sublimation in a florence flask, and was brilliantly white. Fownes does not regard his results as conclusive as to the exact value. (*Phil. Mag.*, (3,) 15, 1839, 62.)

E. MITSCHERLICH: 12.016 (O = 16); 75.1 (O = 100).

Experiments made on the analysis of naphthaline by the ordinary method of organic analysis gave never more than 75.2, and those which seemed most accurate very nearly 75. (Mitscherlich's Lehrbuch der Chemie, 4th ed., 1, 1844, 595.)

Dumas and Stas:
$$12 (O = 16)$$
; 75 $(O = 100)$.

Determined by fourteen experiments on the combustion of carbon in oxygen, the resulting carbon di-oxide being weighed. In five cases natural graphite was employed, and in four graphite from charcoal pig-iron. Both were purified by treatment with acid and heating in chlorine. The necessary oxygen was developed in the combustion-tube from potassic chlorate and cupric oxide. In five experiments diamond was employed, and the oxygen was furnished from a gasometer. The oxygen was displaced by air, especially purified from carbon di-oxide by milk of lime. The products of combustion were collected in tubes filled with pumice stone moistened with sulphuric acid, Liebig potashbulbs and tubes filled with dry potash. The mean of the experiments on graphite gave C = 74.982; those on diamond gave 75.005; the extreme difference was 0.238. The observers point out that the result would not be affected by reduction to vacuum. (Annal. de Chimie et de Physique, (3,) 1, 1841, 5.)

Liebig thinks that potash must have been volatilized, and says that there is no assurance that the oxygen was completely expelled by air. He also points out that the analyses of camphor and benzoic acid, accompanying the investigation, show an excess of carbon for C = 75. (Liebig's Annal., 38, 1841, 195.)

Erdmann and Marchand: 12.009 (O = 16); 75.054 (O = 100).

Erdmann and Marchand repeated Dumas' and Stas' experiments. Five experiments on diamond gave C = 75.028;

31

extreme difference, 0.38. Three experiments on natural and one on artificial graphite gave C = 75.087; extreme difference, 0.13. The number is the mean of all experiments. Erdmann and Marchand adopt 75. Calcium chloride was used in these experiments instead of sulphuric acid to avoid objections as to the possible volatility of the acid. (Erdmann's Journ. für Prak. Chem., 23, 1841, 159.)

Berzelius and Liebig and Redtenbacher: 12.119 (O = 16); 75.741 (O = 100).

Five analyses by Berzelius of the tartrate of lead, the decomposition being effected by heat, gave 62.7431 per cent. plumbic oxide; extreme difference, 0.045. Several analyses of plumbic racemate gave a mean of 62.75 per cent. oxide; extreme difference, 0.05. The salts were prepared by fractional precipitation of plumbic acetate with tartaric and racemic acids respectively. They were dried at 100°. (Poggend. Annal., 19, 1830, 306.) From the analyses of the tartrate Liebig and Redtenbacher calculate C = 75.771, and from the racemate 75.711, taking Pb = 1294.489 and H = 6.2394. (Liebig's Annal., 38, 1841, 137.)

LIEBIG and REDTENBACHER: 12.137 (O = 16); 75.854 (O = 100).

Determined by decomposing known weights of organic salts of silver in a covered crucible by heat and weighing the silver. Five analyses of each of the following salts showed that 18.6113 Ag = 28.8098 acetate; 9.6171 Ag = 16.223 tartrate; 16.2641 Ag = 27.438 racemate; 16.0596 Ag = 25.9019 malate. If Ag = 1351.607 and H = 6.2394, the above value for C follows, with an extreme difference for the 20 analyses of 0.765, (O = 100.) The figures are all calculated for vacuum: [If Ag = 107.93 and H = 1.0025, the average number obtained from the mean of each set of analyses gives C = 12.06865 or 75.429.] The acetate was prepared by partially neutralizing pure acetic acid with ammonia, precipitating with argentic nitrate and recrystallizing the salt from hot aqueous solution. The crystals were dried at 103°. The acetic acid was prepared from plumbic acetate. The tartrate was prepared by adding tartrate of sodium and potassium to a hot (80° to 85°) dilute solution of argentic nitrate till a small permanent precipitate was formed, and cooling the solution. The racemate was prepared from pure acid racemate of ammonium like the tartrate. The malate was prepared from calcium

malate and argentic nitrate. The salt thus obtained was dissolved in nitric acid, and reprecipitated with ammonia added drop by drop, not to complete neutralization, washed and dried. (*Liebig's Annal.*, 38, 1841, 139.)

A. Strecker recalculated Liebig and Redtenbacher's analyses given above, independently of the atomic weight of silver, from the difference in their atomic composition, employing the method of least squares. He found $C=75.415\pm0.061$, or 12.066 ± 0.01 . In the same way, and from the same analyses he calculated the atomic weight of silver at 1348.79, or 107.9032. [The close coincidence between this result and Stas', is certainly worthy of remark.] (Liebig's Annal., 59, 1846, 280.)

Marignac repeated Liebig and Redtenbacher's experiments and got almost the same results, but, by varying the method so as to preclude loss by spirting, different ones.

(Liebig's Annal., 59, 1846, 287.)

Stas had the same experience as Marignac, and also ascribes Liebig and Redtenbacher's high results to loss by spirting. (Bulletin de l'Acad. Roy. des Sciences de Belgique, 16, 1849, 9.)

C. Marignac: 11.986 (O = 16).

Determined by three analyses of the acetate of silver. The salt was decomposed by heat in a tube in such a way that the products of decomposition were forced to pass through porous silver, and loss by spirting was impossible. 100 parts of the salt were found to contain a mean of 64.664 silver, with an extreme difference of 0.005 in vacuo. [If Ag = 107.93, these figures give the above value.] Marignac regards the analysis as a confirmation of Dumas and Stas' determination. The acetate was prepared by solution of argentic carbonate in acetic acid and successive recrystallizations. (Liebig's Annal., 69, 1846, 287; Bibl. Univ., Arch. des Sciences, 1. 1846.)

Strecker believes that the silver in Marignac's determination must have retained carbon. (*Ibid.* 284.)

F. VON WREDE: 12.019 (O = 16); 75.12 (O = 100).

Von Wrede determined the specific gravity of carbon dioxide, taking into consideration its variation from the law of Marriotte. He found it equal to $1.52087 \frac{1+0.0049.p.}{1+at.}$ He also found the specific gravity of oxygen 1.1052 and

83 CARBON.

that of carbonic oxide 0.96779. Comparison gives C = 75.11 to 75.14. (Berzelius' Jahresbericht, 22, 1842, 72.) Berze-

lius adopted this determination.

According to Gmelin-Kraut, 1, (2,) 70, Regnault's value for the specific gravity of oxygen combined with von Wrede's for carbon di-oxide gives C = 12.0037, and with that for carbonic oxide C = 12.0105.

J. S. STAS: 12.005 (O = 16); 75.029 (O = 100).

Determined by passing carbonic oxide over a known weight of pure cupric oxide, and weighing the carbon dioxide formed. Stas got from eight experiments C = 74.993The number taken is the mean of the results, to 75.055. which is misprinted in Stas' paper 75.039.] The carbonic oxide was prepared from oxalic acid by the action of sulphuric acid. It was purified from carbon di-oxide by passing through potash tubes, and from oxygen by passing over hot copper filings, and was kept in a gasometer over water, in which was dissolved a solution of stannous oxide in pot-The cupric oxide was prepared by igniting pure cupric The carbonic acid formed in the experiments was caught in potash and sulphuric acid tubes. The amount of carbon di-oxide weighed was from 23 to 67 grammes. The weighings are reduced to vacuum. (Bulletin de l'Acad. Roy. des Sciences de Belgique, 16, 1849, 9.)

GRAPHON.

B. C. Brodie: 83 (O = 16).

By the action of potassic chlorate and nitric acid on graphite, Brodie obtained a compound of carbon, oxygen and hydrogen containing 11 atoms of carbon, and by the action of heat on this substance two others containing, respectively, 22 and 66 atoms. The first of these is analogous to the hydrated oxide of silicon obtained by Buff and Woehler, if Si = 21. From this fact, and the specific heat of graphite, Brodie concludes that the atomic weight of the graphitic form of C is 33. (Phil. Trans., 149, 1859, 249.) Graham-Otto points out that if Si = 28, graphon must be 44, and that, in that case, the argument from the specific heat loses its applicability.

CERIUM.

The specific heat of metallic cerium, as determined by W. F. Hillebrand, is 0.04479, and the atomic heat 6.18 if the atomic weight is 188. (*Poggend. Annal.*, 158, 1876, 86.)

It is well known that cerium is always accompanied in nature by lanthanium and didymium. The former was discovered in 1839, and the latter in 1843, both by Mosander.

W. HISINGER: 137.93 (O = 16).

According to Hisinger, as reported by Berzelius, the lower oxide of cerium contains 14.821 O per 100 Ce, giving the atomic weight at 574.718 for O = 100, if the lower oxide is regarded as a protoxide. (*Poggend. Annal.*, 8, 1826, 186.)

T. Thomson: 160 (O = 16).

Thomson analysed the sulphate and obtained for cerium the value 625, (O = 100.) [He probably took barium = 70.] (System of Chem., 7th ed., 1, 1831, 466.)

F. J. Otto: 138.91 (O = 16).

According to Gmelin, Otto found in an approximate determination Ce = 578.8, and recorded it in his revised translation of Graham's Chemistry, 1, 1840, 222.

A. Beringer: 138.48 (O = 16).

[Three analyses of cerous chloride with silver give the atomic weight of cerium at 576.375, or 92.22, if Ag = 107.93, and Cl = 35.457. Inconsistent results are given for an analysis of the sulphide.] Three analyses of the sulphate in which the oxide was determined, gave 57.4717 per cent. so-called protoxide, [or Ce = 576.31, or 92.21, if S = 82.0742.] Analysis of the formate gave Ce = 577.04 for C = 75.85. The material for the preparations was ceric oxide obtained from cerite, and purified from lanthanium by digestion with very dilute nitric acid. The lower oxide was assumed to be Ce O. (Liebig's Annal., 42, 1842, 184.)

R. Hermann: 138 (O = 16).

The lower oxide was assumed to be Ce O. 23.523 parts of anhydrous cerous sulphate gave 29.160 parts of barium sulphate, giving Ce = 575, for O = 100, Ba = 856.88, and S = 201.16. The salt was obtained by precipitating basic

sulphate from a sulphuric solution of the cerite oxides, and converting this precipitate into the neutral salt. (Erdmann's Journ. für Prak, Chem., 30, 1843, 184.)

C. Rammelsberg: 137.93 (O = 16).

Hermann states that Rammelsberg experimented on cerium salts free from lanthanium, and got Ce = 574.7, the lower oxide being supposed to contain one atom of oxygen. [I cannot find the original paper.] (Erdmann's Journ. für Prak. Chem., 30, 1843, 184.)

C. Marignac: 141.79 (O = 16).

The result of seven experiments on the titration of cerous sulphate, prepared from basic sulphate, with barium chloride. (Erdmann's Journ. für Prak. Chem., 48, 1849, 406; Bibl. Univ. Arch des Sciences, 8, 265.) Marignac subsequently made experiments which showed these results to be too high from the impurity of the barium sulphate precipitate, (see note to Turner's determination of Barium,) and that the number 575 (for O = 100 and cerous oxide Ce O) was more probable. (Annal. de Chimie et de Physique, (3,) 38, 1853, 148.)

T. Kjerulf: 174.56.

Kjerulf obtained, by three organic analyses of cerium oxalate, Ce = 727.33 on the protoxide theory, O = 100. The salt was prepared by dissolving cerium oxide in oxalic acid. (*Liebig's Annal.*, 87, 1853, 12.) Bunsen points out that this must have been a basic salt. (*Ibid*, 105, 1858, 50.)

R. Bunsen and J. Jegel: 138.192 (O = 16).

The lower oxide was presumed to contain one atom of oxygen. In two experiments cerous sulphate was decomposed with ammonium oxalate. The sulphuric acid thus liberated was determined with barium sulphate; the cerium oxalate precipitate was decomposed by heat with the formation of ceric oxide, which was weighed and the additional oxygen, introduced by heating, determined by iodometric titration. The salt was not anhydrous; the water contents was estimated by difference. The experiments gave respectively 57.49 and 57.46 per cent cerous oxide in the anhydrous salt, or Ce = 576.3 and 575.25 if S = 200. One experiment was made on hydrous cerium oxalate. The cerous oxide was found as before; the water was determined and the

This gave 60.02 oxalic acid was estimated by difference. per cent. cerous oxide, calculated for the anhydrous salt, or The salts were prepared from cerite as fol-Ce = 575.65.lows: the mineral was digested with sulphuric acid, the sulphates formed were leached with water and with dilute nitric acid; this solution was treated with hydrogen sulphide, chlorhydric acid was added and cerium oxalate was precipitated. The oxalate was heated with magnesia to convert the cerium into the higher oxide, which was dissolved in concentrated nitric acid. After diluting the solution, chemically pure basic sulphate was precipitated. the preparation of cerous sulphate and oxalate oxidation was prevented by the action of sulphurous acid. (Liebiq's Annal., 105, 1858, 45.)

C. Rammelsberg: 138.216 (O = 16).

One experiment on the organic analysis of cerium oxalate by heating in a current of oxygen behind copper oxide gave Ce = 575.9, (O = 100), or 92.144, (O = 16), cerous oxide being regarded as $Ce\ O$. Rammelsberg does not adopt his own, but Hermann's determination. (*Poggend. Annal.*, 108, 1859, 44.)

C. Wolf: 136.992 (O = 16).

Determined from experiments on the sulphate, prepared and analyzed as by Bunsen and Jegel. Wolf purified the basic sulphate by solution in nitric acid and reprecipitation in hot water, aided by recrystallizations. He found that the oftener these processes were repeated the smaller was the atomic weight resulting from the analysis. The purifications were repeated until the salt was spectroscopically free from didymium, and was perfectly white, (that employed by other investigators had been yellowish or buff.) The value taken, 45.664, [or \frac{1}{3} of 136.992,] was the smallest and last value reached. The investigation was made in Bunsen's laboratory. (Silliman's Am. Journ., (2,) 46, 1868, 58.)

C. H. Wing: 137.01 (O = 16).

Two experiments were made on the decomposition of hydrous cerium sulphate with oxalic acid, the cerium oxalate being converted into ceric oxide by heat. The amount of cerous oxide in the ceric oxide was calculated according to Wolf's results, giving for the atomic weight of cerium 45.64 and 45.69, S being 32. The cerium was six times recon-

verted into basic sulphate, and repeated recrystallizations were made. The salt was white and spectroscopically pure. The determination was made in Gibbs' laboratory. (Silliman's Amer. Journ. (2,) 49, 1870, 356.)

D. Mendelejeff: 188 (O = 16).

Mendelejeff first suggested raising the atomic weight of cerium from 92 to 138. His reasons were a specific heat determination which he had made with very impure metal, and the fact that the supposed sesquioxide had never been shown to exist. He believes that the atomic weight will be found somewhat below 138, because that is the atomic weight of barium. (Liebig's Annal., suppl., 8, 1871, 186.)

H. Buehrig: 140.648 (O = 16).

Determined from ten analyses of the hydrous oxalate performed by combustion in a current of pure oxygen behind copper oxide. The water was collected in tubes filled with calcic chloride, and the carbonic acid in potash. Five experiments in which the cerium oxide was not determined gave a mean of 94.1304, on the supposition that cerous oxide contains 1 atom of oxygen and that O = 15.96, with an extreme difference of 0.0445. Five determinations in which the cerium was determined as ceric oxide gave 94.2260, with an extreme difference of 0.0431. Carbon was taken at 11.97. The mean result is Ce = 94.1782 for the above mentioned assumptions, [or 140.648 for O = 16, and on the supposition that cerous oxide is a sesqui-compound.] The exalate was prepared from basic nitrate purified by Gibbs' method of oxidation with minium and nitric acid. The salt was spectroscopically pure. (Erdmann's Journ. für Prak. Chem., 120, 1873, 222.)

CHLORINE.

The density of chlorine gas and the specific heat of chlorine compounds leave no doubt that the atomic weight of this element is nearly 35.5. (Gmelin-Kraut, l. c.)

Marcet, Berzelius, Wollaston: 35.28 (O = 16).

Marcet, by experimenting on the calcination of pure marble, and on the saturation of chlorhydric acid with lime, found as the mean of many trials, that 50.77 calcic carbonate are equivalent to 56.1 calcic chloride. Wollaston, taking the equivalent of calcic carbonate at 630, and that of calcium at 255, calculates the equivalent of chlorine at 441 for O = 100. Wollaston cites Berzelius as having obtained the same number by the conversion of plumbic carbonate into chloride. (*Phil. Trans.*, 97, 1807, 301; 104, 1814, 20.)

J. J. Berzelius: 35.412 (O = 16); 221.827 (O = 100).

The molecular weight of potassium chloride was ascertained from four experiments on the decomposition of potassium chlorate, which on being heated lost 89.15 per cent. oxygen. This gives for the chloride 982.567, (O = 100.) 100 parts of potassium chloride were further found equivalent to 192.4 parts argentic chloride, and 100 parts of silver to 132.75 argentic chloride. The value follows. Berzelius in his Lehrbuch accepts Marignac's determination and ascribes the error of the value he had obtained to the imperfect decomposition of that portion of the chlorate which was carried off as dust during the experiment. (Poggend. Annal., 8, 1826, 17; also Lehrbuch der Chemie, 3, 1189, 1191.)

E. Turner: 85.42 (O = 16).

Turner made two experiments on the decomposition of plumbic chloride with argentic nitrate. Assuming the atomic weight of lead at 103.6, and that 100 silver = 132.8 chloride, these analyses gave Cl = 85.48 and 85.48. Turner also decomposed corrosive sublimate with calcic oxide neutralized with nitric acid and precipitated with argentic If mercury = 201, these analyses give a maximum of 85.28, and a minimum of 35.21, of which Turner selects the largest. From calomel treated in the same way, he arrived at the value 85.85. From his experiments on the composition of argentic chloride (and apparently comparison with potassic chloride and chlorate) Turner got 35.45. The mean of the other experiments was 35.35, but Turner considers 85.42 as being the most likely value. plumbum chloride was prepared from the carbonate, and was purified by recrystallization, as was also the corrosive The calomel was "prepared by Mr. Howard," and retained traces of moisture at 300°, which would make the atomic weight derived from its analysis too small. The values are for vacuum. (Phil. Trans., 123, 1833, 529.)

F. Penny: 35.454 (O = 16).

Six experiments on the conversion of silver into nitrate gave 100 Ag = 157.441 nitrate; extreme difference, 0.028. Twelve experiments by three different methods on the conversion of silver into chloride gave 100 Ag = 132.837 chloride. Four series of experiments on the interconversion of potassic chloride, chlorate and nitrate gave for the difference between the molecular weights of the chloride and the nitrate 26.56. Corresponding experiments with sodium salts gave the same difference 26.568. The mean combined with the data for the silver salts gives the molecular weight of argentic chloride at 143.424, and Cl = 35.454. For further details see Penny's determinations of potassium, sodium, nitrogen and silver. The weighings were calculated for vacuum. (Phil. Trans., 129, 1839, 32.*)

R. PHILLIPS: 85.688 (O = 16).

In order to avoid the error possibly incurred by the melting of argentic chloride, etc., Phillips mixed known and nearly equivalent quantities of silver dissolved in nitric acid, or of crystallized argentic nitrate, with ammonium chloride; filtered, washed, and precipitated the comparatively minute amount of chlorine in the filtrate with silver solution. The fusion of this small quantity could cause no loss of importance. Phillips confesses that his ammonium chloride was acid and the only conclusions he draws are that Cl = 36, N = 14, O = 8 and H = 1 may be taken without considerable error if silver is 108. [The method seems to have been original and is nearly that afterwards adopted by Pelouze. The acidity of the ammonium chloride would of course give Cl too high.] (*Phil. Trans.*, 129, 1839, 35.)

C. Marignao: 36.001 (O = 16); 225.007 (O = 100).

Determined by passing chlorhydric acid gas over hot cupric oxide and condensing the water formed. The mean of three experiments was Cl = 450.013; the extreme difference is 0.2 for O = 100. The gas was made from recrys-

^{*}This is one of the most elegant investigations of the kind to be found in chemical literature, though it scarcely receives a mention except from Stas, who accords to it the praise it deserves. Stas' wonderfully exhaustive researches were necessary to prove beyond question that chemistry has a mathematical basis, and that the atomic weights of the elements are incommensurate. Penny's investigation, taken in connection with Stas', shows that the highest degree of accuracy is not incompatible with the simplest means when they are applied with the care and acumen, without which exact results cannot, under any circumstances, be obtained.

tallized sea-salt and concentrated sulphuric acid and was dried by passing through nine tubes filled with sulphuric acid and pummice stone and with calcium chloride. The water was collected in a condenser to which drying tubes were appended. (Paris Comptes Rendus, 14, 1842, 570.)

A. LAURENT: 35.468 (O = 16); 221.672 (O = 100).

Determined by three analyses of chloronaphthalintetrachloride, which he found to contain 58.22; 58.29; 58.28; per cent. Cl. The mean is 58.27 from which the value follows. (Paris Comptes Rendus, 14, 1842, 456.)

(Paris Comptes Rendus, 14, 1842, 456.)
According to Maumené, Laurent confessed that his salt was impure, containing chlorose compounds, in Gerhardt's Comptes Rendus, 1845, 108. (Annal. de Chimie et de Physique, (3,) 18, 1846, 45.)

C. Marignac: 35.37 (O =16); 221.07 (O = 100).

One synthesis of argentic chloride showed that 100 silver equals 32.74 chlorine. Berzelius had found 32.75, which Marignac adopts. Marignac found by six experiments on the decomposition of potassic chlorate by heat, that the molecular weight of potassic chloride was 932.14. He tested the equivalence of potassic and argentic chlorides by precipitating the former with argentic nitrate, filtering without the use of paper through a funnel with a capillary neck. The precipitate was dried and weighed, then melted and reweighed, no loss being observable. 100 potassium chloride gave 192.33 and 192.34 argentic chloride in two experiments, or reduced to vacuum, 192.26. Hence the atomic weight is 442.13. The potassic chloride was prepared by heating chlorate which had been purified by repeated recrystallizations. (Liebig's Annal., 44, 1842, 23.)

C. Marignac: 35.456 (O = 16); 221.6 (O = 100).

In accordance with Pelouze's suggestion, Marignac repeated his determination of the composition of argentic chloride and of the equivalence of potassic and argentic chlorides, retaining the molecular weight of potassic chloride mentioned in the last paragraph. That value was obtained from the mean of six experiments on the decomposition of the chlorate which gave the percentage of oxygen at from 39.155 to 39.167; mean 39.161. Pelouze had got, as the mean of three experiments, 39.157. (Paris Comptes Rendus, 15, 1842, 959.) Marignac made eleven experiments on the equivalence of silver and potassium chloride by Pelouze's

method, a known weight of silver being dissolved in nitric acid and added to a known and nearly equivalent amount of potassic chloride in solution, after which the excess was itrated with decimal standard solution. 100 parts of silver were precipitated by from 69.049 to 69.067, in mean by 69.062 chloride. 100 parts of chloride were precipitated by from 192.33 to 192.37, in mean by 192.348 silver. Five experiments were made on the composition of argentic chloride by dissolving silver in nitric acid, with precautions against loss by spirting, precipitation with chlorhydric acid, washing, drying, melting and weighing in the same vessel. 100 parts of silver gave from 132.825 to 132.844 chloride, mean 132.84. Calculation from these data gives in vacuo Ag = 1349.01; K = 488.94; Cl = 443.20; for O = 100 [or Ag = 107.921; K = 39.115; Cl = 35.456, for O = 16.] (Berzelius' Jahresbericht, 24, 1844, 58; Bibl. Univ., 46, 1843, 350.)

C. GERHARDT: 36 (O = 16).

By heating potassic chlorate in a current of oxygen Gerhardt got, when he took precautions against loss by spirting, a mean of 60.949 chloride, from which he deduces 86 for chlorine without giving further data. (Paris Comptes Rendus, 21, 1845, 1280.) Marignac shows that no data have ever been published which, in connection with Gerhardt's experiments, would give this value for chlorine. He adds further experiments of his own which, without aiming to establish more exactly the true atomic weight, prove it less than 36 (Liebig's Annal., 59, 1846, 284; Bibl. Univ., Arch. des Sciences, 1, 1846.)

E. J. Maumené: 35.462 (O = 16).

Maumené made seven analyses of argentic chloride by reduction in a current of pure hydrogen. Five of these experiments were made with quantities less than 10 grammes, and gave a mean of 100 silver = 32.736 Cl. Two experiments were made with about 30 grammes each, and gave 100 silver equal to 32.86 and 32.853 chlorine. Maumené prefers the latter, and deduces from them for chlorine the value 43.67 or 35.494 taking silver according to his own experiments at 1350.32. [If silver is taken at 107.93 (Stas) the same analyses give 35.462.] (Annal. de Chimie et de Physique. (3,) 18, 1846, 41.)

A. LAURENT: 35.5 (O = 16); 221.88 (O = 100).

A single experiment was made as follows: pure silver was weighed off and placed in a matrass, nitric and chlorhydric acids were added, the liquid was evaporated and the chloride melted. An empty test was carried on at the same time to act as tare. Silver was taken at 1350. (Paris Comptes Rendus, 20, 1849, 5.)

J. DUMAS: 85.5 (O = 16).

Determined by chloridizing different weights of pure silver by heating the metal in a current of chlorine. Experiments on 10 grammes and 20 grammes gave a mean of 35.5055, the difference being 0.013, for chlorine, if silver is 108. (Annal. de Chimie et de Physique, (3,) 55, 1859, 135.)

J. S. Stas: 85.457 (O = 16).

Stas found the atomic weight of chlorine by three inde-

pendent methods:

- (1.) From analysis of argentic chlorate and synthesis of argentic chloride. A known weight of the chlorate was dissolved in water, precipitated with sulphuric acid to secure advantageous division of the salt, and reduced while in suspension by a slow stream of sulphurous anhydride. The chloride was washed, dried, and weighed in the flask in which it was produced. The minute amount of chloride present in the chlorate was collected and taken into consideration, and the wash-water was carefully examined for silver. analyses (of about 140 and 260 grammes) gave for the molecular weight of the chloride 143.383 and 148.407, mean A variety of syntheses of argentic chloride in the 148.895. wet and in the dry way showed that 100 parts silver combined with nearly 32.850 parts chlorine. Stas assumes that none of his syntheses can possibly have given too much chloride and accepts the relation stated. These data give Cl = 35.458.
- (2.) From the mutual relations of potassic chlorate and chloride and argentic chloride, combined with the composition of the last. The chlorate was decomposed either by gentle heat or in the wet way with chlorhydric acid. 100 parts of chlorate were found to contain 60.846 parts chloride as the mean of eight experiments; extreme difference, 0.012, which gives the molecular weight of potassic chloride at 74.59. The relation between potassic and argentic chloride was ascertained by Pelouze's method, (see Marignac's

determination above.) Twenty experiments on quantities of 32 grammes, and less, of silver gave 100 parts Ag = 69.103 parts KCl; extreme difference, 0.008. These data combined with the composition of argentic chloride given above, indicate for chlorine 35.460.

(3.) The composition of argentic nitrate was determined, and the difference between the atomic weights of nitrogen and chlorine. In two experiments silver was dissolved in nitric acid, the solution evaporated to dryness, and the nitrate kept melted until there was no further loss of weight. The result obtained was that 100 silver = 157.484 nitrate; difference, 0.008. From series of experiments on the relation of the chlorides of potassium, sodium, lithium and silver to the nitrates, Stas found the difference between a chloride and a nitrate from 26.586 to 26.591; mean 26.588. These data show that the atomic weight of chlorine lies between 35.455 and 35.460, and confirm the mean of all the determinations of Penny, Marignac, and Stas, 35.457. The silver for this investigation was either distilled or compared with distilled silver; it was found impossible to reduce the amount of silica in the alkaline salts below 0.002 of one per cent., it was therefore determined and allowed for; every possible method of purification by recrystallization and otherwise was resorted to to ensure purity. The weighings are all reduced to vacuum. (Stas, Unters. über Chem. Proport., Leipzig, 1867.)

CHROMIUM.

The specific heat of chromium, as determined from that of the oxide by Kopp, Regnault, and Neumann, corresponds to an atomic heat of from 5.4 to 5.98, if the atomic weight is taken at 52.4. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 56.29 (O = 16); 351.819 (O = 100).

100 parts of plumbic nitrate, on precipitation with potassic chromate, gave 98.772 parts plumbic chromate. The value follows for Pb = 1294.498, and N = 88.518. (Poggend. Annal., 8, 1826, 22.)

T. Thomson: 64 (O = 16); 400 (O = 100).

3.14 grains of metallic chromium, converted into chromic acid by heating with potash and nitre, gave a precipitate of 16.23 grains plumbic chromate. (*Phil. Trans.*, 117, 1827, 159.)

E. Peligot: 52.48 (O = 16); 328 (O = 100).

Peligot reached this value by a careful carbon determination of chromous acetate, produced by precipitating a dilute solution of chromium protochloride with sodium acetate, C = 75. Peligot does not regard the experiment as definitive, the salt possessing but little stability. (Annal. de Chimie et de Physique, (3,) 12, 1844, 527.)

N. J. Berlin: 62.54 (O = 16); 328.39 (O = 100).

Five experiments were made on the decomposition of argentic chromate with chlorhydric acid and alcohol. The silver chloride was washed in the flask in which it was precipitated, treated with aqua regia, melted and weighed without removal. The decanted fluid and the wash-water were evaporated to dryness with excess of ammonia, treated with water and the chromium oxide filtered off, heated to redness and weighed. [Nothing is said of the recovery of any argentic chloride that might have been removed by the decantation.] The value taken is calculated from the comparison of the amounts of argentic chloride and of chromium oxide obtained, Ag = 1349.66; Cl = 443.28. The extreme difference is 1, for O = 100. The argentic chromate was prepared by adding nitrate to a solution of potassic chromate. (Erdmann's Journ. für Prak. Chem., 38, 1846, 145.)

V. A. JACQUELIN: 50.08 (O = 16); 313 (O = 100).

By washing and purifying violet chromium chloride, Jacquelain obtained a substance which he took to be the pure chloride and which was more soluble than the unpurified salt. He analysed it by melting with soda, and arrived at the above number. (Liebig's Annal., 64, 1847, 275; Revue Scient., 14, 198.)

A. Moberg: 63.563 (O = 16); 384.769 (O = 100).

Moberg made twelve experiments on the decomposition of chromium salts by heat. In two cases the sulphate dried at a low red heat was decomposed by strong ignition in a platinum crucible; the results being, 335.65 and 335.29 for chromium. Ten experiments were made on the decomposition of ammonium-chromium-alum which had been dried in a pulverized state for a long time. These determinations gave from 838.965 to 335.739. The value taken is the mean. The alum employed was prepared from pure material, and was repeatedly recrystallized. S = 200; N = 87.5. (Erdmann's Journ. für Prak. Chem., 43, 1848, 115.)

J. Lefort: 52.97 (O = 16).

Determined by fourteen experiments on the precipitation of barium with sulphuric acid from a nitric acid solution of barium chromate. The barium chromate was prepared by precipitating potassium chromate with barium nitrate and drying the precipitate at 250°. [If these analyses are calculated for barium = 137 and S = 32, they give 100 barium chromate = 60.244 barium oxide, extreme difference, 0.26, and the atomic weight as above. Lefort seems to have taken Ba = 136.72. Berlin points out the correction which I have verified.] (Erdmann's Journ. für Prak. Chem., 51, 1850, 261; Journ. de Pharm. et de Chim., 18, 27.)

R. Wildenstein: 63.485 (O = 16).

Determined by thirty-two experiments on the precipitation of barium chloride, desiccated at a red heat, by pure, neutral potassic chromate. The mean of these analyses gave 100 barium chromate = 81.70 barium chloride; extreme difference 0.35. Wildenstein calculates 384.48 without giving the assumption for chlorine. [If Cl = 35.457; Ba = 137, the value follows.] (Erdmann's Journ. für Prak. Chem., 59, 1853, 28.)

F. Kessler: 52.3 (O = 16).

Kessler reached this value by comparing the oxidizing action of potassic chromate with that of potassic chlorate on arsenious acid. Six experiments were made on the oxidizing power of the chromate and twelve on that of the chlorate by a method of titration. By combining the maximum of one with the minimum of the other series, Kessler finds the atomic weight of chromium between 25.93 and 26.40; in mean 26.15, K being = 39.12 and Cl = 35.45. Confirmatory experiments were made on the oxidation of ferrous chloride in the same way. These gave a mean of 26.1. (Poggend. Annal., 113, 1861, 137; 95, 1855, 208.)

M. SIEWART: 52.094 (O = 16).

Determined from the amount of chlorine in sublimed violet chromium chloride. Siewart criticises Kessler's determination and deduces from the latter's data a value 25.02. (Kopp's Jahresbericht, 14, 1861, 240; Halle, Zeitschr. für die Gesammt. Naturwis., 17, 530.)

Kessler points out that the number 25.02 is a misprint in the Jahresbericht, and that Siewart's paper ascribes to him the value 26.02. (Poggend. Ann., 117, 1862, 852.)

COBALT.

The atomic heat of cobalt as determined by Regnault is 6.27 if the atomic weight is assumed at 58.8. (Gmelin-Kraut, l. c.)

E. ROTHOFF: 58.98 (O = 16); 368.65 (O = 100).

269.2 parts of cobalt oxide converted into neutral cobaltous chloride and precipitated with argentic nitrate gave 1029.9 argentic chloride, according to Berzelius' report. (Poggend. Annal., 8, 1826, 185.) Berzelius recalculates this analysis for Cl = 221.64 and Ag = 1349.66, and gets the value taken. (Berzelius' Lehrbuch, 3, 1220.)

R. Schneider: 60.006 (O = 16); 375.04 (O = 100).

Determined from four analyses of the oxalate. The carbon was determined as in organic analysis; the metal by heating a known weight of the salt first in a current of air, then in one of oxygen, and by reduction of the oxide in hydrogen. The mean of the four analyses gave cobalt at 80.003, with an extreme difference of 0.026 for C = 6. The oxalate was prepared by converting the chemically pure cobalt of commerce into roseo-cobaltic chloride, from which the metal was again reduced, then dissolved in chlorhydric acid and carbonate precipitated, which was digested with oxalic acid. (Poggend. Annal., 101, 1857, 398.)

Marignac objects to this determination that the oxalate, being insoluble, may very likely have retained portions of the carbonate which could not be removed by washing.

(Bibl. Univ., Arch. des Sciences, (2,) 1, 1858, 372.)

Schneider answers that he obtained nearly identical results from lots prepared at different times, and that he believes that he has convinced himself that the oxalate contained no carbonate. (*Poggend. Annal.*, 107, 1859, 610.)

Gibbs, reporting Schneider's determination, remarks: "Very numerous and carefully made analyses of the ammonium-cobalt bases, executed in my laboratory, indicate 29.5 as the true equivalent of cobalt." (Silliman's Amer. Journ., (2,) 25, 1858, 438.)

C. Marignac: about 59 (O = 16).

Five experiments were made on the decomposition of cobalt sulphate by heat. This salt can be readily dried without decomposition, and the acid is completely driven off by heat, but the resulting protoxide contains a slight

47

excess of oxygen. In order to remove this excess it was melted under a known weight of an acid silicate of lead. The results for cobalt varied from 29.32 to 29.38. The sulphate was purified by recrystallization. Marignac also experimented on the chloride. The weight of this salt varies greatly with the moisture of the atmosphere when crystallized, and attempts to desiccate it usually result in the formation of some insoluble compound. Three analyses of chloride appearing to contain one molecule of water, and dried at 100°, performed by titration with silver solution, gave cobalt at 29.42 to 29.51. Five experiments were made in the same way on chloride either melted in a current of chlorine or of chlorhydric acid gas, or calcined with ammonium chloride. These determinations gave from 29.36 to 29.42. (Bibl. Univ., Arch. des Sciences, (2,) 1, 1858, 374.) [Marignac, in another investigation in the same volume, takes Ag = 108; Cl = 35.5.]

J. DUMAS: 59 (O = 16).

Determined by five experiments on the titration of cobalt chloride with silver. The mean result for cobalt was 29.542; extreme difference 0.09; Ag = 108; Cl = 85.5. The chloride was prepared by dissolving pure cobalt in aqua regia, evaporating in the presence of excess of chlorhydric acid and heating to redness in a current of chlorhydric acid gas. In two of the determinations cobalt from a different lot, which had been heated in a vacuum was employed. (Annal. de Chimie et de Physique, (8,) 55, 1859, 148.)

W. J. Russell: 58.74 (O = 16).

Determined by fifteen experiments on the reduction of cobalt oxide in hydrogen. The value is the mean; the extreme difference is 0.19. To obtain pure cobalt oxide Claudet's salt was prepared, purified by recrystallization, etc., reduced in hydrogen, the metal dissolved in nitric acid and the resulting salt decomposed by heating in a stream of carbon di-oxide. (Chem. Soc. Journ., (2,) 1, 1863, 57.)

Schneider considers that no sufficient precautions were taken to exclude air in these experiments, and that higher oxides were formed. (*Poggend. Annal.*, 130, 1867, 810.)

E. von Sommaruga: 60 (O = 16).

Determined by seven experiments on the reduction of purpureocobaltic chloride in a current of hydrogen. The mean of the experiments is 29.965; four of them give a

mean 29.996. The extreme difference is 0.098. The salt was prepared by solution of the carbonate in chlorhydric acid, addition of ammonia in excess, exposure to the air, washing of the precipitate with acidulated, then with pure water and drying at 110° . A special examination showed it free from other metals. Sommaruga took Cl = 35.5; N = 14. (Erdmann's Journ. für Prak. Chem., 100, 1867, 118; Sitz.-Bericht der k. k. Akad., 1866.)

C. Winkler: 59 (O = 16).

This value is derived from the mean of five experiments on the precipitation of gold from a solution of neutral crystallized chloride of gold and sodium. The metallic cobalt employed was prepared by the reduction of purpureocobaltic chloride. The latter was made from oxide, and was purified by recrystallization. Gold was assumed at 196. The mean of the results was 29.496; extreme difference, 0.071. (Fresenius' Zeitschr. für Anal. Chem., 6, 1867, 22.)

P. Welesky: 58.98 (O = 16).

Determined from the analysis of cobalti-cyanides, performed by drying the salt at 100°, and heating to redness, first in a current of oxygen then of hydrogen. Four experiments with phenylammonium-cobalti-cyanide gave cobalt at from 29.38 to 29.59. Two experiments with ammonium-cobalti-cyanide gave from 29.46 to 29.55. Mean, 29.48; extreme difference, 0.21. A single experiment by Winkler's method gave 29.42. (Berlin, Bericht der Chem. Ges., 2, 1869, 592.)

W. J. Russel: 58.76 (O = 16).

Determined by the amount of hydrogen set free by the solution of cobalt in hydrochloric acid. The value is the mean of 2 (or 4?) trials. The cobalt employed was that reduced by Russel in his former experiments on the same atomic weight. (Chem. News, 20, 1869, 20.)

R. H. Lee: 59.10 (O = 16).

Determined by analysis of cobalti-cyanide salts. They were decomposed in a crucible by heating from above. The carbon separated was burned off in air and then in oxygen, and the metallic oxide reduced in hydrogen. Six experiments on the strychnine salt gave a mean of 59.05. Six experiments on the brucine salt gave 59.15. Six experi-

ments, made with especial care, on the reduction of purpureo-cobaltic chloride by hydrogen gave 59.09. (Reported by Gibbs. Berlin, Bericht der Chem. Ges., 4, 1871, 789.)

COPPER.

Regnault, Kopp, and others have determined the specific heat of copper. It corresponds to an atomic heat of about 6 if the atomic weight is taken at 63.8. (Gmelin-Kraut, l. c.)

R. Chenevix: F. H. Wollaston: 64 (O = 16); 400 (O = 100.)

Chenevix found 20 parts of oxygen equivalent to 100 parts of copper, whence Wollaston deduces the atomic weight. (Phil. Trans., 104, 1814, 21.)

J. J. Berzelius: 63.296 (O = 16); 395.6 (O = 100).

Determined by two experiments on the reduction of cupric oxide with hydrogen, which gave 395.695 and 395.507. The water was not weighed. (*Poggend. Annal.*, 8, 1826, 182; and *Lehrbuch*, 3, 1216.)

ERDMANN and MARCHAND: 63.456 (O = 16); 396.6 (O = 100.)

Determined by four experiments on the reduction of large quantities of cupric oxide in a current of hydrogen. The hydrogen was displaced by air after the completion of the reduction. The weight of the oxide and of the copper were reduced to vacuum, but not that of the weights employed. To obtain pure cupric oxide, pure vitrol was prepared and electrolytically decomposed. The copper thus obtained was dissolved in nitric acid, and the nitrate decomposed by heat. The value is the mean; the extreme difference is 0.056 for O = 8, or 0.112 for O = 16. (Erdm. Journ. für Prak. Chem., 31, 1844, 389.)

Berzelius points out that these analyses vary among themselves much more than his own. He makes the difference somewhat greater than it really is by neglecting the reduction to vacuum. (*Ibid.*, 37, 1846, 72.)

Hampe shows that these analyses, correctly calculated, give Cu = 63.46. (Zeitschr. für Berg Hütten-und-Sal-Wesen im Preus. St., 21, 1873, 261.)

J. Dumas: 63.5 (O = 16).

Dumas says that experiments on the reduction of cupric oxide and on the sulphidation of copper have shown him that the atomic weight of copper lies between 31.5 and 32, near 31.75, but that his experiments cannot be regarded as decisive. (Annal. de Chimie et de Physique, (8,) 55, 1859, 129.)

MILLON and COMMAILLE: 63.128 (O = 16); 394.55 (O = 100).

These (three) experiments were in most respects a repetition of Erdmann and Marchand's. The value is the mean; the extreme difference is 0.49 for O = 100, or 0.0784 for O = 16. The sulphate was prepared free from iron or zinc by dissolving copper in ammoniacal sulphate or nitrate. The oxide was obtained by heating the nitrate. (Paris Comptes Rendus, 56, 1863, 1249; and 57, 1863, 145.)

Fresenius sees no reason for preferring this to Erdmann and Marchand's value. (Fresenius' Zeitschr. für Anal. Chem., 2, 1863, 474.)

W. Hampe: 63.3296 (O = 16).

In three experiments cupric oxide was reduced in a current of hydrogen with all possible precautions. The hydrogen was displaced by air before weighing, though it was shown by experiment that porous copper does not condense hydrogen. The metal was heated till incipient melting was The reduction and melting were repeated withobserved. out altering the weight. Hampe attempted to control his results by reconverting the metal into oxide, but was unable to effect complete oxidation. The water produced by the reduction was found to be perfectly pure. The mean result was Cu = 31.6696, maximum, 31.6729, minimum, 31.6648. The oxide was prepared from metallic copper. To obtain pure metallic copper, sulphate free from bismuth was electrolytically decomposed, the finely divided metal well washed, then melted, first in a current of carbon di-oxide, afterwards in hydrogen, and then again in carbon di-oxide. From the metal, basic nitrate was formed and from this salt, by heating first in air and then in oxygen, oxide. In two experiments the atomic weight of copper was determined by decomposing cupric sulphate by electrolysis, and weighing The residual fluid was evaporated, and a minute amount of copper, which had escaped decomposition, was

recovered and determined as sulphide. For S=16.037 and 0=8, these experiments gave Cu=31.6577 and 31.66. The value taken is the mean of the two series. All weighings were reduced to vacuum. (Zeitschr. für Berg Hüttenwal Sal.-Wesen im Preus. St., 21, 1873, 260.)

DIDYMIUM.

W. F. Hillebrand found the specific heat of this metal 0.04563, which corresponds to an atomic heat of 6.60 for an atomic weight of 144.78. (*Poggend. Annal.*, 158, 1876, 78.)

C. Marienac: 148.8 (O = 16); 930 (O = 100).

Determined by decomposing disulphate with barium chloride. Assuming the lower oxide as a prot-oxide, he calculated the atomic weight at 620. As Marignac was not confident of the purity of his salt, and subsequently became certain that the method was untrustworthy, details are unnecessary. (Liebig's Annal., 71, 1849, 313.)

C. Marignac: 143.81 (O = 16); 898.8 (O = 100).

Five experiments were made on the sulphate by decomposition with ammonium oxalate. The didymium oxalate was heated to redness, and the resulting oxide weighed. On the assumption that the oxide was protoxide, these determinations gave a mean of 598.2 for Di, with an extreme difference of 2.5. Three experiments were made on the chloride, the insoluble oxychloride, which is unavoidable in drying the salt, being separated. The chlorine was determined with silver, and the Di as in the previous experi-These determinations gave Di at 600.2, with an extreme difference of 5.2 for Cl = 443.2 and S = 200. The salts were prepared from cerite. The cerium was extracted by treatment at first with dilute and afterwards with concentrated nitric acid. The sulphates of Di and La were separated by partial precipitation with oxalic acid and by partial recrystallization. (Annal. de Chimie et de Phys., (3,) 38, 1853, 148.)

R. Hermann: 142.44 (O = 16); 890.25 (O = 100).

In one experiment sulphate which had been heated to a low red heat, was dissolved, decomposed with ammonium

oxalate, the precipitate incinerated and the oxide weighed. The result was Di = 594.46, on the prot-oxide hypothesis, for S = 200. In one experiment the chloride was decomposed with argentic nitrate, oxychloride being filtered off and allowed for, and the argentic chloride weighed. This experiment gave Di = 592.54 for Cl = 443.2. For the preparation of the salt see Lanthanium. (Erdmann's Journ. für Prak. Chem., 82, 1861, 387.)

H. ZSCHIESCHE: About 144 (O = 16).

In five experiments the sulphate was exposed to a white heat until the weight became constant and the oxide on being tested showed no traces of sulphur. The results varied from Di = 46.585 to 48.08, probably, Zschiesche thinks, on account of the presence of La. S = 16. Di was separated from La by the partial precipitation of the nitrates with oxalic acid, the first portion falling being redissolved, and the partial precipitation repeated twenty times. (Erdmann's Journ. für Prak. Chem., 107, 1869, 74.

C. Erk: 142.695 (O = 16).

The sulphate was decomposed with ammonium oxalate, the oxalate incinerated and the oxide weighed. The sulphuric acid was also precipitated as barium salt, and weighed. Three experiments gave a mean of Di = 95.13, on the prot-oxide hypothesis, with an extreme difference of 0.78. The Di salt was found to contain yttrium which was removed by repeated fractional precipitation with sodium sulphate. This re-agent precipitates a double salt of Di and sodium. The purification was continued until the atomic weight became constant. (Kopp's Jahresbericht, 1870, 319, Jena'sche Zeitschr, für Med. und Nat., 6, 299.)

Casselmann thinks that the salt may still have retained yttrium, and Fresenius objects to the barium sulphate determination on the well-known grounds. (Fresenius' Zeitschr, 10, 510.)

D. Mendelejeff: 138 (O = 16).

From the analogy between Di and cerium and other elements, and from the fact that it forms two oxides, Mendelejeff believes that its lower oxide is a sesqui-oxide, and its atomic weight 138. Mendelejeff points out that an error is to be apprehended in the received values from the fact that we have no guarantee of the pureness of Di salts except recrystallization. (Liebig's Annal. Suppl. 8, 1871, 190.)

P. T. CLEVE: 147.01 (O = 16).

Determined by the conversion of didymium oxide into sulphate. The number is the mean of six experiments; extreme difference 0.58. The Di was separated from lanthanium by repeated precipitations of basic nitrate from nitric acid solution, conversion into formate and decomposition of this salt by heat. (Kopp's Jahresbericht, 1874, 259. Bulletin Soc. Chimique, (2,) 21, 246.)

W. F. HILLEBRAND: 144.78 (O = 16).

Determined by one experiment on the conversion of metallic Di into nitrate, and then, by heat, into oxide. The impurities were determined. The metal was reduced electrolytically from the chloride. (*Poggend. Annal.*, 158, 1876, 78.)

ERBIUM.

The physical and chemical analogies of the salts of this element have led Mendelejeff (*Liebig's Annal.*, Suppl. 8, 1871, 195,) and P. T. Cleve (Kopp's Jahresbericht, 1874, 260; Bulletin Soc. Chimique, (2,) 21, 344,) to regard it as triatomic, and its atomic weight as about 170.

M. Delafontaine: 113.04 (O = 16).

M. Delafontaine investigated gadolinite by Mosander's method, and obtained besides yttrium, two substances which he regarded as erbium and terbium. From the sulphates, in which he supposed the metals to exist as protoxides, he determined erbium at 496 and terbium at 471 for O = 100. Popp (Liebig's Annalen, 131, 189,) and Bunsen and Bahr (lbid, 137, 1,) have shown that Mosander's method gives only mixtures. Delafontaine's terbium is thought to have been chiefly the erbium of other chemists. (Liebig's Annal., 134, 1865, 108.)

Bahr and Bunsen: 168.9 (O = 16).

A known weight of erbium oxide was treated with a very slightly excessive quantity of sulphuric acid; the solution evaporated and the excess of acid driven off at as low a temperature as possible. The increase of weight indicates 112.6 for S = 32. The oxide was prepared from gadoli-

nite. The mineral was decomposed with chlorhydric acid, and the earths precipitated with oxalic acid. The oxalates were converted into nitrates, the cerium metals separated with potassic sulphate, and calcium and magnesium with ammonia. If the nitrates of yttrium and erbium are dissolved in boiling water, basic erbium nitrate with some yttrium crystallizes out, leaving yttrium nitrate with some erbium in solution. The process of partial crystallization was continued as long as the atomic weight of the erbium salt continued increasing. Bahr and Bunsen believe, however, that the atomic weight may be some hundredths higher. The salt was spectroscopically free from didymium. (Liebig's Annal., 137, 1866, 2.)

P. T. Cleve and O. M. Hoeglund: 170.55 (O = 16).

Determined from four syntheses of the sulphate, giving 113.7 on the diatomic hypothesis. The oxide was purified by heating the nitrates, etc., according to Berlin. (Blomstrand in Berlin, Ber. der Chem. Ges., 1873, 1467; Bull. Soc. Chimique, 1873, 193 and 289.)

FLUORINE.

Dumas and Peligot and others have determined the vapordensity of a number of fluorine compounds. They correspond to an atomic weight of about 19. (L. Meyer, l. c.)

H. DAVY: 18.86 (O = 16).

Determined by the conversion of Derbyshire spar into sulphate. 100 parts of spar gave a maximum of 175.2 parts calcic sulphate. [If S = 32; Ca = 40; the value follows.] (*Phil. Trans.*, 104, 1814, 64.)

J. J. Berzelius: 18.85 (O = 16).

Determined by conversion of calcic fluoride into sulphate. 100 parts fluoride gave, in mean of three experiments, 175 parts sulphate; extreme difference, 0.2. [If S = 32; Ca = 40; the value follows.] (Poggend. Annal., 8, 1826, 18, and Lehrbuch, 3, 1196.)

P. LOUYET: 19 (O = 16).

Determined by six experiments on the conversion of fluor-spar into calcic sulphate. The mean result was 100 parts spar equal 174.36 sulphate, with an extreme difference of 0.3. Spar from Derbyshire was pulverized, digested with chlorhydric acid, and the foreign matter removed by lutration in water. It was completely dissolved in sulphuric acid, the excess of which was driven off by heat continued till a constant weight was obtained. S = 200; Ca = 250. (Erdmann's Journ. für Prak. Chem., 47, 1849, 104; Annal. de Chim. et de Phys., (3,) 25, 1849, 291.)

E. FREMY.

This chemist says that his analyses essentially confirm Berzelius' determination. (Annal. de Chimie et de Phys., (8,) 47, 1856, 27.)

J. Dumas: 19 (O = 16).

Determined by the conversion of fluorides into sulphates. A single experiment on the conversion of calcic fluoride gave 18.96; two experiments on sodic fluoride, 19.06; and two on potassic fluoride, 18.99. The mean is 19.01; extreme difference, 0.12. Ca = 20; Na = 25; K = 39; S = 16. The alkaline salts were well crystallized and were fused before use. (Annal. de Chim. et de Phys., (3,) 55, 170.)

S. DE LUCCA: 18.96 (O = 16).

Determined by four experiments on the conversion of a pure spar from Gerfalco into sulphate. The extreme difference was 0.15. The decomposition was very difficult. The loss on ignition and the residue left on evaporation of the acid employed were taken into consideration. [S apparently = 16; Ca = 20.] (Paris Comptes Rendus, 51, 1860, 299.)

GALLIUM.

Berthelot has determined the specific heat of gallium at 0.079 corresponding to an atomic heat of 5.52, if the atomic weight is 69.9. (Paris Comptes Rend., 86, 1878, 786.)

L. DE BOISBAUDRAN: 69.9 (O = 16).

This chemist "has prepared several chlorides, [samples of chloride?] several bromides, and several anhydrous iodides of gallium. He has determined the atomic weight of gallium, and found it 69.9, (mean of two experiments.)" (Paris Comptes Rend., 86, 1878, 756.)

GOLD.

Dulong and Petit and Regnault have determined the specific heat of gold. It corresponds to an atomic weight of about 200. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 196.4 (O = 16).

Determined by the amount of mercury necessary to precipitate a known weight of gold from solution of chloride. 142.9 mercury were found equivalent to 98.55 gold. [If Hg = 200, this gives Au = 196.397.] (Poggend. Annal., 8, 1826, 178.)

T. Thomson: 200 (O = 16).

This value is derived from a somewhat inaccurate experiment on the reduction of auric chloride by ferrous sulphate. (*Edinb. Trans. Roy. Soc.*, 11, 1831, 26.)

J. J. Berzelius: 196.73 (O = 16).

Determined by five experiments on the relative amount of gold and of potassic chloride in the residue obtained by heating the double chloride of the two metals in an atmosphere of hydrogen. [Calculated for KCl = 74.594, (Stas,) these experiments give a maximum of 196.79, minimum of 196.63 and a mean of 196.727. The atomic weight derived from the first experiment is misprinted in the Lehrbuch, as is the mean in the Jahresbericht.] (Berzelius' Jahresbericht, 25, 1846, 41; and Lehrbuch, 3, 1845, 1212.)

A. Levol: 196.26 (O = 16).

A known weight of gold was converted into chloride, and this salt decomposed in boiling solution by a current of pure, washed sulphurous acid. The sulphuric acid formed was precipitated as barium salt, and the atomic weight calculated by comparison of the gold employed and the barium sulphate obtained. 1000 gold gave 1782 sulphate. [If the atomic weight of S=32.0742, and that of Ba=137.08, the above value follows.] (Annal. de Chimie et de Phys., (3,) 30, 1850, 355.)

HYDROGEN.

The density of hydrogen as determined by a great number of investigators, especially Regnault, is about 1 of that of oxygen. If oxygen is 16, the atomic weight of hydrogen

is consequently about 1.

The atomic weights of the elements are compared either with that of oxygen or with that of hydrogen. The main advantage of assuming hydrogen as unity is the simplicity of the approximate values expressed in terms of the atomic weight of this element. The hypothesis of Prout has also had much influence in giving currency to this unit. The advantages of oxygen as a standard of comparison consist in the fact that it combines with all the elements, except fluorine, and in the superior accuracy of the determination of its specific gravity. The percentage variation between Regnault's determinations of the specific gravity of hydrogen was thirty-six times as great as occurred in his experiments on oxygen. Unnecessary complication in the approximate values of the atomic weights is as well avoided by assuming oxygen at 16 as by taking hydrogen at 1.

These reasons for the adoption of the atomic weight of oxygen as a standard of comparison appear to me conclusive, and accordingly all values in this paper have been re-

duced to O = 16.

F. H. Wollaston: 1.06 (O = 16); 6.64 (O = 100).

Gay-Lussac and Humboldt having shown that two volumes of hydrogen and one of oxygen form water, and Biot and Arago having determined the specific gravity of these gases, Wollaston calculated the above atomic weight. (*Phil. Trans.*, 104, 1814, 20.)

Berzelius and Dulong: 0.9984 (O = 16); 6.24 (O = 100).

Determined by three experiments on the reduction of cupric oxide by hydrogen. The hydrogen was made from

pure materials, and passed through a solution of litharge in potash, and over a coarse powder of caustic potash before use. The resultant water was caught in calcic chloride and weighed. The determination was also confirmed by experiments on the specific gravity of oxygen and hydrogen. The minimum result for hydrogen was 0.9934, the maximum 1.0086. (Thomson's Annals of Phil., 2, 1821, 48.)

T. Thomson: 1 (O = 16); 6.25 (O = 100).

Thomson found the Sp. Gr. of H=0.0694. Taking that of O as 1.1111 on theoretical grounds (the supposed compound nature of air, etc.,) he calculates the above value. (Erdmann's Journ. für Prak. Chem., 8, 1836, 374; Records of Gen. Sci., R. D. Thomson, 1836, 179.)

J. Dumas: 1.0012 (O = 16); 6.2575 (O = 100).

Determined by nineteen experiments on the reduction of cupric oxide with pure hydrogen. The gas was made from pure materials and was passed through solutions of plumbic nitrate and argentic sulphate, and over potash, and dried with cold sulphuric acid or with phosphoric acid. The weighings of the oxide and of the reduced copper were made in vacuo. [Dumas corrected the results obtained for the air contained in the sulphuric acid, but does not explain how he estimated it, while certain other possible corrections are not mentioned.] The mean of the corrected results is 12.515. The extreme difference is 0.09 for O = 100. Without the correction for absorbed air the mean is 12.533, [or 1.00264]; maximum 12.583; minimum 12.481. (Paris Comptes Rend., 14, 1842, 537.)

ERDMANN and MARCHAND: 1.0016 (O = 16); 6.26 (O = 100).

Determined by eight experiments on the reduction of cupric oxide with hydrogen, the number is the mean of the results. In four of the experiments the correction for vacuum was calculated. These gave H=12.548; extreme difference, 0.067. In four experiments the weighings were made in vacuo. These gave a mean of 12.492, with an extreme difference of 0.015. The oxide employed was either copper scale or was produced from cupric nitrate. The hydrogen was made from pure zinc and sulphuric acid, and was purified with potash in solution and in lumps, mercuric chloride, sulphuric acid, and chloride of calcium. In the

last five experiments the gas was also passed over red-hot copper to remove traces of oxygen.) (Erdmann's Journ. für Prak. Chem., 26, 1842, 461.)

J. S. STAS: 1.0025 (O = 16).

From all the investigations that have been made on the specific gravity of the gases, the composition of water, etc., Stas is inclined to believe that the atomic weight of hydrogen cannot be less than above. Stas found that 100 silver were equivalent to 49.5978 ammonium chloride. [If N=14.044, and Cl=35.457, this relation would give $H\doteq 1.0074$.] (Stas, Untersuch. über. Chem. Prop., Leipzig, 1867.)

J. Thomsen: 1.0025 (O = 16).

Thomsen made three experiments on the oxidation of a known volume of hydrogen by cupric oxide, and five experiments on the combustion of a known volume of hydrogen in oxygen, which proved that 2 litres of hydrogen gave 1.6082 grammes of water under normal conditions, and at latitude 45° . According to Regnault, 1 litre of oxygen and 2 litres of hydrogen would weigh 1.6084 grammes. Hence 1 volume oxygen and 2 volumes hydrogen form water; and if H = 1, O = 15.96, [or if O = 16, H = 1.0025.] (Berlin, Be. der Chem. Ges., 3, 1870, 928.)

INDIUM.

Bunsen found the specific heat of In 0.565 and 0.574, which correspond to an atomic weight of about 114. (Poggend. Annal., 141, 28.)

F. Reich and T. Richter: 111.39 (O = 16).

In one experiment pure indium was dissolved in nitric acid, the oxide precipitated with ammonia and weighed. This experiment gave In = 463.4 for O = 100, and on the supposition that the metal was di-atomic. In a second experiment indium sulphide was dissolved in nitric acid, and the resulting sulphuric acid precipitated with barium chloride. This gave In = 464.9. The number taken is the mean. 8 = 200. The metal was prepared from the oxide. After the removal of lead, etc., with hydrogen sulphide, the oxides

of iron and indium were precipitated with ammonia, the precipitate dissolved in acetic acid and impure indium sulphide reprecipitated. This operation was repeated, and the last traces of iron were removed by partial precipitation with ammonia. (Erdmann's Journ. für Prak. Chem., 92, 1864, 484.)

C. Winkler: 107.754 (O = 16).

Determined by decomposing the nitrate by heat, and weighing the resulting oxide. The mean result of three experiments was In = 35.918 for O = 8, and assuming the univalence of the metal. Extreme difference, 0.079. Metallic indium was prepared by solution of the impure sulphide in chlorhydric acid, precipitation of indium by barium carbonate, solution in sulphuric acid, and precipitation by ammonia of the oxide which was reduced by hydrogen. [This indium seems to have contained iron.] (Erdmann's Journ. für Prak. Chem., 94, 1865, 1.)

C. Winkler: 113.439 (O = 16).

In two experiments the double chloride of gold and sodium was decomposed by pure indium, giving 37.73 and 37.80 for O = 8, and assuming univalence for the metal. In two experiments the nitrate was decomposed by heat, giving In = 37.845 and 37.879. In one experiment the oxide was precipitated from nitric acid solution by ammonia. This experiment gave In = 37.811. The number taken is the mean. The impure indium sulphide was purified as in Winkler's former determination with barium carbonate, but this process requires to be repeated several times. The reduction of the oxide was performed with sodium, the excess of which was removed from the regulus by cupellation in soda. (Erdmann's Journ. für Prak. Chem., 102, 1867, 282.)

R. Bunsen: 113.76 (O = 16).

Determined by converting metallic indium into oxide by means of nitric acid and heat. He seems to regard the experiment only as confirmatory of Winkler's. The metal was the same which served for the determination of the specific heat, and was carefully tested for all impurities. (*Poggend. Annal.*, 141, 1870, 28.)

IODINE.

Dumas determined the specific gravity of iodine vapor. It answers to an atomic weight of about 127. (Annal. de Chim. et de Phys., 33, 1826, 337.)

L. J. GAY-LUSSAC: 123.9 (O = 16).

100 parts of iodine were found equivalent to 26.225 parts of zinc. [If Zn = 65, these figures give the atomic weight at 128.9.] (Poggend. Annal., 14, 1828, 559; Annal. de Chimie, 91, 1814, 5.)

W. PROUT: 126 (O = 16).

Prout found 100 parts of iodine equivalent to 25.8 parts of zinc. [If Zn = 65, this gives I = 125.97.] (Thomson's Annals of Phil., 6, 1815, 323.)

T. Thomson: 124 (O = 16); 775 (O = 100).

Thomson found 20.5 potassic iodide = 19.75 zinc iodide, = 20.75 plumbic nitrate. [If K = 39.1, and plumbic nitrate = 331, the relation given leads to an atomic weight of 124.41.] Thomson thinks that his iodine may have been somewhat impure, as he purified it only by sublimation. (Thomson's System of Chem., 7th ed., 1, 1881, 81.)

J. Dumas: 126.13 (O = 16).

Dumas determined the density of iodine vapor at 8.716 for air = 1. [Referred to the molecular weight of oxygen, this density gives the above number for the atomic weight.] Dumas thinks it probable that it can be more accurately determined by analysis. (Annal. de Chim. et de Phys., 33, 1826, 387.)

J. J. Berzelius: 126.26 (O = 16); 789.14 (O = 100).

Determined by decomposing a known weight of argentic iodide in a current of chlorine, melting the chloride and expelling free chlorine by atmospheric air. The number is the mean of two experiments; difference, 0.01. Ag = 1351.607; Cl = 442.653. The iodide was prepared by precipitation from a solution of potassic iodide with argentic nitrate. The first portion of the precipitate was set aside as possibly contaminated with chlorine. (Poggend. Ann., 14, 1828, 562.)

C. Marignac: 126.844 (O = 16).

In five experiments a known weight of silver was dissolved in nitric acid and precipitated by a known amount of potassic iodide according to Pelouze's modification of Gay-Lussac's method. The mean result was 100 Ag = 153.74 KI in air; extreme difference, 0.14. Stas has recalculated this result for Ag = 107.93, and K = 39.137. The atomic weight so found is, in vacuo, 126.847. In three experiments a known weight of silver was dissolved and precipitated as iodide; mean result, 100 Ag = 217.511 iodide. Extrème difference, 0.04. From these data Stas gets I = 126.84. The iodine was purified by recrystallization as potassic iodate. The methods employed by previous experimenters were ineffectual. (Berzelius' Jahresbericht, 24, 75; Bibl. Univ. de Genève, 46, 1842, 367; also, Stas, Untersuch. über Chem. Prop., 153.)

E. MILLON: 126.07 (O = 16); 787.915 (O = 100).

Three experiments were made on the decomposition of potassic iodate. The mean loss of oxygen was 22.478 per cent; extreme difference, 0.03. If K = 488.94, this gives I = 1580.93. In three experiments argentic iodate, which had been dried for a long time at 200°, was employed, which lost 17.0467 per cent. oxygen; extreme difference, 0.03. If Ag = 1349.01, these data give I = 1570.73. [Berzelius cites this as an atomic weight determination; Millon, however, seems to have regarded it only as a confirmation of Berzelius' number.] Millon prepared pure iodine by passing a current of chlorine through a solution of KI till the precipitated I was redissolved, and reprecipitating with an excess of KI. (Annal. de Chim. et de Phys., (3,) 9, 1843, 407.)

V. A. JACQUELIN: 125.6 (O = 16); 785 (O = 100).

Determined by the analysis of iodic acid with silver. The acid was prepared by the oxidation of iodine with nitric acid of sp. gr. 1.5. The purity of the preparation does not seem to have been tested. Ag = 1351. (Erdmann's Journ. für Prak. Chem., 51, 1850, 458; Annal. de Chim. et de Phys., (3,) 30, 1850, 832.)

J. Dumas: 127 (O = 16).

Determined by the conversion of argentic iodide into chloride in a current of dry chlorine. Two experiments gave 127.04 and 127.01 for Ag = 108; Cl = 35.5. In Gmelin-Kraut's Handbuch these data are recalculated for Ag =

IODINE. 68

107.93 and Cl = 35.457, giving I = 126.941 and 126.928. The argentic iodide used was prepared from zinc iodide which had been prepared from iodine in large crystals. The argentic iodide was fused. (Annal. de Chim. et de Phys., (3,) 55, (3,) 56, (3,) 56, (3,) 56, (3,) 56, (3,) 63.)

J. S. STAS: 126.851.

Stas ascertained the molecular weight of argentic iodide as follows:

In two complete analyses, a known weight of argentic iodate was decomposed by heat in a current of pure, dry nitrogen. The oxygen set free was caught by hot copper and weighed, as well as the residual argentic iodide. In one experiment argentic iodate was dissolved in ammonia, precipitated by sulphuric acid, (to secure advantageous division of the salt,) and reduced while in suspension by a slow current of sulphurous acid. The mean molecular weight reached was 234.779; extreme difference, 0.063. The samples of iodate employed were prepared: (1.) From argentic sulphate and potassic iodate, mixed boiling, the latter in excess, thorough washing and drying in air freed from organic particles; (2.) By the reaction of potassic iodate on argentic hyposulphite. The purity of the salt was carefully tested.

Stas ascertained the composition of argentic iodide as follows:

(1.) A known weight of argentic nitrate was precipitated by hydro-iodic acid and the argentic iodide washed, dried, and weighed in the same vessel. (2.) A known weight of Ag was dissolved in nitric acid, converted into sulphate, dissolved in very dilute sulphuric acid, and precipitated with hydro-iodic acid. The precipitate was washed at temperatures increasing up to 90°. (3.) A known weight of argentic sulphate was allowed to react on a known and nearly equivalent weight of iodine in an aqueous solution of sulphurous and sulphuric acids at 10°, and in the dark, till all the iodine was taken up. The excess of iodine was titrated with silver solution, and the iodide weighed. This method was employed in two experiments. (4) differed from (3) mainly in the conversion of the iodine into ammonium iodide before bringing it into contact with argentic sulphate. Four experiments were made by the last method.

The mean composition of the iodide, as derived from all the experiments, is 100 Ag = 117.5343 iodine. From these data Stas calculates the atomic weight of I at 126.857, and

that of silver at 107.928. [The sum of these weights is not the molecular weight, and this, as well as recalculation of the data, shows that the number is a misprint for 126.851. Stas' results are, therefore, even closer to Marignac's than

his memoir would indicate.]

Most of the experiments were made with iodine prepared by the decomposition of nitric iodide decomposed in a large volume of water at 65°. The iodine was further purified by distillation over barium oxide and by other means. For the preparation of silver see that metal. All possible precautions were observed in the preparation of all reagents and in the conduct of the experiments. (Stas, Untersuch. über Chem. Prop. Leipzig, 1867.)

IRIDIUM.

Regnault determined the specific heat of iridium. It corresponds to an atomic weight of about 198. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 197.19 (O = 16).

Berzelius determined this value from analysis of potassium chloro-iridiate. This salt reduced in hydrogen lost 29 per cent., the same quantity lost by the corresponding platinum salt, (vide platinum.) Berzelius originally calculated the atomic weight of the platinum metals both from the loss of chlorine of these double salts and from the relation between the metal and the potassic chloride left after reduc-In his Lehrbuch he points out the impossibility of complete desiccation, and resorts exclusively to the latter method of calculation. With respect to iridium he merely asserts that its atomic weight is the same as that of platinum, without there, or elsewhere, giving data as to the amounts of iridium and potassic chloride found in the reduced salt. It is, therefore, open to question whether he assumed the identity from the loss on reduction or not. [If Pt = Ir, and if KCl = 74.594, the value fellows; see platinum.] Osmium and iridium were separated by fusion with nitre, solution, and distillation. The residue was fused with potassic chloride and sodium carbonate. On solution the iridium remains behind. This residue was repeatedly roasted and reduced to drive off osmium compounds. The potassium chloro-iridiate was formed from the pure metal. (Poggend. Ann., 13, 1828, 468; Kongl. Vet. Acad. Handl., 1828.)

C. E. CLAUS: W. M. WATTS: 197.6 (O = 16).

Watts recalculated two analyses of potassium chloro-iridiate by Claus from the loss in reduction, and for Cl = 35.457, (Stas.) From one analysis he finds K = 39.87, and Ir = 198.56; from the other K = 89.93, and Ir = 196.62. (Chem. News, 19, 1869, 302.)

IRON.

Regnault, Kopp and others have determined the specific heat of this metal. It corresponds to an atomic weight of about 56. (Gmelin-Kraut, l. c.)

L. J. Thenard: F. H. Wollaston: 66.2 (O = 16); 345 (O = 100).

Thenard determined the composition of the oxide at 22.5 0 and 77.5 Fe, whence Wollaston calculates the value. (Phil. Trans., 104, 1814, 21.)

J. J. Berzelius: 64.27 (O = 16); 839.218 (O = 100).

Determined by repeated experiments on the oxidation of iron, such as is used for piano wire, with nitric acid. The carbon was determined and allowed for. Berzelius in his Lehrbuch shows that the error in this determination lay in the unsuspected presence of soluble silica and on reänalysis he found enough of it to correct the number when taken into account. (Poggend. Ann., 8, 1826, 185.)

G. Magnus: 54.25 (O = 16); 339.06 (O = 100).

Magnus' experiments were made by reducing ferric oxide in a current of hydrogen at about the temperature of boiling mercury. He regarded them simply as comfirmatory of Berzelius' number. (*Poggend. Ann.*, 3, 1825, 84.)

F. Stromeyer: 66.6 (O = 16).

Determined by reducing ferric oxide at a red heat by hydrogen. The oxide is reduced only with great difficulty at a lower temperature. The mean of the experiments gave the oxygen contents at 30.15 per cent., [whence I have calculated the value.] (Poggend. Ann., 6, 1826, 475.)

H. Capitaine: 51.36 (O = 16); 321 (O = 100).

Determined by the peroxidation of galvanically reduced iron and by measuring the hydrogen evolved on the solution of iron in sulphuric acid. (Annal. de Chim. et de Phys., (8,) 2, 1841, 126.)

H. Wackenboder: 66.48 (O = 16).

Wackenroder helped Stromeyer in his reduction of ferric oxide, of which he gives the details. He also describes five experiments of his own, which gave the oxygen contents of ferric oxide at from 30.01 to 30.38. He took no precautions to purify his hydrogen and thinks that the loss of oxygen may have been apparently reduced. [30.195 oxygen corresponds to the above value for Fe.] (Berzelius' Jahresbericht, 24, 1844, 121; Archiv. der Pharm., 36, 1844, 22.)

Svanberg and Norlin: 66.97 (O = 16); 349.809 (O = 100).

In seven experiments a known weight of iron was dissolved in nitric acid and the salt decomposed by heat. The operation was performed in a glass flask. The mean result in vacuo, was 349.104; extreme difference, 0.803. In seven experiments ferric oxide was reduced with purified hydrogen. The mean was Fe = 350.514; extreme difference, 0.735. The number taken is the mean of all the experiments, in vacuo. Berzelius in his Lehrbuch cites the experiments and, by neglecting the reduction to vacuum, gets a slightly different number. He also expresses a preference for the experiments by reduction: (Berzelius' Jahresbericht, 24, 1844, 121; and Poggend. Ann., 62, 1844, 270.)

J. J. Berzelius: 66.05 (O = 16); 350.82 (O = 100).

Berzelius, as a check on the last determination, made two experiments on the oxidation of iron by nitric acid with special precautions against partial reduction. The number is the mean; difference, 0.101. The iron was melted down with glass and magnetic oxide. In his *Lehrbuch* he adopts the mean of these experiments and Svanberg and Norlin's reduction determinations. (*Poggend. Ann.*, 62, 1844, 270.)

ERDMANN AND MARCHAND: 56.016 (O = 16); 350.1 (O = 100).

Erdmann and Marchand made eight experiments on the reduction of ferric oxide in a carefully purified current of

hydrogen. The weighings of the metal were made in vacuo to avoid possible reoxidation in displacing the gas by sir. The number is the mean of the experiments; extreme difference, 1.4 for O = 100. The ferric oxide was prepared by incineration of the oxalate, moistening the residue with nitric acid and reheating. (Erdmann's Journ. für Prak. Chem., 33, 1844, 1.)

L. E. RIVOT: 64.25 (O = 16); 839.01 (O = 100).

Determined by two experiments on the reduction of pure ferric oxide in a current of hydrogen. 100 parts of oxide gave 69.31 and 69.35 parts metallic iron. (Annal. de Chim. et de Phys., (3,) 30, 1850, 188.)

E. MAUMENÉ: 56.0016 (O = 16); 350.01 (O = 100).

Maumené made six experiments by dissolving iron wire in aqua regia, precipitating with ammonia, heating the precipitate to redness and weighing. The number is the mean; extreme difference, 0.34. Maumené had convinced himself by analysis of the extreme purity of the wire. (Erdmann's Journ. für Prak. Chem., 51, 1850, 350.)

J. Dumas: 56.2 (O = 16).

Two experiments on the precipitation of ferric chloride by argentic nitrate gave each 28.1. A single experiment by the same method on ferrous chloride which was slightly yellow, gave 28.1. An experiment made on ferrous chloride, which had been heated in a current of hydrogen and of HCl and was colorless, but contained metallic iron, gave when the admixture was determined, 27.99. Dumas takes Ag = 108; Cl = 85.5, (Annal. de Chim. et de Phys., (3,) 55, 1859, 157.)

LANTHANIUM.

W. F. Hillebrand has determined the specific heat of metallic lanthanium. It corresponds to an atomic heat of 6.23, if the atomic weight is taken at 139. (*Poggend. Ann.*, 158, 1876, 82.)

Several investigations on the atomic weight of lanthanium were made previous to Mosander's announcement of the discovery of didymium. F. J. Otto found it 108.41 shortly after its discovery, and announced it in his translation of

Graham's chemistry. (Gmelin's Handbuch, 5th ed., 1, 46.) Choubine, from analysis of the chloride and of the sulphate, arrived at 108.45. (Erdmann's Journ. für Prak. Chem., 26, 1842, 448.) Rammelsberg determined it from the sulphate, which was rose-colored, at 133.17. (Poggend. Ann., 55, 1842, 65.) R. Hermann found La = 144 from rose-colored sulphate. (Erdmann's Journ. für Prak. Chem., 30, 1843, 198.)

C. G. Mosander: 139.2 (O = 16); 870 (O = 100).

Mosander says that his experiments show the true value to be in the neighborhood of 680, (the metal being assumed bivalent,) but that his salts were not pure, and the determination of little value. (*Poggend. Ann.*, 60, 1843, 301.)

C. Marignac:
$$141.12$$
 (O = 16); 882 (O = 100).

Eleven experiments were made on the decomposition of the sulphate by barium chloride. The results vary greatly. Marignac wrote later (Annal. de Chim et de Phys., (8,) 38, 1853, 148) that experiment had convinced him of the incorrectness of this determination, and that the true value is about 575. (La bivalent.) (Liebig's Ann., 71, 1849, 306.)

M. Holzmann:
$$139.22$$
 (O = 16); 870.15 (O = 100).

In three experiments La sulphate was decomposed by ammonium oxalate. In the filtrate from the precipitated oxalate the sulphuric acid was determined as barium salt. The oxalate was decomposed by heat, and the lanthanium oxide weighed. These experiments gave a mean of 580; extreme difference, 5.6; for bivalent lanthanium. In three experiments the iodate was decomposed by oxalic acid, the oxide determined as before, and the iodine titrated by Bunsen's method. These experiments gave a mean of 580.2; extreme difference, 5.3. S = 200; Ba = 855. In the preparation ration of the salts analyzed the cerium was separated by peroxidation with magnesium oxide and precipitation as basic sulphate. After the removal of yttrium by potassic sulphate, the lanthanium and didymium salts were separated, by making a saturated solution of the sulphates at a temperature of three or four degrees, and gradually raising the temperature. Lanthanium salt then crystallizes out nearly pure. The purification was repeated until the salts were not discolored when heated in an open crucible over the glass-blower's lamp. Bunsen assisted at this investigation. (Erdmann's Journ. für Prak. Chem., 75, 1858, 843.)

C. Czudnowicz: 140.3 (O = 16); 876 (O = 100).

Czudnowicz especially disclaims making this as an atomic weight determination and he adopts Holzmann's value. The salt analysed was the sulphate, and the method the same as that employed by Holzmann. (Erdmann's Journ. für Prak. Chem., 80, 1860, 81.)

R. Hermann: 139.32 (O = 16); 870.75 (O = 100).

Hermann analyzed the carbonate by decomposing it over mercury by sulphuric acid, and measuring the carbon di-oxide liberated. The residue was heated to redness and weighed. This experiment gave La = 580.4, the metal being assumed as bivalent. The carbonate was prepared by precipitating the sulphate with sodium bicarbonate. In three experiments the sulphate was decomposed by ammonium oxalate and the oxide, obtained by incinerating the These analyses gave La = 580.7. oxalate, weighed. one experiment the chloride was analysed with argentic nitrate, giving La = 580.4. The number taken is the mean: extreme difference 2.3. In the preparation of the salts, cerium was separated as basic sulphate, La and didymium were partially separated by crystallization after which a portion of the nearly pure sulphate was precipitated by ammonia, and this precipitate digested with the mother liquor. Didymium sulphate is by this means completely precipitated. 8 = 200; Cl = 443.2; C = 75. Hermann remarks that his former determination was made with impure material. (Erdmann's Journ. für Prak. Chem., 82, 1861, 395.)

H. ZSCHIESCHE: 135.27 (O = 16).

Determined by six experiments on the sulphate. The water was driven off at 280°, and the anhydrous salt exposed to a white heat until the weight became constant, and on being tested, showed no sulphur. The mean result was La = 45.09; extreme difference, 1.15. In preparing the salt from cerite, the cerium was peroxidized with red lead and nitric acid and was precipitated as basic nitrate. The didymium was separated by partial precipitation with oxalic acid and concentration, these operations being repeated as long as the absorption lines of Di were perceptible in the spectroscope. A correction was made for the loss of weight of the crucible, and there was no dust upon its sides. S = 16. (Erdmann's Journ. für Prak. Chem., 104, 1868, 174; 107, 1869, 72.)

C. Erk: 136.39 (O = 16).

Determined by analysis of the sulphate by the method employed by Holzmann. The bases were separated by the methods which Hermann used. Yttrium was also eliminated. Fresenius in his Zeitschrift, 10, 509, objects to the details of the Erk's manipulation of barium sulphate. (Kopp's Jahresbericht, 1870, 319; Jena Zeitschr. für Med. und Nat., 6, 1870, 299.)

D. Mendelejeff: 180 (O = 16).

As La forms but one oxide, the salts of which are not, according to Marignac, isomorphous with those of the lower oxide of didymium, Mendelejeff concludes that it belongs to the same group, but that its oxide is a binoxide, and its atomic weight 180. (Liebig's Ann., Suppl., 8, 1871, 190.)

C. Marignac: 138.75 (O = 16).

By heating the sulphate till all acid was expelled, Marignac, in two experiments, determined La (bivalent) at 92.52 and 92.56; by precipitation with ammonia and heating at 92.24 and 92.48. The sulphate was purified by a great number of partial recrystallizations, and showed only doubtful traces of didymium in the spectroscope. S=16. (Annal. de Chim. et de Phys., (4,) 30, 1878, 67.)

P. T. CLEVE: 139.15 (O = 16).

Determined by the conversion of lanthanium oxide into sulphate. The number is the mean; extreme difference 0.55. The oxide was purified from didymium by repeated partial precipitation from nitric acid solution with ammonia, basic didymium nitrate going down. The lanthanium was finally precipitated with oxalic acid. The oxide was found to be spectroscopically pure by Thalén. (Kopp's Jahresbericht, 1874, 257; Paris Bull. de la Soc. Chim., 21, 196, 246, 344.)

LEAD.

Regnault, Kopp and others have determined the specific heat of lead. It answers to an atomic weight of about 207. (Gmelin-Kraut, l. c.)

J. J. Berzelius and F. H. Wollaston: 207.4 (O = 16); 1295 (O = 100).

Berzelius found 16.5 parts carbon di-oxide equal to 83.5 lead oxide, whence the value, if C=75.4. [If C=12, these figures give lead at 296.67.] Berzelius also determined the composition of the oxide at 7.15 oxygen and 92.85 lead, giving Pb=207.52 or 1297. (Phil. Trans., 104, 1814, 20.)

J. J. Berzelius: 207.12 (O = 16); 1294.498 (O = 100).

Determined by the reduction of a known weight of oxide of lead by hydrogen and the weight of the resultant lead; mean of four nearly coincident experiments. (*Poggend. Ann.*, 8, 1826, 184.

- -Longchamp is credited in some books with an atomic weight determination of lead. He made none, but only speculated on the composition of minium, taking Berzelius' determination as a basis. (Annal. de Chim. et de Phys., 34, 1827, 105.)
 - J. J. Berzelius: 207.078 (O = 16); 1294.242 (O = 100).

This value is the mean of six experiments on the reduction of the oxide in a current of hydrogen. The oxide was produced by the decomposition of the nitrate by heat. As this compound reacts upon Pt, the crucible was lined out with a coating of a very basic nitrate, which prevented the lumps of neutral salt from coming in contact with the crucible. The glass in which the oxide was reduced was not attacked. [The third analysis is miscalculated. It should show an atomic weight of 1295.595. The mean is, therefore, as above, and the extreme difference 2.421.] (Poggend. Ann., 19, 1830, 314.)

J. J. Berzelius: 207.14 (O = 16); 1294.645 (O = 100).

In his *Lehrbuch*, Berzelius selects five analyses made by the above method, three of them the same. These give the above mean, with an extreme difference of 0.704 for O = 100. (*Lehrbuch*, 3, 1219.)

E. Turner: 207.3 (O = 16).

Determined by experiments on the conversion of metallic lead and of oxide of lead into the sulphate by solution in

nitric acid and evaporation with sulphuric acid. In three experiments, Turner found 100 lead = 146.401 sulphate; extreme difference 0.055. Berzelius had found 100 Pb = 146.419 sulphate; extreme difference 0.078. Turner takes the mean of his own and Berzelius' determinations, 146.41. In one experiment Turner found 100 oxide = 185.92 sulphate. Combination of these results gives Pb = 103.6 [or more accurately 103.65.] The oxide was prepared from subnitrate. The lead was prepared from plumbic acetate which was converted into carbonate, then into nitrate, in which form it was recrystallized, then again into carbonate, and reduced by black flux. On testing, it was found perfectly pure. Weighings reduced to vacuum. (Phil. Trans., 123, 1888, 524.)

C. Marignao: 207.04 (O = 16).

Marignac made four experiments on plumbic chloride by Pelouze's modification of the silver titration method. He found Pb = 108.57-.49-.55-.46. The number taken is the mean. The salt was titrated cold, argentic chloride being soluble in hot solutions of plumbic nitrate. The plumbic chloride was purified by recrystallization, and, after being pulverized, was dried at about 200°. According to Marignac there is no difficulty in desiccating it completely at this temperature. Ag = 108; Cl = 35.5. Marignac found it impossible to convert the chloride into the sulphate completely. (Bibl. Univ., Arch. des Sciences, (2,) 1, 1858, 223.)

J. Dumas: 207.1 (O = 16).

From a single experiment on the precipitation of the chloride with argentic nitrate. The chloride used was heated for twelve hours in a current of dry HCl, and the amount of water retained determined. Dumas found it impossible entirely to desiccate the salt without decomposition, drying at 250° does not desiccate it. Cl = 35.5; Ag = 108. (Annal. de Chim. et de Phys., (3,) 55, 1859, 129.)

J. S. Stas: 206.926 (O = 16).

According to the mean of 10 syntheses of plumbic nitrate, 100 lead = 159.9703 nitrate; extreme difference, 0.023. If N = 14.044, this relation gives Pb = 206.918. Stas also made six syntheses of the sulphate, which gave in mean 100 Pb = 146.4275 sulphate; extreme difference, 0.024. If S = 82.0742, this relation gives Pb = 206.934. The syntheses were made in the same way as in the determination of the

atomic weight of silver. The drying of the nitrate could be accomplished only in vacuo and at about 155°. The weighings are for vacuum. The lead used was prepared from commercial acetate by precipitation with metallic lead, of copper, etc., conversion into sulphate, then into carbonate and reduction by potassic cyanide or black flux. (Stas, Untersuch. über Chem. Prop. Leipzig, 1867, 824.)

LITHIUM.

Regnault has determined the specific heat of lithium. It corresponds to an atomic weight of about 7. (Gmelin-

Kraut, L. c.)

The earliest determinations of this constant seem to have been made with a double salt of lithium and potassium, at all events with a very impure material. According to Arfvedson, 420.4 lithium chloride give 1322.4 argentic chloride, whence he deduces as the atomic weight the number 127.757 [or 10.22.] (Poggend. Ann., 8, 1826, 189.) L. N. Vauquelin found 430 lithium sulphate equivalent to 875 barium sulphate. [If S = 32; Ba = 137.08, this relation gives Li = 9.27.] Vauquelin does not describe the preparation of his salt. (Annal. de Chem. et de Phys., 7, 1818, 287.) C. G. Gmelin found Li = 191.21 [or 7.65.] (Poggend. Ann., 15, 480; Gilbert's Ann., 62, 1819, 399.) Kralovanszky by two analyses of the sulphate with barium chloride got Li at from 10.096 to 10.168 (Liebig's Ann., 121, 94; Schweigger's Journ., 54, 1828, 231.) Thomson and Stromeyer also each got similar values. (Thomson's System of Chem., 7th ed., 1, 1881, 420.)

R. Hermann: 6.085 (O = 16); 88.03 (O = 100).

Experiments were made on the carbonate by decomposing it with acid over mercury, and measuring the resultant di-oxide. For C = 75.33, these determinations give Li = 88. Several experiments were also made by analyzing the sulphate with barium chloride. For S = 201.06 and Ba = 856.88, these give Li = 38.05. Hermann precipitated lithium carbonate with ammonium carbonate, and subsequently converted it into sulphate. The chloride was prepared from the phosphate by Berzelius' method. (Poggend. Ann., 15, 1829, 480.)

J. J. Berzelius: 6.633 (O = 16); 40.83 (O = 100).

Berzelius found that 1.874 lithium sulphate gave 3.9985 barium sulphate, and calculated this relation for S=200.75; Ba = 855.29. He also found 4.4545 melted carbonate = 6.653 sulphate, but rejected the analysis. (*Lehrbuch*, 3, 1229, and *Jahresbericht*, 10, 1830, 96.)

R. HAGEN: 6.67 (O = 16).

Hagen precipitated lithium sulphate with barium chloride, and found that 0.852 dry lithium sulphate gave 1.8195 barium sulphate whence he calculates Li = 6.493. [If Ba = 137.08; S = 32; this relation gives Li = 6.57.] (Poggend. Ann., 48, 1889, 363.)

J. W. MALLET: 6.95 (O = 16); 86.89 (O = 100).

In two experiments a known weight of lithium chloride was precipitated by argentic nitrate, and the argentic chloride weighed. In one experiment lithium chloride was titrated with argentic nitrate by Pelouze's method. The number is the mean; the extreme difference is 0.18 for O = 100. Mallet takes Ag = 1849.66; Cl = 443.28. The alkalis were separated from the lithium salt by repeated treatment with ether and alcohol. The salt was examined for impurities, and was fused with a little ammonium chloride to prevent the formation of oxy-chloride. (Silliman's Amer. Journ., (2,) 22, 1856, 849.)

L. Troost: 6.5 (O = 16).

Troost found this number from analysis of the carbonate which had been crystallized from water containing carbon di-oxide and dried at 200°, but does not regard it as definitive. (Annal. de Chim. et de Phys., (8,) 51, 1857, 111.)

J. W. MALLET: 7 (O = 16).

Troost having objected to Mallet's former method of determination, he redetermined it by precipitating the sulphate with a standard solution of barium chloride, the precipitating power of which had been tested on the sulphates of magnesium and sodium. This method was adopted to avoid the well-known imperfections of the sulphur determination. Compared with sodium sulphate the atomic weight of Li was found = 6.92 and 6.95. Compared with magnesium sulphate it was found = 7.07 and 7.09.

Mg = 24; Na = 23. The sulphate was prepared from carbonate, and dried somewhat below a red heat. (Silliman's Amer. Journ., (2,) 28, 1859, 349.)

K. Diehl: 7.026 (O = 16).

Determined by analysis of lithium carbonate with Bunsen's apparatus and in his laboratory. Four experiments; extreme difference, 0.024. C=12. The salt was purified from alkalis by precipitation as carbonate, resolution in acid and reprecipitation, repeated until the sodium line was no longer visible. Diehl found that precipitation of the sulphate with barium chloride gave a nearly constant error on account of the retention of lithium in the precipitate, and led to nearly the same results as Berzelius got. (Liebig's Ann., 121, 1862, 93.)

L. TROOST: 7 (O = 16).

Troost found 1.309 grammes lithium chloride = 4.42 argentic chloride, and 2.75 lithium chloride = 9.3 argentic chloride. From these analyses he deduces the values 7.03 and 6.99. By decomposing the carbonate, dried at 100°, with silicic acid, he found 0.97 carbonate = 0.577 carbon di-oxide and 1.782 carbonate = 1.059 di-oxide, and infers for Li 7 and 7.02. In one experiment the carbonate was converted into sulphate. 1.217 carbonate gave 1.808 sulphate. Troost calculates Li = 7.06. [If Cl = 35.457; Ag = 107.93; C = 12; S = 32; these determinations give, in the same order as above, 7.01; 6.94; 6.98; 7.02; 7.07.] The carbonate was purified by solution in water containing carbon di-oxide, and reprecipitation by boiling, the operation being repeated until the salt was spectroscopically pure. (Paris Comptes Rend., 54, 1862, 366.)

J. S. Stas: 7.022 (O = 16).

According to the mean of three determinations, 100 parts of silver = 39.358 lithium chloride; extreme difference, 0.005. If Ag = 107.93; Cl = 35.457; this ratio gives Li = 7.022. This value is confirmed by experiments on the conversion of the chloride into the nitrate, the results of which give Li = 7.018. The comparison with silver was made according to Pelouze's modification of the silver titration method. The chloride was purified from alkalis, after Preliminary treatment with ether and alcohol, by pouring the dissolved salt into a boiling solution of ammonium car-

bonate containing ammonia in excess. All weighings reduced to vacuum. (Stas, Untersuch. über Chem. Prop., Leipzig, 1867.)

MAGNESIUM.

Regnault and Kopp have each determined the specific heat of this metal. It answers to an atomic weight of about 24. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 26.3 (O = 16); 158.139 (O = 100).

Determined by dissolving magnesium oxide in dilute sulphuric acid, evaporating and heating to incipient redness. 100 oxide were found = 293.985 sulphate. The sulphate was perfectly soluble in water and had therefore lost none of its acid. The oxide was purified by solution in an aqueous solution of carbon di-oxide and reprecipitated by boiling. S = 200.75. (Poggend. Ann., 8, 1826, 188; and Lehrbuch, 3, 1227.)

Marchand and Scheerer recalculated this analysis for S = 200 and reached the value 157.74. They assert that the oxide may have contained alkalis and that the sulphuric acid carries off magnesium sulphate in volatilizing. (Erdmann's Journ. für Prak. Chem., 50, 1850, 892.)

W. Henry: F. H. Wollaston: 23.36 (O = 16); 146 (O = 100).

Henry found that magnesium sulphate contained 88 per cent. magnesium oxide. If S=200 the value follows. (*Phil. Trans.*, 104, 1814, 21.)

— Longchamp: 15.35 (O = 16).

In two experiments, Longchamp found that five parts of crystallized magnesium sulphate are equivalent to 4.91 barium sulphate. [If Ba = 137.08; S = 32, the number follows.] Longchamp found 53 per cent. water which is much too high. According to Marchand and Scheerer, the data for the anhydrous salt give Mg = 97.87, for S = 200; Ba = 856.8, [or 15.74.] (Annal. de Chim. et de Phys., 12, 1819, 265.)

L. J. GAY-LUSSAC: 23.62 (O = 16).

16.205 grammes crystallized sulphate were found equal to 15.345 barium sulphate, and 19.395 magnesium sulphate

to 18.8455 barium sulphate. Calculating from the anhydrous salt Gay-Lussac found from these experiments respectively Mg =147.28 and Mg = 148.09 for Ba = 856.8; S = 200. The salt was found to contain 51.43 water. [Calculated from the anhydrous salt these data give Mg = 23.55 and 23.68. Calculated from the hydrous salt (7 molecules water) the numbers give 24.14 and 24.41, if S = 32; Ba = 137.08.] Gay-Lussac remarks that the sulphate is partially decomposed at a red heat. (Annal. de Chim. et de Phys., 13, 1820, 308.)

T. Scheere: 24.16 (O = 16); 150.97 (O = 100).

Mean of six experiments on the precipitation of the sulphate with barium chloride. Extreme difference, 0.79. S = 200.75; Ba = 855.29. After weighing, the barium sulphate was treated with dilute HCl and the chloride thus extracted allowed for. (*Poggend. Ann.*, 69, 1846, 535.)

T. Scheerer: 24.21 (O = 16); 151.33 (O = 100).

Barium sulphate formed as in the last determination was fused with soda, the barium carbonate dissolved in HCl, and reprecipitated as sulphate. In the filtrate additional magnesia was found. If the error in the former determination was the same, its corrected value would be as above. (Poggend. Ann., 70, 1847, 407.)

Syanberg and Nordenfeldt: 24.72 (O = 16); 154.504 (O = 100).

Four experiments were made on the calcination of the oxalate, and three on the conversion of the magnesia so obtained into sulphate. The oxalate was dried at from 100° to 105° and heated to redness until the weight was constant. The oxide was dissolved in sulphuric acid, evaporated and the excess driven off by heat. The oxalate was prepared from the sulphate by precipitation with sodium carbonate and digestion with oxalic acid. The number is the mean of all experiments; extreme difference, 0.514. 8 = 200.75; C = 75.12; H = 12.48. (Erdmann's Journ. für Prak. Chem., 45, 1848, 473.)

According to Marchand and Scheerer, the data give Mg = 154.27 for S = 200; H = 12.5; C = 75.

MARCHAND and Scheerer: 24.03 (O = 16); 150.19 (O = 100).

Eleven experiments were made in calcining massive magnesium carbonate from Frankenstein, and weighing the

caustic magnesia formed. The carbonate was dried at 800°, and the carbon di-oxide, which passes off above 280°, was caught by caustic baryta solution and determined. The traces of carbon di-oxide not expelled by a yellow heat were set free by solution in chlorhydric acid and also determined as barium carbonate. The silicic acid, etc., were also determined. The mean in air is 150.34; in vacuo as above. Extreme difference, 0.57. C = 75. Eleven other experiments were made with comparatively impure material and less precaution, tending to confirm the above. (Erdmann's Journ. für Prak. Chem., 50, 1850, 409.)

T. SCHEERER: 24 (O = 16); 150 (O = 100).

By separating the neutral sulphates of magnesium and calcium by means of alcohol, Scheerer found that the magnesites used by Marchand and himself contained from one-fourth to one-half per cent. calcium oxide. This correction would make their determination almost exactly 250 or 24. (Liebig's Ann., 110, 1858, 236.)

V. A. JACQUELIN:
$$24.408$$
 (O = 16); 152.55 (O = 100).

Anhydrous, neutral magnesium sulphate, obtained by solution of the oxide in sulphuric acid and heating to redness, gave 33.56 per cent. pure oxide. The method adopted is not described. This oxide by treatment with sulphuric acid gave the original amount of sulphate. If S=200, the number follows. (Annal. de Chim. et de Phys., (3,) 32, 1851, 195.)

A. Macdonnell: 23.9 (O = 16).

Determined from analyses of anhydrous and of crystaltized magnesium sulphate. (*Brit. Assoc. Rep.*, 1852, part 2, 36; and *Kopp's Jahresbericht*, 5, 364.)

J. F. Bahr: 24.77 (O = 16); 154.842 (O = 100).

A known weight of purified magnesium oxide was dissolved in sulphuric acid, evaporated and heated to redness till the weight was constant. The number is the mean of three experiments; extreme difference, 0.515. The oxide was obtained from meteoric olivin. After removal of the heavy metals, the solution was evaporated to dryness with soda, washed and heated to redness. The oxide so obtained was dissolved in acetic acid, oxalic acid was added, the

solution evaporated nearly to dryness, and the oxalate thoroughly washed. Bahr says that the presence of alkalis could not be suspected. S=200. (Erdmann's Journ. für Prak. Chem., 56, 1852, 310; Efversigt af Akad. Færh., 1851, 303.)

Scheerer says that oxide so prepared retains carbonic acid, that sulphate is carried off in heating the sulphate to redness, and that the presence of alkalis is to be suspected. (Erdnam's Journ. für Prak. Chem., 56, 1852, 489.)

J. Dumas: 24.6 (O = 16).

Dumas made eleven experiments on the titration of magnesium chloride with argentic nitrate. He found great difficulty in preparing pure chloride, and does not feel confident of his results. The number is the mean; extreme difference, 0.28. Ag = 108; Cl = 85.5. The chloride was prepared from various salts, but was in all cases finally heated in an atmosphere of HCl. Dumas points out, however, that this process does not remove oxide if present. (Annal. de Chim. et de Phys., (8,) 55, 1859, 129.)

MANGANESE.

Regnault has determined the specific heat of manganese. It corresponds to an atomic weight of about 55. (Gmelin-kraut, l. c.)

J. J. Berzelius: 66.93 (O = 16); 855.787 (O = 100).

By dissolving manganese in nitric acid, evaporating and heating to a low red, Berzelius found 100 Mn = 142.16 oxide. It was not known at the time that the oxide might be partially reduced by this process. (*Poggend. Ann.*, 8, 1826, 185; and *Jahresbericht*, 9, 136.)

J. A. Arfvedson: 56.25 (O = 16); 351.56 (O = 100).

From 1.508 chloride Arfvedson obtained 3.408 argentic chloride. If Ag = 1851.607; Cl = 221.325; the number follows. (Berzelius' Jahresbericht, 9, 1829, 136; Afhandl. i. Fysik., 6, 236.)

E. Turner: 64.9 (O = 16).

Turner analyzed the carbonate in an apparatus similar to Bunsen's. He found 34.72 per cent. carbon di-oxide and 8.427 water. For C=6, he calculates Mn=28.024. By dissolving the protoxide in sulphuric acid, evaporating and heating to redness, he found 9 oxide = 19.01 sulphate. If S=16, this gives Mn=27.96. A second experiment gave 27.93. From 12.47 Mn chloride he obtained 28.42 argentic chloride. [If Cl=35.5, Ag=108; this gives Mn=54.9.] The carbonate was obtained by precipitation with potassium carbonate. The protoxide was obtained by reduction of higher oxides in hydrogen. The chloride was melted in a current of HCl gas. (Edinb. Trans., 11, 1831, 143.)

J. J. Berzelius: 66.34 (O = 16); 845.9 (O = 100).

Berzelius repeated Turner's experiments, taking larger quantities. From the chloride he got from 845.84 to .96; from the sulphate from 846.03 to .29. Ag = 1351.607; Cl = 221.825; S = 201.165. (Berzelius' Jahresbericht, 9, 1830, 136.)

J. J. Berzelius: 66.14 (O = 16); 344.684 (O = 100).

In his Lehrbuch he apparently takes the analyses of the chloride above cited, recalculated for Cl = 221.64; Ag = 1849.66. (Lehrbuch, 3, 1224.)

R. Brandes: 67.06 (O = 16); 356.602 (O = 100).

Determined by analysis of crystallized chloride. The chlorine was determined by precipitation with silver. The Mn was precipitated as carbonate, and converted into oxide by heat. The water was determined by difference, and the composition of the oxide was assumed to be as given by Berzelius, (!) whose values for Ag and Cl were also taken. (Poggend. Ann., 22, 1831, 256.)

K. VON HAUER: 54.98 (O = 16); 848.632 (O = 100).

Determined by nine experiments on the reduction of the sulphate to sulphide by heating the salt in a current of hydrogen sulphide. The reduction was performed in a porcelain tube enclosed in a charcoal fire. Number, mean; extreme difference, 0.34, for 0 = 16. The sulphate was prepared from a pyrolusite containing only silica, iron, and barium. It was reduced to protoxide, dissolved in sulphuric acid, oxidized with nitric acid, precipitated with oxalic

acid, converted into red oxide, dissolved in chlorhydric acid and alcohol, precipitated with ammonium carbonate, dissolved in sulphuric acid, repeatedly heated to redness and recrystallized, and was dried at 300°. Accurate experiments on the reduction of the red oxide proved impracticable on account of the hydroscopicity of the compound. Two experiments on the oxidation of the protoxide, undertaken as a check on the other method, gave 27.486 and 27.527 for O = 8; S = 16. (Erdmann's Journ. für Prak. Chem., 72, 1857, 361; Sitz.-Bericht der k. k. Akad., 1857.)

J. Dumas: 54.96 (O = 16).

Determined by the decomposition of the chloride with argentic nitrate. The number is the mean of five experiments; extreme difference, 0.1 for O=16. Cl=35.5; Ag=108. Dumas had previously made experiments on the reduction of the hyperoxide to protoxide by hydrogen. These gave the atomic weight at from 25.99 to 26.09 for O=8. Dumas believes that a part of the oxide was reduced to metal. The peroxide was prepared from nitrate of the protoxide. (Annal. de Chim. et de Phys., (3,) 55, 1859, 150.)

- RAWACK: 54.02 (O = 16).

Determined, in Schneider's laboratory, by reducing a known weight of red oxide to protoxide in a current of dry hydrogen, and weighing the water formed. The number is derived from the mean of six selected experiments. The extreme difference is 0.22 for O=16. (Poggend. Ann., 107, 1859, 607.)

R. Schneider: 54.038 (O = 16).

The mean result of four analyses of the oxalate by the ordinary method of organic analysis. Extreme difference, 0.04 for O=16. C=12. The oxalate was prepared from chemically pure sulphate by precipitation with sodium carbonate, digestion with oxalic acid, and drying over sulphuric acid. (*Poggend. Ann.*, 107, 1859, 613.)

MERCURY.

The specific heat of mercury in the solid state, as observed by Regnault, and the vapor density, as determined by Dumas, correspond to an atomic weight of slightly above 200. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

FOURCROY AND THENARD, DAVY, WOLLASTON: 200.8 (O = 16); 1255 (O = 100).

Fourcroy and Thenard found 8 O = 100 Hg. Davy found 80 O = 880 Hg, giving Hg = 1266. The latter also found 134 Cl = 380 Hg, which for Cl = 441, gives Hg = 1254. (*Phil. Trans.*, 104, 1814, 21.)

N. G. Sefstroem: 202.58 (O = 16); 1265.822 (O = 100).

Determined by three analyses of the oxide according to which 100 Hg = 7.89, 7.9, and 7.97 O. (Berzelius' Lehrbuch, 3, 1215.)

E. Turner: 200.72 (O = 16).

Turner made a number of determinations of this atomic weight but regarded the value he adopted, 202, only as an approximation. From the oxide, prepared from nitrate, he got 200.77 and 199.97. The compound was decomposed by heat, and the products carried over silver and gold in a narrow tube. Four experiments were made on mercuric chloride which was decomposed by pure calcic oxide, and the Cl precipitated with argentic nitrate. [These analyses recalculated for the Stas' atomic weights of Ag and Cl give 202.079, 201.701, 201.815.] Turner also made two experiments on the reduction of the chloride with stannous chloride, the Hg being collected, dried and weighed. [These experiments recalculated give 199.423 and 199.289.] The mercuric chloride was purified by recrystallization. Weighings reduced to vacuum. (Phil. Trans., 123, 1833, 535.)

ERDMANN AND MARCHAND: 200.14 (0 = 16); 1250.6 (0 = 100).

Determined from the mean of four experiments on the reduction of the oxide in a current of carbon di-oxide. Copper, carbon (from sugar) oxide, and carbon, were introduced in successive layers in a combustion tube. Dry carbon di-oxide was passed through and the mercuric oxide heated. The metal was collected in a receiver to which a tube filled with gold foil was appended. The metal was perfectly clean. Moisture was removed by a stream of dry air after distillation. The oxide was purified by heating it to incipient decomposition the metallic fumes being removed

by a current of dry air. It was tested before being analysed. The extreme difference in the results was 0.8 for O = 100. All weighings in vacuo. (Erdmann's Journ. für Prak. Chem., 51, 1844, 392.)

E. MILLON: 199.94 (O = 16); 1249.63 (O = 100).

Millon made two experiments by heating mercuric chloride with calcic oxide in a current of hydrogen and condensing the metal. The experiments gave 78.87 and 78.82 per cent. mercury. If Cl = 442.64, the value follows. The chloride was dissolved in ether and sublimed. It was perfectly soluble in ether and alcohol, and was well crystallized. (Paris Comptes Rend., 20, 1845, 1291.)

L. Svanberg: 200 (O = 16); 1250 (O = 100).

Svanberg made three experiments by the same method employed by Millon. The mean result was 1248.47; extreme difference, 0.94; but Svanberg shows that there was probably loss, and that the larger the quantity of chloride employed the higher the result. He regards Erdmann and Marchand's result as most probable, but in need of confirmation. Cl = 448.28. (Erdmann's Journ. für Prak. Chem., 45, 1843, 468; Kongl. Vet. Akad. Handl., 1845, 135.)

MOLYBDENUM.

Regnault determined the specific heat of molybdenum. It answers to an atomic weight of about 96. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 95.36 (O = 16); 596.1 (O = 100).

One hundred parts of anhydrous plumbic nitrate, dissolved and precipitated with neutral ammonium molybdate, gave 110.68 parts plumbic molybdate. If Pb = 1294.645, N=87.53, the value follows. Berzelius expresses himself dissatisfied with the accuracy of the determination. (Poggend. Ann., 8, 1826, 23; and Lehrbuch, 3, 1208.)

Syanberg and Struve: 92.13 (O = 16); 575.829 (O = 100).

After trying various methods without getting accordant results, these chemists made ten experiments on the sul-

phide by roasting it first in a current of moist, and then of dry air. Three experiments were excluded as imperfect. The remainder gave a mean of 89.7523 molybdic acid from 100 sulphide; extreme difference, 0.22. The value follows for S = 200. Objections have been made (*Liebig's Ann.*, 68, 211) that the difference in weight between the acid and the sulphate is too small for the purpose of the determination, and that the different analyses give very different atomic weights. The sulphide was prepared by melting together molybdic acid, sulphur, and caustic potash, and leaching the product with water and chlorhydric acid. The sulphide was dried in a current of hydrogen. The molybdic acid was dissolved in ammonia to prove the absence of sulphide. (*Erdmann's Journ. für Prak. Chem.*, 44, 1848, 315.)

N. J. Berlin:
$$91.96$$
 (O = 16); 574.75 (O = 100).

Determined by four analyses of the double mono-sesquimolybdate of ammonium by heating gently with nitric acid in a platinum crucible until only molybdic acid was left. Extreme difference, 8.32 for O = 100; N = 175; H = 12.5. The preparation of the salt is not given. (*Erdmann's Journ.* für Prak. Chem., 49, 1850, 446.)

J. Dumas: 96 (O = 16).

Dumas made five experiments on the reduction of molybdic acid (prepared from the natural sulphide) by means of hydrogen. The reduction was begun at a low temperature in a glass tube, and completed in an unglazed porcelain tube in a reverberatory furnace, where it was kept till several hours heating produced no further alteration in weight. The molybdenum did not assume a metallic appearance. The number is the mean; extreme difference, 0.8 for O = 16. (Annal. de Chim. et de Phys., (3,) 55, 1859, 142.)

M. Delafontaine:
$$92$$
 (O = 16); 575 (O = 100).

This chemist made many experiments in various ways without being able to reach constant results, and only remarks that his experiments indicate Svanberg and Struve's value as the best. (Erdmann's Journ. für Prak. Chem., 95, 1865, 137; Bibl. Univ., Arch. des Sciences, 23, 1865.)

H. Debray: 95.94 (O = 16).

Debray made three experiments on the reduction of molybdic acid. The acid was first converted into the red

oxide in platinum, and at a low temperature, and the small portion of the acid volatilized during this operation was caught and determined. The reduction was completed in a porcelain tube at a white heat. Debray gives his results at 48.03; 48.04; and 47.84. [The analytical data, recalculated, give 95.30; 95.55; 95.73; perhaps on account of misprints. Reduction to vacuum would still further reduce the numbers.] The acid was purified by sublimation in platinum, conversion into ammonium salt, and regeneration by heat. In two experiments ammoniacal solution of molybdic acid was evaporated in the dark with excess of argentic nitrate, the argentic molybdate dissolved out and the excess of silver determined. Debray found 5.510 acid = 7.657silver, and 7.236 acid = 10.847 silver. Hence he calculates M=48 and 47.98. [A little calculation shows that the first data are misprinted. They should read 5.11 acid = The corrected data give for Ag = 107.93; M 7.657 silver. = 96.06 and 95.99. The mean of the recalculated analyses is 95.73.] (Paris Comptes Rend., 66, 1868, 732.)

L. MEYER: 96.10 (O = 16).

Calculated from three analyses of the dichloride, two analyses of the tetrachloride, and two analyses of the pentachloride, made by Leichte and Kempe in Meyer's laboratory. The dichloride was analyzed by heating in a current of hydrogen sulphide, and subsequently in a current of hydrogen. Molybdenum disulphide is the residue. HCl formed was caught in ammonium hydrate and precipitated by argentic nitrate, after the hydrogen sulphide had been driven off by boiling in a flask provided with a con-densing drip-tube. The tetra and pentachloride were decomposed with nitric acid, excess of ammonium hydrate was added, and molybdenum trisulphide precipitated with ammonium sulphide. A weighed portion of the dry precipitate was converted into disulphide by heating in a current of hydrogen. The chlorine of the higher chlorides was determined in the filtrate after precipitation of the trisulphide. By comparing the amount of chloride analyzed with the amount of argentic chloride obtained, Meyer finds in mean M = 95.92; extreme difference, 1.87 for O = 15.96. By comparing the amount of disulphide with that of argentic chloride, M = 95.75; extreme difference, 1.35. By comparing the amount of chloride analyzed with the amount of disulphide obtained for one analysis of tetrachloride and two analyses of pentachloride, he gets M = 95.94; extreme difference, 2.15. The general mean is M = 95.86; extreme

difference, 2.15. Ag = 107.66; S = 31.98; Cl = 35.37; O = 15.96. The specific gravities of the chlorides not having been determined, the weighings are not reduced to vacuum. The pentachloride was prepared from M by heating it in a current of Cl entirely free from air. The metal had been freed from oxide by heating in an atmosphere of HCl. By moderate heating of the pentachloride in dry H, and by distilling pentachloride over the product in dry carbon di-oxide, the trichloride is obtained. The trichloride heated in carbon di-oxide is decomposed into tetrachloride and di-chloride, which latter must be purified with warm dilute nitric acid. (Liebig's Ann., 169, 1874, 360, 344.)

NICKEL.

Regnault has determined the specific heat of nickel. It corresponds to an atomic weight of about 59. (Gmelin-Kraut, l. c.)

E. Rothoff: 59.09 (O = 16); 369.333 (O = 100).

Rothoff converted 188 parts of oxide into chloride, a neutral solution of which gave 718.2 parts argentic chloride. If Cl = 221.64, Ag = 1349.66, the value follows. (Berzelius' Lehrbuch, 3, 1221.)

P. Berthier.

Lassaigne having announced the atomic weight of nickel at 500, (Schweigger's Jahrbuch, 9, 108,) Berthier re-examined the subject and found Rothoff's number confirmed. (Berzelius' Jahresbericht, 5, 1825, 148; Annal. de Chim. et de Phys., 25, 1824, 148.)

ERDMANN AND MARCHAND: 58.2 (O = 16); 365.9 (O = 100).

Determined "with all precaution" by the reduction of the oxide with hydrogen. The results varied from 29.1 to 29.8, but Erdmann has reason to believe the smaller number the more accurate. (Erdmann's Journ. für Prak. Chem., 55, 1852, 202.) NICKEL. 87

H. SAINTE-CLAIRE DEVILLE:

100 parts fused nickel, containing three-tenths per cent. silicon and one-tenth per cent. copper, gave 262 parts anhydrous, yellow nickel sulphate, "corresponding to the atomic weight as given by Berzelius." (Annal. de Chim. et de Phys., (3,) 46, 1856, 182.)

R. Schneider: 58.05 (O = 16); 362.8 (O = 100).

Determined from four analyses of the oxalate. The carbon determinations were made by the ordinary method of organic analysis, because some hydrocarbon forms when the salt is decomposed by heat alone. The metal was determined by heating a known weight of the salt first in air and then in a current of oxygen, and subsequent reduction by hydrogen. In the preparation of the salt the usual precipitate with ammonium sulphide was washed with dilute chlorhydric acid, and the cobalt separated with barium carbonate and chlorine. From the nickel salt obtained the oxalate was precipitated with oxalic acid. The number is the mean of four analyses; extreme difference, 0.082 for O = 8. (Poggend. Ann. 101, 1857, 396.)

C. Marignac: 59 (O = 16).

Marignac made two analyses of the sulphate by decomposing the salt by heat. The decomposition is perfect. avoid errors arising from possible reduction of a portion of the oxide, it was moistened with nitric acid, and recalcined at a moderate temperature. The results obtained were Ni = 29.2 and 29.5. The sulphate was purified by recrystallization. He also made experiments on the chloride by titration with argentic nitrate, according to Pelouze's modification of Gay-Lussac's method. Three such analyses gave from 29.4 to 29.5. In one experiment he also evaporated the nickel nitrate, after filtering off the argentic chloride, and converted it into oxide by heat. This experiment gave Ni = 29.64.The chloride, whether it be distilled or calcined with ammonium chloride, is apt to leave an insoluble residue the weight of which must be deducted. S = 16; Ag = 108; Cl = 35.5. (Bibl. Univ. Arch. des Sciences, (2,) 1, 1858, 375.)

J. Dumas: 59.028 (O = 16).

The number is the mean result of five experiments on the titration of the chloride with argentic nitrate; extreme difference 0.08. Ag = 108; Cl = 85.5. In three cases the nickel chloride was prepared by dissolving fused nickel in aquia regia, repeated evaporation to dryness with HCl, and heating for from twelve to twenty-four hours in a current of HCl gas. In two cases it was produced by passing a current of chlorine over spongy nickel. The chloride analyzed was crystalline and volatile without residue. (Annal. de Chim. et de Phys., (8,) 55, 1859, 149.)

R. Schneider: 58.058 (O = 16).

In consequence of Marignac's criticism (that as nickel oxalate is insoluble it cannot be purified by recrystallization) Schneider repeated his former determination, making special tests for oxalic acid, sodium, and chlorine, with the above result. (Poggend. Ann., 107, 1859, 616.)

W. J. Russell: 58.738 (O = 16).

Determined from the mean of thirteen experiments on the reduction of the oxide in hydrogen. Extreme difference, 0.12 for O=16. The oxide was prepared from three specimens of commercial nickel, which were first converted into pure oxalate and then into nitrate. The oxide was obtained by decomposing the nitrate by a very strong heat. (Journ. Chem. Soc., (2,) 1, 1863, 61.)

Schneider remarks that a portion of the oxide analyzed may have been reduced during the process of decomposing the nitrate. (*Poggend. Ann.*, 130, 1867, 310.) Marignac points out the same danger. (*Bibl. Univ.*, Arch. des Sciences, (2,) 1, 374.)

E. von Sommaruga: 58.026 (O =16).

Determined from the amount of barium sulphate obtained by precipitating the double sulphate of nickel and potassium with barium chloride. The number is the mean of six experiments; extreme difference, 0.168 for O = 8, S = 16; Ba [no doubt] = 68.5; K = 89.2. The salt was prepared by solution of commercial nickel in sulphuric and nitric acid, adding potassic sulphate to the solution, and repeatedly recrystallizing the double sulphate. (Erdmann's Journ. für Prak. Chem., 100, 1867, 115; Sitz.-Ber. der k. k. Akad., 1866.)

C. Winkler: 59.05 (O = 16).

Determined by the amount of gold precipitated from a solution of neutral crystallized potassium chloro-aurate by

a known weight of nickel. The number is the mean of four experiments; extreme difference, 0.186 for O=16, Au=196. The nickel was prepared as follows: commercial nickel carbonate was dissolved in chlorhydric acid, cobalt was repeatedly precipitated with sodium hypochlorite, copper, etc., were removed with hydrogen sulphide, the nickel was precipitated with sodium carbonate, the precipitate dissolved in chlorhydric acid, the chloride sublimed and reduced in a current of hydrogen. (Fresenius' Zeitsch., 6, 1867, 22.)

W. J. Russell: 58.76 (O = 16).

Determined by the amount of hydrogen set free by solution of nickel in chlorhydric acid. The nickel was that obtained in Russell's former determination of the atomic weight of nickel. (Chem. News, 20, 1869, 20.)

R. H. Lee: 58.01 (O = 16).

Determined by analyses of nickel cyanide salts. They were decomposed in a platinum crucible by heat from above. The carbon separated out was burned off first in air and then in oxygen. The metallic oxide was reduced in a current of hydrogen. The mean of six experiments on the strychnine salt gave Ni = 58.04. The mean of six experiments on the brucine salt gave Ni = 57.98. The salts were purified by recrystallization. (Berlin. Bericht der Chem. Gas., 4, 1871, 790.)

NIOBIUM.

The vapor density of the chloride and of the oxychloride, as determined by Deville and Troost, places the atomic weight at about 94. (Paris Comptes Rend., 56, 1863, 891.)

H. Rose: 122 (O = 16).

Rose deduced the atomic weight of niobium from analyses of what he supposed to be the tetrachloride, determining the niobium as niobic acid, and the chlorine as argentic chloride. The results, which varied greatly, indicated the value 97.64. [Marignac having proved that the salt is a pentachloride, this number becomes 122.] Marignac showed

that Rose dealt with a compound containing a large amount of the corresponding tantalium chloride. (*Poggend. Ann.*, 104, 1858, 439.)

Rose; Rammelsberg: 94 (O = 16).

Rose analysed the oxychloride, but did not recognize it as an oxychloride. Rammelsberg calculated the atomic weight from Rose's figures and found that the highest chlorine contents corresponds to an atomic weight of 94. Rose's salt must have been nearly pure as there is no corresponding tantalium compound. (Poggend. Ann., 136, 1869, 853.)

R. Hermann: 104.8 (O = 16).

Hermann deduces this value from analyses of a number of chlorides and sodium salts. The formulas which he gives these compounds are complicated, unlikely, and unsupported by evidence. Marignac has shewn that Hermann's salts contained tantalium. (Erdmann's Journ. für Prak. Chem., 68, 1856, 73.)

O. W. Blomstrand: 95 (O = 16).

Blomstrand made three determinations of the chlorine contents of the pentachloride, getting 64.712 per cent., extreme difference, 0.32. He also made eleven determinations of the niobium in the same compound, weighing it as niobic acid. 100 chloride gave in mean 49.794 acid. The atomic weight calculated from the chlorine contents is 96.67; from the niobic acid, 96.16. Blomstrand also made experiments on sodium niobate which led him to the conclusion that the most probable number is 95. (Gmelin-Kraut, 2, part 2, 73; Acta Univ. Lund., 1864.)

C. Marignao: 94 (O = 16).

Determined from a number of analyses of potassium fluoniobate containing two atoms of potassium. The compound was decomposed by sulphuric acid with which it was evaporated to dryness. The residue was leached with water, the filtrate evaporated and the potassic sulphate melted and weighed. The sulphuric acid remaining with the niobic acid was driven off by heat and the acid weighed. The salt being readily soluble and crystallizing well, can easily be purified from all substances except titanium which Marignac knows no way of separating or determining.

The larger the amount of titanium present, the lower will the atomic weight be; Marignac therefore takes the highest value. (Liebig's Ann., S. 4, 334, 288, 338; Bibl. Univ., Arch. des Sciences, 23, 1865, 25, 1866.)

NITROGEN.

Regnault has determined the specific gravity of nitrogen. It indicates an atomic weight slightly above 14. (Gmelin-Kraut, l. c.)

BIOT and ARAGO; WOLLASTON: 14.03 (O = 16); 87.7 (O = 100).

Biot and Arago found the specific gravities of N and H 0.96918 and 0.07821. If H=13.27 the value follows. [This very accurate value is of course the result of two compensating errors.] (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius; 14.163 (O = 16); 88.518 (O = 100).

Calculated from the specific gravity as determined by Berzelius and Dulong, compared with that of oxygen. By decomposing the nitrate of lead by heat, Berzelius also found N=88.61 for Pb=1294.498. (Poggend. Ann., 8, 1826, 14.)

E. Turner: 14.15 (O = 16).

Determined by experiments on the nitrates of lead, silver, and barium, which were precipitated with sulphuric and hydrochloric acids, and gave respectively N = 14.201; 14.09; 14.17; if Pb = 103.6; Ba = 68.7; Cl = 35.42; S = 16.085; the weighings being reduced to vacuum. The salts were purified by recrystallization. Turner recommends more direct methods. (Phil. Trans., 123, 1838, 537.)

T. Thomson: 14 (O = 16).

From the hypothesis that air is a compound containing four parts of N and one part oxygen, and from the average of various selected determinations of the specific gravity of oxygen, Thomson concludes the specific gravity of oxygen is 1.1111, and that of N 0.9722. These numbers stand

to one another as 16 to 14. (Erdmann's Journ. für Prak. Chem., 8, 1836, 375; Records of General Science, by R. D. Thomson, 1836, 179.)

F. Penny: 14.018 (O = 16).

From the mean of three series of experiments (vide Penny's determination of potassium) it follows that 100 potassic chloride = 135.636 potassic nitrate. Penny found the molecular weight of KCl = 74.527; hence the difference between a chloride and a nitrate is 26.560. Similar experiments were made on the sodium salts. In four experiments 100 sodium chlorate were found = 54.930 chloride; extreme difference, 0.02. In three experiments, 100 sodium chlorate were found = 79.882 sodium nitrate; extreme difference, 0.015. In six experiments 100 sodium nitrate were found = 68.771 chloride; extreme difference, 0.018. In seven experiments 100 chloride were found = 145.416 sodium nitrate; extreme difference, 0.016. These data give sodium chloride = 58.5, and the nitrate = 85.068, or the difference between a chloride and a nitrate = 26.568. Penny found Cl = 35.454. If NO₃-Cl = 26.564, N = 14.018. Weighings for vacuum. (Phil. Trans., 129, 1839, 25.)

L. Svanberg: 13.91 (O = 16).

Determined by four experiments on the decomposition of plumbic nitrate by heat which gave a mean of 67.4016 per cent. oxide; extreme difference, 0.0087. [If Pb = 206.926 (Stas) the value follows.] (Berzelius' Jahresbericht, 22, 1842, 38.)

C. Marignac: 14.02 (O = 16); 87.625 (O = 100).

Marignac made five experiments by dissolving a known weight of silver in nitric acid and melting and weighing the nitrate formed. The silver carried out of the retort by the vapors was precipitated and determined. The mean result was that 100 silver = 157.430 nitrate; extreme difference, 0.046; or, if Ag = 1349.01, N = 87.585. Six experiments were made by the decomposition of a known weight of argentic nitrate with a known weight of potassic chloride by Pelouze's method. Mean, 100 KCl = 227.986 argentic nitrate; extreme difference, 0.18. This gives N = 87.685 if K = 488.94 and Cl = 443.2. Seven experiments by the same method showed that 100 silver dissolved in nitric acid = 49.522 ammonium chloride; extreme difference.

ence, 0.063; Hence N = 87.655. The weighings are reduced to vacuum. (Berzelius' Jahresbericht, 24, 1842, 44; Bibl. Univ. de Genève, 46, 1842, 363.)

T. Anderson: 13.95 (O = 16); 87.204 (O = 100).

Determined by four experiments on the decomposition of plumbic nitrate by heat at a sufficiently low temperature to permit of complete decomposition. The number is the mean; extreme difference, 0.198 for O = 100. Pb = 1294.5. (Annal. de Chim. et de Phys., (3,) 9, 1848, 254.)

J. Pelouze: 14.014 (O = 16); 87.59 (O = 100).

A known weight of argentic nitrate was brought in contact with a known and slightly excessive weight of ammonium chloride and the excess titrated with silver solution. One experiment gave N=175.58; a second gave N=174.78. Ag = 1849.01; Cl = 448.2. The ammonium chloride was purified by sublimation and recrystallization. (Paris Comptes Rend., 20, 1845, 1047.)

P. EINBRODT: 14 (O = 16); 87.5 (O = 100).

Experiments on the decomposition of plumbic nitrate by heat gave N = 87.5 plus a vanishing quantity if Pb = 1294.2239. (*Leibig's Ann.*, 70, 1849, 286.)

J. Dumas: 14 (O = 16).

Determined by experiments on the combustion of ammonia and cyanogen. Particulars not given. C = 6; H = 1. (Annal. de Chim. et de Phys., (3,) 55, 1859, 184.)

J. S. Stas: 14.044 (O = 16).

Stas made seven determinations of the relation between silver and its nitrate by dissolving pure silver in nitric acid, evaporating to dryness and keeping the salt melted until there was no further loss of weight. In two of these experiments the salt was melted in vacuo. The mean result was that 100 Ag = 157.472 nitrate; whence N = 14.040. Later Stas made two more experiments by the same method with all possible precautions to secure accuracy. These gave 100 Ag = 157.484 nitrate and N = 14.042. By the conversion of the chlorides of potassium, sodium, lithium and silver into nitrates, Stas found the difference between a chloride and a nitrate 26.5882. This gives N = 14.045. The weigh-

ings are reduced to vacuum. Cl = 85.457; Ag = 107.98. (Stas, Unters. über Chem. Prop. Leipzig, 1867.)

OSMIUM.

Regnault has determined the specific heat of osmium. It corresponds to an atomic weight of about 199. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 199.04 (O = 16).

Berzelius analyzed potassium chloro-osmate by reduction in a current of hydrogen and solution of the potassium chloride from the residue. 1.3165 grammes of the double salt lost 0.8805 in reduction and the residue was composed of 0.401 potassium chloride and 0.535 osmium. The atomic weight may be calculated either from the chlorine lost or from the relation of the chloride to the metal in the residue. Berzelius preferred the latter as more probably accurate. [If K = 39.187; Cl = 35.457 (Stas;) this relation gives 199.04.] According to W. M. Watts, (Chem. News, 19, 802) the loss of chlorine gives for Stas's values Os = 199.42. Hyperosmic acid was separated from iridium compounds by distilling at a gentle heat. The first portion is perfectly pure. The metal was precipitated from chlorhydric acid solution of hyperosmic acid by mercury and subsequently purified by heating in a current of hydrogen. Potassium chloro-osmate was formed by heating comminuted metal and KCl in a current of chlorine. (Poggend. Ann., 13, 1828, 580; Kongl. Vet. Acad. Handl., 1828.)

E. Fremy: 199.65 (O = 16); 1247.8 (O = 100).

Pure osmium was burned in a current of oxygen and the fumes led over potassic hydrate, by which they are absorbed. An additional potash tube did not increase in weight. Corks were avoided. Number of experiments not given. (Erdmann's Journ. für Prak. Chem., 33, 1844, 409; Journ. de Pharm. et Chim., 1844, 241.)

DEVILLE and DEBRAY: 198 (O = 16).

These chemists determined the vapor density of hyperosmic acid by Dumas' method, finding it 8.89 at 246°,

and 8.87 at 286°. They hence consider it probable that the atomic weight of osmium is the same as that of platinum. The acid was very pure and was prepared by the combustion of metallic osmium in oxygen. (Paris, Comptes Rend., 44, 1857, 1101.)

OXYGEN.

The atomic weight of oxygen is assumed at 16 for the reasons stated under hydrogen, q. v. If hydrogen is taken as unity, 0 = 15.96.

PALLADIUM.

Regnault determined the specific heat of palladium. It corresponds to an atomic weight of about 106. (Gmelin-Kraut, l. c.)

J. J. Berzelius; 106.51 (O = 16).

In his earliest determinations of this constant, Berzelius saturated the metal with sulphur, getting about 711 for S = 201.165; and decomposed the chloride with mercury, getting 704. [711 appears to be a misprint for 714.618 the number given with corresponding data at Poggend., 8, 180.] In this investigation a known weight of potassium chloropalladate was reduced in a current of hydrogen, the weight of the residue determined, the potassium chloride leached from the residue and the metallic palladium weighed. double salt was strongly heated, but not to fusion, in a current of dry air before weighing. It being impossible to desiccate this and the similar platinum-metal salts completely without decomposition, the atomic weight was calculated from the relation between the metal and the KCl. Berzelius found 0.575 Pd = 0.809 KCl, and 0.851 Pd = 1.192 KCl. [If KCl] = 74.594 (Stas) the former gives Pd = 106.086, the latter 106.509.] Berzelius had reason to consider the latter analysis the more accurate. (Poggend. Ann., 13, 1828, 454; Kongl. Vet. Acad. Handl., 1828.)

PHOSPHORUS.

The specific heat of this element, as well as the density of phosphorus and its numerous volatile compounds in the gaseous state, corresponds to an atomic weight slightly above 31. (Gmelin-Kraut, l. c.)

V. Rose; F. H. Wollaston: 35.1 (O = 16).

Wollaston adopted the analysis of Rose, who found that phosphoric anhydride contained 53.28 per cent. oxygen and 46.72 per cent. phosphorus. [This relation gives the above value.] (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius: 31.325 (O = 16).

Berzelius made experiments on the reduction of auric chloride and of argentic sulphate by phosphorus. His results were 0.8115 P = 13.98 Ag; 0.829 P = 8.714 Au; 0.754 P = 7.93 Au. [The first of these analyses is misprinted in the original memoir (Gilbert's Ann., 53, 488).] In the Lehrbuch it is miscalculated as Ruecker has shown. zelius preferred deducing the atomic weight of P from that of silver, because the atomic weight of the latter was more accurately known than that of gold. [If Ag = 107.98, the data give P = 31.825, for Au = 196.67 the latter analyses give P = 31.176 and 81.165.] In all the experiments the precipitated metal was boiled with the solution when the reduction was nearly complete. A trace of gold was observed to precipitate after the experiments were over. The silver was heated to redness before weighing. [J. P. Cooke, Jr., has shown (atomic weight of antimony) that silver is volatile at a red heat. Berzelius must therefore have got too large a result.] The phosphorus was distilled, melted in a glass tube and cooled very slowly, to permit traces of oxides to rise to the surface, and the lower portion of the tube with the phosphorus broken off and instantly (Gilbert's Ann., 53, 1816, 433, and Lehrbuch 3, weighed. **1188.**)

J. Pelouze: 32.024 (O = 16); 200.15 (O = 100).

A known weight of argentic nitrate was brought in contact with a known and slightly excessive weight of phosphorous chloride and the excess titrated. The number of experiments is not given. Ag = 1849.01; Cl = 448.2.

The terchloride was prepared by chloridizing finely divided P with dry chlorine, adding finely divided P, decanting, agitation with tin amalgam and rectification over the same. The fluid was colorless and did not give any precipitate with water. (Paris, Comptes Rend., 20, 1845, 1047.)

V. A. JACQUELIN: 29.83 (O = 16); 186.438 (O = 100).

Determined by experiments on the chlorides of phosphorus with argentic nitrate and plumbic oxide. The results are utterly discordant. (*Paris, Comptes Rend.*, 33, 1851, 698.)

A. Schroetter: 31.0274 (O = 16).

Determined by burning perfectly pure amorphous phosphorus in dry oxygen and weighing the phosphoric anhydride. The number is the mean of 10 experiments; extreme difference, 0.1242. Previous to burning, the phosphorus was heated for a long time in carbon di-oxide or hydrogen. It was burned not in powder but in lumps. (Erdmann's Journ. für Prak. Chem., 63, 1851, 435; Sitz-Bericht der k. k. Akad., 1851.)

B. C. Brodie: 81.81 (O = 16).

Three experiments made by oxidation of phosphorus with aqua regia and determination as magnesium pyrophosphate gave this mean. Brodie seems to regard these determinations only as evidence that the atomic weight needs redetermination. (Journ. Chem. Soc., 5, 1852, 295.)

J. Dumas: 81.08 (O = 16).

Determined by five experiments on the titration of the terchloride with argentic nitrate. The chloride was prepared by the action of dry chlorine on amorphous phosphorus and distillation after the chlorine had been displaced by carbon di-oxide. The portion distilling between 76° and 78° only was used. The number is the mean of the results; extreme difference, 0.08. Ag = 108; Cl = 35.5. (Annal. de Chin. et de Phys., (3,) 55, 1859, 172.)

PLATINUM.

Regnault and Kopp have determined the specific heat of platinum. It answers to an atomic weight of about 197. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 197.19 (O = 16).

Determined by the same method as osmium, q. v., from a single experiment on potassium suroplatinate. 2.135 potassium chloride accompanied 2.822 platinum. [If KCl = 74.594 (Stas,) this gives the above value.] The salt was prepared by precipitating an alcoholic solution of platinum chloride with potassium chloride, washing with alcohol and heating to redness in a current of chlorine. Berzelius remarks that the metal used in his former determinations was impure. (Poggend. Ann., 13, 1828, 468, and Lehrbuch, 3, 1213.)

T. Andrews: 197.88 (O = 16).

Determined by three experiments on potassium chloroplatinate. The salt was dried at 105° in vacuo, decomposed by zinc, the excess of zinc removed by acetic acid, the solution filtered off, and the chlorine titrated. The number is the mean; extreme difference, 0.22. The values assumed for Ag and Cl are not given. They were most likely Marignac's. (Brit. Assoc. Rep., 1852, part 2, 33.)

J. S. Stas made preparations for determining the atomic weight of platinum, but not being able to produce potassium chloroplatinate entirely free from water, and being unacquainted with Bunsen's method of accomplishing this end, resigned the attempt. He made, indeed, three analyses by the same method employed by Berzelius, but unfortunately does not communicate the results. (Stas, Untersuch. über Chem. Prop., Leipzig, 1867, 265.)

POTASSIUM.

Regnault determined the specific heat of potassium. It corresponds to an atomic weight of about 89. (Gmelin-Kraut, l. c.)

M. H. KLAPROTH; F. H. WOLLASTON: 39.517 (O = 16).

Klaproth found that 441 Cl = 591 potassium oxide. Hence Wollaston deduced the value 491 (O = 100) for K. [If Cl = 35.457, this relation gives K = 39.517.] (Phil. Trans., 104, 1814, 20.)

J. J. Berzelius: 39.193 (O = 16); 244.958 (O = 100).

Berzelius found that 100 KCl = 192.4 Ag Cl. If Ag = 1351.607; Cl = 442.65; the above value follows. (*Poggend. Ann.*, 8, 1826, 190.)

F. Penny: 39.073 (O = 16).

Penny made six experiments on the conversion of the chlorate into the chloride. Potassic chlorate was dried at about 105° , dissolved in a flask with HCl, evaporated, dried and weighed. The cake contained some free HCl. It was broken up, pulverized, and a known quantity heated to dull redness but not to fusion, and the HCl expelled allowed for. The mean result was that $100~\rm KCl~O_3=60.823~\rm KCl$; extreme difference, 0.015. This relation gives KCl = 74.527 and if Cl = $35.454~\rm (Penny,)$ the value for K follows. Numerous experiments were also made on the introconversion of the nitrate, the chloride and the chlorate, which established the difference between a chloride and a nitrate, besides confirming the value of K. The salts were purified by recrystallization and were carefully tested for impurities. The weighings are all for vacuum. (*Phil. Trans.*, 129, 1839, 18.)

C. Marignao: 39.2 (O = 16); 245 (O = 100).

By six experiments on the decomposition of the chlorate by heat, 100 chlorate were found to lose 39.161 oxygen; extreme difference 0.012; hence KCl = 932.14. By comparing this value with the molecular weight and the composition of argentic chloride, Cl was calculated at 442.18, leaving for K the number 490. Confirmatory experiments were made on potassic perchloride. The chlorate was purified by recrystallization. The weighings are for vacuum. (Liebig's Ann., 44, 1842, 28.)

C. Marignac: 39.115 (O = 16); 244.47 (O = 100).

Having determined the atomic weight of chlorine from syntheses of argentic chloride, and found it 443.2, the molecular weight of KCl in the last determination, gives K=244.47, for vacuum. Berzelius, by rejecting some analyses and the correction for vacuum, deduces the value 244.429. (Berzelius' Jahresbericht, 25, 1845, 31; Bibl. Univ. de Genève, 46, 1842, 350.)

J. Pelouze: 39.144 (O = 16); 244.65 (O = 100).

A known weight of KCl was brought into contact with a known amount of silver dissolved in nitric acid, the chloride being slightly in excess. This excess was titrated with a decimal solution of silver. The number is the mean of three experiments. Ag = 1349.01; Cl = 443.2. The chloride was prepared by heating the chlorate and recrystallizing the residue. (*Paris Comptes Rend.*, 20, 1845, 1047.)

According to Pelouze, Levol found the molecular weight of KCl 466.245, which combined with Marignac's value of Cl would give K = 244.645 or 39.143. (*Ibid.*)

E. J. MAUMENÉ: 38.96 (O = 16); 243.502 (O = 100.)

The mean of three experiments on the decomposition of KCl with an excess of argentic nitrate showed that 100 KCl = 192.75 AgCl. If Ag = 1850.82 and Cl = 443.67, according to Maumené's determinations, the number follows. The KCl was prepared from the chlorate by heat. To confirm his values for K and Cl, he made seven experiments on the decomposition of the chlorate by heat, and found that 100 chlorate gave 60.791 chloride. An unaccounted for increase in the weight of the flask occurred in these experiments. (Annal. de Chim. et de Phys., (8,) 18, 1846, 41.)

J. S. Stas: 39.187 (O = 16).

According to the mean of seven determinations, 100 parts of KCl dissolved in nitric acid, and evaporated to dryness give 135.6428 parts of nitrate; extreme difference, 0.017. If Cl = 35.457; N = 14.044; the value follows. This value is confirmed by previous experiments which gave 39.130. Potassic chloride, by whatever means it is prepared, still retains silica. Stas, therefore, determined

the amount of silica in the KCl and allowed for it. Weighings for vacuum. (Stas, Untersuch. über Chem. Prop., Leipzig, 1867.)

Stas mentions that Dumas, who was the first to place K at 39, afterwards became convinced that this number was

too low. (*Ibid*, page 318.)

RADCLIFFE.

RHODIUM.

Regnault has determined the specific heat of rhodium. It corresponds to an atomic weight of about 104. (Gmelin-Kraut, l. c.)

J. J. Berzelius: 104.3 (O = 16).

Berzelius made two analyses of dipotassic chlororhodiate. This salt can be completely desiccated in a current of chlorine at a red heat without decomposition. 3.146 grammes gave on reduction in a current of hydrogen 0.930 Cl, and the residue contained 1.804 KCl and 0.912 metallic rhodium. [If KCl = 74.594, Cl = 35.457, (Stas.) the atomic weight of the salt calculated from the Cl contents is 359.881, and that of Rh 104.272. The relation between the Rh and the Cl gives Rh = 104.312.The relation between the KCl and the Rh gives Rh = 104.340. The mean is 104.308.] Berzelius made a second analysis of the crystallized salt in which he determined the water of crystallization. [Under the same suppositions and in the same order, the resulting values for Rh are 106.279; 104.762; 103.708.] In the Lehrbuch only the former analysis is used to deduce the atomic weight. Rhodium was separated from other metals by its insolubility in aqua regia. The double salt was formed by heating finely pulverized Rh in mixture with KCl in a current of chlorine. The double salt was dissolved in water, precipitated with alcohol, washed with alcohol and dried. (Poggend Ann., 13, 1828, 487; Kongl. Vetens. Akad. Handl., 1828.)

In his earlier determination (Rh = 750.68 for O = 100) Berzelius mistook an hydrated oxide for a chloride. (*Ibid.*)

RUBIDIUM.

Kopp determined the specific heat of rubidium chloride. It corresponds to an atomic weight of about 85. (Gmelin-Kraut, l. c.)

Kirchhoff and Bunsen: 85.36 (O = 16).

Determined from the mean of four experiments on the precipitation of the chloride with argentic nitrate. extreme difference was 0.24. Ag = 107.94; Cl = 35.46. An impure mixture of rubidium and potassium chlorides, nearly free from lithium and the earths, was partially precipitated with platinum chloride and the precipitate freed from KCl by repeated boiling with water. The residue was reduced in a current of hydrogen, the rubidium chloride extracted with water, and reprecipitated with platinum chloride. This process was repeated until the potassium lines in the spectrum disappeared. The rubidium was then converted into a mixture of carbonate and oxide, and the caesium separated by extraction with alcohol. The amount of silver precipitated was also tested from time to time and the purification continued till this became constant. (Poggend. Ann., 113, 1861, 339.)

J. PICCARD: 85.41 (O = 16).

Determined by four analyses of rubidium chloride with argentic nitrate. The number is the mean; extreme difference, 0.09. The separation of potassium from rubidium was effected for the different analyses by 6, 7, and 8 successive partial precipitations with platinum chloride, and the separation of caesium by thirty successive extractions of the anhydrous carbonates with warm absolute alcohol. The salt analysed was spectroscopically pure. Ag = 107.94; Cl = 35.46. The experiments were made with Bunsen's assistance. (Erdmann's Journ. für Prak. Chem., 86, 1862, 449.)

L. Grandeau, who is sometimes credited with making a determination of Rb, expressly disclaims doing so. He mentions Bunsen's value as the true atomic weight and says that his analyses of the sulphate, undertaken to test its purity, led him to adopt the number 85.4; apparently for brevity's sake. (Annal. de Chim. et de Phys., (3,) 67, 1863, 227.)

R. Godeffroy: 85.476 (O = 16).

Determined by four analyses of rubidium chloride prepared and analysed exactly as Godeffroy determined cosium, q. v.; extreme difference, 0.04. Cl = 85.5; Ag = 108. (Liebig's Am., 181, 1877, 189.)

RUTHENIUM.

Bunsen has determined the specific heat of ruthenium. It corresponds to an atomic weight of about 104. (Gmelin-Kraut, l. c.)

C. E. CLAUS: 104.57 (O = 16).

Determined from three analyses of potassium chlororutheniate by the same method Berzelius had employed for other platinum metals. Claus found an average of 28.783 per cent. Ru; extreme difference 0.48, and 41.063 KCl; extreme difference, 0.51. [If K=39.137, Cl=35.457; this composition gives Ru=104.57. The weighings as given in the memoir are misprinted.] Claus also determined the chlorine with silver; the results were such as to show that the salt was not anhydrous, though it had been dried at 200° in an atmosphere of Cl. The salt was prepared by the evaporation of a solution of ruthenium and potassic hydrate in aqua regia, solution of other chlorides of Ru in dilute HCl, and removal of basic compounds by mechanical concentration in water. Claus later takes the atomic weight of Ru = 104. In this memoir he puts it at 651.387 $(0=100,)\ 104.22$ (0=16,) without mentioning the values of K and Cl. (Poggend. Ann., 65, 1845, 218.)

SELENIUM.

Regnault determined the specific heat of selenium, which accords with an atomic weight of about 79. (Gmelin-Kraut, l.c.)

J. J. Berzelius: 79.23 (O = 16).

Berzelius found that 100 Se absorb 179 dry chlorine gas, and that the product was exactly decomposed by water into chlorhydric acid and selenious acid. [If Cl = 35.457 (Stas) the value follows.] (Poggend. Ann., 8, 1826, 21.)

F. SACC: 78.55 (O = 16); 490.98 (O \doteq 100).

Sacc's experiments are very discordant. He made three experiments on the reduction of a known weight of selenious acid with ammonium bisulphite and chlorhydric acid. The mean result was Se = 490.38; extreme difference, 5.5. In four experiments barium seleniate was decomposed by heating to redness with sulphuric acid in excess. The salt was found to contain 41.95 selenious acid; extreme difference 0.04. For Ba = 856.877 the resulting value is 491.49. The selenium was purified by solution in nitric acid, evaporation and sublimation, and by reduction with HOl and Selenious acid was prepared by ammonium bisulphite. oxidation with nitric acid. Barium seleniate was prepared by precipitation of barium nitrate with sodium seleniate and heating to redness. Sacc regards 490.3 as the most probable value of Se. (Annal. de Chim. et de Phys., (8,) 31, 1851, 119.)

A. Schroetter: 78.6 (O = 16).

Details not given. (Kopp's Jahresbericht, 4, 1851, 318; Sitz.-Bericht der k. k. Acad., 6, 1851, 214.)

ERDMANN AND MARCHAND: 78.6 (O = 16); 492.5 (O = 100).

Determined from experiments on mercuric selenide by the same methods employed for the determination of S, q. v. Three experiments gave 71.726, 71.731, 71.741, per cent. mercury. (Erdmann's Journ. für Prak. Chem., 55, 1852, 202.)

J. Dumas: 76.46 (O = 16).

Determined by seven experiments on the chloridation of selenium. The chloride was condensed in a tube cooled to —20°, after which the escaping gases were led through other tubes filled with asbestos. The extreme difference in the results was 0.46. Cl = 35.5. (Annal. de Chim. et de Phys., (3,) 55, 1859, 129.)

O. Pettersson and G. Ekman: 79.08 (O = 16).

Determined by five analyses of selenious acid. A warm solution of the acid was acidified with chlorhydric acid, precipitated with sulphurous acid and the selenium collected on a glass filter. Many precautions are necessary in the precipitation and drying. The value is the mean; extreme difference, 0.04. (Berlin, Bericht der Chem. Gesell., 9, 1876, 1212; in extenso in the Acta of the Scientific Soc. of Upsala.)

SILICON.

The vexed question of the composition of silicic acid has been settled by H. F. Weber, who found that the specific heat of this element becomes nearly constant above 200° and that the atomic heat is 5.8 for Si = 28. (Poggend. Ann., 154, 1875, 575.)

J. J. Berzelius: 29.63 (O = 16); 185.19 (O = 100).

100 parts of silicon, which had been heated to redness, and freed from silicic acid by hydrofluoric acid, gave 208 parts silicic acid, whence the value. Berzelius also made analyses of barium fluosilicide from which he calculated the oxygen 'contents of the acid at 51.975. This gives for the atomic weight of Si 29.58. (Poggend. Ann., 8, 1826, 20; and Lehrbuch, 3, 1200.)

J. Pelouze: 28.46 (O = 16); 177.88 (O = 100).

A known weight of perfectly pure silver, dissolved in nitric soid, was brought in contact with a known and slightly excessive weight of silicon tetrachloride and the excess titrated with decimal silver solution. The value is derived from the mean of two experiments; difference 0.76 for O = 100; Cl = 448.2, Ag = 1849.01. The chloride was prepared by Ebelmen; it was perfectly transparent, volatilized without residue, and had been dried for a long time in a vacuum. (Paris, Comptes Rend., 20, 1845, 1047.)

J. Dumas: 28.02 (O = 16).

Determined from the mean of two experiments on the tetrachloride which was weighed off in a glass bulb and

introduced, so enclosed, into a vessel centaining water. The bulb was broken and the chlorine contents titrated with argentic nitrate. The difference between the experiments was 0.2 for O = 16, Ag = 108, Cl = 35.5. The chloride was repeatedly rectified; its boiling point was 59°. (Annal. de Chim. et de Phys., (3,) 55, 1859, 129.)

J. Schiel: 28.01 (O = 16).

Determined by two analyses of the tetrachloride. The salt was decomposed with a slight excess of ammonium hydrate and the chlorine titrated with argentic nitrate. The atomic weights of Cl and Ag used are not stated. Schiel found 0.6738 silicic chloride = 2.277 argentic chloride, and 1.3092 silicic chloride = 4.418 argentic chloride. [For Ag = 107.93, Cl = 35.457, these data give Si = 28.13, and 27.89.] (Liebig's Ann., 120, 1861, 94.)

SILVER.

Dulong and Petit, Regnault and others have determined the specific heat of silver and found it in accord with an atomic weight of about 108. (Gmelin-Kraut, l. c.)

MARCET; DAVY; WENZEL; WOLLASTON.

Wollaston in his table of equivalents mentions that Marcet found 441 Cl = 1350 silver, and Davy the same quantity of chlorine = 1860 silver. Wenzel found 200 sulphur = 1860 silver. (Phil. Trans., 104, 1814, 21.)

J. J. BERZELIUS: 108.129 (O = 16); 675.804 (O = 100).

Berzelius found that 100 silver gave 182.75 argentic chloride. Taking Cl = 221.825 he calculates Ag = 1351.607. He expresses uncertainty whether or no this value should not be reduced to one half. (*Poggend. Ann.*, 8, 1826, 180.)

E. Turner: 108 (O = 16).

Turner determined the composition of argentic chloride at 100 silver to 132.8 chloride. These numbers are for

107

vacuum. If Cl = 35.42 (Turner) the value follows. (*Phil. Trans.*, 123, 1833, 536.)

F. Penny: 107.97 (O = 16).

Penny made six experiments on the conversion of silver The silver was dissolved in cold nitric acid, into nitrate. the solution evaporated, and the nitrate fused all in one flask and with precautions against loss by spiriting. found 100 Ag = 157.441 nitrate; extreme difference, 0.028. In five experiments the nitrate from the preceding determinations was converted into chloride, by means of chlorhydric acid, in the same flask, dried, fused, and weighed. Penny could detect no decomposition in fusion. He found 100 Ag = 132.8372 chloride; extreme difference, 0.01. In two experiments silver was dissolved in nitric acid, precipitated with chlorhydric acid, evaporated and fused, giving 132.830 and 132.838. The mean of all seven experiments is 132.836. Penny takes 132.837. From the relations of the chlorides, chlorates, and nitrates of potassium and sodium, Penny had determined the difference between the atomic weights of a chloride and a nitrate at 26.565. gives the molecular weight of argentic chloride at 143.424 and Ag = 107.97. The silver used, as well as the water and the acids, were carefully tested for impurities and a minute amount of solid residue in the twice distilled water and in the acids was allowed for. The weighings were all reduced to vacuum. (*Phil. Trans.*, 129, 1839, 27.)

C. Marignac: 108 (O = 16); 675 (O = 100).

Silver was dissolved in nitric acid and precipitated with chlorhydric acid. One experiment, reduced to vacuum, gave 100 silver = 132.74 chloride, which Marignac considered confirmatory of Berzelius' value, 132.75. He therefore adopted the latter number. 100 potassic chloride were found to produce 192.26 argentic chloride, in two experiments, the difference between which was 0.01. By analysis, by means of heat, of potassic chlorate, Marignac had found the molecular weight of the chloride 932.14, these relations give the molecular weight of argentic chloride at 1792.13 and the atomic weight of silver at 1350. The potassic chloride was prepared by heating the chlorate and cooling the resulting chloride over sulphuric acid. (Liebig's Ann., 44, 1842, 23.)

C. Marignac: 107.922 (O = 16); 674.505 (O = 100).

Marignac redetermined the relation between silver and potassic chloride by Pelouze's method. He found 100 Ag = 69.062 KCl in six experiments, the extreme difference between which was 0.018. In five experiments he found 100 KCl = 192.348 Ag; extreme difference 0.04. He also redetermined the composition of argentic chloride. The silver was dissolved in a long-necked flask and the fumes passed into a second flask containing water. Solution being effected, the water from the second flask was added to the contents of the first, and the whole precipitated with HCl. The chloride was washed, dried, melted and weighed in the same flask. The result was 100 Ag = 182.84 chloride; extreme difference 0.019. Combination of these data with Marignac's old value for the molecular weight of KCl, 932.14, gives Ag = 1349.01. All weighings reduced to vacuum. Berzelius revised the result by throwing out one experiment and by rejecting the correction for vacuum. He thus got Ag = 1849.66. (Berzelius' Jahresbericht, 24, 58; 25, 81; Bibl. Univ. de Genève, 46, 1842, 850.)

In opposition to Prout's hypothesis, Marignac cites his analyses of argentic acetate, in which the escaping gases were forced to pass over porous silver. They gave in three experiments 64.664 silver from 100 acetate; extreme difference 0.005. If C=75, this gives Ag=1349.6. He. also found 100 Ag=157.455 nitrate. [If N=87.5, this gives Ag=1348.88.] He also found 100 Ag=49.556 ammonium chloride. (Liebig's Ann., 59, 284; Bibl. Univ. de Genève, 1846.)

LIEBIG and REDTENBACHER; STRECKER: 107.903 (O = 16); 674.395 (O = 100).

Strecker recalculated Liebig and Redtenbacher's analyses of argentic acetate, tartrate, racemate and malate by the method of least squares, and from the difference in the atomic composition of these salts. He obtained for Ag the value 1348.79. Vide Carbon. (Liebig's Ann., 59, 1846, 280.)

E. J. MAUMENÉ: 108.026 (O = 16); 675.16 (O = 100).

In four experiments argentic oxalate was mixed with sand in a flask and decomposed by heat in a current of air. The products of decomposition were passed over cupric oxide, and through drying tubes and potash tubes. In five experiments the acetate was treated in the same way, but not mixed with sand. The mean result was Ag = 1850.82; extreme difference 0.77. Maumené found it very difficult to purify the oxalate, which showed traces of nitric acid after 100 washings. (Annal. de Chim. et de Phys., (8,) 18, 1846, 41.)

J. S. Stas: 107.93 (O = 16).

Thirteen syntheses of argentic iodide, performed by bringing hydroiodic acid in contact with argentic sulphate or nitrate, gave 100 Ag = 117.5848 iodine. Three analyses of argentic iodate, performed by decomposition by heat in a current of nitrogen or by reduction of the salt, while in suspension, by a current of sulphurous anhydride, gave AgI = 284.779. Hence Ag = 107.928. Four syntheses of the bromide, performed by bringing hydrobromic acid in contact with argentic sulphate, gave 100 Ag = 74.0805 Br.Two analyses of argentic bromate, by reduction while in suspension with sulphurous anhydride, gave Ag Br = 187.87. Hence, Ag = 107.921. Seven syntheses of argentic chloride, three of them by combustion of silver in chlorine, three by precipitation with HCl, and one by precipitation with ammonium chloride, gave 100 Ag = 32.8445 Cl. Stas adopts the number 32.85 on the supposition that no excess of chlorine was possible. The chloride was fused. Two analyses of the chlorate, accomplished by heat or by evaporation with chlorhydric acid, gave Ag Cl = 143.395. Hence Ag = 107.987. Five syntheses of the sulphide, performed by heating silver in a current of sulphur vapor, or of hydrogen sulphide, gave 100 Ag = 114.8522 argentic Six analyses of the sulphate by reduction in a current of hydrogen, showed that 100 sulphate contained 69.203 silver, hence Ag = 107.920, [107.926? vide Sulphur.] From analysis of potassium chlorate, Stas had determined the molecular weight of KCl at 74.59. By twenty-four determinations he found 100 Ag = 69.103 KCl, hence Ag = 107.943. The silver was prepared either by Levol's method or by decomposing an ammoniacal solution of argentic nitrate with a mixture of ammonium sulphite and a copper salt. The metal was heated to the boiling point until the sodium line disappeared and the metallic fumes were a pale blue. To test its purity, it was compared with distilled silver. See Stas's determinations of Cl, Br, I, S,

and K. All weighings reduced to vacuum. (Slas, Untersuch. über Chem. Prop., Leipzig, 1867.)

SODIUM.

The specific heat of sodium has been determined by Regnault and indicates an atomic weight of about 28. (Gmelin-Kraut, l. c.)

H. DAVY; F. H. WOLLASTON: 23.28 (O = 16); 145.5 (O = 100).

Davy found that 134 Cl combine with 88 Na to form sodium chloride. If Cl = 441, the value follows. (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius: 23.164 (O = 16).

Berzelius found that 100 Na Cl = 244.6 Ag Cl. [If Ag Cl = 143.387, (Stas,) the value follows.] (*Poggend. Ann.*, 8, 1826, 189.)

F. Penny: 23.046 (O = 16).

Penny made four experiments on the conversion of the chlorate into the chloride by means of HCl. A known weight of the salt was dissolved in a flask in the acid and evaporated, dried and weighed without removal. The sodium chloride was not fused. The mean result was that 100 chlorate equals 54.930 chloride; extreme difference, 0.02. This relation gives the molecular weight of the chloride at 58.5. Penny had found the atomic weight of Cl = 35.454; hence the value for Na. [If Cl = 35.457 (Stas,) Na = 23.043. Stas himself found 23.048.] The sodium chlorate was prepared by precipitating potassium chlorate with sodium bitartrate, and purifying the sodium chlorate by recrystallization. The weighings are for vacuum. (Phil. Trans., 129, 1839, 25.)

J. Pelouze: 22.97 (O = 16); 143.59 (O = 100).

A known weight of perfectly pure silver was dissolved in nitric acid, and brought in contact with a known and slightly excessive weight of sodium chloride, and the excess titrated with decimal silver solution. The mean result of three experiments was that 100 Ag = 51.141 Na Cl; extreme difference, 0.083. The value follows for Ag = 1349.01; Cl = 443.2. The sodium chloride was prepared either from sodium sulphate and barium chloride, or from sodium carbonate and chlorhydric acid, or from a very pure rock salt. It was repeatedly recrystallized and was dried at 200° or melted. (Paris Comptes Rend., 20, 1845, 1047.)

J. Dumas: 23.011 (O = 16).

Determined from the mean of seven experiments on the titration of sodium chloride with argentic nitrate; extreme difference, 0.09. Ag = 108; Cl = 35.5 [Dumas gives the mean as 23.014 instead of 23.0114.] For five experiments Na Cl recrystallized ten times and melted was employed. For two experiments (giving an average of 28.036) the residue from the incineration of the acetate was used to prepare Na Cl, which was recrystallized four times and melted. (Annal. de Chim. et de Phys., (3,) 55, 1859, 129.)

J. S. Stas: 23.043 (O = 16).

According to the mean of 10 determinations 100 Ag = 54.2078 Na Cl; extreme difference 0.0038. The sodium chloride was found to contain a minute quantity of silicic acid which reduces the result from Na = 28.049 to 23.045 for Ag = 107.93; Cl = 35.457. According to the mean of five determinations 100 Na Cl = 145.4526 sodium nitrate; extreme difference 0.025. If N = 14.044 this gives Na = 23.045. The lowest determination gives Na = 23.042. The sodium chloride was purified by recrystallization and in part by conversion into sodium chloroplatinate. The weighings are for vacuum. (Stas, Untersuch. über Chem. Prop., Leipzig, 1867.)

STRONTIUM.

Regnault determined the specific heat of strontium chloride. It corresponds to an atomic weight of about 87.5. (Gmelin-Kraut, L. c.)

M. H. KLAPROTH; F. H. WOLLASTON: 94.4 (O = 16); 590 (O = 100).

Klaproth found 42 sulphuric anhydride = 58 strontium oxide; whence the value for S = 200. (*Phil. Trans.*, 104, 1814, 20.)

F. STROMEYER; 87.34 (O = 16); 545.929 (O = 100).

According to Berzelius, Stromeyer found that 100 strontium chloride = 181.25 argentic chloride; whence the value, for Ag = 1849.66; Cl = 221.64. (Berzelius' Lehrbuch, 3, 1229.) In Gilbert's Ann., 54, 1816, 251, Stromeyer refers to this analysis as by V. Rose. Stromeyer himself found 0.5 grm. carbonate = 75.5394 c. c. carbon di-oxide [which gives Sr = 88.26 if 1000 c. c. carbon di-oxide weigh 1.96483 grm.] Stromeyer calculated Sr = 552.28 for O = 100.

—. SALVETAT: 88 (O = 16); 550 (O = 100).

Determined from the loss of weight of strontium carbonate by calcination and on driving off carbon di-oxide with sulphuric acid. Details not given. (*Paris Comptes Rend.*, 17, 1848, 818.)

J. Pelouze: 87.68 (O = 16); 548.02 (O = 100).

A known weight of perfectly pure silver was brought in contact with a known and slightly excessive amount of strontium chloride and the excess titrated with decimal silver solution. The number is the mean of two experiments; extreme difference, 0.2. Ag = 1849.01; Cl = 443.2. The chloride was purified by recrystallization and was dried at 200° or below redness. (Paris Comptes Rend., 20, 1047.)

C. Marignac: 87.54 (O = 16).

Marignac made experiments on three different preparations of strontium chloride, (1,) (2,) (8.) Compared with silver by Pelouze's method it was found that ten grammes strontium chloride = (1) 8.103; (2) 8.099; (3) 8.101 silver. The same strontium chloride converted into sulphate gave (1) 6.887; (2) 6.8855; (3) 6.884 sulphate. In both these series of experiments the strontium was weighed as airdried, hydrous, crystalline chloride. Comparison gives Sr

= (1) 43.79; (2) 43.82; (3) 43.77. In each experiment of the latter series the water was determined by driving it off at a red heat. It was proved that the chloride does not undergo decomposition at this temperature, and the water contents was found to vary no more than 0.0005 of the total weight. In three more experiments the water was determined, and the anhydrous salt analysed by Pelouze's method giving (1) 43.77; (2) 43.74; (3) 43.76. Ag = 108; Cl = 35.5; S = 16. The chloride was prepared (1) from the chemically pure chloride of commerce by precipitating barium with sulphuric acid, separation of lime by precipitation of the strontium chloride by HCl gas and washing with chlorhydric acid. The purity was tested by the solubility of a portion converted into sulphate. The chloride was finally redissolved and precipitated with alcohol. (2) was prepared from (1) by a repetition of the same process. (3) was prepared from (2) by recrystallization. (Bibl. Univ., Arch. des Sciences, (2,) 1, 1858, 220.)

J. Dumas: 87.52 (O = 16).

Determined from the mean of six experiments on the analysis of strontium chloride with argentic nitrate. The extreme difference was 0.14, Cl = 35.5; Ag = 108. The salt was purified by boiling with sulphuric acid, and precipitation with and recrystallization from chlorhydric acid. These processes were in some cases several times repeated. The pure salt was fused in a current of HCl gas. (Annal. de Chim. et de Phys., (8), 65, 1859, 129.)

SULPHUR.

Deville and Troost and others have determined the density of sulphur in the gaseous form. It corresponds to an atomic weight of about 32. The specific heat of sulphur also agrees moderately well with this value. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

J. J. Berzelius; F. H. Wollaston: 32 (O = 16); 200 (O= 100).

According to Wollaston, Berzelius found that plumbic sulphide was composed of 86.64 lead and 13.36 S. Hence the value, for lead = 1295. (*Phil. Trans.*, 104, 1814, 20.)

J. J. Berzelius: 32.19 (O = 16) 201.165 (O = 100).

A known weight of lead was dissolved in pure nitric acid, precipitated with sulphuric acid and evaporated. The mean result of four experiments was that 100 Pb = 146.44 sulphate. The variation was only in the fifth figure. If lead = 1294.498 the value follows. [If this relation is recalculated with Stas's atomic weight of lead, S = 82.096.] (Poggend. Ann. 8, 1826, 16.)

E. Turner: 32.17 (O = 16).

Determined from syntheses of plumbic and baric sulphates. The former gave 16.083, the latter, 16.087. Ba = 68.7, Pb = 103.6. The numbers are for vacuum. Vide Barium and Lead. (Phil. Trans., 123, 1833, 539.)

T. Thomson: 32 (O = 16); 200 (O = 100).

This chemist found the specific gravity of sulphurous acid in mean of two experiments, 2.22216, almost exactly double 1.1111 which he takes (on utterly untenable grounds) for the specific gravity of oxygen. (Erdmann's Journ. für Prak. Chem., 8, 1836, 870; Records of General Science by R. D. Thomson, 1836, 179.)

Erdmann and Marchand: 32.004 (O = 16); 200.026 (O = 100).

Determined by four experiments on the decomposition of mercuric sulphide by copper, in a current of carbon dioxide, the mercury being caught in a cold receiver. The mean composition was found to be for vacuum 86.211 mercury and 13.789 sulphur, extreme difference, 0.017 Hg. If Hg = 1250.6, the value follows. In purifying the sulphide it was first heated to drive off excess of sulphur and then sublimed three times, the first and last portions of the sublimate being rejected. (Erdmann's Journ. für Prak. Chem., 31, 1844, 396.)

J. J. Berzelius: 32.12 (O = 16); 200.75 (O = 100).

Berzelius' former value, 201.165, is changed by the new value for lead, 1294.645 to 200.8017. Three new experiments were made by gently heating argentic chloride in a current of hydrogen disulphide. The mean of three experiments gives 8 = 200.706; extreme difference 0.11. Cl = 443.38, Ag = 1349.66. (Berzelius' Jahresbericht, 25, 1845, 37, and Lehrbuch, 3, 1185.)

H. STRUVE: 32.002 (O = 16).

Determined by six experiments on the reduction of a known weight of argentic sulphate in a current of hydrogen. The number is the mean; extreme difference, 0.146. Ag = 108. The sulphate was prepared by precipitating the nitrate with an excess of sulphuric acid, and drying at a high temperature. (Liebig's Ann., 80, 1851, 203; Berzelius' Jahresbericht, 30, 20.)

J. Dumas: 32.0196 (O = 16).

Determined by five experiments on the combustion of silver in sulphur vapor. The number is the mean; extreme difference, 0.054. Ag = 108. The sulphur was purified by repeated distillation. The silver was heated to redness in a current of sulphur vapor, the excess of sulphur being afterwards distilled off in a current of carbon di-oxide. (Annal. de Chim. et de Phys., (3,) 55, 1859, 147.)

J. S. Stas: 32.0742 [?] (O = 16).

According to the mean of six analyses of argentic sulphate by decomposition in a current of hydrogen at as low a temperature as possible, 100 sulphate yield 69.203 [more exactly 69.20317] silver; extreme difference, 0.012. Five syntheses of the sulphide, performed by heating silver in a current of sulphur vapor or hydrogen disulphide, showed that 100 silver = 114.8522 sulphide; extreme difference, 0.005. By comparing these figures, which are for vacuum, Stas deduces S = 32.0742; Ag = 107.920. [There seems to be a trifling error in this calculation. The weighings seem to be correct, for the means correspond to the details given. As given, the numbers indicate S = 32.058; 6. The latter is almost identical with Stas's Ag = 107.926. mean value, 107.930.] The sulphate was prepared by the action of sulphuric acid on argentic nitrate, or by solution of silver in sulphuric acid. The salt was heated above the boiling point of sulphuric acid. (Stas, Unters. über Chem. Prop., Leipzig, 1867.)

TANTALIUM.

Deville and Troost have determined the vapor density of tantalium chloride. It agrees with an atomic weight of 182. (Paris Comptes Rend., 64, 1867, 294.)

J. J. Berzelius: 167.74 (O = 16).

Berzelius decomposed the sulphide in dry chlorine gas and decomposed the resulting chloride with water. 99.75 parts sulphide yielded 89.35 tantalic acid. On the supposition that the acid contains three atoms of oxygen Berzelius calculates the atomic weight at 1148.365 for 8 = 200.75. [If the acid contains five atoms of oxygen the value becomes 167.74.] (Poggend. Ann., 4, 1825, 14, and Lehrbuch, 3, 1209.)

Rose denies that the sulphide formed, as Berzelius prepared it, by heating tantalium in carbon disulphide vapor is a constant compound. (*Poggend. Ann.*, 99, 580.) Marignac, however, shows that Berzelius, Rose and Hermann, obtained constant results from its analysis, from 89.50 to 90 acid from 100 sulphide. If Ta = 182, the sulphide would give 90.24 acid. (*Liebig's Ann.*, S, 4, 1866, 358.)

H. Rose: 172 (O = 16).

Out of twelve analyses of the chloride, in which both the chlorine and the tantalic acid were determined, Rose selected two in which the agreement was best. [These analyses calculated for Ag = 107.93, Cl = 35.457, give Ta = 171.96.] The chloride was prepared from tantalic acid especially freed from tungsten and tin by mixing with carbon, drying in carbon di-oxide, and heating in a current of chlorine in which the salt was allowed to cool. Excess of chlorine was expelled by dry air, and the salt was hermetically sealed in glass. Rose supposed the acid to contain two atoms of oxygen and therefore deduces the value 859.81. (O = 100). (Poggend. Ann., 99, 1856, 75.)

Marignac seems to prove that the material with which Rose dealt contained niobium. He states that the chlorides of the two elements cannot be separated from one another, and that there are no characteristics by which their purity can be decided. (*Liebig's Ann.*, S, 4, 1866, 352.)

R. HERMANN:

Hermann made many analyses of tantalium salts to which, however, he ascribes quite incomprehensible formulas. Marignac has shown that his methods were utterly inadequate to produce pure preparations. He assumes two atoms of tantalium and three atoms of oxygen in the acid and gives the atomic weight as 645. (O=100.) (Erdmann's Journ. für Prak. Chem., 70, 1857, 198.)

C. Marignac: 182 (O = 16).

Berzelius', Rose's and Marignac's analyses of the double fluoride of tantalium and potassium show that the fluorine is combined with Ta and potassium in proportions of two to five. The salt has also exactly the crystal form of the niobium salt. Hence the acid is a ditantalic pentoxide. Four experiments were made on this salt by drying at 100°, moistening with sulphuric acid and heating gradually till the excess of acid was driven off. The potassic sulphate was leached out, evaporated, melted and weighed, and the tantalic acid heated to redness and weighed. mean potassic sulphate contents was found to be 44.29 per cent; extreme difference, 0.15. The mean amount of tantalic acid obtained was 56.59; extreme difference, 0.25. If K = 39, these data give Ta = 182.3. Four analyses were also made of the ammonium salt. This contained traces of potassium which were determined and allowed for in each case. The mean amount of tantalic acid obtained was 65.25 per cent; extreme difference, 0.34. This gives Ta = 182, the number which Marignac adopts. The salts were obtained by dissolving tantalic acid, which had not been heated to redness, in fluohydric acid, adding potassic or ammonic hydrate and purifying by recrystallization. These salts are much less soluble than the corresponding niobium and titanium salts. (Liebig's Ann., S. 4, 1866, 234.)

TELLURIUM.

Regnault and Kopp have each determined the specific heat of tellurium and found it in accord with an atomic weight of about 128. (Gmelin-Kraut, l. c.)

J. J. Berzelius 129.03 (O = 16); 806.452 (O = 100).

A known weight of metallic tellurium was oxidized with nitric acid, the excess of acid being driven off by heat. It was found that 100 Te gave 124.8 tellurious acid. (*Poggend. Ann.*, 8, 1826, 24.)

J. J. Berzelius: 128.28 (O = 16); 801.76 (O = 100).

Determined as before but with purer material. Three experiments were made, which gave 802.838, 801.786, 801.74. Berzelius took the mean of the latter two. The tellurium was prepared from tetradymite by heating with potassium carbonate and olive oil in a closed crucible, dissolving the potassium telluride so formed in water free from air, precipitating the tellurium by a current of air and distilling it in a current of hydrogen. (Poggend. Ann., 32, 1834, 16.)

K. von Hauer: 128.06 (O = 16).

Determined from the mean of five experiments on the precipitation of bromine with argentic nitrate from the double bromide of potassium and tellurium. The bromine contents was found to be 69.9236 per cent., for Ag = 108.1; Br = 80; extreme difference 0.172. If K = 89.2, the value follows. The salt was prepared by mixing tellurium and potassic bromide in atomic proportions, adding water and bromine, heating to drive off excess of bromine and repeated recrystallization. (Erdmann's Journ. für Prak. Chem., 73, 1858, 98; Sitz-Bericht der k. k. Acad.)

J. Dumas: 129 (O = 16).

No details are given. (Annal. de Chim. et de Phys., (3,) 55, 1859, 129.)

THALLIUM.

Regnault determined the specific heat of thallium. It agrees with an atomic weight of 204. (Gmelin-Kraut, l. c.)

A. Lamy: 204 (O = 16).

Three analyses of the chloride with argentic nitrate gave

a mean of 204; extreme difference 1.2. An experiment on the precipitation of the sulphate with barium nitrate gave 204.3. [The atomic weights used were probably those accepted by Dumas.] The salts were purified by recrystallization. (Annal. de Chim. et de Phys., (3,) 67, 1863, 411.)

W. Crookes: 202.96 (O = 16).

These determinations were made from the sulphate, which was prepared with great care. By decomposing the sulphate with potassic iodide and weighing the thallic iodide formed, the atomic weight was found at 202.73; by precipitation with barium nitrate, 203.55; with chlorhydric acid and alcohol, thallic chloride being weighed, 201.85; from the amount of sulphate produced from a known weight of metal, 203.1; by precipitation with platinum chloride, 203.56. The values taken for Cl, I, etc., are not given; [they were probably those accepted by Dumas.] (Erdmann's Journ. für Prak. Chem., 92, 1864, 277; Chem. News.)

H. Werther: 204 (O = 16).

In five experiments Werther decomposed thallic iodide with potassic hydrate and zinc, both perfectly pure, and precipitated the iodine with silver. The mean result of these experiments was Tl = 204.4; extreme difference 1.7. [The value assumed for I is not stated. One experiment, which gave exactly 204, according to Werther, recalculated for Ag = 107.93; I = 126.85 gives Tl = 203.63.] Three experiments were made by decomposing the iodide with ammoniacal solution of argentic nitrate and weighing the argentic iodide formed. These determinations gave Tl = 203.47; extreme difference 0.3. The preparation of the iodide is not given. (Erdmann's Journ. für Prak. Chem., 92, 1864, 136.)

M. Hebberling: 203.94 (O = 16).

Hebberling made three experiments on the sulphate by precipitation with barium chloride, which gave in mean Tl = 204.13; extreme difference 0.2. He also made two experiments on the chloride by precipitation with argentic nitrate. These gave 203.8 and 203.5. The atomic weights assumed are not stated. [If Ag = 107.93; Cl = 35.457; the first analysis of the chloride gives Tl = 203.44. The data for the second analysis are misprinted. If a probable correction of a single figure is made, the data give Tl = 10.000

203.026.] The salts were purified by recrystallization. (Liebig's Ann., 134, 1865, 11.)

W. Crookes: 204.155 (O = 16).

Determined by experiments on the solution of metallic thallium in nitric acid and evaporation to dryness. The number is the mean of ten experiments; extreme difference, 0.038. The balance stood in a partial vacuum, and the weighings were made at two different pressures and calculated for Very elaborate precautions were taken throughvacuum. out. Crookes also mentions determinations made with barium nitrate, but gives no data. The thallium was prepared in seven different lots by the reduction of as many different salts which had been purified by recrystallization &c. The metal was fused in lime. The reagents were expecially prepared by methods similar to those of Stas. Crookes took N = 14.009, O = 15.96, and calculated for Tl the value 203.642. [If O = 16, the value becomes 204.155.] (Phil. Trans., 163, 1873, 277.)

THORIUM.

From the isomorphism existing between thorium, tin, and titanium, and from the similarity of thorium to zirconium, Delafontaine and Marignac believe the oxide to contain two atoms of oxygen. (Liebig's Ann., 131, 100.) Neither the specific heat of this element nor the vapor density of any of its compounds has been determined so far as I know.

J. J. Berzelius; 238 (O = 16); 1887.72 (O = 100).

From the sulphate, precipitated by heating a solution of the salt and redissolved in cold water, Berzelius got the values 748.493 and 735.713 by precipitating with barium chloride. He also analysed the double sulphate of potassium and thorium. From the relation between the sulphuric acid and the thorium oxide found, the atomic weight would seem to be 750.63, while the relation between the potassic sulphate obtained, and the amount of oxide gives 740.6. These numbers are calculated on the supposition that the oxide contains a single atom of oxygen. Ba =

855.29, 8 = 200.75, K = 488.856. (Poggend. Ann., 16, 1829, 398, and Lehrbuch, 3, 1224.)

J. J. Chydenius: 236.64 (O = 16).

This chemist analysed the sulphate, the double sulphate of potassium and thorium, the oxalate, the acetate and the formate, getting results which vary from 228.52 to 243.76. He averages with his own results analyses made by Berzelius and by Berlin, which, however, alter the result inappreciably. According to Delafontaine, the methods employed for purification are ineffectual. Chydenius assumes a single atom of oxygen in the oxide. (Poggend. Ann., 119, 1863, 55.)

N. J. Berlin: 231.64 (O = 16).

Chydenius reports two analyses of the oxalate by Berlin which gave for thorium 57.87 and 57.95, or 281.48 and 231.80. (*Poggend. Ann.*, 119, 1863, 56.)

M. Delafontaine: 281.5 (O = 16).

Determined from analyses of the sulphate. Fourteen experiments on the decomposition of this salt, by the heat of a strong double-draught lamp, gave a mean of 52.51 per cent. oxide; extreme difference, 0.88. In three experiments the sulphur contents of the salt was determined by precipitation with barium chloride after the sulphate had been decomposed with ammonium oxalate. The mean amount of sulphuric anhydride so found was 31.92 per cent.; extreme difference, 0.78. Three experiments on the water contents gave 15.68 per cent; extreme difference, 0.21. The sum of these means is 100.11. The value of thorium was calculated from the relation of the oxide to the sulphuric anhydride for S = 32, Ba = 137. The salt was prepared from thorite and from orangite by decomposition with sulphuric acid and recrystallization of the sulphate The purification was continued with the help of heat. until the crystals and the mother liquor had exactly the same composition. Marignac assisted at this investigation. (Liebig's Ann., 131, 1864, 100.)

P. T. CLEVE: 233.88 (O = 16).

Cleve made six analyses of the anhydrous sulphate, getting in mean Th = 233.8; extreme difference, 1.36. From

analyses of the oxalate he got 283.97; extreme difference, 0.6. (Kopp's Jahresbericht, 1874, 261; Bull. Soc. Chim., (2,) \$1, 116.)

TIN.

Regnault and Kopp have each determined the specific heat of tin. It agrees with an atomic weight of about 118. Dumas, Cahours and others have determined the vapor density of volatile tin compounds with a similar result. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

J. J. Berzelius; 117.647 (O = 16); 735.294 (O = 100).

Berzelius determined this value by oxidizing pure tin foil by means of nitric acid and weighing the oxide. He found 100 tin = 127.2 stannic acid. (*Poggend. Ann.*, 8, 1826, 184.)

G. J. MULDER: 116.112 (O = 16); 725.7 (O = 100).

Two experiments were made by oxidizing tin with nitric acid, evaporating, drying, and heating to redness. They gave each 100 tin = 127.56 stannic acid; whence the value. All possible precautions are said to have been taken. The metal was prepared by the reduction of pure oxide with soot and a flux. (Erdmann's Journ. für Prak. Chem., 48, 1849, 85; Scheikundige Onderzoek., 5. Deel, 260.)

C. L. Vlaanderen: about 118. (O = 16).

Determined from experiments on the oxidation and reduction of tin and stannic acid in vessels of various materials. The experiments regarded as the most accurate were made on the reduction of the acid in a current of hydrogen in porcelain vessels. The acid had been heated in platinum. These experiments gave 59.04 and 59.12. Stannic acid heated in glass or porcelain was found to retain nitric acid. (Kopp's Jahresbericht, 11, 1858, 188; Mulder, Scheikundige Verh. en Onderzoek., 2. Deel, 150.)

J. Dumas: 118.08 (O = 16).

Two experiments were made on the oxidation of pure tin by nitric acid. The stannic acid being heated white hot in platinum vessels gave for the atomic weight 59.1, The tin employed was prepared from pure chloride. Two experiments on the titration of the chloride with argentic nitrate gave 59.06 and 59.03. Ag = 108, Cl = 35.5. (Annal. de Chim. et de Phys., (3,) 55, 1859,

TITANIUM.

The specific heat of titanic acid has been determined by Regnault and by Kopp, and indicates an atomic weight of about 50. Dumas determined the vapor density of the tetrachloride at 6.886. [If the molecular weight of O = 32, and if Cl = 35.457, this gives Ti = 56.025.] (Gmelin-Kraui, l. c., and Poggend. Ann., 9, 1827, 441.)

H. Rose: 61.17 (O = 16).

Determined by roasting titanium sulphide and weighing the titanic acid formed. The highest result obtained was 1.017 sulphide from 0.757 acid. This result Rose adopted on the supposition that an excess was impossible. For S=201.16 these data give Ti=62.25 (O=16); 389.1 (O=100). [If S=32, Ti=61.17.] The sulphide was prepared by heating titanic acid in a current of carbon disulphide. (Gilbert's Ann., 73, 1828, 135.)

Rose subsequently expressed the opinion that the sulphide employed in this analysis was impure, and contained undecomposed titanic acid, but afterwards came to the conclusion that it was perfectly pure, accounting for the variation of the results from those he obtained later by the theory that the sulphide and the oxide of this element, like those of tantalium, were entirely dissimilar compounds. Marignac has shown that tantalium sulphide is of normal constitution. (Poggend. Ann., 99, 1856, 576.)

H. Rose: 48.28 (O = 16).

Titanium chloride was decomposed with water, titanic acid precipitated by ammonic hydrate, and the chlorine Precipitated from the filtrate with argentic nitrate. Taking Ag = 1351.607, Cl = 221.325; Rose calculated the chlorine contents in four experiments at from 74.43 to 74.53 per

cent; mean 74.46 and Ti at 303.686. According to Gmelin-Kraut, these analyses recalculated for Stas's values give Ti = 48.28. The chloride was prepared by the action of chlorine on a mixture of titanic acid and carbon, and was rectified four or five times over potassium and mercury. It was clear and developed no chlorine on decomposition with water. (Poggend. Ann., 15, 1829, 145.)

C. G. Mosander: 47.33 (O = 16); 295.81 (O = 100).

Mosander determined the oxygen contents of titanic acid at from 39.83 to 40.82 per cent.; mean 40.427. Mosander never described the method of analysis. [The oxygen contents was probably determined from the chloride, for the above data give Ti = 294.7, while Berzelius records the determination as having given 295.81.] (Poggend. Ann., 19, 1830, 212, and Berzelius' Lehrbuch, 3, 1211.)

J. PIERRE: 60.36 (O = 16).

Determined by three experiments on the titration of the chloride with argentic nitrate by Pelouze's method. Pierre does not give the values taken for Cl and Ag. He calculates the atomic weight of Ti at 314.69. [If Ag = 107.93, Cl = 35.457; his data give Ti = 314.75 (O = 100); 50.36 (O = 16), with an extreme difference in the latter case of 0.08.] He made two other determinations giving lower results, but it was found that the chloride employed was slightly decomposed by contact with air. The chloride was prepared from artificial titanic acid which was free from iron, and was further purified by fractional distillation. (Annal. de Chim. et de Phys., (3,) 20, 1847, 257.)

• A. Demoly: 56.512 (O = 16).

Determined by experiments on the tetrachloride. The salt was decomposed with water, the titanic acid precipitated by ammonic hydrate, and the chlorine precipitated in the filtrate, after the excess of ammonic hydrate had been volatilized and the solution acidified. Both precipitates were weighed. Demoly calculates the atomic weight of Ti at 350, without mentioning what values he accepted for silver and chlorine. [If Ag = 107.93, Cl = 35.457; the atomic weight, calculated from the argentic chloride, is 353.2 (O = 100); or 56.512 (O = 16), with an extreme difference in the three experiments of 0.88 for O = 16.] The chloride was prepared from rutile by preliminary conver-

sion into nitride, &c. It was purified by rectification over mercury and potassium. (Liebig's Ann., 72, 213; Laurent and Gerhardt, Comptes Rend., 1849, 325.)

TUNGSTEN.

Regnault has determined the specific heat of tungsten, and Roscoe the vapor density of the chloride. These experiments place the atomic weight of tungsten at about 184. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

J. J. Berzelius: 189.26 (O = 16); 1183.355 (O = 100).

A weighed quantity of tungstic acid was reduced in a current of hydrogen, again weighed, then re-oxidized and reweighed. The number is the mean result of the two operations. The number is given in Berzelius' Lehrbuch as 1188.36 with the data, which are also given in Poggend. Am., 8, 23. It is pointed out in Graham-Otto that this value must be misprinted, an observation which I have verified. (Poggend. Ann., 4, 1825, 152.)

Berzelius made an earlier determination than the foregoing by the oxidation of the sulphide, getting 1207. He points out the source of error in this experiment arising from the formation of irreducible sulphate. (Berzelius'

Jahresbericht, 5, 1825, 121.)

R. Schneider: 184.12 (O = 16); 1150.78 (O = 100).

Schneider made five experiments on the reduction of tungstic acid with hydrogen in a porcelain tube heated by a charcoal fire. These analyses gave the mean contents of the acid at 79.316 tungsten per hundred; extreme difference, 0.096. This composition corresponds to an atomic weight of 1150.39. He also made three experiments on the combustion of tungsten, getting a mean of 79.327 tungsten per 100 acid; extreme difference, 0.005, or an atomic weight of 1151.17. The value taken is the mean. The tungstic acid was prepared by decomposing ammoniotungstic sulphide with chlorhydric acid, washing the precipitate with acid, solution in ammonia, reprecipitation with chlorhydric acid, and so on until a perfectly pure product was obtained. The tungstic acid was finally dried and

heated to redness. (Erdmann's Journ. für Prak. Chem., 50, 1850, 163.)

R. F. MARCHAND: 184.1 (O = 16); 1150.6 (O = 100).

Determined from two experiments on the reduction of tungstic acid in a current of hydrogen, and two experiments on the combustion of tungsten. These determinations were made in the same manner as and at the same time with Schneider's. The extreme difference was 3.5 for O = 100. (Liebig's Ann., 77, 1851, 263.)

J. B. von Borck: 183.816 (O = 16); 1148.85 (O = 100).

Determined by seven experiments on the reduction of tungstic acid at a white heat by hydrogen, and by two experiments on the combustion of tungsten. The number is the mean; extreme difference, 10.38 for O = 100. The tungstic acid was prepared from Wolframite by fusing the mineral with potassium carbenate, solution in water containing alcohol, precipitation with calcic chloride and decomposition of the calcic tungstate with chlorhydric acid. The tungstic acid so produced was converted into ammonium salt which, on decomposition, yields a compound free from iron and manganese. (Erdmann's Journ. für Prak. Chem., 54, 1851, 254.)

A. RICHE: 174 (O = 16).

This value was reached by five determinations of the amount of water produced by the reduction of tungstic acid in a current of hydrogen, which gave a mean of 87.07; extreme difference, 1.78. The tungstic acid was obtained by heating the ammonium salt, or by the decomposition of the oxychloride produced by heating tungstic acid and carbon in a current of chlorine. (Annal. de Chim. et de Phys., (3,) 50, 1857, 10.)

J. Dumas: 184 (O = 16).

Dumas made six experiments on the reduction of tungstic acid in hydrogen at a high temperature in a nacelle of unglazed porcelain, and two experiments on the titration of the chloride with argentic nitrate. The extreme difference between the results was 0.69 for O=8. The acid was pre-

pared by gently heating the ammonium salt in a muffle. (Annal. de Chim. et de Phys., (3,) 55, 1859, 144.)

F. A. Bernoulli: 186.8 (O = 16); 1167.5 (O = 100).

Berneulli made five experiments on the reduction of tungstic acid by hydrogen in a porcelain tube at a very high temperature, two experiments on the amount of water formed in reduction, and four experiments on the oxidation of tungsten. The mean result was W = 93.41; extreme difference, 0.75. [If experiment 9, in which oxidation seems to have taken place, is left out, the mean becomes 93.35; extreme difference, 0.18.] The tungstic acid was prepared from ammonium tungstate which had been boiled for several days with nitric acid. The tungstic acid was One part of it was green, another part heated to redness. The determinations from the different colored acids did not differ, and Bernoulli considers them isomeric modifications of the same compound. There appear to be misprints in the data given. (Poggend. Ann., 111, 1860, 599.)

C. Scheibler: 184 (O = 16).

Scheibler reached this value by five determinations of the water contents (9 molecules) of barium metatungstate. From determinations of the barium and the tungsten in the same compound Scheibler reached other values, but he regards the water determination as the most trustworthy. (Erdmann's Journ. für Prak. Chem., 83, 1861, 328.)

E. Zettnow: 183.952 (O = 16).

Determined from analyses of ferrous tungstate and argentic tungstate. A known weight of ferrous tungstate was melted with sodium carbonate and the mass dissolved. The ferric hydrate was thoroughly washed, dissolved in chlorhydric acid, reduced to ferrous chloride with zinc of known composition, and titrated with potassic permanganate in several measured portions. Four such series of experiments were made, and gave a mean of 92.038 for W; extreme difference, 0.33. The ferrous tungstate was prepared by melting pure anhydrous sodium tungstate with ferrous chloride and sodium chloride, dissolving, separating impurities, crystallizing, washing the crystals with water, chlorhydric acid and sodium carbonate. The argen-

tic tungstate was decomposed with nitric acid and titrated with sodium chloride or decomposed with hot sodium chloride solution, the argentic chloride being weighed. Five experiments gave a mean of 91.915 for W; extreme difference, 0.13. The argentic tungstate was prepared by the precipitation of sodium tungstate with argentic nitrate, thorough washing and drying in yellow light. The permanganate solution was prepared according to Mohr and tested with ammonio-ferrous sulphate. Fe = 28, Ag = 108. (Poggend. Ann., 130, 1867, 30.)

H. E. ROSCOE: 184.04 (O = 16).

Determined by reducing tungstic acid in a current of hydrogen, by reoxidizing the metal, and by reducing the chloride in a current of hydrogen, the chlorhydric acid being condensed and estimated as argentic chloride. In the experiments on the acid, that compound was reduced, and reoxidized three times with almost identical results. The mean of the second and third reductions of the same sample gave W = 183.84. In the experiments on the chloride, the chlorine and the tungsten were each determined, and gave a mean of 184.25 for Cl = 35.5. The tungstic acid was prepared by the decomposition of the chloride, washing and heating to redness in a platinum vessel. It was canary yellow. The chloride was prepared from pure tungsten. (Liebig's Ann., 162, 1872, 366.)

URANIUM.

No certainty exists as to the relation between the equivalent and the atomic weight of uranium. The latter is commonly accepted as about 120. Mendelejeff gives grounds for supposing it to be 240, (Liebig's Ann., S. 8, 1871, 178,) and L. Meyer regards it as probably 180, a value which accords well with the specific heat of the black oxide as observed by Regnault. (Gmelin-Kraut, l. c.) For the purposes of this paper it seems best to retain the customary value.

J. A. Arfvedson: 128.6 (O = 16).

Determined by experiments on the reduction of uranoso-

uranic oxide and on the oxidation of uranous oxide. By combustion of uranous oxide in oxygen he found in two experiments that 100 oxide combined with 3.695 and with 3.73 oxygen. From the reduction of the green oxide he found that 100 uranous oxide combine with 3.67 oxygen. He deduces as the mean 3.688. Regarding uranous oxide as the metal, Arfvedson calculated the atomic weight at 2711.36. [If the lower oxide is a protoxide, the data give 128.6 for O = 16.] The uranous oxide was prepared from pitchblende by solution in aqua regia, precipitation of heavy metals with hydrogen sulphide, precipitation with ammonic hydrate, solution in ammonium carbonate to remove iron, reprecipitation, heating to redness, washing with chlorydric acid to remove impurities, and reduction in hydrogen. (Poggend. Ann., 1, 1824, 254.)

E. Peligot: 119.128 (O = 16).

In two experiments the amount of carbon in the acetate was found to be 11.27 and 11.3; mean 11.285. In one experiment the uranic oxide was determined at 67.8 per cent. [From these data the above value follows.] Peligot takes 120 or 750, C = 75. The preparation of the salt is not given. Peligot mentions the oxalate and gives analyses, but does not deduce an atomic weight from them. (Annal. de Chim. et de Phys., (8,) 5, 1842, 39.)

J. J. EBELMEN: 118.86 (O = 16); 742.875 (O = 100).

Ebelmen made six experiments on the reduction of the oxalate to uranous oxide by hydrogen and heat. The value follows with an extreme difference of 0.65 for C = 75; H = 12.5. All the weighings were reduced to vacuum. To obtain pure oxalate, the nitrate was precipitated by oxalic acid and this preparation decomposed by heat. The oxide thus obtained was digested with chlorhydric acid, washed, dissolved in nitric acid, recrystallized, and precipitated with oxalic acid. The oxalate was dried at 100°. According to Rammelsberg the reduction of the oxalate is accompanied by the separation of carbon which remains with the oxide. (Annal. de Chim. et de Phys., (8,) 5, 1842, 189.)

Berzelius, Arfvedson, Marchand: 128.4 (O = 16); 802.49 (O = 100).

While Arfvedson was making his first determination, Berzelius also made an experiment on the combustion of uranous oxide getting 103.685 uranic from 100 uranous oxide. Marchand (Erdmann's Journ. für Prak. Chem., 23, 1841, 498) got in the same way 103.668. The average of the combustion experiments of all three chemists is 103.694, whence Berzelius calculates the value. (Berzelius' Jahresbericht, 22, 1842, 113.) Peligot and Rammelsberg, as well as Marchand, point out faults in this method, such as the probable condensation of hydrogen in the protoxide and the tendency to form higher oxides. (Poggend. Ann., 59, 1843, 4.)

C. RAMMELSBERG.

This chemist made experiments on the reduction by hydrogen of the green oxide, prepared in various ways, and got results varying from 580.4 to 767.6 for O=100. (Poggend. Ann., 59, 1843, 9.) By precipitation of uranous chloride with silver he reached the number 787.5 for Cl=442.65. The chlorine contents found varies in three experiments from 73.89 to 74.46. The chloride was prepared by heating uranous oxide in an atmosphere of chlorine. (Poggend. Ann., 56, 1842, 821.)

J. Wertheim: 119.42 (O = 16); 746.86 (O = 100).

Determined by three experiments on the decomposition of the double acetate of uranium and sodium. The mean loss of acetic acid by heating the salt to redness was 82.477 per cent.; extreme difference, 0.036. The number follows for C = 75, H = 6.25, Na = 390.9. [In Poggend. Ann., 57, 484, an abstract is given of a paper read before the academy (of Berlin?) by Mitscherlich, in which he states that Wertheim's experiments above described give 740.512. Berzelius in his Jahresbericht, 23, 137, makes or quotes the same statement, so also does Rammelsberg, Poggend. Ann., 59, 4, and it has been repeated elsewhere. I have recalculated the data given by Wertheim and find the results correctly deduced in his own report. For Na = 23.043 (Stas); the data give U = 119.53.] The salt was prepared from uraninite by solution in nitric acid, precipitation with hydrogen sulphide, evaporation of the filtrate to dryness, solution in hot water, crystallization and recrystallization, heating the crystals to drive off nitric acid, solution in acetic acid, digestion with sodium carbonate and recrystallization. (Erdmann's Journ. für Prak. Chem., 29, 1843, 209.)

C. Rammelsberg: about 120 (O = 16).

Determined in six experiments, undertaken at Berzelius' suggestion, by treating uranous oxide with nitric acid and sulphuric acid and weighing the sulphate. It is very difficult to weigh the uranous oxide which constantly increases in weight. Two experiments were made on the green oxide, which was prepared either by heating uranous oxide, or the nitrate, in air. Two experiments were made on magnesium uraniate by dissolving the compound in nitric acid and heating to redness. The compound was found unstable in character. One experiment was made by heating the double acetate of uranium and sodium and three experiments by heating the double acetate of barium and The results obtained varied from 633.17 to 753.76. Rammelsberg considers the determinations confirmatory of Wertheim's and Ebelmen's. (Poggend. Ann., 66, 1845, 95.)

E. Peligor: 120 (O = 16); 750 (O = 100).

Determined by combustion of the oxalate in a current of air, both the carbonic acid and the green oxide of uranium being weighed. At first Peligot got only 780 as the atomic weight by this process, but by repeating the recrystallization of the salt until determinations gave constant results, he got a mean of 750. He says that he came to the same value by comparing the amount of uranic oxide obtained from the acetate with the weight of the salt employed. (Paris Comptes Rend., 22, 1846, 487.)

VANADIUM.

Roscoe has determined the vapor density of vanadium chloride. It agrees with an atomic weight of about 51. (L. Meyer, l. c.)

J. J. Berzelius: 52.47 (O = 16).

Berzelius made four experiments on the relation between the higher and the lower oxides of vanadium, three by reduction with hydrogen at a very high temperature and one by oxidation. He supposed the higher oxide to have the formula VO₂, and the lower VO, and consequently got for the atomic weight the number 855.84 (O = 100). R. Schneider has shown that the data as given by Berzelius are discordant, (Poggend. Ann., 88, 819,) a fact of small importance in view of the succeeding investigation. The higher oxide analyzed by Berzelius was produced by gently heating the ammonium salt. (Poggend. Ann., 22, 1831, 14; Kongl. Vet. Akad. Handl., 1831.)

Roscoe examined some ammonium vanadate which Berzelius had sent Faraday and found that it contained phos-

phorus. (Liebig's Ann., S, 6, 1868, 93.)

H. E. Roscor: 51.33 (O = 16).

Roscoe made four experiments on the reduction of vanadic acid $(\nabla_2 O_5)$ in carefully purified hydrogen. The acid was prepared from ammonium vanadate. To free this compound from phosphorus and silicic acid it was powdered, decrepitated with sodium in an iron crucible, washed with water and with chlorhydric acid, re-oxidized with nitric acid, chloridized in a current of chlorine, the chloride rectified and decomposed with water. The acid so obtained was dried, moistened with sulphuric acid, exposed to the fumes of fluohydric acid for ten days and melted. This pure acid was first heated for several hours in dry air and afterwards in hydrogen. The mean result of four experiments was V = 51.871; extreme difference, 0.228. Nine experiments were made on the titration of the chloride by Pelouze's method. Eight experiments were also made on the analysis of the chloride with argentic nitrate by the ordinary method. The mean of the seventeen experiments on the chloride gives the contents in chlorine at 61.276 per cent.; extreme difference, 0.69. This composition indicates an atomic weight of 51.29. Roscoe takes Cl = 35.457, Ag = The vanadium chloride was purified by rectification over sodium in a current of carbon di-oxide. The reagents were prepared according to Stas. (Liebig's Ann., S, 6, 1868, 86.)

Roscoe mentions atomic weight determinations by Czudnowicz as giving 55.35. This chemist, however, did not calculate an atomic weight from his analyses, but used that obtained by Berzelius. (*Poggend. Ann.*, 120, 1868, 17.)

YTTRIUM.

The composition of yttrium oxide is not definitely settled. Mendelejeff concludes from the general behavior of its compounds that it is a sesqui-oxide. As, however, all the chemists who have made atomic weight determinations of this element have considered it a prot-oxide, I shall assume it to be so and the atomic weight, therefore, about 60.

J. J. Berzelius:
$$64.29$$
 (O = 16); 401.84 (O = 100).

This determination was made before the discovery of erbium and can scarcely be correct. The value was reached by analysis of the sulphate with barium chloride. Ba = 856.88, S = 201.165. (Poggend. Ann., 8, 1826, 186; 10, 1827, 841.)

N. J. Berlin: 59.7 (O = 16).

According to Blomstrand in Berlin, Ber. der Chem. Ges., 1873, 1467. I can find no other record of this determination which probably appeared in the Forhandl. ved de Skandinaviske Naturforsk, 1860, 448.

0. Popp:
$$68 (0 = 16)$$
,

The mean of four analyses of the sulphate showed that 40.15 oxide were equivalent to 38.28 sulphuric anhydride, giving a molecular weight for the oxide of 42.015; extreme difference, 0.018. The yttrium was precipitated with sublimed oxalic acid, the free acid being afterwards neutralized with ammonia. The sulphuric acid was precipitated with barium chloride in the filtrate with precautions. Popp, who denies the existence of erbium and terbium, separated yttrium from the cerite oxides by precipitation with barium carbonate, yttrium remaining in solution, S = 16, Ba = 68.5. (Liebig's Ann., 131, 1864, 183.)

M. Delafontaine: about 64 (O = 16).

Delafontaine does not pretend that this number is exact. It is derived from analyses of the sulphate. His method of separation was essentially Mosander's, which was proved by Popp and by Bunsen and Bahr to give impure salts. (Liebig's Ann., 134, 1865, 108.)

Bahr and Bunsen: 61.7 (O = 16).

Determined by saturating the oxide with sulphuric acid as in the determination of erbium, q.v. Partial recrystallization does not produce pure yttrium nitrate, but only concentrates traces of didymium in the salt. Didymium must be separated with potassic sulphate. Erbium nitrate is more easily decomposed by heat than yttrium nitrate. The nitrates were therefore partially decomposed, yttrium nitrate dissolved out and the process repeated until there was no trace of erbium or didymium visible in the spectroscope. The mean of two determinations gave Y = 30.85; difference, 0.1. S = 16. (Liebig's Ann., 137, 1866, 21.)

M. Delafontaine: 58.5 (O = 16).

Determined by three experiments on the sulphate which gave in mean 48.28 per cent. oxide for S = 32. [In the Jahresbericht this determination is reported as giving Y = 74.5. Yttrium is apparently a misprint for yttrium oxide.] The yttrium salt seems to have been prepared according to the method of Bahr and Bunsen. (Kopp's Jahresbericht, 1866, 184; Bibl. Univ., Arch. des Sciences, (2), 25, 1866, 112.)

P. T. CLEVE AND O. M. HOEGLUND: 59.7 (O = 16).

Determined by analysis of the sulphate. The oxide was purified by heating the nitrates, etc., according to N. J. Berlin. (Blomstrand, in Berlin, Bericht der Chem. Ges., 1873, 1467; Bihang till Vet. Akad. Handl, 1873, B. 1, 3, No. 8.)

ZINC.

The specific heat of zinc has been determined by Regnault and others. The vapor density of volatile organic compounds has been determined by Frankland and others. These experiments agree in placing the atomic weight at about 65. (Gmelin-Kraut, l. c.; L. Meyer, l. c.)

GAY-LUSSAC, BERZELIUS, WOLLASTON: 65.547 (O = 16); 409.67 (O = 100).

In his experiments on the oxidation of zinc Gay-Lussac

zinc. 135

found that 100 Zn = 24.41 oxygen. This value is repeatedly cited in his memoir. (Gilbert's Ann., 30, 1811, 297; Mêmoire D'Arceuil, 2, 174.) Wollaston gives the same figures on Gay-Lussac's authority. (Phil. Trans., 104, 1814, 21.) Wollaston calculates from these data Zn = 410, (O = 100)Berzelius in each of two experiments got 100 Zu = 124.4(Gilbert's Ann., 37, 1811, 460.) In Poggend. Ann., 8, 1826, 184, as well as in his Lehrbuch, Berzelius cites Gay-Lussac as having found 100 Zn = 24.8 oxygen. states that his own determinations were in perfect accordance with these figures, and calculates from them the stomic weight of zinc at 403.226 or 64.52, and this was the accepted value for many years. I cannot find any other determinations by either of these chemists, and am obliged to suppose that there was a mistake made in recording the data from which Berzelius made his calculations; if so, it is remarkable that neither Berzelius nor the other chemists who determined this value perceived it; for the question was reopened during Berzelius' life, and A. Erdmann made his determination at Berzelius' request.

V. A. JACQUELIN: 66.24 (O = 16); 414 (O = 100).

This number was reached by measuring the amount of hydrogen developed by a known weight of zinc from sulphuric acid on the supposition that the specific gravity of hydrogen is 0.0624. The results seem to have been inconsistent. Subsequently Jacquelain arrived at the same number by oxidizing an impure zinc of known composition. (Paris Comptes Rend., 14, 1842, 636; and Annal. de Chim. et de Phys., (3,) 7, 1843, 204.)

P. A. FAVRE:
$$66$$
, $(O = 16)$; 412.5 $(O = 100)$.

Favre made four experiments on the combustion of zinc oxalate, the carbon di-oxide being collected and its weight compared with that of the oxide. The mean result was Zn=412.66; extreme difference, 1.11. C=75. He also made three experiments by passing the hydrogen developed by a known weight of zinc over cupric oxide, the water being caught. These experiments gave in mean Zn=412.16; extreme difference, 0.65 for H=12.5. (Annal. de Chim. et de Phys., (8), 10, 1844, 163.)

A. Erdmann; 65.05 (O = 16); 406.591 (O = 100). Determined by oxidizing pure zinc with nitric acid, and

driving off the acid by heating the salt in a porcelain crucible. Platinum is attacked. The number is the mean of four experiments; extreme difference, 0.698. The zinc was prepared by mixing pure oxide with carbon, and distiling in a current of hydrogen. (Berzelius' Jahresbericht, 24, 1844, 182; Efversigt af Kongl. Vet. Akad. Handl., 1, 3.)

ZIRCONIUM.

Deville and Troost have determined the vapor density of the chloride. It agrees with an atomic weight of about 90. (L. Meyer, l. c.)

J. J. Berzelius: 89.6 (O = 16).

In one experiment the sulphate was decomposed with ammonic hydrate, the oxide weighed and the sulphuric acid precipitated with barium chloride. In five experiments the sulphate was decomposed at a white heat, ammonium carbonate being added at the close of the operation. The mean result was that 100 parts of sulphuric anhydride unite with 75.853 parts of zirconium oxide; extreme difference, 0.23. Berzelius deduces the value 840.08 for O = 100, S = 201.165; on the supposition that the oxide contains three atoms of oxygen. [Being a binoxide, this relation gives Zr = 89.6 for O = 16.] The sulphate seems to have been prepared by dissolving the oxide in sulphuric acid and expelling the excess of acid by heat. (Poggend. Ann., 4, 1825, 126.)

R. HERMANN:

This chemist made some experiments on the chloride getting in three determinations a mean of 839.45 for O = 100 and on the tri-oxide supposition. The extreme difference was 20.1. Cl = 443.65. The chloride was produced by heating the oxide with carbon in a current of chlorine. Hermann adopts not his own but Berzelius' determination. (Erdmann's Journ. für Prak. Chem., 31, 1844, 77.)

C. Marignac: 90 (O = 16).

Determined from analyses of potassium fluo-zirconiate. The salt was decomposed with sulphuric acid, the excess of acid driven off by heat, the residue weighed, the potassic sulphate leached out with water, and the residue again weighed. Marignac does not pretend that the determination is accurate. The results gave from 45.01 to 45.48. He thinks that some potassic sulphate may have escaped solution, and therefore takes the minimum. K = 39, S = 16.

According to Marignac, Deville also found the atomic weight of zirconium somewhat higher than Berzelius by analysis of the chloride with which he determined the vapor density. (Annal. de Chim. et de Phys., (3,) 60, 1860, 257.)

				·	
·					
		·			
					!
·	·			,	
					1
					1
					.
			·		

APPENDIX.

DETERMINATIONS BY T. THOMSON.

In Thomson's Annals of Philosophy, volumes 16 and 17, 1820-21. Thomson published a series of papers descriptive of experiments undertaken for the purpose of verifying Prout's hypothesis. His method consisted in mixing reagents in what he considered equivalent proportions, and after precipitation examining portions of the supernatant liquid for an excess of each of the salts supposed to neutralize one another. In all except four cases, either the salt analyzed was a sulphate and the precipitant barium chloride, or the determination was dependent upon such an analysis; yet although Thomson took barium = 70, in no instance was he able to detect either barium or sulphuric acid in the residual solution when the quantity of the re-agents corresponded to the atomic weights which he adopts. Comparison of his results with those reached by more accurate experimenters will make this exact neutralization appear impossible, nor were his contemporaries able to repeat his experiments successfully. Thomson's determinations are, as such, utterly valueless, yet as they were for many years extensively accepted in English and American scientific literature they are inserted here for reference. In the following table Thomson's numbers are multiplied, when necessary, for the sake of comparison with the values now accepted.

DETERMINATIONS INVOLVING BARIUM = 70.

Arsenic76	Magnesium24
Barium140	Manganese56 Nickel52
Bismuth216	Nickel52
Calcium40	Nitrogen14
Carbon12	Phosphorus32
Chlorine36	Potassium40
Chromium56	Silver110
Cobalt52	Sodium24
Copper64	Strontium88°
Iron56	Sulphur32
Lead208	Zinc68

189

THOMSON FURTHER DETERMINED-

Antimony at 132 by oxidation.

Boron at 12 from analysis of borax.

Mercury at 200 by conversion of the oxide into chloride.

Tin at 116 by oxidation with nitric acid.

REDUCTION OF WEIGHINGS TO VACUUM.

In discussing the analyses recorded in the foregoing pages, or in reconciling atomic weight determinations by various chemists, it may be found convenient to employ the following table. The maximum error involved is less than 0.01 per cent. or 0.1 milligram per gram.

GRAM WEIGHTS BEING OF BRASS, FRACTIONS OF PLATINUM.

For substances the sp. gr. of which exceeds 6.1; no correction is necessary.

For substances the sp. gr. of which is less than 6.1:—

To correct the entire grams; multiply their number by the correction in the table opposite the sp. gr. of the substance, found in the first column, and add the product to the observed weight.

To correct the fractions of a gram, multiply the correction opposite the sp. gr. of the substance, found in the third column of the table, by the first two decimal figures of the observed weight, if the sp. gr. of the substance is less than 3, and by the first decimal only, if the sp. gr. exceeds 3, and add the product to the observed weight.

ALL WEIGHTS USED BEING OF PLATINUM.

For substances the sp. gr. of which exceeds 7.8, no correction is necessary.

For substances the sp. gr. of which is less than 7.8:—Multiply the correction opposite the sp. gr. of the substance, found in the *third* column, by the number of grams, tenths and hundredths observed, if the sp. gr. falls short of 3, or by the number of grams and tenths, if the sp. gr. exceeds 3, and add the product to the observed weight.

The table shows within what limits it is necessary to know the sp. gr.

(Weights of pecific Gravi	Brass) for ty between—	Correction per Gram. Error $< \frac{1}{10}$ Mg.	(Weights of F Specific Grav	
27.788 and	1 11.064	-0.000 067 gram.		
11.064	6.904	0.000 000	51.766 an	d 18.568
6.904	5.019	+0.000 067	18.568	7.807
5.019	8.948	0.000 188	7.807	5.480
8.948	8.247	0 000 200	5.480	4.222
3.247	2.759	0.000 267	4.222	8.488
2.759	2.899	0 000 888	8.488	2.893
2.899	2.122	0.000 400	2.898	2,500
2,122	1.908	0.000 467	2.500	2.201
1.903	1.724	0.000 588	2.201	1.965
1.724	1.576	0.000 600	1.965	1.776
1.576	1.452	0.000 667	1.776	1.619
1,452	1.877	0.000 788	1.619	1.488
1.877	1.254	0 000 800	1.488	1.877
1.254	1.174	0.000 867	1.877	1.281
1.174	1.103	0.000 988	1.281	1.197
1.103	1.041	0.001 000	1.197	1.124
1.041	0.985	0.001 067	1.124	1.059
		0.001 188	1.059	1.002
		0.001 200	1.002	0.950

(Sill. Amer. Jour., 16, 1878, 265; Liebig's Ann., 195, 1879, 222.)



INDEX TO AUTHORITIES.

Acta Universitatis Lundensis, 90.

Afhandlingar i Fysik, etc., 79.

Annales de Chimie et de Physique, 8, 10, 12, 18, 15, 19, 20, 21, 22, 24, 28, 30, 85, 40, 41, 42, 48, 47, 50, 51, 55, 56, 61, 62, 68, 65, 66, 67, 71, 72, 78, 74, 77, 78, 79, 81, 84, 86, 87, 88, 98, 97, 100, 102, 104, 106, 109, 111, 118, 115, 118, 119, 128, 124, 126, 127, 129, 136, 187.

Berlin, Bericht der Deutschen Chemischen Gesellschaft, 18, 48, 49, 54, 59, 89, 105, 184.

Berzelius' Lehrbuch der Chemie, 7, 12, 14, 17, 18, 22, 28, 26, 27,28, 29, 88, 46, 49, 54, 56, 71, 74, 76, 80, 82, 88, 86, 98, 105, 112, 114, 116, 120, 121, 124.

Berzelius' Jahresbericht über die Fortschritte der Chemie, etc., 8, 41, 56, 62, 65, 66, 74, 78, 79, 80, 86, 92, 98, 100, 108, 114, 115, 125, 130, 186.

Bibliothèque Universelle de Genève, Archives des Sciences, 15, 16, 22, 27, 82, 41, 46, 47, 62, 72, 84, 87, 88, 91, 98, 100, 108, 118, 184.

Bihang till Vetenskaps Akademien Handlingar, 184. (See Kongliga Vetenskaps Akademien.)

British Association Reports, 15, 19.

Bulletin de la Société Chimique, 58, 54, 122.

Bulletin de l'Académie Royale des Sciences, etc., de Belgique, 82, 88.

Chemical News, 25, 48, 65, 89.

Edinburgh Royal Society Transactions, 56, 80.

Erdmann's Journal für Praktische Chemie, 18, 28, 24, 27, 28, 29, 31, 35, 37, 44, 45, 48, 49, 52, 55, 58, 59, 60, 62, 67, 68, 69, 76, 77, 78, 79, 81, 88, 86, 88, 90, 92, 94, 96, 102, 104, 114, 117, 119, 122, 126, 127, 130, 186.

Fresenius' Zeitschrift für Analytische Chemie, 48, 50, 52, 89.

Forhandlinger ved de Skandinaviske Naturforskeres, 188.

Gilbert's Annalen der Physik, etc., 96, 128, 185.

Gmelin-Kraut, Handbuch der Chemie, 7, 9, 12, 18, 18, 19, 20, 21, 28, 26, 87, 48, 46, 49, 56, 62, 64, 65, 68, 78, 76, 79, 81, 88, 86, 90, 91, 94, 95, 96, 98, 101, 102, 108, 106, 110, 111, 118, 117, 118, 122, 128, 125, 128, 184.

Halle, Zeitschrift für die Gesammten Naturwissenschaften, 45.

Jenaische Zeitschrift für Medicin und Naturwissenschaft, 52.

Journal de Pharmacie et de Chimie, 94.

Journal of the Chemical Society, 47, 88, 97.

Klatzo, Ueber die Constitution der Beryllerde, 18.

Kongliga Vetenskaps Akademien Handlingar, 21, 83, 94, 95, 101.

Kopp's Jahresbericht über die Fortschritte der Chemie, 45, 52, 58, 78, 104, 122, 184.

Laurent and Gerhardt's Comptes Rendus Mensuels, etc., 125.

Liebig's Annalen der Chemie, etc., 15, 26, 28, 29, 80, 81, 82, 85, 86, 87, 40, 41, 48, 51, 52, 58, 54, 68, 78, 75, 78, 86, 91, 98, 99, 108, 106, 107, 115, 116, 117, 120, 121, 125, 126, 128, 182, 183, 184.

Meyer, L., Moderne Theorien der Chemie, 7, 9, 12, 20, 28, 81, 118, 122.

Mitscherlich's Lehrbuch der Chemie, 80.

Œfversigt af Vetenskaps Akademien Færhandlingar, 79, 186. (See Kongliga Vetenskaps Akademien.)

Otto's German Translation of Graham's Chemistry, 88, 84, 67.

Paris Comptes Rendus, 8, 18, 15, 20, 26, 40, 42, 50, 55, 56, 58, 75, 88, 85, 89, 98, 95, 97, 100, 105, 111, 112, 116, 181, 185.

Pelouze, Traité de Chimie, 8.

Philosophical Magazine, 8, 17, 22, 80.

Philosophical Transactions of the Royal Society, 14, 26, 28, 88, 87, 28, 39, 48, 49, 54, 56, 65, 71, 72, 76, 82, 91, 92, 96, 99, 106, 107, 110, 112, 118, 114, 120.

Poggendorff's Annalen der Physik, etc., 9, 10, 11, 12, 18, 14, 17, 18, 19, 21, 24, 25, 26, 29, 88, 86, 88, 48, 45, 46, 49, 51, 58, 54, 59, 60, 61, 64, 65, 63, 67, 68, 91, 94, 95, 98, 99, 101, 102, 108, 104, 105, 106, 110, 114, 116, 118, 121, 122, 128, 124, 125, 127, 128, 129, 180, 181, 182, 188, 185, 186.

Proceedings of the American Academy of Arts and Sciences, 11, 12.

Schweigger's Journal für Chemie und Physik, 28, 86.

Scheikundige Verhandelingen en Onderzoekingen, 122.

Silliman's American Journal of Science, 25, 26, 87, 46, 74, 75.

Sitzungs-Bericht der k. k. Akademie zu Wien, 48, 81, 88, 97, 104, 118.

Stas, Untersuchungen über Chemischen Proportionen und Atomgewichte, 28, 48, 59, 62, 64, 78, 76, 94, 98, 101, 110, 111, 115.

Stockholm Akademien Handlingar, 18. (See Kongliga Vetenskaps Akademien.)

Thomson's System of Chemistry, 7, 14, 17, 20, 28, 24, 29, 61, 78.

Thomson's Annals of Philosophy, 58, 61, 189.

Thomson, R. D., Records of General Science, 92, 114.

Zeitschrift für Berg Hütten-und Salinen-Wesen im Preussichen Staate, 49, 51.

INDEX TO ATOMIC WEIGHT DETERMINATIONS.

	•	
ALLEN. (JOHNSON and)	BERZELIUS, J. J.	Borck, J. B. von
Cæsium 25	Chlorine 37, 38	Tungsten126
Aluminium 7	Chromium 43	Boron 19, 140
Anderson, T.	Copper 49	Brandes, R.
Nitrogen 93	Fluorine 54	Manganese 80
ANDREWS, T.	Gold 56	Brodie, B. C.
Barium 15	Iodine 61	Graphon 33
Platinum 98	Iridium 64	Phosphorus 97
Antimony 9, 140	Iron 65, 66	Bromine 20
ARAGO. (BIOT and)	Lead 71	Buehrig, H.
Carbon 28	Lithium 74	Cerium 37
Nitrogen 91	Magnesium 76	Bunsen, R. W.
ARFVEDSON, J. A.	Manganese 79, 80	Cæsium 25
Lithium 73	Molybdenum 83	Indium 60
Manganese 79	Nickel 87	Bunsen, R.W., and J. JEGEL.
Uranium128, 129	Nitrogen 91	Cerium 35
Arsenic 12 AWDEJEW.	Osmium 94	BUNSEN, R. W. (J. F. BAHR
	Palladium 95	and)
Beryllium 17	Phosphorus 96	Erbium 53
Russ T P	Platinum 98	Yttrium134
BAHR, J. F.	Potassium 99	BUNSEN. (KIRCHOFF and) Cæsium 24
Magnesium 78	Rhodium101	Rubidium102
BAHR, J. F., and R. W. BUNSEN.	Selenium104	Rubidium102
Erbium 53	Silicon 105 Silver 106	Cadmium 23
Yttrium 134	Sodium110	Cæsium 24
BALARD, A. J.	Sulphur113, 114	Calcium 26, 139
Bromine 20	Tantalium 116	CAPITAINE, H.
Barium 13	Tellurium118	Iron 66
BERINGER, A.	Thorium120	Carbon28
Cerium 34	Tin122	Cerium 34
BERLIN, N. J.	Tungsten 125	CHENEVIX, R.
Chromium 44	Uranium129	Copper 49
Molybdenum 84	Vanadium131	Chlorine 37, 139
Thorium121	Yttrium 133	CHOUBINE.
Yttrium 122	Zinc 134	Lanthanium 68
Bernoulli, F. A.	Zirconium 136	Chromium 43
Tungsten 127		CHYDENIUS, J. J.
BERTHIER, P.	BERZELIUS and DULONG.	Thorium121
Nickel 86	Hydrogen 57	CLAUS; C. E.
Beryllium 16	Nitrogen 91	Iridium 65
DERZELIUS, J. J.	BERZELIUS and LIEBIG.	Ruthenium103
Aluminium 7	Carbon 31	CLEVE, P. T.
Antimony 9	BIOT and ARAGO.	Didymium 53
Arsenic 12	Carbon 28	Erbium 53
Barium 14	Nitrogen 91	Lanthanium 70
Beryllium 17	Bismuth 18, 139	Thorium121
Boron	BLOMSTRAND, C. W.	CLEVE, P. T., and O. M.
Bromine 21	Niobium 90	Hoegland.
Calcium 26, 28	Boisbaudran, L. DE	Erbium 54
Carbon 29	Gallium 56	Yttrium134
10		145

INDEX.

Cobalt 46, 139	Dumas, J.	GAY-LUSSAC, L. J.
COMMAILLE. (MILLON and)	Magnesium 79	Zinc134
Copper 50	Manganese 81	GERHARDT, C.
Соокт, J. Р., Jr.	Molybdenum 84	Chlorine 41
Antimony II	Nickel 87	GIBBS, W.
Copper 49, 139	Nitrogen 93	Cerium 37
CROOKES, W. Thallium119, 120	Phosphorus 97 Potassium 100	Cobalt 46 GMELIN, C. G.
Czudnowicz, C.	Selenium104	Lithium 73
Lanthanium 60	Silicon105	GMELIN-KRAUT.
	Sodium 111	Boron 19
DAVY, H.	Strontium113	Carbon 33
Fluorine 54	Sulphur 115	fodine 62
Mercury 82	Tellurium118	GODEFFROY, R.
Silver106	Tin122	Cæsium 25
DAVY, J. Sodium110	Tungsten126 Dumas and Stas.	Rubidium103
DEBRAY, H.	Carbon 30	Gold 56 Graphon 33
Molybdenum 84	Carbon	Crapton 33
DEBRAY. (DEVILLE and)	EBELMEN, J. J.	Hagen, R.
Osmium 94	Uranium129	Lithium 74
DELAFONTAINE, M.	EINBRODT, P.	Hampe, W.
Erbium 53 Molybdenum 84	Nitrogen 93	Copper50
	EKMAN, G. (O. PETTERSSON	HAUER, K. VON
Thorium121	and)	Cadmium 23
Yttrium 133, 134	Selenium105	Manganese 80 Tellurium118
DEMOLY, A. Titanium124	Erbium 53 ERDMANN, A.	HEBBERLING, M.
Deville, H. Sainte-Claire	Zinc135	Thallium119
Nickel 87	ERDMANN and MARCHAND.	HENRY, W.
DEVILLE and DEBRAY.	Calcium 27	Magnesium 76
Osmium 94	Carbon 30	HERMANN, R.
DEVILLE. (WOEHLER and)	Copper 49	Cerium 34
Boron 19	Hydrogen 58 Iron 66	Didymium 51
DEXTER, W. P.	Iron 66	Lanthanium 68, 69
Antimony 10	Mercury 82 Nickel 86	Lithium 73 Niobium 90
Didymium 51 Diehl, K.	Selenium104	Tantalium117
Lithium 75	Sulphur114	Zirconium136
DULONG. (BERZELIUS and)	ERK, C.	HILLEBRAND, W. F.
Hydrogen 57	Didymium 52	Didymium 53
Nitrogen 91	Lanthanium 70	HISINGER, W.
Dumas, J.		Cerium 34
Aluminium 8	FAVRE, P. A.	HOEGLUND, O. M. (P. T.
Antimony 10 Arsenic 12, 13	Zinc135	CLEVE and) Erbium 54
Barium 16	Fluorine 54 FOURCROY and THENARD.	Yttrium134
Bismuth 19	Mercury 82	HOLZMANN, M.
Bromine 22	Fownes, G.	Lanthanium 68
Cadmium 24	Carbon 30	Hydrogen 57
Calcium 26, 28	FREMY, E.	
Carbon 29	Fluorine 55	Indium 59
Chlorine 42	Osmium 94	Iodine 61
Cobalt 47	FREMY. (PELOUZE and) Aluminium 8	Iridium 64 Iron 65, 139
Copper 50 Fluorine 55	Aummum 8	ISNARD.
Hydrogen 58	Gallium 55	Aluminium 8
Iodine 61, 62	GAY-LUSSAC, L. J.	
Iron 67	Iodine 61	JACQUELAIN, V. A.
Lead 72		Chromium 44


INDEX.


JACQUELAIN, V. A.	LIEBIG, J.	MARIGNAC, C.
Iodine 62		Silver 107, 108
Magnesium 78	LIEBIG and REDTENBACHER.	StrontiumII
Phosphorus 97		Tantalium II
Zinc135	Silver 108	Zirconium136
JEGEL, J. (R. BUNSEN and)	LIEBIG. (BERZELIUS and)	Mather, W. W.
Cerium 35	Carbon 31	
JOHNSON and ALLEN.	Lithium 73	Maumené, E. J.
	Loewig, C.	Chlorina
Cæsium 25		Chlorine 41
	Bromine 21	Iron 6
KEMPE. (LEICHTE and)	LONGCHAMP.	Potassium100
Molybdenum 85	Magnesium 76	Silver108
Kessler, F.	LOUYET, P.	MENDELEJEFF, D.
Antimony 10	Fluorine 55	Cerium 37
	LUCCA, S. DE	Didymium 52
Arsenic 13 Chromium 45	Fluorine 55	Erbium 53
Vergrand Drivers	1	Lanthanium 70
Kirchhoff and Bunsen.	MACDONNELL, A.	Uranium128
Cæsium 24	Magnesium 78	MERCER.
Rubidium 102	Magnesium 76, 139	Cæsium 25
KJERULF, T.	MAGNUS, G.	Mercury 81, 140
Cerium 35	Iron 65	MEYER, L.
KLAPROTH, M. H.	MALLET, J. W.	
Potassium 99		Molybdenum 85
Strontium 112	Lithium 74	Uranium128
KLAPROTH. (WOLLASTON	Manganese 79, 139	MILLON, E.
and)	MARCET.	Iodine 62
Barium 13	Chlorine 37	Mercury 83
KLATZO, G.	Silver106	MILLON and COMMAILLE.
Beryllium 18	Marchand, R. F.	Copper50
KRALOVANSZKY.	Tungsten126	MITSCHERLICH, E.
	Uranium129	Carbon 30
Lithium 73	MARCHAND and Scheerer.	Moberg, A.
KRAUT. (GMELIN—)	Magnesium 76	Chromium 44
Boron 19	MARCHAND. (ERDMANN	Molybdenum 83
Carbon 33	and)	Mosander, C. G.
Iodine 62	Calcium 27	Lanthanium 68
1	Carbon 30	Titanium124
LAGERHJELM, P.	Copper 49	MULDER, G. J.
Bismuth 18	Hydrogen 58	Tin122
LAMY, A.	Iron 66	
Thallium 118	Mercury 82	Nickel 86, 139
Lanthanium 67	Nickel 86	Niobium89
LASSAIGNE.	Selenium104	Nitrogen 91
	Sulphur114	Nordenfeldt. (Svanberg
Nickel 86	Manager C	and)
LAURENT, A.	MARIGNAC, C.	
Boron 20	Barium 15, 16	Magnesium 77
Chlorine 40, 42	Bromine 21	NORLIN. (SVANBERG and)
Lead 70, 139	Calcium 27	Iron 66
Lee, R. H.	Carbon 32	
Cobalt 48	Cerium 35	Odling, W.
Nickel 89	Chlorine 39, 40	Aluminium 8
LEFORT, J.	Cobalt	Osmium 94
Chromium 45	Didymium 51	OTTO, F. J.
LEICHTE and KEMPE.	Iodine 62	Cerium 34
Molybdenum 85	Lanthanium 68	Lanthanium 67
LENSSEN, E.		Oxygen 95
Cadmium 24	Nickel 87	,,
Levol, A.		Palladium 95
Gold 56	Nitrogen 92	PELIGOT, E.
Potassium100	Potassium 99, 100	Chromium 44
10(422)(11(1) 100	10tassium 99, 100	

PELIGOT, E.	KOSCOE, H. E.	STAS, J. S.
Uranium 129, 131	Tungsten128	Chlorine 42
PELOUZE, J.	Vanadium 132	Hydrogen so
	Does U	Hydrogen 59 Lodine 63
Arsenic 12		100me 03
Barium 15	Niobium 89, 90	Lead 72
Chlorine 40	Tantalium 116	Lithium 75
Nitrogen 93	Titanium123	Nitrogen 93
Phosphorus 96	Rose, H., and Weber.	Potassium100
Determinen 90	Antimonia	Cil
Potassium100	Antimony 9	Silver109
Silicon105	Rose, V.	Sodium111
Sodium 110	Phosphorus 96	Sulphur 115
Strontium112	Strontium112	STAS, J. S. (J. DUMAS and)
PELOUZE and FREMY.	ROTHOFF, E.	Carbon 30
		Conserve A
Aluminium 8	Cobalt 46	STRECKER, A.
PENNY, F.	Nickel 86	Carbon 32
Chlorine 39	Rubidium 102	Silver108
Nitrogen 92	Russell, W. I.	STROMEYER, F.
Potassium 99	Cobalt 47, 48	Cadmium 23
	Nickel 88, 89	Iron 65
Silver107		T 1011 05
Sodium110	Ruthenium103	Lithium 73
PETTERSSON, O., and G.	_	Strontium112
EKMAN.	SACC, F.	Strontium
Selenium105	Selenium104	STRUVE, H.
PHILLIPS, R.	SALVETAT.	Barium 15
Chladas		
Chlorine 39	Barium 15	Sulphur 115
Phosphorus 96, 139	Calcium 26	STRUVE, H., (L. SVANBERG
Piccard, J.	Strontium112	and)
Rubidium 102	Scheerer, T.	Molybdenum 83
PIERRE, J.	Magnesium 77, 78	Sulphur 113, 139
Titonium		Correspond
Titanium124	SCHEERER, T. (MARCHAND	Syanberg, L.
Platinum 98	and)	Mercury 83
Popp, O.	Magnesium 76, 77	Nitrogen 92
Yttrium 133	SCHEIBLER, C.	SVANBERG and NORDEN-
Potassium 98, 139	Tungsten127	FELDT.
PROUT, W.	SCHIEL, J.	Magnesium -
		Magnesium 77
Iodine 61	Silicon106	SVANBERG and NORLIN.
	Schneider, R.	Iron 66
Rammelsberg, C.	Antimony 9	SVANBERG and STRUVE.
Cerium 35, 36	Bismuth 19	Molybdenum 83
Lanthanium 68	Cobalt 46	,
		Tontolium
Niobium 90	Manganese 81	Tantalium116
Uranium130, 131	Nickel87, 88	Tellurium117
RAWACK.	Tungsten125	Thalén.
Manganese 81	SCHROETTER, A.	Lanthanium 70
REDTENBACHER. (LIEBIG	Phosphorus 97	Thallium118
and)	Selenium 104	
		Menaku, L. J., and F. H.
Carbon 31	SEFSTROEM, N. G.	THENARD, L. J., and F. H. WOLLASTON. Iron
Silver108	Mercury 82	Iron 65
REGNAULT.	Selenium 103	THENARD L. J., (FOURCEOY
Hydrogen 57	SIEWART, M.	and)
Nitrogen 91	Chromium 45	Mercury 82
Prior F and T Discomen		
REICH, F., and T. RICHTER.	Silicon105	THOMSEN, J.
Indium 59	Silver106, 139	Hydrogen 59
Riche, A.	Sodium 111, 139	THOMSON, T.
Tungsten126	SOMMARUGA, E. VON.	Aluminium 7
RICHTER, T. (F. REICH and)	Cobalt 47	Antimony140
Indium 59		·
		Amonic
D 39	Nickel 88	Arsenic 130
Rivot, L. E.	Nickel 88 Stas, J. S.	Arsenic139 Barium14, 139
RIVOT, L. E. Iron 67	Nickel 88 STAS, J. S.	Arsenic 139 Barium 14, 139 Beryllium 17
Rivot, L. E.	Nickel	Arsenic 130

INDEX.

THOMSON, T.	TURNER, E.	WOEHLER and DEVILLE.
Boron 20, 140	Nitrogen 91	Boron 19
Cadmium 23	Silver 106	Graphon 33
Calcium139	Sulphur 114	Wolf, C.
Carbon29, 139	•	Cerium 36
Cerium 34	Unger, B.	Wollaston, F. H.
Chlorine 139	Antimony 11	Calcium 26
Chromium43, 139	Uranium 128	
Cobalt 139		Chlorine 37
Copper139	Vanadium131	Copper 49
Gold 56	VAUQUELAIN, L. N.	Hydrogen 57
Hydrogen 58	Lithium 73	Magnesium 76
Iodine 61	VLAANDEREN, C. L.	Mercury 82
Iron139	Tin122	Nitrogen 91
Lead139		Phosphorus 96
Lithium 73		Potassium 99
Magnesium 139		Silver106
Manganese139	Iron 66	Sodium 110
Mercury140	Wallace, W.	Strontium112
Nickel 139	Bromine 22	Sulphur 113
	WATTS, W. M.	Zinc 134
Phosphorus139		WOLLASTON and KLAPROTH.
Potassium139	Osmium 94	Barium 13
Silver139	WEBER. (H. ROSE and)	WOLLASTON. (BERZELIUS
Sodium 139	Antimony 9	and)
Strontium139	WEEREN, J.	Lead 71
Salphur 114	Beryllium 17	WOLLASTON. (L. J. THE-
Tin140	WELESKY, P.	NARD and)
Zinc139	Cobalt 48	Iron 65
Thorium120	WENZEL.	WREDE, F. VON
Tin122, 140	Silver106	Carbon 32
TISSIER, C.	WERTHEIM, J.	
Aluminium 8	Uranium130	
Titanium 123	WERTHER, H.	Yttrium133
TROOST, L.	Thallium119	
Lithium74, 75	WILDENSTEIN, R.	
Tungsten125	Chromium 45	ZETTNOW, E.
Turner, E.	WING, C. H.	Tungsten 127
Barium 14	Cerium 36	Zinc134, 139
Chlorine 38	WINKLER, C.	Zirconium136
Lead 71	Cobalt 48 Indium 60 Nickel 88	Zschiesche, H.
Manganese 80	Indium 60	Didymium 52
Mercury 82	Nickei 88	Lanthanium 69





. • .

SMITHSONIAN MISCELLANEOUS COLLECT

CONSTANTS OF NATU

PART V.

Á RECALCULATION

-0F-

THE ATOMIC WEIGH

BY

FRANK WIGGLESWORTH CLARKE, S. B.,

Professor of Chemistry and Physics in the University of Cincir



WASHINGTON: SMITHSONIAN INSTITUTION. 1882.



ADVERTISEMENT.

The present publication is one of a series devoted to the discussion and more precise determination of various "Constants of Nature;" and forms the Fifth contribution to that subject published by this Institution.

The First number of the series, embracing tables of "Specific Gravities" and of Melting and Boiling Points of Bodies, prepared by the same author, Prof. F. W. Clarke, was published in 1873. The Fourth part of the series, comprising a complete digest of the various "Atomic Weight" determinations of the chemical elements published since 1814, commencing with the well-known "Table of Equivalents" by Wollaston, (given in the Philosophical Transactions for that year,) compiled by Mr. George F. Becker, was published by the Institution in 1880. The present work which may be regarded as practically supplementary to that digest, (or perhaps rather as the memoir to which that digest is introductory,) comprises a very full discussion and re-calculation of the "Atomic Weights" from all the existing data, and the assignment of the most probable value to each of the elements.

The manuscript of the work was presented to the Institution in its completed form by Prof. F. W. Clarke, the cost of publication only being at the expense of the Smithsonian fund.

SPENCER F. BAIRD, Secretary of Smithsonian Institution.

WASHINGTON, January, 1882.




TABLE OF CONTENTS.

troduction	
rmulæ for the Calculation of Probable Error	
. Oxygen	
2. Silver, Potassium, Sodium, Chlorine, Bromine, Iodine, and Sulphu	
3. Nitrogen	
4. Carbon	
5. Barium	
6. Strontium	
7. Calcium	
8. Lead	
9. Fluorine	
IO. Phosphorus	
II. Boron	
12. Silicon	
13. Lithium	
14. Rubidium	
15. Cæsium	
16. Thallium	
17. Glucinum	
18. Magnesium	
19. Zinc	
20. Cadmium	
21. Mercury	
22. Chromium	
23. Manganese	
24. Iron	
25. Copper	
26. Molybdenum	
27. Tungsten	
28. Uranium	
29. Aluminum	
o Cold	
Nickel and Cobalt	
Selenium	
. TallarinM	
Vanadium	
A	
Bismuth	
DISTRICT	

CONTENTS.

	PAGE.
38. Tin	204
39. Titanium	207
40. Zirconium	212
41. Thorium	214
42. Gallium	218
43. Indium	219
44. Cerium	220
45. Lanthanum	229
46. Didymium	236
47. The Yttrium Group. Scandium, Yttrium, Ytterbium, Erbium, Ter-	
bium, Phillipium, Decipium, Thulium, Samarium, etc.	240
48. Columbium (Niobium)	247
49. Tantalum	248
50. Platinum	249
51. Osmium	254
52. Iridium	255
53. Palladium	256
54. Rhodium	258
55. Ruthenium	259
Appendix	261

INTRODUCTION.

In the autumn of 1877 the writer began collecting data relative to the determinations of atomic weights, with the purpose of preparing a complete resumé of the entire subject, and of recalculating all the estimations. The work was fairly under way, the material was collected and partly discussed, when I received from the Smithsonian Institution a manuscript by Professor George F. Becker, entitled "Atomic Weight Determinations: a Digest of the Investigations Published since 1814." This manuscript, which has lately been issued as Part IV of the "Constants of Nature," covered much of the ground contemplated in my own undertaking. It brought together all the evidence, presenting it clearly and thoroughly in compact form; in short, that portion of the task could not well be improved ¹Pon. Accordingly, I decided to limit my own labors to a critical recalculation of the data; to combine all the figures ^uPon a common mathematical basis, and to omit everything Which could as well be found in Professor Becker's "Digest."

At the very beginning of my work certain questions confronted me. Should I treat the investigations of different individuals separately, or should I combine similar data together in a manner irrespective of persons? For example, ought I, in estimating the atomic weight of silver, to take Stas' work by itself, Marignac's work by itself, and so on, and then average the results together; or should I rather combine all series of figures relating to the composition of Potassium chlorate into one mean value, and all the data concerning the composition of silver chloride into another mean, and, finally, compute from such general means the constant sought to be established? The latter plan was finally adopted; in fact, it was rendered necessary by the method of least squares, which method was alone adequate to supply me with good processes for calculation.

The mode of discussion and combination of results was briefly as follows. The formulæ employed are given in another chapter. I began with the ratio between oxygen and hydrogen; in other words, with the atomic weight of oxygen referred to hydrogen as unity. Each series of experiments was taken by itself, its arithmetical mean was found, and the probable error of that mean was computed. Then the several means were combined according to the appropriate formula, each receiving a weight dependent upon its probable error. The general mean thus established was taken as the most probable value for the atomic weight of oxygen, and, at the same time, its probable error was mathematically asssigned.

Next in order came a group of elements which were best discussed together, namely, silver, chlorine, potassium, sodium, bromine, iodine, and sulphur. For these elements there were data from thirteen experimenters. All similar figures were first reduced to common standards, and then the means of individual series were combined into general Thus all the data were condensed into twenty ratios, from which several independent values for the atomic weight of each element could be computed. The probable errors of these values, however, all involved the probable error of the atomic weight of oxygen, and were, therefore, higher than they would have been had the latter element not entered into consideration. Here, then, we have suggested a chief peculiarity of this whole revision. atomic weight of each element involves the probable errors of all the other elements to which it is directly or indirectly referred. Accordingly, an atomic weight determined by reference to elements whose atomic weights have been defectively ascertained will receive a high probable error, and its weight, when combined with other values, will be relatively For example, an atomic weight ascertained by direct comparison with hydrogen will, other things being equal. have a lower probable error than one which is referred to hydrogen through the intervention of oxygen; and a metal whose equivalent involves only the probable error of oxygen will be more exactly known than one which depends upon the greater errors of silver and chlorine. These points will appear more clearly evident in the subsequent actual discussions.

But although the discussion of atomic weights is ostensibly mathematical, it cannot be purely so. Chemical considerations are necessarily involved at every turn. In assigning weights to mean values I have been, for the most part, rigidly guided by mathematical rules; but in some cases I have been compelled to reject altogether series of data which were mathematically excellent, but chemically worthless because of constant errors. In certain instances there were grave doubts as to whether particular figures should be included or rejected in the calculation of means; there having been legitimate reasons for either procedure. Probably many chemists would differ with me upon such points of judgment. In fact, it is doubtful whether any two chemists, working independently, would handle all the data in precisely the same way, or combine them so as to produce exactly the same final results. Neither would any two mathematicians follow identical rules or reach identical conclusions. In calculating the atomic weight of any element those values are assigned to other elements which have been determined in previous chapters. Hence a variation in the order of discussion might lead to slight differences in the final results.

As a matter of course the data herein combined are of very unequal value. In many series of experiments the weighings have been reduced to a vacuum standard; but in most cases chemists have neglected this correction altogether. In a majority of instances the errors thus introduced are slight; nevertheless they exist, and interfere more or less with all attempts at a theoretical consideration of the results. For example, they affect seriously the investigation of Prout's hypothesis, and are often great enough to account for seeming exceptions to it. Such questions as these will be considered in the appendix.

Another serious source of error affecting many of the re-

sults was not discovered until recently. A large number of computations had been actually finished, involving, among other things, the greater part of Stas' work, when Dumas published his investigation upon the occlusion of oxygen by silver. Here it was shown that a very great number of atomic weight determinations must have been vitiated by constant errors, which, though constant for each series, were probably of different magnitude in different series, and, therefore, could not be systematically corrected for. At the time of the announcement of this discovery of Dumas my work was so far under way that I thought it best to complete my discussion without reference to it, and then to study its influence in the appendix. In the chapter upon aluminum, however, it will be noted that Mallet eliminated this error in great part from his experimental results.

Necessarily, this work omits many details relative to experimental methods, and particulars as to the arrangements of special forms of apparatus. For such details original memoirs must be consulted. Their inclusion here would have rendered the work unwarrantably bulky. There is such a thing as over-exhaustiveness of treatment, which is equally objectionable with under-thoroughness.

Of course, none of the results reached in this revision can be considered as final. Every one of them is liable to repeated corrections. To my mind the real value of the work, great or little, lies in another direction. have been brought together and reduced to common standards, and for each series of figures the probable error has Thus far, however much my methods been determined. of combination may be criticized, I feel that my labors will The ground is now cleared, in a measure, have been useful. for future experimenters; it is possible to see more distinctly what remains to be done; some clues are furnished as to the relative merits of different series of results. I hope to be able, from time to time, as new determinations are published, to continue the task here begun, and perhaps, also, to add, in the near future, some data of my own establishing.

In addition to the usual periodicals the following works

- have been freely used by me in the preparation of this volume:
- BERZELIUS, J. J. Lehrbuch der Chemie. 5 Auflage. Dritter Band. SS. 1147-1231. 1845.
- Van Geuns, W. A. J. Præve eener Geschiedenis van de Æquivalentgetallen der Scheikundige Grondstoffen en van hare Soortelijke Gewigten in Gasvorm, voornamelijk in Betrekking tot de vier Grondstoffen der Bewerktuigde Natuur. Amsterdam, 1853.
- MULDER, E. Historisch-Kritisch Overzigt van de Bepalingen der Æquivalent-Gewigten van 13 Eenvoudige Ligchamen. Utrecht, 1853.
- MULDER, L. Historisch-Kritisch Overzigt van de Bepalingen der Æquivalent-Gewigten van 24 Metalen. Utrecht, 1853.
- OUDEMANS, A. C., Jr. Historisch-Kritisch Overzigt van de Bepaling der Æquivalent-Gewigten van Twee en Twintig Metalen. Leiden, 1853.
- STAS, J. S. Untersuchungen über die Gesetze der Chemischen Proportionen über die Atomgewichte und ihre gegenseitigen Verhältnisse. Uebersetzt von Dr. L. Aronstein. Leipzig, 1867.

The four Dutch monographs above cited are especially valuable. They represent a revision of all atomic weight data down to 1853, as divided between four writers.

FORMULÆ FOR THE CALCULATION OF PROBABLE ERROR.

Although the ordinary formula for the probable error of an arithmetical mean is familiar to all physicists, it is perhaps best to reproduce it here, as follows:

(1.)
$$\epsilon = \pm .6745 \sqrt{\frac{S}{n (n-1)}}$$

Here n represents the number of observations or experiments in the series, while S is the sum of the variations of the individual results from the mean.

In combining several arithmetical means, representing several series, into one general mean each receives a weight indicated by its probable error; greater as the latter becomes less, and *vice versa*. Let A, B, C, etc., be such mean results, and a, b, c, their probable errors respectively. Then the general mean is determined by this formula:

(2.)
$$M = \frac{\frac{A}{a^3} + \frac{B}{b^3} + \frac{C}{c^2}}{\frac{1}{a^3} + \frac{1}{b^3} + \frac{1}{c^2}} \dots$$

For the probable error of this general mean we have:

(3.)
$$m = \frac{1}{\sqrt{\frac{1}{a^2} + \frac{1}{b^2} + \frac{1}{c^2}}} \dots$$

In the calculation of atomic and molecular weights the following formulæ have been employed. For assistance in connection with them my thanks are due to Professors H. T. Eddy and E. W. Hyde of the University of Cincinnati.

Using, as before, capital letters to represent known quantities and small letters for their probable errors respectively,

we have for the sum or difference of two quantities, A and B:

$$\epsilon = \sqrt{a^1 + b^2}$$

For the product of A multiplied by B the probable error is

(5.)
$$\epsilon = \sqrt{(Ab)^2 + (Ba)^2}$$

For the product of three quantities, ABC:

(6.)
$$\epsilon = \sqrt{(BCa)^2 + (ACb)^2 + (ABc)^2}$$

For a quotient, $\frac{B}{A}$, the probable error becomes

Given a proportion, A:B::C:x, the probable error of the fourth term is as follows:

(8.)
$$\epsilon = \frac{\sqrt{\left(\frac{BCa}{A}\right)^2 + (Cb)^2 + (Bc)^2}}{A}$$

This formula is used in nearly every atomic weight calculation, and is, therefore, exceptionally important. Rarely a more complicated case arises in a proportion of this kind:

$$A:B::C+x:D+x$$

In this proportion the unknown quantity occurs in two terms. Its probable error is found by this expression, and is always large:

(9.)
$$\epsilon = \sqrt{\frac{(C-D)^2}{(A-B)^4}} \left(B^2 a^2 + A^2 b^2 \right) + \frac{B^2 c^2 + A^2 d^2}{(A-B)^2}$$

When several independent values have been calculated for an atomic weight they are treated like means, and combined according to formulæ (2) and (3.) Each final result

is, therefore, to be regarded as the general mean of all reliable determinations. This method of combination may not be the best one theoretically possible, but it seemed to be the only one practically available. The data are too imperfect to warrant the use of much more elaborate processes of discussion.

		·		
•				
,				
	·			
	•			

RECALCULATION OF THE ATOMIC WEIGHTS.

OXYGEN.

The ratio between oxygen and hydrogen is the foundation upon which the entire system of atomic weights depends. Hence, the accuracy of its determination has, from the beginning, been recognized as of extreme importance. A trifling error here may become cumulative when repeated through a moderate series of other ratios.

Leaving out of account the earliest researches, which have now only a historical value, we find that three methods have been employed for fixing this important constant. First, the synthesis of water, effected by passing hydrogen gas over red hot oxide of copper. Secondly, the exact determination of the relative density of the two gases. Thirdly, by weighing the quantity of water formed upon the direct union of a known volume of hydrogen with oxygen.

The first of these methods has been employed in three leading investigations, namely, by Dulong and Berzelius,* by Dumas, and by Erdmann and Marchand. The essential features of the method are in all cases the same. Hydrogen gas is passed over heated oxide of copper, and the water thus formed is collected and weighed. From this weight and the loss of weight which the oxide undergoes, the exact composition of water is readily calculated. Dulong and Berzelius made but three experiments, with the following results for the percentages of oxygen and hydrogen in in water:

0.	H.
88.942	11.058
88.809	11.191
88.954	11.046

Thomson's Annals of Philosophy, July, 1821, p. 50.

These figures, rather roughly determined, and by no means exact enough to meet the requirements of modern science, give a mean value of 16.021 for the atomic weight of oxygen. As the weighings were not reduced to a vacuum, this correction was afterwards applied by Clark,* who showed that these syntheses really make O = 15.894; or, in Berzelian terms, if O = 100, H = 12.583.

In 1842 Dumast published his elaborate investigation upon the composition of water. The first point was to get pure hydrogen. This gas, evolved from zinc and sulphuric acid, might contain oxides of nitrogen, sulphur dioxide, hydrosulphuric acid, and arsenic hydride. These impurities were removed in a series of wash bottles; the H₂S by a solution of lead nitrate, the HAs by silver sulphate, and the others by caustic potash. Finally, the gas was dried by passing through sulphuric acid, or, in some of the experiments, over phosphorus pentoxide. The copper oxide was thoroughly dried, and the bulb containing it was weighed. By a current of dry hydrogen all the air was expelled from the apparatus, and then, for ten or twelve hours, the oxide of copper was heated to dull redness in a constant stream of the gas. The reduced copper was allowed to cool in an atmosphere of hydrogen. The weighings were made with the bulbs exhausted of air. The following table gives the results:

Column A contains the symbol of the drying substance. B gives the weight of the bulb and copper oxide. C, the weight of bulb and reduced copper. D, the weight of the vessel used for collecting the water. E, the same, plus the water. F, the weight of oxygen. G, the weight of water formed. H, the crude equivalent of H when O = 10,000. I, the equivalent of H, corrected for the air contained in the sulphuric acid employed. This correction is not explained, and seems to be questionable.

^{*} Philosophical Magazine, 3d series, 20, 341.

[†] Compt. Rend., 14, 537.

									0	X	Y	} E	N.	•									
	ï	1240.6	1248.0	1247.3	1249.0	1254.6	1255.0	1253.3	1249.0	1255.1	1248.9	1249.0	1250.8	1254.8	1256.2	1252.2	1249.1	1255.1	1254.7	1248.0		1251.5	
	H.	1250.5	1249.0	1248.1	1250.6	1256.2	1256.3	1254.6	1250.0	1258.3	1250.4	1251.2	1253.3	1257.7	1258.1	1255.8	1250.6	1257.3	1257.5	1248.8		1253.3	1
	ij	14.827	22.905	23.053	50.50	85.960	49.047	39.178	51.623	67.586	58.320	59.078	67.282	66.869	58.360	63.577	41.390	38.458	36.175	34.677		Means	_
\	8 4	13.179	20.362	20.495	57.004	76.364	43.571	34.811	45.887	60.031	51.838	52.508	59.789	62.090	51.838	56.483	36.789	34.162	32.133	30.827	-		
	ri	495.634	511.132	462.764	948.323	973.291	916.206	878.482	876.244	890.246	799.417	933.910	998.700	752.273	799.455	1128.319	920.030	926.275	924.837	912.539			
	D.	480.807	488.227	439.711	884.190	887.331	867.159	839.304	824.624	822.660	741.095	874.832	931.487	682.374	741.097	1064.762	878.640	887.817	888.662	877.862			
	Ü	278.806	324.186	296.175	568.825	728.182	490.155	627.104	566.738	844.612	590.487	535.137	613.492	598.765	590.487	881.362	719.563	720.000	727.632	716.825	_		
	B.	291.985	344.548	316.671	625.829	804.546	533.726	661.915	612.625	904.643	642.325	587.645	673.280	660.855	642.325	937.845	756.352	754.162	759.762	747.652		•	
	¥	H, SO,		3	P.O.	H, SO,			P,0,		H,SO,	P.O.	***************************************	II,SO,		***************************************	P ₂ O ₅		***************************************	***************************************			

In the sum total of these nineteen experiments, 840.161 grammes of oxygen form 945.439 grammes of water. This gives, in percentages, for the composition of water, oxygen 88.864; hydrogen, 11.136. Hence the atomic weight of oxygen, calculated in mass, is 15.9608. In the following column the values are given as deduced from the individual data given under the headings F and G:

```
15.994
16.014
16.024
15.992
15.916
15.916
15.943
16.000
15.892
15.995
15.984
15.958
15.902
15.987
15.926
15.992
15.904
15.900
16.015
```

Mean, 15.9607, with a probable error of \pm .0070.

In calculating the above column several discrepancies were noted, probably due to misprints in the original memoir. On comparing columns B and C with F, or D and E with G, these anomalies chiefly appear. They were detected and carefully considered in the course of my own calculations; and, I believe, eliminated from the final result.

The paper by Erdmann and Marchand* followed closely after that of Dumas. The method of research was essentially the same as that of the latter chemist, varying only in points of comparatively unimportant detail. The results are given in two series, in one of which the weighings were

^{*} Journ. f. Prakt. Chem., 1842, bd. 26, s. 461.

not actually made in vacuo, but were, nevertheless, reduced to a vacuum standard. The second series represents actual vacuum weighings. The quantity of water formed in each experiment, was from 41.664 to 95.612 grammes. I give below only the percentages of oxygen and hydrogen in water as deduced from Erdmann and Marchand's data:

First Series.

O.	H.
88.836	11.164
88.821	11.179
88.874	11.126
88.868	11.132

Second Series.

Ο.	H.
88.887	11.113
88.898	11.102
88.895	11.105
88.899	11.101

Hence, the atomic weight of oxygen is, as follows:

First Series.	Second Series.
15.915	15.997
15.891	16.015
15.976	16.010
15.966	16.016
	
Mean, 15.9369, ± .0138	Mean, 16.0005, ± .0030

The effect of discussing these two series separately is somewhat startling. It gives to the four experiments in Erdmann and Marchand's second group a weight vastly greater than their other four and Dumas' nineteen taken together. For so great a superiority as this there is no adequate reason; and it is highly probable that it is due almost entirely to fortunate coincidences, rather than to greater accuracy of work. We will, therefore, treat Erdmann and Marchand's experiments as one series, giving all equal weight, and then combine them with the results obtained by Dumas. We now have—

By DumasBy Erdmann and Marchand	
General mean	0 = 15.0642 + .0060

In discussing the relative density of oxygen and hydrogen gases we need only consider the more modern researches of Dumas and Boussingault, and of Regnault. As the older work has some historical value, I may in passing just cite its results. For the density of hydrogen we have .0769, Lavoisier; .0693, Thomson; .092, Cavendish; .0732, Biot and Arago; .0688, Dulong and Berzelius. For oxygen there are the following determinations: 1.087, Fourcroy, Vauquelin, and Séguin; 1.103, Kirwan; 1.128, Davy; 1.088, Allen and Pepys; 1.1036, Biot and Arago; 1.1117, Thomson; 1.1056, De Saussure; 1.1026, Dulong and Berzelius; 1.106, Buff; 1.1052, Wrede.*

In 1841 Dumas and Boussingault† published their determinations of gaseous densities. For hydrogen they obtained values ranging from .0691 to .0695; but beyond this mere statement they give no details. For oxygen three determinations were made, with the following results:

If we take the two extreme values given above for hydrogen, and regard them as the entire series, they give us a mean of .0693, $\pm .00013$.

This mean hydrogen value, combined with the mean oxygen value, gives for the atomic weight of the latter element the number 15.9538, $\pm .031$.

Regnault's researches, published four years later,‡ were of

^{*} For Wrede's work, see Berzelius' Jahresbericht for 1843. For Dulong and Berzelius, see the paper already cited. All the other determinations are taken from Gmelin's Handbook, Cavendish edition, v. 1, p. 279.

[†] Compt. Rend., 12, 1005. Compare also with Dumas, Compt. Rend., 14, 537.

[‡] Compt. Rend., 20, 975.

OXYGEN. 7

1

a more satisfactory kind. Indeed, they are among the classics of physical science; and probably approach as near to absolute accuracy as is possible for experiment.

For hydrogen three determinations of density gave the following results:

For oxygen four determinations were made, but in the first one the gas was contaminated by traces of hydrogen, and the value obtained, 1.10525; was, therefore, rejected by Regnault as too low. The other three are as follows:

Now, combining the hydrogen and oxygen series, we have for the atomic weight of oxygen, 15.9628, ± .0044.*

Upon combining the result of Regnault's work with that from Dumas and Boussingault's we get the following value:

This result, it will be seen, agrees remarkably well with that obtained in the experiments upon the synthesis of water.

^{*}Since these computations were made, Professor John Le Conte has called my attention to the existence of slight numerical errors in Regnault's own reductions. As corrected by Le Conte, Regnault's figures give 1.105612 for the density of oxygen, and 0.069269 for that of hydrogen. Hence the atomic weight of O becomes 15.9611, instead of 15.9628. The difference is slight, but still it ought not to be ignored. All the computations in the body of this work, having been finished before I received Professor Le Conte's figures, must stand, nevertheless, as they are. For further details Le Conte refers to Phil. Mag., (4,) 27, p. 29, 1864; and also to the Smithsonian Report for 1878, p. 428.

The third method indicated at the beginning of this discussion has been recently employed in part by J. Thomsen* of Copenhagen. Unfortunately this chemist has not published the details of his work, but only the end results. These serve to confirm the values for oxygen fixed by other methods, but they cannot well be included in the systematic discussion. Partly by the oxidation of hydrogen over heated copper oxide, and partly by its direct union with oxygen, Thomsen finds that at the latitude of Copenhagen, and at sea level, one litre of dry hydrogen at 0° and 760 mm. pressure will form .8041 gramme of water. According to Regnault, at this latitude, level, temperature, and pressure, a litre of hydrogen weighs .08954 gramme. these data, O = 15.9605. It will be seen at once that Thomsen's work depends in great part upon that of Regnault, and yet that it affords an admirable reinforcement of the latter.

It is now plain, in conclusion, that all the different lines of research point to an atomic weight for oxygen a little below 16.00. Five distinct investigations confirm each other wonderfully. Upon combining the values obtained by the two chief methods we get the following final results:

```
From synthesis of water...... O = 15.9642, \pm .0060
From gaseous densities..... O = 15.9627, \pm .0043
```

In the general mean the atomic weight of oxygen becomes 15.9633, with a probable error of \pm .0035.†

^{*} Ber. d. Deutsch. Chem. Gesellschaft, 1870, s. 928.

[†] Le Conte's correction of Regnault's figures introduced here would make 0 == 15.9622, instead of 15.9633. Difference, .0011.

SILVER, POTASSIUM, SODIUM, CHLORINE, BROMINE, IODINE, AND SULPHUR.

The atomic weights of these seven elements depend upon each other to so great an extent that they can hardly be considered independently. Indeed, chlorine, potassium, and silver have always been mutually determined. ratio between silver and chlorine, the ratio between silver and potassium chloride, and the composition of potassium chlorate, these three atomic weights were first accurately fixed. Similar ratios, more recently worked out by Stas and others, have rendered it desirable to include bromine, iodine, sulphur, and sodium in the same general discussion.

Several methods of determination will be left altogether out of account. For example, in 1842 Marignac* sought to fix the atomic weight of chlorine by estimating the quantity of water formed when hydrochloric acid gas is passed over heated oxide of copper. His results were wholly inaccurate, and need no further mention here. A little later Laurent† redetermined the same constant from the analysis of a chlorinated derivative of naphthalene. This method did not admit of extreme accuracy, and it presupposed a knowledge of the atomic weight of carbon; hence it may be properly disregarded. Maumené's analyses of the oxalate and acetate of silver gave good results for the atomic weight of that metal; but they also depend for their value upon our knowledge of carbon, and will, therefore, be discussed further on with reference to that element.

Let us now consider the ratios upon which we must rely for ascertaining the atomic weights of the seven elements in question. After we have properly arranged our data we may then discuss their meaning. First in order we may

^{*}Compt. Rend., 14, 570. Also, Journ. f. Prakt. Chem., 26, 304. †Compt. Rend., 14, 456. Journ. f. Prakt. Chem., 26, 307.

[†] Ann. d. Chim. et d. Phys., (3,) 18, 41. 1846.

conveniently take up the percentage of potassium chloride obtainable from the chlorate.

The first reliable series of experiments to determine this percentage was made by Berzelius.* All the earlier estimations were vitiated by the fact that when potassium chlorate is ignited under ordinary circumstances a little solid material is mechanically carried away with the oxygen gas. Minute portions of the substance may even be actually volatilized. These sources of loss were avoided by Berzelius, who devised means for collecting and weighing this trace of potassium chloride. All the successors of Berzelius in this work have benefitted by his example; although for the methods by which loss has been prevented we must refer to the original papers of the several investigators. In short, then, Berzelius ignited potassium chlorate, and determined the percentage of chloride which remained. Four experiments gave the following results:

```
60.854
60.850
60.850
60.851
```

Mean, 60.851, with a probable error of ± .0006

The next series was made by Penny,† in England, who worked after a somewhat different method. He treated potassium chlorate with strong hydrochloric acid in a weighed flask, evaporated to dryness over a sand bath, and then found the weight of the chloride thus obtained. His results are as follows, in six trials:

^{*} Poggend. Annalen, 1826, bd. 8, s. 1.

[†] Phil. Transactions, 1839, p. 20.

In 1842 Pelouze* made three estimations by the ignition of the chlorate, with these results:

60.843 60.857 60.830 Mean, 60.843, ± .0053

Marignac, in 1842,† worked with several different recrystallizations of the commercial chlorate. He ignited the salt, with the usual precautions for collecting the material carried off mechanically, and also examined the gas which was evolved. He found that the oxygen from 50 grammes of chlorate contained chlorine enough to form .003 gramme of silver chloride. Here are the percentages found by Marignac:

In chlorate once crystallized	60.845
In chlorate once crystallized	60.835
In chlorate twice crystallized	60.833
In chlorate twice crystallized	60.844
In chlorate three times crystallized	60.839
In chlorate four times crystallized	60.839

Mean, 60.8392, $\pm .0013$

In the same paper Marignac describes a similar series of experiments made upon potassium perchlorate, KClO₄. In three experiments it was found that the salt was not quite free from chlorate, and in three more it contained traces of iron. A single determination upon very pure material gave 46.187 per cent. of oxygen and 53.813 of residue.

In 1845 two series of experiments were published by Gerhardt.† The first, made in the usual way, gave these results:

60.871 60.881 60.875 Mean, 60.8757, ± .0020

^{*} Compt. Rend., 15, 959.

[†] Ann. d. Chem. u. Pharm., bd. 44, s. 18.

[‡] Compt. Rend., 21, 1280.

In the second series the oxygen was passed through a weighed tube containing moist cotton, and another filled with pumice stone and sulphuric acid. Particles were thus collected which in the earlier series escaped. From these experiments we get—

These last results were afterwards sharply criticized by Marignac,* and their value seriously questioned.

The next series, in order of time, is due to Maumené.† This chemist supposed that particles of chlorate, mechanically carried away, might continue to exist as chlorate, undecomposed; and hence that all previous series of experiments might give too high a value to the residual chloride. In his determinations, therefore, the ignition tube, after expulsion of the oxygen, was uniformly heated in all its parts. Here are his percentages of residue:

60.788 60.790 60.793 60.791 60.785 60.795 60.795 Mean, 60.791, ± .0009

The question which most naturally arises in connection with these results is, whether portions of chloride may not have been volatilized, and so lost.

Closely following Maumené's paper there is a short note by Faget,‡ giving certain mean results. According to this chemist, when potassium chlorate is ignited slowly, we get

^{*} Supp. Bibl. Univ. de Genéve, Vol. I.

[†] Ann. d. Chim. et d. Phys., (3,) 18, 71. 1846.

[‡] Ann. d. Chim. et d. Phys., (3,) 18, 80. 1846.

60.847 per cent. of residue. When the ignition is rapid, we get 60.942. As no detailed experiments are given, these figures can have no part in our discussion.

Last of all we have two series determined by Stas.* In the first series we have the results obtained by igniting the chlorate. In the second series the chlorate was reduced by strong hydrochloric acid, after the method followed by Penny:

```
First Series.
60.8380
60.8395
60.8440
60.8473
60.8450

Mean, 60.84276, ± .0012

Second Series.
60.850
60.853
60.844

Mean, 60.849, ± .0017
```

In these experiments every conceivable precaution was taken to avoid error and ensure accuracy. All weighings were reduced to a vacuum standard; from 70 to 142 grammes of chlorate were used in each experiment; and the chlorine carried away with the oxygen in the first series was absorbed by finely divided silver and estimated. It is difficult to see how any error could have crept in.

Now, to combine these different series of experiments.

```
Penny,
           ----- 60.8225, ± .0014
Pelouze.
        46
             --- 60.843, ± .0053
        66
Marignac,
           ----- 60.8392, ± .0013
Gerhardt, 1st
           ---- 60.8757, ± .0020
     2d
             ----- 60.9487, ± .0011
Maumené.
              _____ 60.791, ± .0009
Stas,
     ıst
             ----- 60.8428, ± .0012
     2d
   General mean, from all nine series, representing forty
      experiments _____ 60.846, ± .00038
```

^{*} See Aronstein's Translation, p. 249.

This value is exactly that which Stas deduced from both of his own series combined, and gives great emphasis to his wonderfully accurate work. It also finely illustrates the compensation of errors which occurs in combining the figures of different experimenters.

Similar analyses of silver chlorate have been made by Marignac and by Stas. Marignac's figures I have not been able to find,* and Stas gives but two experiments. The following are his percentages of oxygen in silver chlorate:†

For the direct ratio between silver and chlorine there are seven available series of experiments. Here, as in many other ratios, the first reliable work was done by Berzelius.1

He made three estimations, using each time twenty grammes of pure silver. This was dissolved in nitric acid. In the first experiment the silver chloride was precipitated and collected on a filter. In the second and third experiments the solution was mixed with hydrochloric acid in a flask, evaporated to dryness, and the residue then fused and weighed without transfer. One hundred parts of silver formed of chloride:

^{*}Since all the calculations were finished I have secured a copy of Marignac's figures. They are as follows: The third column gives the percentage of O in AgClO₂.

24.510 gr	m. Ag(ClO, gave	18.3616 AgCl.	25.103
25.809	44	**	19.3345 "	25.086
30.306	"	44	22.7072 "	25.074
28.358	46	• •	21.2453 "	25.082
28.287	"	46	21.1833 "	25.113
57.170	"	66	42.8366 "	25.072

Mean, 25.088, ± .0044

The introduction of these figures into the subsequent calculations could not produce any appreciable result. They would practically vanish from the general mean. However, they serve here as confirmation of Stas' work.

[†] Aronstein's Translation, p. 214.

[†] Thomson's Annals of Philosophy, 1820, v. 15, p. 89.

Turner's work* closely resembles that of Berzelius. Silver was dissolved in nitric acid and precipitated as chloride. In experiments one, two, and three the mixture was evaporated and the residue fused. In experiment four the chloride was collected on a filter. A fifth experiment was made, but has been rejected as worthless.

The results were as follows: In a third column I put the quantity of AgCl proportional to 100 parts of Ag.

```
28.407 grains Ag gave 37.737 AgCl. 132.844
41.917 " 55.678 " 132.829
40.006 " 53.143 " 132.837
30.922 " 41.070 " 132.818

Mean, 132.832, ± .0038
```

The same general method of dissolving silver in nitric acid, precipitating, evaporating, and fusing without transfer of material was also adopted by Penny.† His results for 100 parts of silver are as follows, in parts of chloride:

```
132.836
132.840
132.830
132.840
132.840
132.830
132.838

Mean, 132.8363, ± .0012
```

In 1842 Marignac‡ found that 100 parts of silver formed 132.74 of chloride, but gave no available details. Later,||

^{*} Phil. Transactions, 1829, 291.

[†] Phil. Transactions, 1839, 28.

¹ Ann. Chem. Pharm., 44, 21.

See Berzelius' Lehrbuch, 5th Ed., Vol. 3, pp. 1192, 1193.

in another series of determinations, he is more explicit, and gives the following data: The weighings were reduced to a vacuum standard.

79.853 gr	m. Ag ga	ve 106.080 A	AgCl.	Ratio, 132.844
69.905	"	92.864	66	132.843
64.905	66	86.210	66	132.825
92.362	"	122.693	"	132.839
99.653	44	132.383	**	132.844
				Mean, 132.839, ± .0024

The above series all represent the synthesis of silver chloride. Maumené* made analyses of the compound, reducing it to metal in a current of hydrogen. His experiments make 100 parts of silver equivalent to chloride:

```
132.734
132.754
132.724
132.729
132.741
Mean, 132.7364, ± .0077
```

By Dumast we have the following estimations:

Finally, there are seven determinations by Stas,‡ made with his usual accuracy and with every precaution against error. In the first, second, and third, silver was heated in chlorine gas, and the synthesis of silver chloride thus effected directly. In the fourth and fifth silver was dissolved in nitric acid, and the chloride thrown down by passing hydrochloric acid gas over the surface of the solution. The whole was then evaporated in the same vessel, and the chloride fused, first in an atmosphere of hydrochloric acid,

^{*} Ann. d. Chim. et d. Phys., (3,) 18, 49. 1846.

[†] Ann. Chem. Pharm., 113, 21. 1860.

[‡] Aronstein's Translation, p. 171.

and then in a stream of air. The sixth synthesis was similar to these, only the nitric solution was precipitated by hydrochloric acid in slight excess, and the chloride thrown down was washed by repeated decantation. All the decanted liquids were afterwards evaporated to dryness, and the trace of chloride thus recovered was estimated in addition to the main mass. The latter was fused in an atmosphere of HCl. The seventh experiment was like the sixth, only ammonium chloride was used instead of hydrochloric acid. From 98.3 to 399.7 grammes of silver were used in each experiment, the operations were performed chiefly in the dark, and all weighings were reduced to vacuum. In every case the chloride obtained was beautifully white. The following are the results in chloride for 100 of silver:

```
132.841
132.843
132.849
132.846
132.848
122.8417
Mean, 132.8445, ± .0008
```

We may now combine the means of these seven series, representing in all thirty-three experiments. One hundred parts of silver are equivalent to chlorine, as follows:

Berzelius	32.757, ± .0190
Turner	$32.832, \pm .0038$
Penny	32.8363, ± .0012
Marignac	$32.839, \pm .0024$
Maumené	32.7364, ± .0077
Dumas	$32.8755, \pm .0044$
Stas	$32.8445, \pm .0008$
General mean	32.84I8. + .0006

Here, again, we have a fine example of the evident compensation of errors among different series of experiments. We have also another tribute to the accuracy of Stas, since this general mean varies from the mean of his results only within the limits of his own variations.

The ratio between silver and potassium chloride, or, in other words, the weight of silver in nitric acid solution which can be precipitated by a known weight of KCl, has been fixed by Marignac and by Stas. Marignac,* reducing all weighings to vacuum, obtained these results. In the third column I give the weight of KCl proportional to 100 parts of Ag.

8 grm	. Ag =	3.2626	KCl:		69.067		
_	"	15.001	"		69.050		
	66	15.028	**		69.066		
	46	15.131	"		69.063		
	"	15.216	"		69.063		
	"	17.350	46		69.063		
				Mean	60.062	_	~

Mean, 69.062, $\pm .0017$

Stas' experiments upon this ratio may be divided into two series.† In the first series the silver was slightly impure, but the impurity was of known quantity, and corrections could therefore be applied. In the second series pure silver was employed. The potassium chloride was from several different sources, and in every case was purified with the utmost care. From 10.8 to 32.4 grammes of silver were taken in each experiment, and the weighings were reduced to vacuum. The method of operation was, in brief, as follows: A definite weight of potassium chloride was taken, and the exact quantity of silver necessary, according to Prout's hypothesis, to balance it was also weighed out. metal, with suitable precautions, was dissolved in initric acid, and the solution mixed with that of the chloride. After double decomposition the trifling excess of silver remaining in the liquid was determined by titration with a normal solution of potassium chloride. One hundred parts of silver required the following of KCl:

^{*} See Berzelius' Lehrbuch, 5th edition, Vol. 3, pp. 1192, 1193.

[†] Aronstein's Translation, pp. 250-257.

```
First Series.
       69.105
       69.104
       69.103
       69.104
       69.102
Mean, 69.1036, ± .0003
   Second Series.
       69.105
       69.099
       69.107
       69.103
       69.103
      69.105
       69.104
       69.099
       69.1034
       69.104
      69.103
       69.102
       69.104
       69.104
      69.105
      69.103
       69,101
       60.105
       69.103
Mean, 69.1033, ±.0003
```

Now, combining the three series, with their thirty experiments, we get the following:

General mean	69.1032,	±	.0002
Stas, 1st series		_	•
Marignac			•

The quantity of silver chloride which can be formed from a known weight of potassium chloride has also been determined by Berzelius, Marignac, and Maumené. Berzelius* found that 100 parts of KCl were equivalent to 194.2 of

^{*} Poggend. Annal., 8, 1. 1826.

AgCl; a value which, corrected for weighings in air, becomes 192.32. This experiment will not be included in our discussion.

In 1842 Marignac* published two determinations, with these results from 100 KCl:

192.33 192.34

Mean, corrected for weighing in air, 192.26, ± .003

In 1846 Marignac† published another set of results, as follows. The weighings were reduced to vacuum. The usual ratio is in the third column.

17.034	grm. KCl gave	32.761	AgCļ.		192.327
14.427	66	27.749	"		192.341
15.028	44	28.910	"		192.374
15.131	"	29.102	66		192.334
15.216	66	29.271	**		192.370
				3.5	
	_			Mean	T02 240 - 006

Three estimations of the same ratio were also made by Maumené, 1 as follows:

```
10.700 grm. KCl gave 20.627 AgCl. 192.776
10.5195 " 20.273 " 192.716
8.587 " 16.556 " 192.803

Mean, 192.765, ± .017
```

The three series of ten experiments in all foot up thus:

These figures show clearly that the ratio which they represent is not of very high importance. It might be rejected altogether without impropriety, and is only retained for the

^{*}Ann. Chem. Pharm., 44, 21. 1842. • † Berzelius' Lehrbuch, 5th Ed., Vol. 3, pp. 1192, 1193.

[‡] Ann. d. Chim. et d. Phys., (3,) 18, 41. 1846.

sake of completeness. It will obviously receive but little weight in our final discussion.

In estimating the atomic weight of bromine the earlier experiments of Balard, Berzelius, Liebig, and Löwig may all be rejected. Their results were all far too low, probably because chlorine was present as an impurity in the materials employed. Wallace's determinations, based upon the analysis of arsenic tribromide, are tolerably good, but need not be considered here. In the present state of our knowledge, Wallace's analyses are better fitted for fixing the atomic weight of arsenic, and will, therefore, be discussed with refference to that element.

The ratios with which we now have to deal are closely similar to those involving chlorine. In the first place there are the analyses of silver bromate by Stas.* In two careful experiments he found in this salt the following percentages of oxygen:

There are also four analyses of potassium bromate by Marignac.† The salt was heated, and the percentage loss of oxygen determined. The residual bromide was feebly alkaline. We cannot place much reliance upon this series. The results are as follows:

When silver bromide is heated in chlorine gas, silver chloride is formed. In 1860 Dumast employed this method

^{*}Aronstein's Translation, pp. 200-206.

[†] See E. Mulder's Overzigt, p. 117; or Berzelius' Jahresbericht, 24, 72.

¹ Ann. Chem. Pharm., 113, 20.

for estimating the atomic weight of bromine. His results are as follows: In the third column I give the weight of AgBr equivalent to 100 parts of AgCl.

2.028 gm	n. AgBr gave	1.547	AgCl.	131.092
4.237	"	3.235	46	130.974
5.769	66	4.403	46	131.024
				Mean, 131.030, ± .023

This series is evidently of but little value.

But the two ratios upon which, in connection with Stas' analyses of silver bromate, the atomic weight of bromine chiefly depends are those which connect silver with the latter element directly and silver with potassium bromide.

Marignac,* to effect the synthesis of silver bromide, dissolved the metal in nitric acid, precipitated the solution with potassium bromide, washed, dried, fused, and weighed the product. The following quantities of bromine were found proportional to 100 parts of silver:

74.072 74.055 74.066

Mean, reduced to a vacuum standard, 74.077, ± .003

Much more elaborate determinations of this ratio are due to Stas.† In one experiment a known weight of silver was converted into nitrate, and precipitated in the same vessel by pure hydrobromic acid. The resulting bromide was washed thoroughly, dried, and weighed. In four other estimations the silver was converted into sulphate. Then a known quantity of pure bromine, as nearly as possible the exact amount necessary to precipitate the silver, was transformed into hydrobromic acid. This was added to the dilute solution of the sulphate, and, after precipitation was complete, the minute trace of an excess of silver in the clear supernatant fluid was determined. All weighings were re-

^{*} E. Mulder's Overzigt, p. 116. Berzelius' Jahresbericht, 24, 72.

[†] Aronstein's Translation, pp. 154-170.

duced to a vacuum. From these experiments, taking both series as one, we get the following quantities of bromine corresponding to 100 parts of silver:

	74.0830
	74.0790
	74.0795
	74.0805
	74.0830
Mean,	74.081, ± .0006

Combining this with Marignac's result, 74.077, \pm .003, we get as a general mean the value 74.0809, \pm .0006.*

The ratio between silver and potassium bromide was first accurately determined by Marignac.† I give, with his weighings, the quantity of KBr proportional to 100 parts of Ag:

2.131 g	rm. Ag	= 2.351	KBr.	110.324
2.559	"	2.823	"	110.316
2.447	66	2.700	44	110.339
3.025	"	3.336	"	110.283
3.946	"	4-353	44	110.314
11.569	"	12.763	44	110.321
20.120	66	22.191	"	110.293

Mean, corrected for weighing in air, 110.343, ± .005

Stas,‡ working in essentially the same manner as when he fixed the ratio between potassium chloride and silver, obtained the following results:

^{*0.} W. Huntington, in his paper upon the atomic weight of cadmium, (Amer. Acad. Proc., 1881,) gives three analyses and three syntheses of silver bromide. These give a mean value of Ag: Br:: 100: 74.064. This figure I record here in order that other chemists may not overlook the work of Mr. Huntington, although it came out too late for use in my own calculations.

[†] E. Mulder's Overzigt, p. 116. Berzelius' Jahresbericht, 24, 72.

[‡] Aronstein's Translation, pp. 334-347.

```
110.361

110.360

110.342

110.346

110.338

110.360

110.336

110.344

110.332

110.343

110.357

110.334
```

Mean, 110.3463, ± .0020

Combining this with Marignac's mean result, 110.343, \pm .005, we get a general mean of 110.3459, \pm .0019.

The ratios upon which we must depend for the atomic weight of iodine are exactly parallel to those used for the determination of bromine.

To begin with, the percentage of oxygen in potassium iodate has been determined by Millon.* In three experiments he found:

Millon also estimated the oxygen in silver iodate, getting the following percentages:

The analysis of silver iodate has also been performed with extreme care by Stas.† From 76 to 157 grammes were used

^{*} Ann. d. Chim. et d. Phys., (3,) 9, 400. 1843.

[†] Aronsteins' Translation, pp. 179-200.

in each experiment, the weights being reduced to a vacuum standard. As the salt could not be prepared in an absolutely anhydrous condition, the water expelled in each analysis was accurately estimated and the necessary corrections applied. In two of the experiments the iodate was decomposed by heat, and the oxygen given off was fixed upon a weighed quantity of copper heated to redness. Thus the actual weights, both of the oxygen and the residual iodide, were obtained. In a third experiment the iodate was reduced to iodide by a solution of sulphurous acid, and the oxygen was estimated only by difference. In the three percentages of oxygen given below the result of this analysis comes last. The figures for oxygen are as follows:

```
16.976
16.972
16.9761
Mean, 16.9747, ± .0009
```

This, combined with Millon's series above cited, gives us a general mean of $16.9771, \pm .0009$.

The ratio between silver and potassium iodide seems to have been determined only by Marignac,* and without remarkable accuracy. In five experiments 100 parts of silver were found equivalent to potassium iodide as follows:

```
1.616 grm. Ag = 2.483 KI.
                                    Ratio, 153.651
2.503
        66
                 3.846 "
                                          153.665
          46
                 5.268 "
3.427
                                          152.720
          "
                 3.290 "
2.141
                                          153.667
10.821
                16.642 "
                                          153.794
```

Mean, 153.6994, ± .0178

The synthesis of silver iodide has been effected by both Marignac and Stas. Marignac, in the paper above cited, gives these weighings. In the last column I add the ratio between iodine and 100 parts of silver:

```
      15.000 grm. Ag gave 32.625 AgI.
      117.500

      14.790 " 32.170 " 117.512

      18.545 " 40.339 " 117.519
```

Mean, corrected for weighing in air, 117.5335, \pm .0036

^{*} Berzelius' Lehrbuch, 5th Ed., 3, 1196.

Stas* in his experiments worked after two methods, which gave, however, results concordant with each other and with those of Marignac.

In the first series of experiments Stas converted a known weight of silver into nitrate, and then precipitated with pure hydriodic acid. The iodide thus thrown down was washed, dried, and weighed without transfer. By this method 100 parts of silver were found to require of iodine:

117.529 117.536

Mean, 117.5325, ± .0024

In the second series a complete synthesis of silver iodide from known weights of iodine and metal was performed. The iodine was dissolved in a solution of ammonium sulphite, and thus converted into ammonium iodide. The silver was transformed into sulphate and the two solutions mixed. When the precipitate of silver iodide was completely deposited the supernatant liquid was titrated for the trifling excess of iodine which it always contained. As the two elements were weighed out in the ratio of 127 to 108, while the atomic weight of iodine is probably a little under 127, this excess is easily explained. From these experiments two sets of values were deduced; one from the weights of silver and iodine actually employed, the other from the quantity of iodide of silver collected. From the first set we have of iodine for 100 parts of silver:

117.5390 117.5380 117.5318 117.5430 117.5420 117.5300

Mean, 117.5373, ± .0015

From the weight of silver iodide actually collected we

^{*} Aronstein's Translation, pp. 136, 152.

get as follows. For experiment number three in the above column there is no equivalent here:

```
117.529

117.531

117.539

117.538

117.530

Mean, 117.5334, ± .0014
```

Now, combining these several sets of results, we have the following general mean:

One other comparatively unimportant iodine ratio remains for us to notice. Silver iodide, heated in a stream of chlorine, becomes converted into chloride; and the ratio between these two salts has been thus determined by Berzelius and by Dumas.

From Berzelius* we have the following data: In the third column I give the ratio between AgI and 100 parts of AgCl.

```
5.000 grm. AgI gave 3.062 AgCl. 163.292
12.212 " 7.4755 " 163.360
Mean, 163.326, ± .023
```

Dumas't results were as follows:

```
3.520 grm. AgI gave 2.149 AgCl. 163.793
7.011 " 4.281 " 163.770
Mean, 163.782, ± .008
```

General mean from the combination of both series, 163.733, $\pm .0076$.

We now come to the ratios connecting sulphur with silver

^{*} Ann. d. Chim. et d. Phys., (2,) 40, 430. 1829.

[†] Ann. Chem. Pharm., 113, 28. 1860.

and chlorine. Other ratios have been applied to the determination of the atomic weight of sulphur, but they are hardly applicable here. The earlier results of Berzelius were wholly inaccurate, and his later experiments upon the synthesis of lead sulphate will be used in discussing the atomic weight of lead. Erdmann and Marchand determined the amount of calcium sulphate which could be formed from a known weight of pure Iceland spar; and later they made analyses of cinnabar, in order to fix the value of sulphur by reference to calcium and to mercury. Their results will be applied in this discussion towards ascertaining the atomic weights of the metals just named. For our present purposes only three ratios need be considered.

First in order let us take up the composition of silver sulphide, as directly determined by Dumas, Stas, and Cooke. Dumas'* experiments were made with sulphur which had been thrice distilled and twice crystallized from carbon disulphide. A known weight of silver was heated in a tube in the vapor of the sulphur, the excess of the latter was distilled away in a current of carbon dioxide, and the resulting silver sulphide was weighed.

I subjoin Dumas' weighings, and also the quantity of Ag.S proportional to 100 parts of Ag, as deduced from them:

```
9.9393 grm. Ag = 1.473 S. Ratio, 114.820

9.962 " 1.4755 " " 114.811

30.637 " 4.546 " " 114.838

30.936 " 4.586 " " 114.824

30.720 " 4.554 " . " 114.824
```

Mean, 114.8234, ± .0029

Dumas used from ten to thirty grammes of silver in each experiment. Stas,† however, in his work, employed from sixty to two hundred and fifty grammes at a time. Three of Stas' determinations were made by Dumas' method, while in the other two the sulphur was replaced by pure sulphu-

^{*} Ann. Chem. Pharm., 113, 24. 1860

[†] Aronstein's Translation, p. 179.

retted hydrogen. In all cases the excess of sulphur was expelled by carbon dioxide, purified with scrupulous care. Impurities in the dioxide may cause serious error. The five results come out as follows for 100 parts of silver:

```
114.854

114.853

114.854

114.851

114.849

————

Mean, 114.8522, ± .0007
```

The experiments made by Professor Cooke* with reference to this ratio were only incidental to his elaborate researches upon the atomic weight of antimony. They are interesting. however, for two reasons: they serve to illustrate the volatility of silver, and they represent, not syntheses, but reductions of the sulphide by hydrogen. Cooke gives three series In the first the silver sulphide was long heated of results. to full redness in a current of hydrogen. Highly concordant and at the same time plainly erroneous figures were obtained; the error being eventually traced to the fact that some of the reduced silver, although not heated to its melting point, was actually volatilized and lost. The second series, from reductions at low redness, are decidedly better. In the third series the sulphide was fully reduced below a visible red heat. Rejecting the first series we have from Cooke's figures in the other two the subjoined quantities of sulphide corresponding to 100 parts of silver:

```
7.5411 grm. Ag. S lost .9773 grm. S. Ratio, 114.889
5.0364
                       .6524
                               66
                                              114.882
              66
                                "
                                         66
2.5815
                                              114.886
                       .3345
              66
                                64
                                              114.892
2.6130
                       .3387
                                              114.891
2.5724
                       .3334
                                       Mean, 114.888, ± .0012
1.1357 grm. Ag<sub>2</sub>S lost .1465 S.
                                       Ratio, 114.810
                       .1670 "
1.2936
                                              114.823
                                       Mean, 114.8165, ± .0044
```

^{*} Proc. American Acad. of Arts and Sciences, v. 12. 1877.

Now, combining all four series, we get the following results:

Dumas	
Stas	114.8522, ± .0007
Cooke's 2d	114.888, ± .0012
" 3d	114.8165, ± .0044
General mean	114.8581, ± .0006

Here again we encounter a curious and instructive compensation of errors, and another evidence of the accuracy of Stas.

The percentage of silver in silver sulphate has been determined by Struve and by Stas. Struve* reduced the sulphate by heating in a current of hydrogen, and obtained these results:

```
5.1860 grm. Ag<sub>2</sub>SO<sub>4</sub> gave 3.5910 grm. Ag.
                                                  69.244 per cent.
                 66
                           4.1922
                                                  69.243
 8.6465
                 66
                             5.9858
                                        "
                                                  69.228
                 66
                                       "
                                                             46
                             8.0608
11.6460
                                                  69.215
                 66
                                        "
                                                             "
9.1090
                             6.3045
                                                  69.212
 9.0669
                             6.2778
                                                  69.239
```

Mean, 69.230, ± .004

Stas,† working by essentially the same method, with from 56 to 83 grammes of sulphate at a time, found these percentages:

```
69.200
       69.197
       69.204
       69.209
       69.207
       69.202
Mean, 69.203, ± .0012
```

Combining this mean with that from Struve's series we get a general mean of 69.205, $\pm .0011$.

^{*} Ann. Chem. Pharm., 80, 203. 1851.

[†] Aronstein's Translation, pp. 214-218.

The third and last sulphur ratio with which we have now to deal is one of minor importance. When silver chloride is heated in a current of sulphuretted hydrogen the sulphide is formed. This reaction was applied by Berzelius* to determining the atomic weight of sulphur. He gives the results of four experiments; but the fourth varies so widely from the others that I have rejected it. I have reason to believe that the variation is due, not to error in experiment, but to error in printing; nevertheless, as I am unable to track out the cause of the mistake, I must exclude the figures involving it entirely from our discussion.

The three available experiments, however, give the following results: The last column contains the ratio of silver sulphide to 100 parts of chloride.

```
6.6075 grm. AgCl gave 5.715 grm. Ag<sub>2</sub>S. 86.478
9.2323 " 7.98325 " 86.471
10.1775 " 8.80075 " 86.472
```

Mean, 86.4737, \pm .0015

We have also a single determination of this value by Svanberg and Struve.† After converting the chloride into sulphide they dissolved the latter in nitric acid. A trifling residue of chloride, which had been enclosed in sulphide, and so protected against change, was left undissolved. Hence a slight constant error probably affects this whole ratio. The experiment of Svanberg and Struve gave 86.472 per cent. of silver sulphide derived from 100 of chloride. If we assign this figure equal weight with the results of Berzelius, and combine, we get a general mean of 86.4733, \pm .0011.

For sodium there are but two ratios of any definite value for present purposes. The early work of Berzelius we may disregard entirely, and confine ourselves to the consideration of the results obtained by Penny, Pelouze, Dumas, and Stas.

^{*} Berzelius' Lehrbuch, 5th Ed., Vol. 3, p. 1187.

⁺ Journ. für Prakt. Chem., 44, 320. 1848.

The percentage of oxygen in sodium chlorate has been determined only by Penny,* who used the same method which he applied to the potassium salt. Four experiments gave the following results:

The ratio between silver and sodium chloride has been fixed by Pelouze, Dumas, and Stas. Pelouze† dissolved a weighed quantity of silver in nitric acid, and then titrated with sodium chloride. Equivalent to 100 parts of silver he found of chloride:

By Dumast we have seven experiments, with results as follows: The third column gives the ratio between 100 of silver and NaCl.

2.0535 gr	m. NaCl =	3.788 gr	m. Ag.		54.211	
2.169	44	4.0095	44		54.097	
4-3554	46	8.0425	"		54.155	
6.509	44	12.0140	"		54.178	
6.413	"	11.8375	44		54.175	
2.1746	"	4.012	44		54.202	
5.113	"	9.434	"		54.187	
				Mean	E4 172	aaa6

Mean, 54.172, ± .0096

Stas, || applying the method used in establishing the similar ratio for potassium chloride, and working with salt from

^{*} Phil. Transactions, 1839, p. 25.

[†] Compt. Rend., 20, 1047. 1845.

[‡] Ann. Chem. Pharm., 113, 31. 1860.

^{||} Aronstein's Translation, p. 274.

six different sources, found of sodium chloride equivalent to 100 parts of silver:

```
54.2093
54.2088
54.2070
54.2070
54.2070
54.2060
54.2076
54.2081
54.2083
54.2089
```

Mean, 54.2078, ± .0002

Now, combining these three series, we get the following result:

```
Pelouze_____ 54.141, ± .0063
Dumas _____ 54.172, ± .0096
Stas _____ 54.2078, ± .0002
    General mean..... 54.2076, ± .0002
```

Here the work of Stas is of such superior excellence that the other series might be completely rejected without appreciably affecting our calculations.

We have now before us the data establishing, with greater or less accuracy, twenty different ratios relating to the atomic weights of the seven elements under discussion. In these we are to discuss the results of about two hundred and fifty separate experiments. Before beginning upon our calculations we will tabulate our ratios, and number them for convenient future reference. Of course it will be understood that the probable errors given below relate to the last term of each proportion:

```
(3.)
     66
(4.)
     66
            AgClO<sub>3</sub> ----- 25.0795, ± .0010
(5.)
     66
            (6.)
            AgIO_8 ...... 16.9771, \pm .0009
(7.)
         Ag in Ag_2SO_4 ..... 69.205, \pm .0011
(8.)
```

```
(9.) Ag: NaCl:: 100: 54.2076, \pm.0002

(10.) Ag: KCl:: 100: 69.1032, \pm.0002

(11.) Ag: KBr:: 100: 110.3459, \pm.0019

(12.) Ag: KI:: 100: 153.6994, \pm.0178

(13.) Ag: Cl:: 100: 32.8418, \pm.0006

(14.) Ag: Br:: 100: 74.0809, \pm.0006

(15.) Ag: I:: 100: 117.5345, \pm.0009

(16.) Ag: Ag<sub>3</sub>S:: 100: 114.8581, \pm.0006

(17.) KCl: AgCl:: 100: 192.294, \pm.0029

(18.) AgCl: AgBr:: 100: 131.030, \pm.023

(19.) AgCl: AgI:: 100: 86.4733, \pm.0011
```

Now, from ratios 1 to 7 inclusive, we can at once, by applying the known atomic weight of oxygen, deduce the molecular weights of seven haloid salts. Let us consider the first calculation somewhat in detail.

Potassium chlorate yields 39.154 per cent. of oxygen and 60.846 per cent. of residual chloride. For each of these quantities the probable error is \pm .00038. The atomic weight of oxygen is 15.9633, \pm .0035, so that the value for three atoms becomes 47.8899, \pm .0105. We have now the following simple proportion: 39.154:60.846::47.8899:x, = the molecular weight of potassium chloride, = 74.4217. The probable error being known for the first, second, and third term of this proportion, we can easily find that of the fourth term by the formula given in our introduction. It comes out \pm .0164. By this method we obtain the following series of values, which may conveniently be numbered consecutively with the foregoing ratios:

```
(21.) KCl, from (1,) = 74.4217, ± .0164

(22.) KBr, " (2,) = 119.117, ± .0962

(23.) KI, " (3,) = 165.210, ± .0529

(24.) NaCl, " (4,) = 58.366, ± .0137

(25.) AgCl, " (5,) = 143.062, ± .0320

(26.) AgBr, " (6,) = 187.453, ± .0432

(27.) AgI, " (7,) = 234.195, ± .0530
```

With the help of these molecular weights we are now able to calculate eight independent values for the atomic weight of silver:

```
from (10) and (21,) Ag = 107.696, \pm .024
First,
             (11) " (22,) " = 107.948, \pm .087
Second,
             (12) " (23,) " = 107.488, \pm .037
Third,
             (9) " (24,) " = 107.671, \pm .025
Fourth.
Fifth,
             (13) " (25,) " = 107.694, \pm .024
             (14) " (26,) " = 107.681, \pm .025
Sixth,
             (15) " (27,) " = 107.659, \pm .024
Seventh.
             (8) " (16,) " = 107.712, \pm .025
Eighth,
             General mean, " = 107.675, ± .0096
```

It is noticeable that six of these values agree very well. The second and third, however, diverge widely from the average, but in opposite directions; they have, moreover, high probable errors, and consequently little weight. Of these two, one represents little and the other none of Stas' work. Their trifling influence upon our final results becomes curiously apparent in the series of silver values given a little further along.

When we consider closely, in all of its bearings, any one of the values just given, we shall see that for certain purposes it must be excluded from our general mean. example, the first is derived partly from the ratio between silver and potassium chloride. From this ratio, the atomic weight of one substance being known, we can deduce that of the other. We have already used it in ascertaining the atomic weight of silver, and the value thus obtained is included in our general mean. But if from it we are to determine the molecular weight of potassium chloride, we must use a silver value derived from other sources only, or we should be assuming a part of our result in advance. other words, we must now use a general mean for silver from which this ratio with reference to silver has been re-Hence the following series of silver values, which are lettered for reference:

```
A. General mean from all eight...... 107.675, ± .0096
B.
                rejecting the first ..... 107.671, \pm .0105
C.
                           second _____ 107.671, ± .0097
D.
                           third _____ 107.679, ± .0100
                     **
E.
                           fourth _____ 107.675, ± .0104
         "
F.
                     "
                           fifth ..... 107.671, ± .0105
         "
                     "
G.
                           sixth _____ 107.674, ± .0104
         "
H.
                     "
                          seventh_____ 107.678, ± .0105
                     66
                           eighth _____ 107.679, ± .0104
I.
```

These values are essentially the same, both in magnitude and in weight. For all practical purposes any one of them is as good as any other. Still, on theoretical grounds, it may be well to keep them distinct and separate in the remainder of this discussion.

We are now in a position to determine more closely the molecular weights of the haloid salts which we have already been considering.

For silver chloride, still employing the formula for the probable error of the last term of a proportion, we get the following values:

Subtracting from this the atomic weight of silver, 107.675, \pm .0096, we get for the atomic weight of chlorine, Cl = 35.370, \pm .014.

For silver bromide we have these results:

Hence, using the general mean for silver as above, $Br = 79.768, \pm .019$.

Silver iodide comes out as follows:

Hence $I = 126.557, \pm .022$.

For the molecular weight of sodium chloride we have:

Hence, if chlorine = 35.370, $\pm .014$, then Na = 22.998, $\pm .011$.

For potassium chloride:

For potassium bromide we get:

And for potassium iodide:

Now, taking the molecular weights of these three potassium salts in connection with the atomic weights just found for chlorine, bromine, and iodine, we get these values for potassium:

Finally, the three sulphur ratios give us three estimates for the atomic weight of sulphur. In the third of these I have applied the "A" value for silver and the general mean for silver chloride:

We may now appropriately compare the results of this

discussion with the atomic weights deduced by Stas from his own experiments only. His values are given under two headings: one for oxygen = 16, the other for O = 15.96. As we have been using the figure 15.9633 for oxygen, here is at the outset a discrepancy. Starting from this value we found:

Ag = 107.675,
$$\pm$$
 .0096
Cl = 35.370, \pm .014
Br = 79.768, \pm .019
I = 126.557, \pm .022
Na = 22.998, \pm .011
K = 39.019, \pm .012
S = 31.984, \pm .012

If we assume 16 to be the true figure for oxygen, we get the following results, which I have placed in a column parallel with the values found by Stas:

The New Values.	Stas.	Differences.
Silver 107.923	107.930	.007
Chlorine 35.451	35-457	.006
Bromine 79.951	79.952	100.
Iodine 126.848	126.850	.002
Sodium 23.051	23.043	.009
Potassium 39.109	39.137	.028
Sulphur 32.058	32.074	.016

These differences are insignificant. No other criticism could more severely test the character of Stas' work, or more definitely illustrate his magnificent accuracy of manipulation.

NITROGEN.

The atomic weight of nitrogen has been determined from the density of the gas, from the ratio between ammonium chloride and silver, and from the composition of certain nitrates.

Upon the density of nitrogen a great many experiments have been made. In early times this constant was determined by Biot and Arago, Thomson, Dulong and Berzelius, Lavoisier, and others. But all of these investigations may be disregarded as of insufficient accuracy; and, as in the case of oxygen, we need consider only the results obtained by Dumas and Boussingault, and by Regnault.

Taking air as unity, Dumas and Boussingault* found the density of nitrogen to be—

For hydrogen, as was seen in our discussion of the atomic weight of oxygen, the same investigators found a mean of .0693, \pm .00013. Upon combining this with the above nitrogen mean, we find for the atomic weight of the latter element, $N = 14.026, \pm .0295$.

By Regnault† much closer work was done. He found the density of nitrogen to be as follows:

```
.97148
.97148
.97154
.97155
.97108
.97108
Mean, .97137, ± .000062
```

^{*} Compt. Rend., 12, 1005. 1841. † Compt. Rend., 20, 975. 1845.

For hydrogen, Regnault's mean value is .069263, \pm .000019. Hence, combining as before, N = 14.0244, \pm .0039.*

The value found by combining both series of experiments is $N = 14.0244, \pm .0039$.

In discussing the more purely chemical ratios for establishing the atomic weight of nitrogen, we may ignore, for the present, the researches of Berzelius, of Anderson, and of Svanberg. These chemists experimented chiefly upon lead nitrate, and their work is consequently now of greater value for fixing the atomic weight of lead. Their results will be duly considered in the proper connection further on.

The ratio between ammonium chloride and silver has been determined by Pelouze, by Marignac, and by Stas. The method of working is essentially that adopted in the similar experiments with the chlorides of sodium and potassium.

For the ammonium chloride equivalent to 100 parts of silver, Pelouzet found:

Mean, 49.5365, ± .013

Marignac‡ obtained the following results. The usual ratio for 100 parts of silver is given also:

8.063	grm. Ag =	3.992 gm	m. NH ₄ Cl.	49.510
9.402	46	4.656	44	49.521
10.339	**	5.120	**	49.521
12.497	4	6. 191	**	49.540
11.337	**	5.617	**	49.546
11.307	**	5-595	46	49.483
4.326	44	2.143	44	49.538

Mean, 49.523, ± .0055

^{*} Professor Le Conte, in his corrections of Regnault's calculations, already cited in a foot note to the chapter on oxygen, finds for the density of nitrogen the value 0.971346. Hence N = 14.0225. This correction is very slight, but it should be considered in any future revision of the atomic weights.

[†] Compt. Rend., 20, 1047. 1845.

¹ Berzelius' Lehrbuch, 5th Ed., 3d v., 1184, 1185.

But neither of these series can for a moment compare with that of Stas.* He used from 12.5 to 80 grammes of silver in each experiment, reduced his weighings to a vacuum standard, and adopted a great variety of precautions to ensure accuracy. He found for every 100 parts of silver the following quantities of NH₄Cl:

```
49.600
49.599
49.598
49.597
49.593
49.597
49.597
49.602
49.597
49.598
49.592
```

Mean, 49.5973, ± .0005

Now, combining these three series, we get:

```
Pelouze 49.5365, ± .013

Marignac 49.523, ± .0055

Stas 49.5973, ± .0005

General mean 49.597, ± .0005
```

The quantity of silver nitrate which can be formed from a known weight of metallic silver has been determined by Penny, by Marignac, and by Stas. Penny† dissolved silver in nitric acid in a flask, evaporated to dryness without transfer, and weighed. One hundred parts of silver thus gave of nitrate:

```
157.430

157.437

157.458

157.440

157.430

157.455

Mean, 157.4417, ± .0033
```

^{*} Aronstein's Translation, pp. 56-58. † Phil. Trans., 1839.

Marignac's* results were as follows. In the third column they are reduced to the common standard of 100 parts of silver:

68.987	grm. Ag gave	108.608 grm.	AgNO ₃ .	157.433
57.844	**	91.047	"	157.401
66.436	66	104.592	"	157-433
70.340	66	110.718	"	157.404
200.000	"	314.894	"	157-447

Mean, 157.4236, ± .0061

Stas,† employing from 77 to 405 grammes of silver in each experiment, made two different series of determinations at two different times. The silver was dissolved with all the usual precautions against loss and against impurity, and the resulting nitrate was weighed, first after long drying without fusion just below its melting point; and again, fused. Between the fused and the unfused salt there was in every case a slight difference in weight, the latter giving a maximum and the former a minimum value.

In Stas' first series there are eight experiments; but the seventh he himself rejects as inexact. The values obtained for the nitrate from 100 parts of silver are given below in two columns, representing the two conditions in which the salt was weighed. The general mean given at the end I have deduced from the means of the two columns considered separately:

	Onfusea.		rusea.
	157.492		157.474
	157.510		157.481
	157.485		157.477
	157.476		157.471
	157.478		157.470
	157.471		157.463
	157.488		157.469
Mean,	157.4857	Mean,	157.472
	General mean	, 157.474, ±	.0014

22.6....

^{*} Berzelius' Lehrbuch, 5th Ed., 3, pp. 1184, 1185.

[†] Aronstein's Translation, pp. 305 and 315.

In the later series there are but two experiments, as follows:

	Unfused.		Fused.
	157.4964		157.488
	157.4940		157.480
Mean,	157.4952	Mean,	157.484
	General	mean, 157.486. +	.0003

Now, to combine all four sets of results:

Penny	157.4417, ± .0033
Marignac	157.4236, ± .0061
Stas, 1st series	
Stas, 2d series	157.4860, ± .0003
	
General mean	157.470. + .0003

For the direct ratio between silver nitrate and silver chloride there are two series of estimations. A weighed quantity of nitrate is easily converted into chloride, and the weight of the latter ascertained. In two experiments Turner* found of chloride from 100 parts of nitrate:

Penny,† in five determinations, found the following percentages:

The general mean from both series is 84.3743, \pm .0025.

The ratio directly connecting silver nitrate with ammonium chloride has been determined only by Stas.‡ The

^{*} Phil. Trans., 1833, 537.

[†] Phil. Trans., 1839.

[‡] Aronstein's Translation, p. 309.

usual method of working was followed; namely, nearly equivalent quantities of the two salts were weighed out, the solutions mixed, and the slight excess of one estimated by titration. In four experiments 100 parts of silver nitrate were found equivalent to chloride of ammonium as follows:

The similar ratio between potassium chloride and silver nitrate has been determined by both Marignac and Stas.

Marignac* gives the following weights. I add the quantity of KCl proportional to 100 parts of AgNO₈:

1.849 g	rm. KCl =	4.218 gr	m. AgNO ₃ .	43.836
2.473	44	5.640	"	43.848
3.317	46	7.565	. "	43.847
2.926	44	6.670	44	43.868
6. 191	**	14.110	44	43.877
4.351	**	9.918	"	43.870

Mean, 43.858, ± .0044

Stas'† results are given in three series, representing silver nitrate from three different sources. In the third series the nitrate was weighed in vacuo, while for the other series this correction was applied in the usual way. For the KCl equivalent to 100 parts of AgNO₃ Stas found:

```
First Series.

43.878

43.875

43.875

43.874

Mean, 43.8755, ± .0005
```

^{*} Berzelius' Lehrbuch, 5th Ed., 3d vol., 1184, 1185.

[†] Aronstein's Translation, p. 308.

```
Second Series.

43.864
43.869
43.876

Mean, 43.8697, ± .0023

Third Series.
43.894
43.878
43.885

Mean, 43.8857, ± .0031
```

Combining all four series we have:

Mar	igna	c	43.858,	± .0044	
Stas	, 1st	serie	s	43.8755,	± .0005
46	2 d	44		43.8697,	± .0023
**	3 d	44		43.8857,	± .0031
	G	e ner	al mean	43.8715.	+ .0004

There have also been determined by Penny and by Stas a series of ratios connecting the alkaline chlorides and chlorates with the corresponding nitrates. One of these, relating to the lithium salts, will be studied further on with reference to that metal.

The general method of working upon these ratios is due to Penny.* Applied to the ratio between the chloride and nitrate of potassium it is as follows: A weighed quantity of the chloride is introduced into a flask which is placed upon its side and connected with a receiver. An excess of pure nitric acid is added, and the transformation is gradually brought about by the aid of heat. Then, upon evaporating to dryness over a sand bath, the nitrate is brought into weighable form. The liquid in the receiver is also evaporated, and the trace of solid matter which had been mechanically carried over is recovered and also taken into account. In another series of experiments the nitrate was taken, and by pure hydrochloric acid converted into chloride; the process being the same. In the following columns of figures I

^{*} Phil. Trans., 1839.

have reduced both series to one standard; namely, so as to express the number of parts of nitrate corresponding to 100 of chloride:

```
First Series .- KCl treated with HNO2.
               135.639
               135.637
               135.640
               135.635
               135.630
               135.640
               135.630
         Mean, 135.636, ± .0011
Second Series .- KNO3 treated with HCl.
               135.628
               135.635
               135.630
               135.641
               135.630
               135.635
               135.630
         Mean, 153.633, ± .0011
```

Stas* results are as follows:

```
135.643

135.638

135.647

135.649

135.645

135.655

Mean, 135.6453, ± .0014
```

These figures by Stas represent weighings in the air. Reduced to a vacuum standard this mean really becomes 135.6423.

Now, combining, we have:

^{*} Aronstein's Translation, p. 270.

By the same general process Penny* determined how much potassium nitrate could be formed from 100 parts of chlorate. He found as follows:

```
82.505
82.497
. 82.498
82.500
———
Mean, 82.500, ± .0012
```

For 100 parts of sodium chlorate he found of nitrate:

For the ratio between the chloride and nitrate of sodium Penny made two sets of estimations as in the case of potassium salts. The subjoined figures give the amount of nitrate equivalent to 100 parts of chloride:

```
First Series.—NaCl treated with HNO<sub>3</sub>.

145.415

145.408

145.420

145.424

145.410

145.418

145.420
```

Second Series .- Na NO2 treated with HCl.

Mean, 145.4164, ± .0015

145.419 145.391 145.412 145.415 145.412 145.412 Mean, 145.410, ± .0026

^{*} Phil. Trans., 1839.

Stas* gives the following series:

```
145.453
145.468
145.465
145.469
```

Mean, after reducing to vacuum standard, 145.4526, ± .0030

Combining, we have as follows:

```
Penny, 1st series ______ 145.4164, ± .0015
" 2d " ______ 145.410, ± .0026
Stas _____ 145.4526, ± .0030

General mean _____ 145.4185, ± .0012
```

We have now, apart from the determinations of gaseous density, nine ratios, representing one hundred and fourteen experiments from which to calculate the atomic weight of nitrogen. Let us first collect and number these ratios:

- (1.) Ag: $AgNO_3$:: 100: 157.479, \pm .0003
- (2.) $AgN\ddot{O}_3$: AgCl:: 100: 84.3743, \pm .0025
- (3.) $AgNO_8$: KCl:: 100: 43.8715, \pm .0004
- (4.) $AgNO_3$: NH_4Cl :: 100: 31.488, \pm .0006
- (5.) Ag: NH₄Cl:: 100: 49.597, ± .0005
- (6.) KCl: KNO_3 :: 100: 135.6363, \pm .0007
- (7.) KClO₃: KNO₃:: 100: 82.500, ± .0012
- (8.) NaCl: NaNO₃:: 100:145.4185, $\pm .0012$
- (9.) $NaClO_3 : NaNO_3 :: 100 : 79.8823, \pm .0029$

From these ratios we are now able to deduce the molecular weight of ammonium chloride and of the three nitrates named in them. For these calculations we may use the already determined atomic weights of silver, oxygen, potassium, sodium, and chlorine, and the molecular weights of silver chloride and sodium chloride. These two molecular weights involve, respectively, the most probable values for silver, sodium, and chlorine. We cannot, however, appropriately use the directly determined molecular weight of potassium chloride, since the most probable value for the

^{*} Aronstein's Translation, p. 278.

atomic weight of potassium is only in part derived from that salt. The following are the values which we shall employ:

Now, from ratio number five we can get the molecular weight of ammonium chloride, $NH_4Cl = 53.4048$, $\pm .0048$, and N = 14.0336, $\pm .0153$.

From ratio number four an independent value for nitrogen can be calculated, namely, $N = 14.0330, \pm .015$.

For the molecular weight of silver nitrate three values are deducible, namely:

Hence $N = 13.9840, \pm .0174$.

The molecular weight of potassium nitrate is twice calculable, as follows:

And $N = 13.9774, \pm .0216$.

So also for sodium nitrate we have:

And N = 13.9906, $\pm .0163$.

We have now before us six estimates of the atomic weight of nitrogen. It only remains for us to combine these after

the usual method, as follows, in order to obtain the most probable value:

ı.	From	a specific gravity of $N_{}N = 14.0244$, ± .0	039
2.	"	ammonium chloride " = 14.0336	o. ± .o	153
3.	44	ratio number four " = 14.0330), ± .0	150
4.	66	silver nitrate " = 13.9840), ± .0	174
5.	44	potassium nitrate " = 13.9774	, ± .0	216
6.	64	sodium nitrate = 13.9900	ό, ± .ο	163
	G	General mean " - 14 0010	· - ^	025

If oxygen is 16, this becomes 14.0291. Stas found N = 14.044. The difference is .015, showing a remarkably close agreement.

CARBON.

Although there is a large mass of material relating to the atomic weight of carbon, much of it may be summarily set aside as having no value for present purposes. The density of carbon dioxide, which has been scrupulously determined by many investigators,* leads to no safe estimate of the constant under consideration. The numerous analyses of hydrocarbons, like the analyses of naphthalene by Mitscherlich, Woskresensky, Fownes, and Dumas, give results scarcely more satisfactory. In short, all the work done upon the atomic weight of carbon before the year 1840 may be safely rejected as unsuited to the present requirements of exact science. As for methods of estimation we need consider but three, as follows:

First.—The analysis of organic salts of silver.

Second.—The determination of the weight of carbon dioxide formed by the combustion of a known weight of carbon.

^{*} Notably by Lavoisier, Biot and Arago, De Saussure, Dulong and Berzelius, Buff, Von Wrede, Regnault, and Marchand. For details, Van Geuns' monograph may be consulted.

51

Third.—The method of Stas, by the combustion of carbon monoxide.

The first of these methods, which is also the least accurate, was employed by Liebig and Redtenbacher* in 1840. They worked with the acetate, tartrate, racemate, and malate of silver, making five ignitions of each salt, and determining the percentage of metal. From one to nine grammes of material were used in each experiment.

In the acetate the following percentages of silver were found:

After applying corrections for weighing in air this mean becomes 64.6065.

In the tartrate the silver came out as follows:

Or, reduced to a vacuum, 59.2806

In the racemate we have:

```
59.290
59.292
59.287
59.283
59.284
-----
Mean, 59.2872, ± .0012
Or, corrected, 59.2769
```

^{*}Ann. Chem. Pharm., 38, 137. Mem. Chem. Soc., 1, 9. Phil. Mag., (3,) 19, 210.

And from the malate:

61.996
61.972
62.015
62.059
62.011
Mean, 62.0106, ± .0096
Or, corrected, 62.0016

Now, applying to these mean results the atomic weights already found for oxygen and silver, we get the following values for carbon:

From the acetate ______C = 12.0306, ± .0047

" tartrate _____" = 12.0356, ± .0064

" racemate ____" = 12.0413, ± .0063

" malate ____" = 12.0408, ± .0054

General mean ____" = 12.0363, ± .0028

Now these results, although remarkably concordant, are by no means unimpeachable. They involve two possible sources of constant error, namely, impurity of material and the volatility of the silver. These objections have both been raised by Stas, who found that the silver tartrate, prepared as Liebig and Redtenbacher prepared it, always carried traces of the nitrate, and that he, by the ignition of that salt, could not get results at all agreeing with theirs. In the case of the acetate a similar impurity would lower the percentage of silver, and thus both sources of error would reinforce each other and make the atomic weight of carbon come out too high. With the three other salts the two sources of error act in opposite directions, although the volatility of the silver is probably far greater in its influence than the impurity. Even if we had no other data relating to the atomic weight of carbon, it would be clear from these facts that the results obtained by Liebig and Redtenbacher must be decidedly in excess of the true figure.

A different method of dealing with organic silver salts was adopted by Maumené,* in 1846, for the purpose of estab-

^{*} Ann. d. Chim. et d. Phys., (3,) 18, 41.

CARBON. 53

lishing, by reference to carbon, the atomic weight of silver. We will simply reverse his results and apply them to the atomic weight of carbon. He effected the combustion of the acetate and the oxalate of silver, and, by weighing both the residual metal and the carbon dioxide formed, he fixed the ratio between these two substances. In the case of the acetate his weighings show that for every gramme of metallic silver the weights of CO₂ were produced, which are shown in the third column:

8.083 g	rm. Ag =	6.585 gr	m. CO ₂ .	.8147
11.215	44	9.135	"	.8136
14.351	44	11.6935	"	.8148
9.030	66	7.358	46	.8148
20.227	66	16.475	"	.8145
				Mean, .81448

The oxalate of silver, ignited by itself, decomposes too violently to give good results; and for this reason it was not used by Liebig and Redtenbacher. Maumené, however, found that when the salt was mixed with sand the combustion could be tranquilly effected. The oxalate employed, however, with the exception of the sample represented in the last experiment of the series, contained traces of nitrate, so that these results involve slight errors. For each gramme of silver the appended weights of CO₂ were obtained:

14.299 gm	m. Ag:	= 5.835 g	rm. CO ₂ .	.4081
17.754	"	7.217	**	.4059
11.550	46	4.703	"	.4072
10.771	"	4. 387	44	.4073
8.674	66	3.533	66	.4073
11.4355	66	4.658	**	.4073
				Mean, .40718

Now, one of these salts being formed by a bivalent and the other by a univalent acid, we have to reduce both to a common standard. Doing this, we have the following results for the ratio between the atomic weight of silver and the molecular weight of CO_2 ; if Ag = 1.00,

Here the slight error due to the impurity of the oxalate becomes of such trifling weight that it practically vanishes.

From these data, if Ag = 107.675, $\pm .0096$, $CO_2 = 43.8485$, $\pm .0086$.

Hence $C = 11.9219, \pm .0111$.

As has already been said, the volatility of silver renders all the foregoing results more or less uncertain. Far better figures are furnished by the combustion of carbon directly, as carried out by Dumas and Stas* in 1840 and by Erdmann and Marchand† in 1841. In both investigations weighed quantities of diamond, of natural graphite, and of artificial graphite were burned in oxygen, and the amount of dioxide produced was estimated by the usual methods. The graphite employed was purified with extreme care by treatment with strong nitric acid and by fusion with caustic alkali. I have reduced all the published weighings to a common standard, so as to show in the third column the amount of oxygen which combines with a unit weight (say one gramme) of carbon. Taking Dumas and Stas' results first in order we have from natural graphite:

1.000	grm.	C gave	3.671	grm. CO ₃ .		2.6710
.998		44	3.660	44		2.6673
.994		46	3.645	"		2.6670
1.216		44	4.461	**		2.6686
1.471		"	5-395	u ,		2.6676
					Mean,	2.6683, ± .0005

With artificial graphite:

.992 gr	m. C ga	ve 3.642 g	rm. CO ₂ .	2.6714
.998	"	3.662	"	2.6682
1.660	44	6.085	"	2.6654
1.465	44	5.365	"	2.6744
				Mean, 2.66985, ± .0013

^{*} Compt. Rend., 11, 991-1008. Ann. Chim. Phys., (3,) 1, 1.

[†] Journ. f. Prakt. Chem., 23, 159.

And with diamond:

```
.708 grm. C gave 2.598 grm. CO.
                                       2.6695
.864 " 3.1675
                                      2.6661
         66
                                       2,6628
                        "
1.219
               4.465
         66
                        "
                                       2.6680
1.232
                4.519
         66
                        "
1.375
                5.041
                                       2.6662
```

Mean, 2.6665, ± .0007

Erdmann and Marchand's figures for natural graphite give the following results:

In one experiment 1.8935 grm. of artificial graphite gave 6.9355 grm. CO₂. Ratio for O, 2.6628. This, combined with the foregoing series, gives a mean of 2.6636, \pm .0007.

With diamond they found:

```
.8052 grm. gave 2.9467 grm. CO<sub>2</sub>.
                                             2.6596
1.0858
        66
                 3.9875
                                             2.6632
           "
                            "
1.3557
                 4.9659
                                             2.6629
           "
                            "
1.6305
                 5.97945
                                             2.6673
                                             2.6653
.7500
                 2.7490
                                      Mean, 2.6637, ± .0009
```

Now, combining all these series, we get the following result:

Hence, if O = 15.9633, $\pm .0035$, C = 11.973, $\pm .0030$.

Another very exact method for determining the atomic weight of carbon was employed by Stas* in 1849. Carefully purified carbon monoxide was passed over a known weight

[#] Bull. Acad. Bruxelles, 1849, (1,) 31.

of copper oxide at a red heat, and both the residual metal and the carbon dioxide formed were weighed. The weighings were reduced to a vacuum standard, and in each experiment a quantity of copper oxide was taken representing from eight to twenty-four grammes of oxygen. The method, as will at once be seen, is in all essential features similar to that usually employed for determining the composition of water. The figures in the third column, deduced from the weights given by Stas, represent the quantity of carbon monoxide corresponding to one gramme of oxygen:

9.265 gm	m. O =	= 25.483 (CO ₂ .	1.75046
8.327	**	22.900	"	1.75010
13.9438	**	38.351	"	1.75040
11.6124	"	31.935	66	1.75008
18.763	46	51.6055	"	1.75039
19.581	44	53.8465	66	1.74994
22.515	46	61.926	44	1.75043
24.360	"	67.003	"	1.75053
				Mean, 1.75029, ± .00005

Hence the molecular weight of carbon monoxide is $27.9404, \pm .0062$. And $C = 11.9771, \pm .0071$.

Now, in order to complete our discussion, we must combine the four values we have found for carbon:

```
1. By Liebig and Redtenbacher... C = 12.0363, ± .0028
2. By Maumené's figures ....... " = 11.9219, ± .0111
3. By combustion of carbon..... " = 11.9730, ± .0030
4. By Stas' method ...... " = 11.9771, ± .0071

General mean...... " = 12.0021, ± .0019
```

But values one and two are hardly reliable enough to be included in our final estimate. They involve dangerous constant errors, and ought, therefore, to be disregarded. Rejecting them altogether, and taking a general mean from values three and four, we get for the most probable figure for the atomic weight of carbon, C = 11.9736, $\pm .0028$. If oxygen is 16, then carbon becomes 12.0011. In other words, the ratio between oxygen and carbon is almost exactly 16 to 12.

BARIUM.

For determining the atomic weight of barium we have a series of six ratios, established by the labors of Berzelius, Turner, Struve, Pelouze, Marignac, and Dumas. Andrews* and Salvetat,† in their papers upon this subject, gave no details nor weighings; and, therefore, their work may be properly disregarded. First in order in point of importance, if not first chronologically, is the ratio between silver and anhydrous barium chloride, as determined by Pelouze, Marignac, and Dumas.

Pelouze, in 1845, made the three subjoined estimations of this ratio, using his well known volumetric method. A quantity of pure silver was dissolved in nitric acid, and the amount of barium chloride needed to precipitate it was accurately ascertained. In the last column I give the quantity of barium chloride proportional to 100 parts of silver:

3.860 grn	ı.BaCl, p	pt. 4.002 gr	m. Ag.	96.452
5.790	"	6.003	"	96.452
2.895	66	3.001	"	96.468
				 _

Mean, 96.4573, ± .0036

Essentially the same method was adopted by Marignac|| in 1848. His experiments were made upon four samples of barium chloride, as follows. A, commercial barium chloride, purified by recrystallization from water. B, the same salt, calcined, redissolved in water, the solution saturated with carbonic acid, filtered, and allowed to crystallize. C, the preceding salt, washed with alcohol, and again recrystallized. D, the same, again washed with alcohol. For 100 parts of silver the following quantities of chloride were required:

^{*} Chemical Gazette, October, 1852.

[†] Compt. Rend., 17, 318.

¹ Compt. Rend., 20, 1047. Journ. für Prakt. Chem., 35, 73.

^{||} Arch. d. Sci. Phys. et Nat., 8, 271.

Dumas* employed barium chloride prepared from pure barium nitrate, and took the extra precaution of fusing the salt at a red heat in a current of dry hydrochloric acid gas. Three series of experiments upon three samples of chloride gave the following results:

1.7585 gr	m.BaCl ₂ =	1.826 g	ran. Ag.	Ratio, 96.303
3.842	"	3.988	"	96.339
2.1585	44	2.2405	"	96.340
4.0162	"	4.168	"	96.358
				
				Mean, 96.3325, ± .0068

^{*} Ann. Chem. Pharm., 113, 22. 1860. Ann. Chim. Phys., (3,) 55, 129.

59

Series B.

1.6625	grm. BaCl ₂ =	1.727	grm. Ag.	Ratio,	96.265
2.4987	66	2.594	6 "		96.304
3.4468	66	3.579	**		96.306
4.0822	66	4.239	5 "		96.290
4.2062	44	4.368	3 "		96.289
4.4564	44	4.629	44		96.271
8.6975	66	9.031	44		96.307
				Mean,	96.2902, ± .0043
			Series C.		
2.2957	grm. BaCl ₂ =	2.383	grm. Ag.	Ratio,	96.316
4.1372	"	4.293			96.371

4.2662 " 4.430 " 96.303 4.4764 " 4.647 " 96.329 5.6397 " 5.852 " 96.372

Mean, 96.3382, $\pm .0096$

We have now eight series of experiments upon this ratio, representing thirty distinct estimations. Combining, we get a general mean as follows:

Pelouze	96.4573, ± .0036
Marignac, A	96.3543, ± .0033
" B	96.3540, ± .0013
" C	96.3605, ± .0017
" D	96.3670, ± .0057
Dumas, A	$96.3325, \pm .0068$
" B	96.2902, ± .0043
" C	96.3382, ± .0096
General mean	96.3596, ± .0009

The ratio between silver and crystallized barium chloride has also been fixed by Marignac.* The usual method was employed, and two series of experiments were made; in the second of which the water of crystallization was determined previous to the estimation. Five grammes of chloride were taken in each determination. The following quantities of BaCl₂·2H₃O correspond to 100 parts of silver:

^{*} Journ. f. Prakt. Chem., 74, 212. 1858.

,	
A .	В.
113.109	113.135
113.135	113.122
113.097	113.060
	
Mean, 113.114, ± .0074	Mean, 113.106, ± .0154

The general mean from both series is 113.113, \pm .0067.

The direct ratio between the chlorides of silver and barium was early established both by Berzelius* and Turner.† Berzelius found that 100 parts of dry barium chloride gave of silver chloride:

Turner made five experiments, with the following results:

137.45 137.54 137.70 137.62 137.64

Of these, Turner regards the fourth and fifth as the most exact. These give a mean of 137.63, \pm .007, while the other three are in mean 137.563, \pm .049. Combining Berzelius' figures with those of Turner, we get as follows:

General mean	137.841, ± .c	047
" 4, 5	137.63, ± .c	07
Turner, I, 2, 3	$137.563, \pm .0$	49
Berzelius	138.07, ± .0	07

Incidentally to some of his other work Marignac‡ determined the percentage of water in crystallized barium chloride. Two sets of three experiments each were made, the first upon five grammes and the second upon ten grammes of salt. The following are the percentages obtained:

^{*} Poggend. Annal., 8, 177. † Phil. Trans., 1829, 291.

[†] Journ. f. Prakt. Chem., 74, 212. 1858.

A.	В.
14.790	14.80
14.796	14.81
14.800	14.80
Mean, 14.795, ± .0019	Mean, 14.803, ± .002
General mean of both series	14 700 + 0014

General mean of both series, 14.799, ± .0014

The ratio between barium nitrate and barium sulphate has been determined only by Turner.* According to his experiments 100 parts of sulphate correspond to the following quantities of nitrate:

For the similar ratio between the sulphate and the chloride there are experiments by Turner, Berzelius, Struve, and Marignac. Turner† found that 100 parts of chloride ignited with sulphuric acid gave 112.19 parts of sulphate. By the common method of precipitation and filtration a lower figure was obtained, because of the slight solubility of the sulphate. This point bears directly upon many other atomic weight determinations.

Berzelius,‡ treating barium chloride with sulphuric acid, obtained the following results in BaSO₄ for 100 parts of BaCl₂:

Struve, || in two experiments, found:

```
* Phil. Trans., 1833, 538.

† Phil. Trans., 1829, 291.

‡ Poggend. Annal., 8, 177.

|| Ann. Chem. Pharm., 80, 204. 1851.
```

Marignac's* three results are as follows:

8.520 grm.	BaCl ₂ gave	9.543	BaSO ₄ .	Ratio, 112.007
8.519	46	9.544	**	112.032
8.520	"	9.542	44	111.995
				Mean, 112.011, ± .0071

Rejecting Turner's single result as unimportant, we may combine the other series:

```
Berzelius 112.175, ± .0034
Struve 112.0938, ± .0018
Marignac 112.011, ± .0071
General mean 112.106, ± .0015
```

The data from which we are to calculate the atomic weight of barium may now be tabulated as follows:

- (1.) Ag₂: BaCl₂:: 100: 96.3596, ± .0009
- (2.) $Ag_2 : BaCl_2.2H_2O :: 100 : 113.113, \pm .0067$
- (3.) BaCl₂: 2AgCl:: 100: 137.841, ± .0047
- (4.) Per cent. of H₂O in BaCl₂.2H₂O, 14.799, ± .0014
- (5.) $BaSO_4: BaN_2O_6::112.028, \pm .014$
- (6.) BaCl₂: BaSO₄:: 100: 112.106, ± .0015

From these ratios, with the aid of the atomic weights already established, we can immediately calculate four independent values for the molecular weight of BaCl₂:

We have here an interesting example of the compensation of constant errors. Ratios (2) and (4) both represent work done by Marignac upon barium chloride containing water of crystallization. If now, as is not improbable, the salt contained a trifling excess of water, the molecular weight of barium chloride as calculated from (2) would come out too high, while on the other hand the result from ratio (4) would err in the opposite direction. In point of fact, the

^{*} Journ. f. Prakt. Chem., 74, 212. 1858.

63

two results in the present calculation nearly compensate each other, and, on account of their relatively high probable errors, they exert but an unimportant influence upon the general mean.

In conclusion, we have three independent values for the atomic weight of barium:

If O = 16, then Ba = 137.007. In other words, the ratio between oxygen and barium is almost an exact ratio between two whole numbers.

In the above discussion it will at once be noticed that the second and third values for Ba have very high probable errors, and that they therefore exert almost no influence upon the general mean. This fact by no means renders them worthless however, for, at the lowest estimate, they are useful in confirmation of the better determinations. It is also highly probable that the method of discussion, rigidly carried out, does not do them absolute justice.

STRONTIUM.

The ratios which fix the atomic weight of strontium resemble in general terms those relating to barium, only they are fewer in number and represent a comparatively small amount of work. The early experiments of Stromeyer,* who measured the volume of CO₂ evolved from a known weight of strontium carbonate, are hardly available for the present discussion. So also we may exclude the determination by Salvétat,† who neglected to publish sufficient details.

Taking the ratio between strontium chloride and silver first in order, we have series of figures by Pelouze and by Dumas. Pelouze‡ employed the volumetric method already described under barium, and in two experiments obtained the subjoined results. In another column I append the ratio between SrCl, and 100 parts of silver:

Dumas, by the same general method, made sets of experiments with three samples of chloride which had previously been fused in a current of dry hydrochloric acid. His results, expressed in the usual way, are as follows:

			Series A.		
3.137 gr	m. SrCl ₂	= 4.280 g	rm. Ag.	Ratio, 73.2	944
1.982	46	2.705	44	73.2	717
3.041	"	4. 142	66	73-4	186
3.099	44	4.219	"	73-4	534
				Mean, 73.3	 595, ± .0303

^{*} Schweigg. Journ., 19, 228. 1816.

[†] Compt. Rend., 17, 318. 1843.

[†] Compt. Rend., 20, 1047. 1845.

[|] Ann. Chim. Phys., (3,) 55, 29. 1859. Ann. Chem. Pharm., 113, 34.

Series R.

3.356 grm.	SrCl ₂ =	4-574	grm.	Ag.	Ratio,	73.3713	
6.3645	"	8.667	"			73-4327	
7.131	"	9.712	"			73.4246	•
					Mean,	73.4095.	± .0130

Series C.

7.213 g	rm. SrCl ₂ =	= 9.811 g	rm. Ag.	Ratio,	73.5195	
2.206	**	3.006	"		73.3866	
4.268	**	5.816	"		73.5529	
4.018	46	5-477	"		73.3613	
				Mean.	72.4551.	+ .0321

Combining, we have:

Pelouze		73.4781, ±	.0050
Dumas,	A	73·3595, ±	.0303
44	B	73.4095, 士	.0130
"	C	73.4551, ±	.0321
(General mean	73.4655. +	.6046

The foregoing figures apply to anhydrous strontium chloride. The ratio between silver and the crystallized salt, SrCl₂.6H₂O, has also been determined in two series of experiments by Marignac.* Five grammes of salt were used in each estimation, and, in the second series, the percentage of water was first determined. The quantities of the salt corresponding to 100 parts of silver are given in the last column:

Series A.

Series B.

^{*} Journ. Prakt. Chem., 74, 216. 1858.

In the same paper Marignac gives two sets of determinations of the percentage of water in crystallized strontium chloride. The first set, corresponding to "B" above, comes out thus:

In the second set ten grammes of salt were taken at a time, and the following percentages were found:

General mean, from both series, 40.575, ± .0015

The chloride used in the series of estimations last given was subsequently employed for ascertaining the ratio between it and the sulphate. Converted directly into sulphate, 100 parts of chloride yield the quantities given in the third column:

```
5.942 grm. SrCl<sub>2</sub> gave 6.887 grm. SrSO<sub>4</sub>. 115.932

5.941 " 6.8855 " 115.949

5.942 " 6.884 " 115.927

Mean, 115.936, ± .004
```

Now, to sum up the ratios and calculate the atomic weight of strontium.

```
(1.) Ag: SrCl_2:: 100: 73.4655, \pm .0046
(2.) Ag: SrCl_2:6H<sub>2</sub>O:: 100: 123.470, \pm .006
```

(3.) Per cent. of H_2O in $SrCl_2.6H_2O$, 40.575, $\pm .0015$

(4.) SrCl₂: SrSO₄:: 100: 115.936, ± .004

We now have the molecular weight of SrCl,, as follows:

And for the atomic weight of strontium itself we have two values, as follows:

If O = 16, then Sr = 87.575.

CALCIUM.

For determining the atomic weight of calcium we have sets of experiments by Berzelius, Erdmann and Marchand, and Dumas. Salvétat* also has published an estimation, but without the details necessary to enable us to make use of his results. I also find a reference† to some work of Marignac; which, however, seems to have been of but little importance. The earlier work of Berzelius was very inexact as regards calcium, and it is not until we come down to the year 1842 that we find any material of decided value.

The most important factor in our present discussion is the composition of calcium carbonate, as worked out by Dumas and by Erdmann and Marchand.

In 1842 Dumast made three ignitions of Iceland spar, and determined the percentages of carbon dioxide driven off and of lime remaining. The impurities of the material were also determined, the correction for them applied, and the weighings reduced to a vacuum standard. The percentage of lime came out as follows:

```
56.12
56.04
56.06
————
Mean, 56.073, ± .016
```

^{*} Compt. Rend., 17, 318. 1843. † See Oudeman's monograph, p. 51.

[‡] Compt. Rend., 14, 537. 1842.

About this same time Erdmann and Marchand* began their researches upon the same subject. Two ignitions of spar, containing .04 per cent. of impurity, gave respectively 56.09 and 56.18 per cent. of residue; but these results are not exact enough for us to consider further. Four other results obtained with artificial calcium carbonate are more noteworthy. The carbonate was precipitated from a solution of pure calcium chloride by ammonium carbonate, was washed thoroughly with hot water, and dried at a temperature of 180°. With this preparation the following residues of lime were obtained:

It was subsequently shown by Berzelius that calcium carbonate prepared by this method retains traces of water even at 200°, and that minute quantities of chloride are also held by it. These sources of error are, however, in opposite directions, since one would tend to diminish and the other to increase the weight of residue.

In the same paper there are also two direct estimations of carbonic acid in pure Iceland spar, which correspond to the following percentages of lime:

In a still later paper† the same investigators give another series of results based upon the ignition of Iceland spar. The impurities were carefully estimated, and the percentages of lime are suitably corrected:

^{*} Journ. für Prakt. Chem., 26, 472. 1842.

[†] Journ. für Prakt. Chem., 31, 269. 1844.

```
4.2134 grm. CaCO<sub>3</sub> gave 2.3594 grm. CaO.
                                              55.997 per cent.
                                    66
                66
15.1385
                         8.4810
                                               56.022
                                                        "
                         13.1958
                                               56.031
23.5503
23.6390
                         13.2456
                                               56.032
                                               56.044
42.0295
                         23.5533
49.7007
                         27.8536
                                               56.042
                                        Mean, 56.028, ± .0047
```

Six years later Erdmann and Marchand* published one more result upon the ignition of calcium carbonate. They found that the compound began giving off carbon dioxide below the temperature at which their previous samples had been dried, or about 200°, and that, on the other hand, traces of the dioxide were retained by the lime after ignition. These two errors do not compensate each other, since both tend to raise the percentage of lime. In the one experiment now under consideration these errors were accurately estimated, and the needful corrections were applied to the final The percentage of residual lime in this case came This agrees tolerably well with the figures out 55.998. found in the direct estimation of carbonic acid, and, if combined with those two, gives a mean for all three of 56.006, $\pm .0043.$

Combining all these series we get the following result:

```
      Dumas
      56.073, ± .016

      Erdmann and Marchand
      56.006, ± .007

      " " 56.028, ± .0047

      " " 56.006, ± .0043

      General mean
      56.0198, ± .0029
```

For reasons given above this mean is probably vitiated by a slight constant error, which makes the figure a trifle too high.

In the earliest of three papers by Erdmann and Marchand there is also given a series of determinations of the ratio between calcium carbonate and sulphate. Pure Iceland

^{*} Journ. für Prakt. Chem., 50, 237. 1850.

spar was carefully converted into calcium sulphate, and the gain in weight noted. One hundred parts of spar gave of sulphate:

136.07 136.06 136.02 136.06

Mean, 136.0525, ± .0071

In 1843 the atomic weight of calcium was redetermined by Berzelius,* who investigated the ratio between lime and calcium sulphate. The calcium was first precipitated from a pure solution of nitrate by means of ammonium carbonate, and the thoroughly washed precipitate was dried and strongly ignited in order to obtain lime wholly free from extraneous matter. This lime was then, with suitable precautions, treated with sulphuric acid, and the resulting sulphate was weighed. Correction was applied for the trace of solid impurity contained in the acid, but not for the weighing in air. The figures in the last column represent the percentage of weight gained by the lime upon conversion into sulphate:

```
      1.80425 grm. CaO gained
      2.56735 grm.
      142.295

      2.50400
      " 3.57050 " 142.592

      3.90000
      " 5.55140 " 142.343

      3.04250
      " 4.32650 " 142.202

      3.45900
      " 4.93140 " 142.567
```

Mean, 142.3998, ± .0518

Last of all we have the ratio between calcium chloride and silver, as determined by Dumas.† Pure calcium chloride was first ignited in a stream of dry hydrochloric acid, and the solution of this salt was afterwards titrated with a silver solution in the usual way. The CaCl, proportional to 100 parts of Ag is given in a third column:

^{*} Journ. für Prakt. Chem., 31, 263. Ann. Chem. Pharm., 46, 241.

[†] Ann. Chim. Phys., (3,) 55, 129. 1859. Ann. Chem. Pharm., 113, 34.

2.738	grm. CaCl ₂ =	= 5.309 g	rm. Ag.	51.573
2.436	**	4.731	44	51.490
1.859	44	3.617	44	51.396
2.771	46	5.3885	u	51.424
2.240	"	4.3585	"	51.394

Mean, 51.4554, ± .0230.

We have now four ratios to calculate from, as follows:

- (1.) Per cent. of CaO in CaCO₃, 56.0198, ± .0029
- (2.) CaO: SO₃:: 100: 142.3998, ± .0518
- (3.) CaCO₈: CaSO₄:: 100: 136.0525, ± .0071
- (4.) Ag: CaCl₂:: 100: 51.4554, ± .0230

These give us the subjoined values for calcium:

If O = 16, then Ca = 40.082.

A glance at the above figures will show that, if, as is probable, the value deduced from the composition of calcium carbonate is a trifle too high, the general mean must be too high also. It is, therefore, interesting to see what result the very latest of Erdmann and Marchand's experiments will lead to. They found, after taking every precaution, in a single experiment that calcium carbonate yielded 55.998 per cent. of lime. From this we get Ca = 39.905; or, if O = 16, Ca = 39.997. It is possible, then, that "Prout's law" may hold good for calcium.

LEAD.

For the atomic weight of lead we have to consider experiments made upon the oxide, chloride, nitrate, and sulphate. The researches of Berzelius upon the carbonate and various organic salts need not now be considered, nor is it worth while to take into account any work of his done before the year 1818. The results obtained by Döbereiner* and by Longchamp† are also without special present value.

For the exact composition of lead oxide we have to depend upon the researches of Berzelius. His experiments were made at different times through quite a number of years; but were finally summed up in the last edition of his famous "Lehrbuch.". In general terms his method of experiment was very simple. Perfectly pure lead oxide was heated in a current of hydrogen, and the reduced metal weighed. From his weighings I have calculated the percentages of lead thus found and given them in a third column:

Earlier Results.

8.045 grm	. PbO g	ave 7.4675 g	rm. Pb.	92.8217 p	er cent.
14.183	44	13.165	"	92.8224	64
10.8645	44	10.084	44	92.8160	44
13.1465	44	12.2045	**	92.8346	44
21.9425	44	20.3695	**	92.8313	66
11.159	44	10.359	"	92.8309	• •
		Late	st.		
6.6155	44	6. 141	44	92.8275	44
14.487	44	13.448	44	92.8280	46
14.626	44	13.5775	**	92.8313	¥

Mean, 92.8271, ± .0013

For the synthesis of lead sulphate we have data by Berzelius, Turner, and Stas. Berzelius, whose experiments

^{*} Schweig. Journ., 17, 241. 1816.

[†] Ann. Chim. Phys., 34, 105. 1827.

[‡] Bd. 3, s. 1218.

^{||} Lehrbuch, 5th Ed., 3, 1187.

73

were intended rather to fix the atomic weight of sulphur, dissolved in each estimation ten grammes of pure lead in nitric acid, then treated the resulting nitrate with sulphuric acid, brought the sulphate thus formed to dryness, and weighed. One hundred parts of metal yield of PbSO₄:

Turner,* in three similar experiments, found as follows:

In these results of Turner's absolute weights are implied.

The results of Stas' syntheses,† effected after the same general method, but with variations in details, are as follows.

Corrections for weighing in air were applied:

```
146.443

146.427

146.419

146.432

146.421

146.423

Mean, 146.4275, ± .0024
```

Combining, we get the subjoined result:

Turner, in the same paper, also gives a series of syntheses of lead sulphate, in which he starts from the oxide instead

^{*} Phil. Trans., 1833, 527-538. † Aronstein's Translation, 333.

of from the metal. One hundred parts of PbO, upon conversion into PbSO, gained weight as follows:

	35.84		
	35.71		
	35.84		
	3 5· 7 5		
	35.79		
	35.78		
	35.92		
Mean,	35.804,	\pm	.018

These figures are not wholly reliable. Numbers one, two, and three represent lead oxide contaminated with traces of nitrate. The oxide of four, five, and six contained traces of minium. Number seven was free from these sources of error, and, therefore, deserves more consideration. The series as a whole undoubtedly gives too low a figure; and this error would tend to slightly raise the atomic weight of lead.

Still a third series by Turner establishes the ratio between the nitrate and the sulphate; a known weight of the former being in each experiment converted into the latter. One hundred parts of sulphate represent of nitrate:

```
109.312
109.310
109.300
Mean, 109.307, ± .002
```

In all these experiments by Turner the necessary corrections were made for weighing in air.

For the ratio between lead chloride and silver we have a series of results by Marignac and one experiment by Dumas. There are also unavailable data by Turner and by Berzelius.

Marignac,* applying the method used in his researches upon barium and strontium, and working with lead chloride which had been dried at 200°, obtained these results.

^{*} Journ. für Prakt. Chem., 74, 218. 1858.

The third column gives the ratio between PbCl, and 100 parts of Ag:

```
4.9975 grm. PbCl<sub>2</sub> = 3.8810 grm. Ag. 128.768
4.9980 " 3.8835 " 128.698
5.0000 " 3.8835 " 128.750
5.0000 " 3.8860 " 128.667
```

Mean, 128.721, ± .016

Dumas,* in his investigations, found that lead chloride retains traces of water even at 250° , and is sometimes also contaminated with oxychloride. In one estimation 8.700 grammes PbCl, saturated 6.750 of Ag. The chloride contained .009 of impurity; hence, correcting, Ag: PbCl,:: 100: 128.750. If we assign this figure equal weight with those of Marignac, we get as the mean of all, 128.7266, \pm .013. The sources of error indicated by Dumas, if they are really involved in this mean, would tend slightly to raise the atomic weight of lead.

The synthesis of lead nitrate, as carried out by Stas,† gives excellent results. Two series of experiments were made, with from 103 to 250 grammes of lead in each determination. The metal was dissolved in nitric acid, the solution evaporated to dryness with extreme care, and the nitrate weighed. All weighings were reduced to the vacuum standard. In series A the lead nitrate was dried in an air current at a temperature of about 155°. In series B the drying was effected in vacuo. 100 of lead yield of nitrate:

```
A.

159.973

159.975

159.975

159.968

159.973

Mean, 159.9743, ± .0012
```

^{*} Ann. Chem. Pharm., 113, 35. 1860. † Aronstein's Translation, 326.

B. 159.970 159.964 159.959 159.965

Mean, 159.9645, ± .0015

Mean from both series, 159.9704, ± .0010

There still remain to be noticed two sets of experiments upon lead nitrate, which were originally intended to establish the atomic weight of nitrogen. Lead nitrate was carefully ignited and the residual oxide weighed. The first series, bearing Svanberg's name,* gives simply the percentage of oxide found, as follows:

The second series is by Anderson,† and gives the weighings upon which the percentages rest. The latter come out thus:

5.19485 gm	n. PbN ₂ O ₆ gave	3.5017	grm. PbO.	67.4071 p	er cent.
9.7244	"	6.5546	46	67.4037	44
9.2181	44	6.2134	**	67.4044	44
9.6530	44	6.5057	44	67.3957	44

Mean, 67.4027, ± .0016

It will at once be seen that these series are identical; the discordance between the first figures of the two being undoubtedly due to some misprint in the weighings of the Anderson set. How it happens that the same work has been published by two separate authors I will not attempt to explain; neither will I undertake to determine which of the two is really entitled to credit.

^{*} Journ. für Prakt. Chem., 27, 381. 1842. † Ann. Chim. Phys., (3,) 9, 254. 1843.

77

We have now seven ratios upon which to base our computations:

LEAD.

```
(1.) Per cent. of Pb in PbO, 92.8271, ± .0013
(2.) Per cent. of PbO in PbN<sub>2</sub>O<sub>6</sub>, 67.4016, ± .0014
(3.) Pb: PbSO<sub>4</sub>:: 100: 146.4262, ± .0023
(4.) PbO: PbSO<sub>4</sub>:: 100: 135.804, ± .018
(5.) PbSO<sub>4</sub>: PbN<sub>2</sub>O<sub>6</sub>:: 100: 109.307, ± .002
(6.) Pb: PbN<sub>2</sub>O<sub>6</sub>:: 100: 159.9704, ± .0010
(7.) Ag: PbCl<sub>4</sub>:: 100: 128.7266, ± .013
```

Discussing these separately, we get an equal number of values for the atomic weight of lead:

If O = 16, this becomes Pb = 207.079.

In the above discussion are included several values which diverge widely from this general mean, and which, for other reasons, are probably erroneous. Although but one of these carries much weight, it is as well to exclude them, and to base our computations upon the others. If, now, we reject the second, fourth, and fifth values, we get for the atomic weight of lead, Pb = 206.471, $\pm .021$. If O = 16, this becomes Pb = 206.946.

From the synthesis of the nitrate Stas found 206.918, and from the sulphate, 206.934. The agreement of these values with our own general mean is certainly very close.

FLUORINE.

The atomic weight of fluorine has been determined only by one general method, namely, by the conversion of fluorides into sulphates. Excluding the early results of Davy,* we have only to consider the experiments of Berzelius, Louyet, Dumas, and DeLuca, with reference to the fluorides of calcium, sodium, potassium, barium, and lead.

The ratio between calcium fluoride and sulphate has been determined by the four investigators above named, and by one general process. The fluoride is treated with strong sulphuric acid, the resulting sulphate is ignited, and the product weighed. In order to ensure complete transformation special precautions are necessary; such, for instance, as repeated treatment with sulphuric acid, and so on. For details like these the original papers must be consulted.

The first experiments in chronological order are those of Berzelius,† who operated upon an artificial calcium fluoride. He found, in three experiments, for one part of fluoride the following of sulphate:

Louyet's researches‡ were much more elaborate than the foregoing. He began with a remarkably concordant series of results upon fluor spar, in which one gramme of the fluoride yielded from 1.734 to 1.737 of sulphate. At first he regarded these as accurate, but he soon found that particles of spar had been coated with sulphate, and had therefore escaped action. In the following series this source of error

was guarded against.

^{*} Phil. Trans., 1814, 64.

[†] Poggend. Annal., 8, 1. 1826.

[‡] Ann. Chim. Phys., (3,) 25, 300. 1849.

Starting with fluor spar, Louyet found of sulphate as follows:

A second series, upon artificial fluoride, gave:

Dumas* published but one result for calcium fluoride. 495 grm. gave .864 grm. sulphate, the ratio being 1:1.7455.

De Luca† worked with a very pure fluor spar, and published the following results. The ratio between CaSO₄ and one gramme of CaF₂ is given in the third column:

.9305 gm	m. CaF, ga	ive 1.630 gm	m. CaSO ₄ .	1.7518
.836	66	1.459	**	1.7452
.502	46	.8755	66	1.7440
.3985	44	.6945	**	1.7428

If we include Dumas' single result with these, we get a mean of 1.7459, $\pm .0011$.

Upon combining all these series, we get as follows:

Berzelius	1.7500,	± .0004
Louyet, 1st series	1.7437,	± .0003
" 2d "	1.7417,	± .0004
De Luca and Dumas	1.7459,	1100. ±
General mean	1.74493.	+ .0002

For the ratio between the two sodium salts we have experiments by Dumas and by Louyet. According to Louyet one gramme of NaF gives of Na₂SO₄:

^{*} Ann. Chem. Pharm., 113, 28. 1860.

[†] Compt. Rend., 51, 299. 1860.

[‡] See the papers already quoted.

The weighings published by Dumas are as follows:

The general mean of both series is 1.6863, $\pm .0004$.

Dumas also gives experiments upon potassium fluoride. The quantity of sulphate formed from one gramme of fluoride is given in the last column:

The ratios for the fluorides of lead and of barium are due entirely to Louyet. One gramme BaF, gave of BaSO₄:

With the lead fluoride a new method of treatment was adopted. The salt was fused, powdered, dissolved in nitric acid, and precipitated by dilute sulphuric acid. The evaporation of the fluid and the ignition of the sulphate was then effected without transfer. Five grammes of fluoride were taken in each operation, yielding of sulphate:

We now have five ratios to calculate from, as follows:

```
(1.) CaF_1: CaSO_4:: 1.0: 1.74493, \pm .0002
(2.) NaF: Na_2SO_4:: 1.0: 1.6863, \pm .0004
(3.) KF: K_2SO_4:: 1.0: 1.4991, \pm .0007
(4.) BaF_1: BaSO_4:: 1.0: 1.3310, \pm .0004
(5.) PbF_2: PbSO_4:: 5.0: 6.1783, \pm .0002
```

From these we get five values for F:

-		
From	(1) =	18.926, ± .009
44	(2)= =	19.050, ± .014
66	(3) ================================	18.975, ± .032
64	(4)	18.993, ± .033
44	(5) =	19.092, ± .016
	General mean "=	18.084. + .0065

If O = 16, this becomes 19.027.

Before leaving the subject of fluorine we must notice two possible sources of error beyond the always to be considered one of impurities in the materials employed. First, an incomplete conversion of a fluoride into a sulphate would lead to results tending to raise the atomic weight of fluorine. On the other hand, the value for fluorine which has most weight is that derived from calcium fluoride. But it was shown under calcium that the atomic weight determined for that metal was probably a trifle too high. This error, introduced into our fluorine calculations, tends to lower our final results. These two errors, then, if they really exist, will, in part at least, compensate each other.

PHOSPHORUS.

The material from which we are to calculate the atomic weight of phosphorus is by no means abundant. Berzelius, in his Lehrbuch,* adduces only his own experiments upon the precipitation of gold by phosphorus, and ignores all the earlier work relating to the composition of the phosphates. These experiments we will consider with reference to gold.

Pelouze,† in a single titration of phosphorus trichloride with a standard solution of silver, obtained a wholly erroneous result; and Jacquelain,‡ in his similar experiments, did even worse. Schrötter's criticism upon Jacquelain sufficiently disposes of the latter.||

There are, in short, but two investigations upon the atomic weight of phosphorus which have any value for present purposes, namely, the researches of Schrötter and of Dumas. These chemists worked with different materials and by different methods, and yet obtained beautifully concordant results.

Schrötters burned pure amorphous phosphorus in dry oxygen, and weighed the pentoxide thus formed. One gramme of P yielded P₂O₅ in the following proportions:

Hence $P = 30.9562, \pm .0074$.

^{* 5}th Ed., 1188. † Compt. Rend., 20, 1047. ‡ Compt. Rend., 33, 693. || Journ. für Prakt. Chem., 57, 315. ½ Journ. für Prakt. Chem., 53, 435. 1851.

Dumas* prepared pure phosphorus trichloride by the action of dry chlorine upon red phosphorus. The portion used in his experiments boiled between 76° and 78°. This was titrated with a standard solution of silver in the usual manner. Dumas publishes weights, from which I calculate the figures given in the third column, representing the quantity of trichloride proportional to 100 parts of silver:

```
      1.787 grm. PCl<sub>3</sub> = 4.208 grm. Ag.
      42.4667

      1.466 " 3.454 " 42.4435

      2.056 " 4.844 " 42.4443

      2.925 " 6.890 " 42.4528

      3.220 " 7.582 " 42.4690
```

Mean, 42.4553, $\pm .0036$

Hence $P = 31.0314, \pm .0467$.

Now, combining these two values, we have:

If O = 16, this becomes 31.0292.

The fact here noticeable, that Dumas' figures give a value for P slightly higher than that deduced from those of Schrötter, may be accounted for upon the supposition that the phosphorus trichloride contained traces of oxychloride. Such an impurity would tend to raise the apparent atomic weight of phosphorus, and its occurrence is by no means improbable.

^{*} Ann. Chem. Pharm., 113, 29. 1860.

BORON.

The atomic weight of this element has been determined by Berzelius and by Laurent, and calculated by Dumas from some experiments by Deville.

Berzelius* based his determination upon three concordant estimations of the percentage of water in borax. Laurent† made use of two similar estimations, and all five may be properly put in one series, thus:

Hence $B = 10.943, \pm .023$.

Dumas't calculations were based on Deville's analyses of the chloride and bromide of boron, which give the ratios between AgCl and BCl₃, and between AgBr and BBr₃. Reducing the weighings to a common standard, 100 parts of silver chloride correspond to the quantities of boron trichloride given in the third column:

Hence $B = 10.808, \pm .174$.

With the bromide, 2.446 BBr₃ gave 5.496 AgBr. If we assign this experiment equal weight with one in the chloride series, and include the probable error of Br, $\mathbf{B} = 10.964, \pm .364$.

The three values combine as follows:

^{*} Poggend. Annal., 8, 1. 1826. † Journ. für Prakt. Chem., 47, 415. 1849.

[†] Ann. Chem. Pharm., 113, 31. 1860.

From borax	$B = 10.943, \pm .023$
From BCl,	" == 10.808, ± .174
From BBr ₈	" = 10.964, ± .364
General mean	" = 10.041, + .023

If O = 16, B = 10.966.

Further investigation of the atomic weight of boron is evidently desirable.

SILICON.

Although Berzelius* attempted to ascertain the atomic weight of silicon, first by converting pure Si into SiO₂, and later from the analysis of BaSiF₆, his results were not satisfactory. We need only consider the estimations of Pelouze, Schiel, and Dumas.

Pelouze,† experimenting upon silicon tetrachloride, employed his usual method of titration with a solution containing a known weight of silver. One hundred parts of Ag gave the following equivalencies of SiCl₄:

Hence Si = 28.408.

Essentially the same method was adopted by Dumas.‡ Pure SiCl₄ was weighed in a sealed glass bulb, then decomposed by water, and titrated. The results for 100 Ag are given in the third column:

Hence Si = 28.117.

^{*} Lehrbuch, 5 Aufl., 3, 1200.

[†] Compt. Rend., 20, 1047. 1845.

[‡] Ann. Chem. Pharm., 113, 31. 1860.

Dumas and Pelouze's series combine as follows:

General mean	39.4265, ± .0071
Dumas	39·377, ± ·014
Pelouze	$39.4447, \pm .0083$

Hence $SiCl_4 = 169.810, \pm .034$.

Schiel,* also studying the chloride of silicon, decomposed, it by ammonia. After warming and long standing it was filtered, and in the filtrate the chlorine was estimated as AgCl. One hundred parts of AgCl correspond to the quantities of SiCl₄ given in the last column:

Hence $SiCl_4 = 169.437$, $\pm .080$, and Si = 27.957. Combining the values for $SiCl_4$ we have this result:

Hence $Si = 28.195, \pm .066$; or, if O = 16, Si = 28.260.

It will be observed that all of these determinations rest upon the composition of SiCl₄, a compound for which it would not be easy to guarantee absolute purity. All the errors likely to occur in the determination of the atomic weight would be plus errors, so that the value deduced above is almost certainly too high.

^{*} Ann. Chem. Pharm., 120, 94.

LITHIUM.

The earlier determinations of the atomic weight of lithium by Arfvedson, Stromeyer, C. G. Gmelin, and Kralovanzky were all erroneous, because of the presence of sodium compounds in the material employed. The results of Berzelius, Hagen, and Hermann were also incorrect, and need no further notice here. The only investigations which we need to consider are those of Mallet, Diehl, Troost, and Stas.

Mallet's experiments* were conducted upon lithium chloride, which had been purified as completely as possible. In two trials the chloride was precipitated by nitrate of silver, which was collected upon a filter and estimated in the ordinary way. The figures in the third column represent the LiCl proportional to 100 parts of AgCl:

```
7.1885 grm. LiCl gave 24.3086 grm. AgCl. 29.606
8.5947 " 29.0621 " 29.574
```

In a third experiment the LiCl was titrated with a standard solution of silver. 3.9942 grm. LiCl balanced 10.1702 grm. Ag, equivalent to 13.511 grm. AgCl. Hence 100 AgCl = 29.563 LiCl. Mean of all three experiments, 29.581, $\pm .0087$.

Diehl,† whose paper begins with a good resumé of all the earlier determinations, describes experiments made with lithium carbonate. This salt, which was spectroscopically pure, was dried at 130° before weighing. It was then placed in an apparatus from which the carbon dioxide generated by the action of pure sulphuric acid upon it could be expelled, and the loss of weight determined. From this loss the following percentages of CO₂ in Li₂CO₃ were determined:

^{*}Silliman's Amer. Journal, November, 1856. Chem. Gazette, 15, 7.

[†] Ann. Chem. Pharm., 121, 93.

Diehl's investigation was quickly followed by a confirmation from Troost.* This chemist, in an earlier paper,† had sought to fix the atomic weight of lithium by an analysis of the sulphate, and had found a value not far from 6.5; thus confirming the results of Berzelius and of Hagen, who had employed the same method. But Diehl showed that the BaSO₄ precipitated from Li₂SO₄ always retained traces of Li, which were recognizable by spectral analysis, and which accounted for the error. In the later paper Troos made use of the chloride and the carbonate of lithium, both spectroscopically pure. The carbonate was strongly ignited with pure quartz powder, thus losing carbon dioxide, which loss was easily estimated. The subjoined results were obtained:

This combined with Diehl's mean, 59.417, $\pm .006$, gives a general mean of 59.420, $\pm .0057$.

The lithium chloride employed by Troost was heated in a stream of dry hydrochloric acid gas; of which the excess, after cooling, was expelled by a current of dry air. The salt was weighed in the same tube in which the foregoing operations had been performed, and the chlorine was then estimated as silver chloride. The usual ratio between LiCl and 100 parts of AgCl is given in the third column:

This combined with Mallet's mean, 29.581, \pm .0087, gives a general mean of 59.584, \pm .0075.

Finally, we come to the work of Stas,‡ which was exe-

^{*} Zeit. Anal. Chem., 1, 402.

[†] Annales d. Chim. et d. Phys., 51, 108.

[‡] Aronstein's Translation, 279-302.

89

cuted with his usual wonderful accuracy. In three titrations, in which all the weights were reduced to a vacuum standard, the following quantities of LiCl balanced 100 parts of pure silver:

In a second series of experiments, intended for determining the atomic weight of nitrogen, LiCl was converted into LiNO₃. The method was that employed for a similar purpose with the chlorides of sodium and of potassium. One hundred parts of LiCl gave of LiNO₃:

We have now the following ratios from which to deduce the atomic weight of lithium:

```
(1.) AgCl: LiCl:: 100: 29.584, \pm .0075
(2.) Ag: LiCl:: 100: 39.358, \pm .001
(3.) LiCl: LiNO<sub>3</sub>:: 100: 162.5953, \pm .0025
(4.) Per cent. of CO<sub>3</sub> in Li<sub>2</sub>CO<sub>3</sub>, 59.420, \pm .0057
```

Hence two values for the molecular weight of LiCl:

For lithium itself we get three values:

If O = 16, then Li = 7.0235. Stas himself gives 7.022 as his determination. Difference, .0015.

RUBIDIUM.

The atomic weight of rubidium has been determined by Bunsen, Piccard, and Godeffroy; but only from analyses of the chloride.

Bunsen,* employing ordinary gravimetric methods, estimated the ratio between AgCl and RbCl. His rubidium chloride was purified by fractional crystallization of the chloroplatinate. He obtained the following results, to which, in a third column, I add the ratio between RbCl and 100 parts of AgCl:

```
One grm. RbCl gave 1.1873 grm. AgCl. 84.225
" 1.1873 " 84.225
" 1.1850 " 84.388
" 1.1880 " 84.175

Mean, 84.253, ± .031
```

The work of Piccard† was similar to that of Bunsen. In weighing, the crucible containing the silver chloride was balanced by a precisely similar crucible, in order to avoid the correction for displacement of air. The filter was burned separately from the AgCl, as usual; but the small amount of material adhering to the ash was reckoned as metallic silver. The rubidium chloride was purified by Bunsen's method. The results, expressed according to the foregoing standard, are as follows:

```
1.1587 grm. RbCl = 1.372 AgCl + .0019 Ag.
                                             84.300
           66
                  1.6632 "
1.4055
                               .0030 "
                                             84.303
                  1.1850 "
100.1
           "
                               .0024 "
                                             84.245
           66
                   1.7934 "
                               .0018 "
1.5141
                                             84.313
                                       Mean, 84.290, ± .0105
```

Godeffroy,‡ starting with material containing both rubidium and cæsium, separated the two metals by fractional

^{*} Zeit. Anal. Chem., 1, 136. Poggend. Annal., 113, 339. 1861.

[†] Journ. für Prakt. Chem., 86, 454. 1862. Zeit. Anal. Chem., 1, 518.

[‡] Ann. Chem. Pharm., 181, 185. 1876.

CÆSIUM. 91

crystallization of their alums, and obtained salts of each spectroscopically pure. The nitric acid employed was tested for chlorine and found to be free from that impurity, and the weights used were especially verified. In two of his analyses of RbCl the AgCl was handled by the ordinary process of filtration. In the other two it was washed by decantation, dried, and weighed in a glass dish. The usual ratio is appended in the third column:

```
1.4055 grm. RbCl gave 1.6665 grm. AgCl.
1.8096
              44
                       2,1461
                               66
                                           84.320
2.2473
              "
                       2.665
                                 "
                                           84.326
              "
2.273
                       2.6946
                                 44
                                           84.354
                                    Mean, 84.3345, ± .0051
```

Combining the three series, we get the following result:

Hence Rb = 85.251, $\pm .018$. If O = 16, Rb = 85.529.

CÆSIUM.

The atomic weight of cæsium, like that of rubidium, has been determined from the analysis of the chloride. The earliest determination, by Bunsen,* was incorrect, because of impurity in the material employed.

In 1863 Johnson and Allen published their results.† Their material was extracted from the lepidolite of Hebron, Maine, and the cæsium was separated from the rubidium as bitartrate. From the pure cæsium bitartrate cæsium chloride was prepared, and in this the chlorine was estimated as

^{*} Zeit. Anal. Chem., 1, 137.

[†] Amer. Journ. Sci. and Arts, (2,) 35, 94.

silver chloride by the usual gravimetric method. Reducing their results to the convenient standard adopted in preceding chapters, we have, in a third column, the quantities of CsCl equivalent to 100 parts of AgCl:

1.8371 gr	n. CsCl gave	1.5634 g	rm. AgCl.	117.507
2.1295	46	1.8111	46	117.580
2.7018	66	2.2992	44	117.511
1.56165	61	1.3302	"	117.399
			Mea	n, 117.499, ± .025

Shortly after the results of Johnson and Allen appeared a new series of estimations was published by Bunsen.* His cæsium chloride was purified by repeated crystallizations of the chloroplatinate, and the ordinary gravimetric process was employed. The following results represent, respectively, material thrice, four times, and five times purified:

```
1.3835 grm. CsCl gave 1.1781 grm. AgCl. Ratio, 117.435
1.3682 " 1.1644 " 117.503
1.2478 " 1.0623 " 117.462

Mean, 117.467, ± .013
```

Godeffroy's work† was, in its details of manipulation, sufficiently described under rubidium. In three of the experiments upon cæsium the silver chloride was washed by decantation, and in one it was collected upon a filter. The results are subjoined:

```
1.5825 grm. CsCl gave 1.351 grm. AgCl. Ratio, 117.135
1.3487 " 1.1501 " 117.265
1.1880 " 1.0141 " 117.148
1.2309 " 1.051 " 117.107

Mean, 117.164, ± .023
```

We may now combine the three series to form a general mean:

^{*} Poggend. Annal., 119, 1. 1863. † Ann. Chem. Pharm., 181, 185. 1876.

Hence Cs = 132.583, $\pm .024$; or, if O = 16, Cs = 132.918.

THALLIUM.

The atomic weight of this interesting metal has been fixed by the researches of Lamy, Werther, Hebberling, and Crookes. Lamy and Hebberling investigated the chloride and sulphate; Werther studied the iodide; Crooke's experiments involved the synthesis of the nitrate. The last mentioned work was so thorough and admirable that the other researches are included here only for the sake of historical completeness.

Lamy* gives the results of one analysis of thallium sulphate and three of thallium chloride. 3.423 grammes Tl₂SO₄ gave 1.578 grm. BaSO₄; whence 100 parts of the latter are equivalent to 216.920 of the former. In the thallium chloride the chlorine was estimated as silver chloride. The following results were obtained. In the third column I give the amount of TlCl proportional to 100 parts of AgCl:

```
3.912 grm. TlCl gave 2.346 grm. AgCl. 166.752
3.000 " 1.8015 " 166.528
3.912 " 2.336 " 167.466
```

Mean, 166.915, ± .1905

Hebberling's† work resembles that of Lamy. Reducing his weighings to the standards adopted above, we have from his sulphate series, as equivalent to 100 parts of BaSO₄, the amounts of Tl₂SO₄ given in the third column:

^{*} Zeit. Anal. Chem., 2, 211. 1863.

[†] Ann. Chem. Pharm., 134, 11. 1865.

1.4195 g	rm. Tl ₂ SO ₄ gave	.6534 grm. BaSO ₄ .	217.248
1.1924	46	.5507 "	216.5 24
.8560	** •	-3957 "	216.325
			Mean, 216,600

Including Lamy's single result, as of equal weight, we get a mean of $216.754. \pm .1387$.

From the chloride series we have these results, with the ratio stated as usual:

Lamy's mean was 166.915, \pm .1905. Both means combined give a general mean of 166.555, \pm .0865.

Werther's* determinations of iodine in thallium iodide were made by two methods. In the first series TlI was decomposed by zinc and potassium hydroxide, and in the filtrate the iodine was estimated as AgI. One hundred parts of AgI correspond to the amounts of TlI given in the last column:

.720 gm	m. TlI ga	ve .51 gr	m. AgI.	141.176
2.072	44	1.472	46	140.761
.96 0	44	.679	"	141.384
. 385	46	.273	"	141.026
1.068	44	-759	46	140.711
				Mean, 141.012, ± .085

In the second series the thallium iodide was decomposed by ammonia in presence of silver nitrate, and the resulting AgI was weighed. Expressed according to the foregoing standard the results are as follows:

```
1.375 grm. TlI gave .978 grm. AgI. Ratio, 140.593

1.540 " 1.095 " 140.639

1.380 " .981 " 140.673

Mean, 140.635, ± .016
```

General mean of both series, 140.648, $\pm .016$.

^{*} Journ. für Prakt. Chem., 92, 128. 1864.

From the foregoing results three values for the atomic weight of thallium are calculable:

In 1873 Crookes,* the discoverer of thallium, published his final determination of its atomic weight. His method was to effect the synthesis of thallium nitrate from weighed quantities from absolutely pure thallium. No precaution necessary to ensure purity of materials was neglected; the balances were constructed especially for the research; the weights were accurately tested and all their errors ascertained; weighings were made partly in air and partly in vacuo, but all were reduced to absolute standards; and unusually large quantities of thallium were employed in each experiment. In short, no effort was spared to attain as nearly as possible absolute precision of results. The details of the investigation are too voluminous, however, to be cited here; the reader who wishes to become familiar with them must consult the original memoir. Suffice it to say that the research is a model which other chemists will do well to copy.

The results of ten experiments by Professor Crookes may be stated as follows. In a final column we may state the quantity of nitrate producible from 100 parts of thallium. The weights given are in grains:

Thallium.	$TINO_3 + Glass.$	Glass Vessel.		Ratio.
497-9 729 95	1121.851852	472.557319		130.3875
293.193507	1111.387014	729.082713		130.3930
288. 562777	971.214142	594.949719		130.3926
324.963740	1142.569408	718.849078		130.3900
183.790232	1005.779897	766. 133831		130.3912
190.842532	997.334615	748.491271		130.3920
195.544324	1022.176679	767.203451		130.3915
201.816245	1013.480135	750.332401		130.3897
295.683523	1153.947672	768.403621		130.3908
299.203036	1159.870052	769.734201		130.3917
		N	iean,	130.3910, ± .00034

^{*} Philosophical Transactions, 1873, p. 277.

Hence, using the atomic weights and probable errors previously found for N and O, Tl = 203.715, $\pm .0365$. If O = 16, Tl = 204.183.

Crookes himself, using 61.889 as the molecular weight of the group NO₃, gets the value T1 = 203.642; the lowest value in the series being 203.628, and the highest 203.666; an extreme variation of 0.038. This is extraordinary accuracy for so high an atomic weight, at least as far as Crookes' work is concerned. But its value depends in reality upon the accuracy of other chemists in fixing the atomic weights of N and O; a slight variation in either of the latter constants producing a large variation here. What . Crookes really has done has been to fix with almost absolute certainty the ratio between Tl and NO₂. If the latter group should have the molecular weight 62, in accordance with Prout's hypothesis, then Tl = 204.008. words, the ratio thus fixed by Crookes is almost exactly represented by two whole numbers, and supports Prout's hypothesis in a very decided way. Crookes himself seems to have overlooked this fact, for he regards his results as militating against the hypothesis in question.

GLUCINUM.

The atomic weight of glucinum is at present much in doubt; our knowledge of it depending upon the unsettled question whether the oxide is GlO or Gl₂O₃. The formula GlO agrees with Mendelejeff's law, and is advocated by Reynolds,* Lothar Meyer,† and Brauner.‡ The symbol Gl₂O₃, on the other hand, is favored by Nilson and Pettersson,|| and by Humpidge.§ Humpidge, Meyer, and Brauner

^{*} Phil. Mag., (5,) 3, 38. 1877. Chem. News, 42, 273. 1880.

[†] Ber. der Deutsch. Chem. Gesell., 13, 1780. 1880. Also, 11, 576. 1879.

[†] Phil. Mag., (5,) 11. Jan., 1881.

^{||} Berichte, 11, 381 and 906. 1879. Also, 13, 2035. 1880.

[&]amp; Chem. News, 42, 261. 1880.

offer only theoretical discussions of the subject; Reynolds and Nilson and Pettersson have determined the specific heat of the metal, but give opposed results. In the following calculations the simpler formula will be assumed, not as a finality, but because of its accordance with the system of Mendeleieff.

The data from which we are to calculate the atomic weight of glucinum have been determined by Awdejew, Weeren, Klatzo, Debray, and Nilson and Pettersson. Berzelius'* single experiment on the sulphate may be left out of account.

Awdejew,† whose determination was the earliest of any value, analyzed the sulphate. The sulphuric acid was thrown down as barium sulphate; and in the filtrate, from which the excess of barium had been first removed, the glucina was precipitated by ammonia. The figures which Awdejew publishes represent the ratio between SO₃ and GlO, but not absolute weights. As, however, his calculations were made with SO₃ = 501.165, and Ba probably = 855.29, we may add a third column showing how much BaSO₄ is proportional to 100 parts of GlO:

SO ₃ .	G10.	Ratio.
4457	1406	921.242
4531	1420	927.304
7816	2480	915.903
12880	4065	920.814

Mean, 921.316, ± 1.577

The same method was followed by Weeren and by Klatzo, except that Weeren used ammonium sulphide instead of ammonia for the precipitation of the glucina. Weerent gives the following weights of GlO and BaSO₄. The ratio is given in a third column, just as with the figures by Awdejew:

^{*}Poggend. Annal., 8, 1. † " 56, 106. 1842. † " 92, 124. 1854.

G10.	BaSO.	Ratio.	
.3163 grm.	2.9332 grm,	927.031	
.2872 "	2.6377 "	918.419	
.2954 "	2.7342 "	925.592	
.5284 "	4.8823 "	902.946	•
		_	

Mean, 918.497, ± 3.624

Klatzo's* figures are as follows, with the third column added by the writer:

G10.	BaSO.	Ratio.
.2339 grm.	2.1520 grm.	920.052
.1910 "	1.7556 "	919.162
.2673 "	2.4872 "	930.490
.3585 "	3.3115 "	923.710
.2800 "	2.5842 "	922.989
		_

Mean, 923.281, ± 1.346

Combining these series into a general mean, we get the subjoined result:

Hence GlO = 25.224, $\pm .269$.

Debray† analyzed a double oxalate of glucinum and ammonium, $Gl(NH_4)_2C_4O_8$. In this the glucina was estimated by calcination, after first converting the salt into nitrate. The following percentages were found:

The carbon was estimated by an organic combustion. I give the weights, and put in a third column the percentages of CO, thus obtained:

^{*} Zeitschrift für Anal. Chem., 8, 523. 1869.

[†] Ann. de Chim. et de Phys., (3,) 44, 37. 1855.

Salt.	CO_2 .	Per cent. CO2.
.600 grm.	.477 grm.	79.5 0 0
.603 "	.478 "	79.270
.600 "	. •477 "	79.500
		

Mean, 79.423, ± .052

Calculating the ratio between CO, and GlO, we have for the molecular weight of the latter, GlO = 25.220, \pm .180. The agreement between this result and the one previously deduced from the sulphate is certainly very striking.

Last of all and best of all we come to the determinations recently published by Nilson and Pettersson.* These chemists sought to use the sublimed chloride of glucinum, but found it to contain traces of lime derived from a glass tube. They finally resorted to the sulphate as the most available salt for their purposes. This, which they write $Gl_2(SO_4)_3$ 12H₂O, and which we formulate as $GlSO_4.4H_2O$, yields pure glucina upon strong ignition. The subjoined percentages of glucina were thus obtained:

Hence GlO = 25.048, and Gl = 9.085, \pm .0055. If O = 16, Gl = 9.106. If SO₈ = 80, then Gl = 9.096.

If the oxide is Gl₂O₃, then the value Gl = 9.085, $\pm .0055$ becomes Gl = 13.628, $\pm .0082$.

It would be easy enough to combine this value for Gl with those derived from the experiments of the investigators previously cited, but it is hardly worth while. All the other estimations have such high probable errors that they would practically vanish from the general mean. Their influence would hardly extend to the third decimal place, and they may therefore be neglected.

^{*} Compt. Rend., 91, 168. 1880.

MAGNESIUM.

There is perhaps no common metal of which the atomic weight has been subjected to closer scrutiny than that of magnesium. The value is low, and its determination should, therefore, be relatively free from many of the ordinary sources of error; it is extensively applied in chemical analysis, and ought consequently to be accurately ascertained. Strange discrepancies, however, exist between the results obtained by different investigators; so that the generally accepted figure cannot be regarded as absolutely free from doubt.

The determinations of Berzelius* and other early chemists need not be here considered. Nor does the estimation made by Macdonnell† deserve more than a passing mention. He puts the atomic weight of magnesium at 23.9, but gives no details concerning his method of determination. The researches which we have to consider are those of Scheerer, Svanberg and Nordenfeldt, Jacquelain, Bahr, Marchand and Scheerer, and Dumas.

Scheerer's method of investigation was exceedingly simple.‡ He merely estimated the sulphuric acid in anhydrous magnesium sulphate, employing the usual process of precipitation as barium sulphate. He gives no weighings, but reports the percentages of SO_3 thus found. In his calculations, O = 100, $SO_3 = 500.75$, and BaO = 955.29. It is easy, therefore, to recalculate the figures which he gives, so as to establish what his method really represents, viz., the ratio between the sulphates of barium and magnesium.

Thus revised, his four analyses show that 100 parts of MgSO₄ yield the following quantities of BaSO₄:

^{*} Lehrbuch, 5 Aufl., Bd. 3, s. 1227.

[†] British Association Report, 1852, part 2, p. 36.

[†] Poggend. Annal., 69, 535. 1846.

	Per cent. SO3.		
193.575	66.573		
193.677	· 66.608		
193.767	66.639		
193.631	66.592	-1	O JEE
			* *** # * # #

Mean, 193.6625, ± .0274

Hence, using the atomic weights deduced in previous chapters for Ba, S, and O, Mg = 24.544, \pm .0311. In a subsequent note* Scheerer shows that the barium sulphate of the foregoing experiments carried down with it magnesium salts in such quantity as to make the atomic weight of magnesium 0.39 too low. Corrected, Mg becomes = 24.545.

The work of Bahr, of Jacquelain, and in part that of Svanberg and Nordenfeldt, also relates to the composition of magnesium sulphate. Jacquelain's experiments were as follows.† Dry magnesium sulphate was prepared by mixing the ordinary hydrous salt to a paste with sulphuric acid, and calcining the mass in a platinum crucible over a spirit lamp to constant weight and complete neutrality of reaction. This dry sulphate was weighed and intensely ignited three successive times. The weight of the residual MgO having been determined, it was moistened with sulphuric acid and recalcined over a spirit lamp, thus reproducing the original weight of MgSO₄. Jacquelain's weighings for these two experiments show that 100 parts of MgO correspond to the quantities of MgSO₄ given in the last column:

1.466 grm.	MgSO4 gave	.492 grm.	MgO.	297.968
.492 "	MgO "	1.466 "	MgSO ₄ .	297.968

Jacquelain also made one estimation of sulphuric acid in the foregoing sulphate as BaSO₄. His result, (1.464 grm. MgSO₄ = 2.838 grm. BaSO₄,) reduced to the standard adopted in dealing with Scheerer's experiments, give for 100 parts of MgSO₄, 193.852 BaSO₄. If this figure be given equal weight with a single experiment in Scheerer's series,

^{*} Poggend. Annal., 70, 407.

[†] Ann. d. Chim. et Phys., 3 serie, 32, 202.

and combined with the latter, the mean will be 193.700, \pm .0331. From this the atomic weight of magnesium becomes 24.244, \pm .033. This again, corrected according to Scheerer for the magnesium salts carried down by the barium sulphate, becomes 0.39 higher, or Mg = 24.283. Of course this correction, determined by Scheerer for a single experiment, can only be a rough approximation in a mean like the foregoing. It is better than no correction at all, the character of the error involved being known.

Bahr's* work resembles in part that of Jacquelain. This chemist converted pure magnesium oxide into sulphate, and from the increase in weight determined the composition of the latter salt. From his weighings 100 parts of MgO equal the amounts of MgSO₄ given in the third column:

```
1.6938 grm. MgO gave 5.0157 grm. MgSO<sub>4</sub>. 296.122
2.0459 " 6.0648 " 296.437
1.0784 " 3.1925 " 296.040
```

Mean, 296.200, ± .0815

About four years previous to the investigations of Bahr the paper of Svanberg and Nordenfeldt† appeared. These chemists started with the oxalate of magnesium, which was dried at a temperature of from 100° to 105° until it no longer lost weight. The salt then contained two molecules of water, and upon strong ignition it left a residue of MgO. The percentage of MgO in the oxalate comes out as follows:

```
7.2634 grm. oxalate gave 1.9872 grm. oxide. 27.359 per cent.
6.3795 " 1.7464 " 27.375 "
6.3653 " 1.7418 " 27.364 "
6.2216 " 1.7027 " 27.368 "

Mean, 27.3665, ± .0023
```

In three of these experiments the MgO was treated with H₂SO₄, and converted, as by Jacquelain and by Bahr in their later researches, into MgSO₄. One hundred parts of MgO gave of MgSO₄ as follows:

^{*} Journ. für Prakt. Chem., 56, 310. 1852. † Journ. für Prakt. Chem., 45, 473. 1848.

```
1.9872 grm. MgO gave 5.8995 grm. MgSO<sub>4</sub>. 296.875

1.7464 " 5.1783 " 296.513

1.7418 " 5.1666 " 296.624

Mean, 296.671, ± .072
```

We have now for this ratio between MgO and MgSO₄ three series; not at all concordant. We may combine them, assigning to each of Jacquelain's two results a weight corresponding to one of Bahr's:

```
Jacquelain ______ 297.968, ± .0999
Bahr ______ 296.200, ± .0815
Svanberg and Nordenfeldt _____ 296.671, ± .072

General mean _____ 296.806, ± .0475
```

In 1850 the elaborate investigations of Marchand and Scheerer* appeared. These chemists undertook to determine the composition of some natural magnesites, and, by applying corrections for impurities, to deduce from their results the sought for atomic weight. The magnesite chosen for the investigation was, first, a yellow, transparent variety from Snarum; second, a white opaque mineral from the same locality; and, third, a very pure quality from Frank-In each case the impurities were carefully determined; but only a part of the details need be cited here. Silica was of course easily corrected for by simple subtraction from the sum of all of the constituents; but iron and calcium, when found, having been present in the mineral as carbonates, required the assignment to them of a portion of the carbonic acid. In the atomic weight determinations the mineral was first dried at 300°. The loss in weight upon ignition was then carbon dioxide. It was found, however, that even here a correction was necessary. Magnesite, upon drying at 300°, loses a trace of CO2, and still retains a little water; on the other hand, a minute quantity of CO. The CO, expelled at 300° remains even after ignition. amounted in one experiment to .054 per cent.; that retained after calcination to .055 per cent. Both errors tend in the

^{*} Journ. für Prakt. Chem., 50, 385.

same direction, and increase the apparent percentage of MgO in the magnesite. On the yellow mineral from Snarum the crude results are as follows, giving percentages of MgO, FeO, and CO₂ after eliminating silica:

CO ₂ .	$M_{\mathcal{G}}O$.	FeO.
51.8958	47.3278	.7764
51.8798 •	47-3393	.7809
51.8734	47.3154	.8112
51.8875	47.3372	· 775 3
	Mean, 47.3299 ± .0037	

After applying corrections for loss and retention of CO,, as previously indicated, the mean results of the foregoing series become—

The ratio between the MgO and the CO₂, after correcting for the iron, will be considered further on.

Of the white magnesite from Snarum but a single analysis was made, which, for present purposes may be ignored. Concerning the Frankenstein mineral three series of analyses were executed. In the first series the following results were obtained:

8.996 gn	m. CO ₂	= 8.2245 g	rm. MgO.	47.760 pe	r cent. MgO.
7.960	44	7.2775	46	47.761	44
9.3265	"	8.529	44	47.767	46
7.553	46	6.9095	44	47.775	64
				Mean, 47.766, =	± .0022

This mean, corrected for loss of CO, in drying, becomes 47.681. I give series second with corrections applied:

6.8195	grm. MgCO ₃ gave	e 3.2500	grm. MgO.	47.658 p	er cent.
11.3061	"	5.3849	"	47.628	**
9.7375	**	4.635	"	47.599	44
12.3887	44	5.9033	",	47.650	44
32.4148	44	15.453	44	47.674	**
38.8912	44	18.5366	**	47.663	46
26.5223	**	12.6445	44	47.675	44
			Mean,	47.650,	± .0069

The third series was made upon very pure material, so that the corrections, although applied, were less influential. The results were as follows:

4.2913 gr	m.MgCO3 g	gave 2.0436 gi	m. MgO.	47.622]	per cent.
27.8286	46	13.2539	44	47.627	44
14.6192	44	6.9692	46	47.672	"
18.3085	**	8.7237	" .	47.648	44
			Mean	. 47.642.	+ .0077

In a supplementary paper* by Scheerer, it was shown that an important correction to the foregoing data had been overlooked. Scheerer, re-examining the magnesites in question, discovered in them traces of lime, which had escaped notice in the original analyses. With this correction the two magnesites in question exhibit the following mean composition:

	Snarum.	Frankenstein.
CO,	52.131	52.338
MgO	46.663	47-437
CaO	.430	.225
FeO	.776	
	100.000	100.000

Correcting for lime and iron, by assigning each its share of CO_2 , the Snarum magnesite gives as the true percentage of magnesia in pure magnesium carbonate, the figure 47.624. To this, without serious mistake, we may assign the weight indicated by the probable error, \pm .0037; the quantity previously deduced from the percentages of MgO given in the uncorrected analyses.

From the Frankenstein mineral, similarly corrected, the final mean percentage of MgO in MgCO₃ becomes 47.628. This, however, represents three series of analyses, whose combined probable errors may be properly assigned to it. The combination is as follows:

^{*} Ann. d. Chem. und Pharm., 110, 240.

third column:

Result, \pm .0020, probable error of the general mean.

We may now combine the results obtained from both magnesites:

Snarum mineral ____ Per cent. MgO, 47.624, ± .0037
Frankenstein mineral _ " 47.628, ± .0020
General mean _ " 47.627, ± .0018

The last investigation upon the atomic weight of magnesium which we have to consider is that of Dumas.* Pure

magnesium chloride was placed in a boat of platinum, and ignited in a stream of dry hydrochloric acid gas. The excess of the latter having been expelled by a current of dry carbon dioxide, the platinum boat, still warm, was placed in a closed vessel and weighed therein. After weighing, the chloride was dissolved and titrated in the usual manner with a solution containing a known quantity of pure silver. The weighings which Dumas reports give, as proportional to

100 parts of silver, the quantities of MgCl, stated in the

2.203 grm. MgCl₂ = 4.964 grm. Ag. 44.380 5.678 2.5215 " 44.408 2.363 66 5.325 44.376 3.994 9.012 44.319 44 66 5.834 2.578 44.189 2.872 66 66 6. 502 44.171 " " 2.080 4.710 44.161 ** 66 2.214 5.002 44.262 2.086 66 4.722 " 44.176 66 1.688 3.823 44 44.154 44.276 1.342 3.031

There are now before us the following ratios, from which to deduce the sought-for atomic weight:

Mean, 44.261, ± .020

^{*} Ann. Chem. Pharm., 113, 33. 1860.

```
(1.) MgSO<sub>4</sub>: BaSO<sub>4</sub>:: 100: 193.700, ± .0331

(2.) MgO: MgSO<sub>4</sub>:: 100: 296.806, ± .0475

(3.) Per cent. of MgO in oxalate, 27.3665, ± .0023

(4.) Per cent. of MgO in carbonate, 47.627, ± .0018

(5.) Ag: MgCl<sub>2</sub>:: 100: 44.261, ± .020
```

From these we find three values for the molecular weight of MgO:

We have also three values for the atomic weight of magnesium:

Or, if O = 16, Mg becomes = 24.159.

In this general mean all the determinations are included, good or bad. Dumas' result is unquestionably wrong; the error, probably, being due to the presence of oxychloride in the MgCl, which was used. It is doubtful whether any precautions could have eliminated that error. If we take only Marchand and Scheerer's work on magnesium carbonate as having positive value, we shall get from their analyses the following result, viz: Mg = 23.959, \pm .0046. Or, if O = 16, this becomes 24.014. The atomic weight of magnesium, therefore, varies from the whole number 24, only within the ordinary limits of experimental error.

ZINC.

The several determinations of the atomic weight of zinc are by no means closely concordant. The results obtained by Gay-Lussac* and Berzelius† were undoubtedly too low, and may be disregarded here. We need consider only the work done by Jacquelain, Favre, and Axel Erdmann.

In 1842 Jacquelain published the results of his investigations upon this important constant.\(\frac{1}{2}\) In two experiments a weighed quantity of zinc was converted into nitrate, and that by ignition in a platinum crucible was reduced to oxide. In two other experiments sulphuric acid took the place of nitric. As the zinc contained small quantities of lead and iron, these were estimated, and the necessary corrections applied. From the weights of metal and oxide given by Jacquelain the percentages have been calculated:

Nitric Series.

9.917 gr	m. Zn gav	e 12.3138 grm. ZnO.	80.536 pe	r cent. Zn.
9.809	44	12.1800 "	80.534	44
		Sulphuric Series.		
2.398	**	2.978 grm. ZnO.	80.524	44
3.197	**	3.968 "	80.570	44

Mean of all four, 80.541, ± .007

Hence $Zn = 66.072, \pm .028$.

The method adopted by Axel Erdmann || is essentially the same as that of Jacquelain, but varies from the latter in certain important details. First, pure zinc oxide was prepared, ignited in a covered crucible with sugar, and then, to complete the reduction, ignited in a porcelain tube in a current of hydrogen. The pure zinc thus obtained was converted into oxide by means of treatment with nitric acid and sub-

^{*} Mémoire d'Arceuil, 2, 174.

[†] Gilb. Annal., 37, 460.

[‡] Compt. Rend., 14, 636.

^{||} Poggend. Annal., 62, 611. Berz. Lehrb., 3, 1219.

zinc. 109

sequent ignition in a porcelain crucible. Erdmann's figures give us the following percentages of metal in the oxide:

Hence $Zn = 64.9045, \pm .019$.

If we combine the results of Jacquelain with those of Erdmann, we get a mean percentage of zinc, 80.324, \pm .0032; and an atomic weight of Zn = 65.168, \pm .018. The reason for the discordance between the two experimenters will be considered further along.

Favre * employed two methods of investigation. First, zinc was dissolved in sulphuric acid, the hydrogen evolved was burned, and the weight of water thus formed was determined. To his weighings I append the ratio between metallic zinc and 100 parts of water:

Hence $Zn = 65.803, \pm .020$.

The second method adopted by Favre was to burn pure zinc oxalate, and to weigh the oxide and carbonic acid thus produced. From the ratio between these two sets of weights the atomic weight of zinc is easily deducible. From Favre's weighings, if CO₂ = 100, ZnO will be as given in the third column below:

7.796 grm. ZnO = 8.365 grm. CO₂. 93.198
7.342 " 7.883 " 93.137
5.2065 " 5.588 " 93.173
Mean, 93.169,
$$\pm$$
 .012
Hence Zn = 65.8395, \pm .022.

* Ann. Chim. Phys., (3,) 10, 163. 1844.

A fourth combustion of the oxalate is omitted from the above series, having been rejected by Favre himself. In this the oxide formed was contaminated by traces of sulphide.

The four values for zinc now before us are so discordant that a combination of them after the usual method can have only a trifling significance. The following is the result thus obtained:

```
From Jacquelain's figures ... Zn = 66.072, ± .028
From Favre's water series ... " = 65.803, ± .020
From Favre's oxalate series ... " = 65.8395, ± .022
From Erdmann's figures ... " = 64.9045, ± .019

General mean ... " = 65.557, ± .011
```

It will be seen that three of these values agree tolerably well, placing the atomic weight of zinc in the neighborhood of 66, while the other is, in round numbers, about a unit lower. This lower figure, however, has the smallest probable error, and it will be found also, upon careful consideration, that it is less likely than the others to be vitiated by experimental inaccuracies. Both chemically and mathematically it is the best.

Upon comparing Erdmann's results with those of Jacquelain two points are worth noticing: first, Erdmann worked with purer material than Jacquelain, although the latter applied corrections for the impurities which he knew were present; secondly, Erdmann calcined his zinc nitrate in a porcelain crucible, while Jacquelain used platinum. In the latter case it has been shown that portions of zinc may become reduced and alloy themselves with the platinum of the crucible. Hence a lower weight of oxide from a given quantity of zinc, a higher percentage of metal, and an increased atomic weight. This source of constant error has undoubtedly affected Jacquelain's experiments, and vitiated his results. In Erdmann's work no such errors seem to be present.

Over Favre's experiments Erdmann's have the important merit of simplicity. In the latter it is difficult to detect sources of error; in the former it is easy. In Favre's water series it was essential that the hydrogen should first be thoroughly dried before combustion, and then that every trace of water formed should be collected. A trivial loss of hydrogen or of water would tend to increase the apparent atomic weight of zinc.

In the combustion of the zinc oxalate equally great difficulties are encountered. Here a variety of errors are possible, such as are due, for example, to impurity of material, to imperfect drying of the carbon dioxide, and to incomplete collection of the latter. It may not be easy to prove that such errors actually did creep into Favre's work, and yet their possibility hinders us from absolutely accepting his results.

All things considered, then, Erdmann's determination of the atomic weight of zinc is the one most entitled to credit, and must be taken for the present in lieu of the general mean deduced from all four of the values. This determination, Zn = 64.9045, $\pm .019$, becomes, if O = 16, 65.054.

CADMIUM.

The earliest determination of the atomic weight of this metal was by Stromeyer, who found that 100 parts of cadmium united with 14.352 of oxygen.* With our value for the atomic weight of oxygen these figures make Cd = 111.227. This result has now only a historical interest.

The more modern estimates of the atomic weight of cadmium are four in number, by v. Hauer, Lenssen, Dumas, and Huntington. Of these that by v. Hauer† comes first in chronological order. He heated pure anhydrous cadmium sulphate in a stream of dry hydrogen sulphide, and weighed the cadmium sulphide thus obtained. His results

^{*} See Berz. Lehrbuch, 5th Ed., 3, 1219.

⁺ Journ. für Prakt. Chem., 72, 350. 1857.

were as follows, with the percentage of CdS in CdSO, therefrom deduced:

7.7650 gm	. CdSO, gav	ve 5.3741 g	rm. CdS.	69.209	per cent.
6.6086	"	4.5746	66	69.222	44
7.3821	".	5.1117	66	69.245	44
6.8377	44	4.7336	44	69.228	"
8.1956	**	5.6736	44	69.227	66
7.6039	66	5.2634	46	69.220	"
7.1415	66	4.9431	66	69.217	44
5.8245	**	4.0335	"	69.251	44
6.8462	44	4.7415	"	69.257	"

Mean, 69.231, ± .0042

Lenssen* worked upon pure cadmium oxalate, handling. however, only small quantities of material. This salt, upon ignition, leaves the following percentages of oxide:

```
.5128 grm. oxalate gave .3281 grm. CdO.
                                              63.982 per cent.
                       .4193
                                              63.996
.4017
                       .2573
                                              64.053
```

Mean, 64.010, ± .014

Dumas dissolved pure cadmium in hydrochloric acid, evaporated the solution to dryness, and fused the residue in hydrochloric acid gas. The cadmium chloride thus obtained was dissolved in water and titrated with a solution of silver after the usual manner. From Dumas' weighings I calculate the ratio between CdCl, and 100 parts of silver:

2.369 gm	ı. CdCl, =	2.791	rm. Ag.	84.880
4.540	"	5.348	61	84.892
6.177	66	7.260	"	85.803
2.404	16	2.841	44	84.618
3.5325	66	4.166	44	84.794
4.042	"	4.767	`**	84.791
				Mean, 84.843, ± .026

Latest of all comes Huntington's twork, done under the direction of Professor J. P. Cooke. Bromide of cadmium

^{*} Journ. für Prakt. Chem., 79, 281.

[†] Ann. Chem. Pharm., 113, 27. 1860.

[†] Proc. Amer. Acad., 1881.

was prepared by dissolving the carbonate in hydrobromic acid, and the product, dried at 200°, was purified by sublimation in a porcelain tube. Upon the compound thus obtained two series of experiments were made.

In one series the bromide was dissolved in water, and a quantity of silver not quite sufficient for complete precipitation of the bromine was then added in nitric acid solution. After the precipitate had settled, the supernatant liquid was titrated with a standard solution of silver containing one gramme to the litre. The precipitate was washed by decantation, collected by reverse filtration, and weighed. To the weighings I append the ratio between CdBr, and 100 parts of silver bromide:

1.5592 grm	a. CdBr, g	ave 2.1529 gr	m. AgBr.	Ratio, 72.423
* 3.7456	"	5.1724	66	72.415
2.4267	"	3.3511	66	72.415
* 3.6645	"	5.0590	"	72.435
* 3.7679	**	5.2016	"	72.437
2.7938	"	3.8583	66	72.410
* 1.9225	**	2.6552	66	72.405
3-4473	66	4-7593	66	72.433
				Mean. 72 4216

Mean, 72.4216, ± .0028

The second series was like the first, except that the weight of silver needed to effect precipitation was noted, instead of the weight of silver bromide formed. In the experiments marked with an asterisk, both the amount of silver required and the amount of silver bromide thrown down were determined in one set of weighings. The third column gives the CdBr, proportional to 100 parts of silver:

* 3.7456	grm. CdBr ₂ =	2.9715	grm. Ag.	126.051
5.0270	44	3.9874	"	126.072
* 3.6645	**	2.9073	44	126.045
* 3.7679	66	2.9888	44	126.067
* 1.9225	**	1.5248	44	126.082
2.9101	66	2.3079	44	126.093
3.6510	46	2.8951	"	126.110
3.9782	44	3.1551	"	126.088

Mean, 126.076, ± .0052

From the first seriesCdBr, = 271.498,
$$\pm$$
 .032
From the second series ... " = 271.505, \pm .027
General mean ... " = 271.502, \pm .0215

Hence $Cd = 111.966, \pm .043$.

According to Huntington's own calculations these experiments fix the ratio between silver, bromine, and cadmium as Ag: Br: Cd::108:80:112.31. This result militates strongly against Prout's hypothesis.

Upon combining all the determinations we get the following result:

Or, if O = 16, then Cd = 112.092.

It will be seen that Dumas and Huntington's determinations, both made with haloid salts of cadmium, agree with wonderful closeness, and so confirm each other. On the other hand, v. Hauer's data give a value for the atomic weight of cadmium which is much lower. Apparently, v. Hauer's method was good, and the reason for the discrepancy remains to be discovered. Until it is ascertained I prefer to use the above mean value for Cd, rather than to adopt one investigation and reject the others.

MERCURY.

In dealing with the atomic weight of mercury we may reject the early determinations by Sefström* and a large part of the work done by Turner.† The latter chemist, in addition to the data which will be cited below, gives figures

^{*} Sefström. Berz. Lehrb., 5th Ed., 3, 1215. Work done in 1812.

[†] Phil. Trans., 1833, 531-535.

to represent the percentage composition of both the chlorides of mercury; but these results are neither reliable nor in proper shape to be used.

First in order we may consider the percentage composition of mercuric oxide, as established by Turner and by Erdmann and Marchand. In both investigations the oxide was decomposed by heat, and the mercury was accurately weighed. Gold leaf served to collect the last traces of mercurial vapor.

Turner gives four estimations.* Two represent oxide obtained by the ignition of the nitrate, and two are from commercial oxide. In the first two the oxide still contained traces of nitrate, but hardly in weighable proportions. A comparison of the figures from this source with the others is sufficiently conclusive on this point. The third column represents the percentage of mercury in HgO:

```
144.805 grains Hg = 11.54 grains O.
                                          92.619 per cent.
125.980
                  10.08
                                          92.592
            "
                           "
                                                   "
173.561
                   13.82
                                          92.625
            46
                    9.101 "
                                                   "
114.294
                                          92.620
                                    Mean, 92.614, ± .0050
```

In the experiments of Erdmann and Marchand† every precaution was taken to ensure accuracy. Their weighings, reduced to a vacuum standard, give the subjoined percentages:

```
82.0079 grm. HgO gave 75.9347 grm. Hg. 92.594 per cent. 51.0320 " 47.2538 " 92.597 " 84.4996 " 78.2501 " 92.604 " 44.6283 " 41.3285 " 92.606 " 118.4066 " 109.6408 " 92.597 "
```

Mean, 92.5996, ± .0015

Combining, we have:

```
Turner ______ 92.614, ± .0050
Erdmann and Marchand _____ 92.5996, ± .0015
General mean _____ 92.601, ± .0014
```

^{*} Phil. Trans., 1833, 531-535.

[†] Journ. für Prakt. Chem.; 31, 395. 1844.

With a view to establishing the atomic weight of sulphur Erdmann and Marchand also made a series of analyses of pure mercuric sulphide. These data are now best available for discussion under mercury. The sulphide was mixed with pure copper and ignited; mercury distilling over and copper sulphide remaining behind. Gold leaf was used to retain traces of mercurial vapor, and the weighings were reduced to vacuum:

```
34.3568 grm. HgS gave 29.6207 grm. Hg.
                                      86.215 per cent. Hg.
        66
                                      86.206
                                              46
24.8278
               21.40295 "
                                                "
37.2177
            66
                   32.08416
                                      86.207
                    69.6372 · "
                                                "
80.7641
           66
                                      86.223
                                Mean, 86.2127, ± .0027
```

For the percentage of mercury in mercuric chloride we have data by Turner, Millon, and Svanberg. Turner,* in addition to some precipitations of mercuric chloride by silver nitrate, gives two experiments in which the compound was decomposed by pure stannous chloride, and the mercury thus set free was collected and weighed. The results were as follows:

```
44.782 grains Hg = 15.90 grains Cl. 73.798 per cent. 73.09 " 25.97 " \frac{73.784}{\text{Mean}} " Mean, \frac{73.791}{73.791}, \pm .005
```

Millon† purified mercuric chloride by solution in ether and sublimation, and then subjected it to distillation with lime. The mercury was collected as in Erdmann and Marchand's experiments. Percentages of metal as follows:

```
73.87
73.81
73.83
73.87
Mean, 73.845, ± .010
```

Svanberg,‡ following the general method of Erdmann

^{*} Phil. Trans., 1833, 531-525. † Ann. Chim. Phys., (3,) 18, 345 1846. † Journ. für Prakt. Chem., 45, 472. 1848.

and Marchand, made three distillations of mercuric chloride with lime, and got the following results:

```
      12.048 grm. HgCl, gave
      8.889 grm. Hg.
      73.780 per cent.

      12.529 " 9.2456 " 73.794 " 73.810 " 73.810 " 73.795, ± .006

      Mean. 73.795, ± .006
```

Combining these series we have:

General mean	73.798, ± .0034
Svanberg	$73.795, \pm .006$
Millon	$73.845, \pm .010$
Turner	73.791, ± .005

In this mean Turner's figures undoubtedly receive undue weight, for, on experimental grounds, they are probably inferior to both of the other series. It is better, however, that the general mean should remain as it is, than that I should deal arbitrarily with any of the data.

We now have three figures to calculate from:

```
Per cent. of Hg in HgO....... 92.601, ± .0014
" HgS...... 86.2127, ± .0027
" HgCl<sub>2</sub> ..... 73.798, ± .0034
```

These give us three values for the atomic weight of mercury and a general mean as follows:

If O = 16, then this becomes 200.171.

CHROMIUM.

Concerning the atomic weight of chromium there has been much discussion, and many experimenters have sought to establish the true value. The earliest work upon it hav-

ing any importance was that of Berzelius,* in 1818 and 1826, which led to results much in excess of the correct figure. His method consisted in precipitating a known weight of lead nitrate with an alkaline chromate and weighing the lead chromate thus produced. The error in his determination arose from the fact that lead chromate, except when thrown down from very dilute solutions, carries with it minute quantities of alkaline salts, and so has its apparent weight notably increased. When dilute solutions are used, a trace of the precipitate remains dissolved, and the weight obtained is too low. In neither case is the method trustworthy.

In 1844 Berzelius' results were first seriously called in question. The figure for chromium deduced from his experiments was somewhat over 56; but Peligot† now showed, by his analyses of chromous acetate and of the chlorides of chromium, that the true number was near 52.5. Unfortunately, Peligot's work, although good, was published with insufficient details to be useful here. For chromous acetate he gives the percentages of carbon and hydrogen, but not the actual weights of salt, carbon dioxide, and water from which they were calculated. His figures vary considerably moreover; enough to show that their mean would carry but little weight when combined with the more explicit data furnished by other chemists.

Jacquelain's work we may omit entirely. He gives an atomic weight for chromium which is notoriously too low, and prints none of the numerical details upon which his result rests. The researches which particularly command our attention are those of Berlin, Moberg, Lefort, Wildenstein, Kessler, and Siewert.

Among the papers upon the atomic weight under consideration that by Berlin is one of the most important. His starting point was normal silver chromate; but in one ex-

^{*} Schweigg. Journ., 22, 53, and Poggend. Annal., 8, 22.

[†] Compt. Rend., 19, 609 and 734; 20, 1187; 21, 74.

[†] Compt. Rend., 24, 679. 1847.

[|] Journ. für Prakt. Chem., 37, 509, and 38, 149. 1846.

periment the anhydrochromate Ag, Cr, O, was used. These salts, which are easily obtained in a perfectly pure condition, were reduced in a large flask by means of hydrochloric acid and alcohol. The chloride of silver thus formed was washed by decantation, dried, fused, and weighed without transfer. The united washings were supersaturated with ammonia, evaporated to dryness, and the residue treated with hot water. The resulting chromic oxide was then collected upon a filter, dried, ignited, and weighed. The results were as follows:

```
4.6680 grm. Ag<sub>2</sub>CrO<sub>4</sub> gave 4.027 grm. AgCl and 1.0754 grm. Cr<sub>2</sub>O<sub>3</sub>.
                                              **
                                 2.983
                                                              .7960
2.5060
                                  2.1605
                                                  "
                                                              .5770
                                                                             "
                                  1.8555
2.1530
                                                  66
                                                                             "
                                                               .4945
4.3335 grm. Ag<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> gave 2.8692
                                                              1.5300
```

From these weighings three values are calculable for the atomic weight of chromium. The three ratios upon which these values depend we will consider separately; taking first that between the chromic oxide and the original silver salt. In the four analyses of the normal chromate the percentages of Cr₂O₃ deducible from Berlin's weighings are as follows:

And from the single experiment with Ag, Cr, O, the percentage of Cr, O, was 35.306.

For the ratio between Ag₂CrO₄ and AgCl, putting the latter at 100, we have for the former:

In the single experiment with anhydrochromate 100 AgCl is formed from 151.035 Ag, Cr, O₇.

Finally, for the ratio between AgCl and Cr.O., the five experiments of Berlin give, for 100 parts of the former, the following quantities of the latter:

```
26.705
      26.685
      26.707
      26.650
      26.662
Mean, 26.682, ± .0076
```

These results will be discussed in connection with the work of other investigators at the end of this chapter.

In 1848 the researches of Moberg* appeared. His method simply consisted in the ignition of anhydrous chromic sulphate and of ammoniacal chrome alum, and the determination of the amount of chromic oxide thus left as residue. In the sulphate, Cr₂(SO₄), the subjoined percentages of Cr.O. were found. The brackets indicate two different samples of material, to which, however, we are justified in ascribing equal value:

```
.542 grm. sulphate gave .212 grm. Cr2O2.
                                                39.114 per cent.
1.337
              44
                                    "
                          .523
                                                39.117
.5287
               44
                                    ..
                          .207
                                                39.153
                                    "
                                                39.303
1.033
                          .406
                                                39.286
 .868
                          .341
```

Mean, 39.1946, ± .0280

From the alum, (NH₄), Cr, (SO₄), 24H, O, we have these percentages of Cr.O.. The first series represents a salt long dried under a bell jar at a temperature of 18°. The crystals taken were clear and transparent, but may possibly have lost traces of water, which would tend to increase the atomic weight found for chromium. In the second series the salt was carefully dried between folds of filter paper, and

^{*} Journ. für Prakt. Chem., 43, 114.

[†] This objection is suggested by Berlin in a short note upon Lefort's paper. Journ. für Prakt. Chem., 71, 191.

results were obtained quite near those of Berlin. Both of these series are discussed together, neither having a remarkable value:

1.3185	grm. alum	gave .213	grm. Cr ₂ O ₃ .	16.155 p	er cent.	1
.7987	64	.129	66	16.151	64	1
1.0185	44	. 1645	66	16.151	•6	١
1.0206	66	.1650	46	16.167	"	1
.8765	44	.1420	44	16.201	46	I
.7680	44	.1242	66	16.172	44	ì
1.6720	44	.2707	44	16.190	46	1
.5410	46	.0875	44	16.174	46	1
1.2010	44	. 1940	66	16.153	46	l
1.0010	66	.1620	66	16.184	"	1
.7715	46	.1235	66	16.007	44	ì
1.374	44	.2200	66	16.012	"	}

Mean, 16.143, ± .0125

The determinations made by Lefort* are even less valuable than those by Moberg. This chemist started out from pure barium chromate, which, to thoroughly free it from moisture, had been dried for several hours at 250°. The chromate was dissolved in pure nitric acid, the barium thrown down by sulphuric acid, and the precipitate collected upon a filter, dried, ignited, and weighed in the usual manner. The natural objection to the process is that traces of chromium may be carried down with the sulphate, thus increasing its weight. In fact, Lefort's results are somewhat too high. Calculated from his weighings, 100 parts of BaSO₄ correspond to the amounts of BaCrO₄ given in the third column:

1.2615	grm. BaCrO ₄ gave	1.1555	grm. B	aSO ₄ .	109.174
1.5895	44	1.4580	"	1	109.019
2.3255	66	2.1340	"	1	108.974
3.0390	66	2.7855	"	1	109.101
2.3480	44	2.1590	"	1	08.754
1.4230	65	1.3060	14	1	08.708
1.1975	44	1.1005	"	1	08.814
3.4580	44	3.1690	**	1	109.119
2.0130	. .	1.8430	"	1	109.224

^{*} Journ. für Prakt. Chem., 51, 261. 1850.

3.5570 grm	. BaCrO ₄ gave	3.2710 g	rm. BaSO4.	108.744
1.6470	44	1.5060	46	109.363
1.8240	**	1.6725	46	109.058
1.6950	"	1.5560	66	108.933
2.5960	£4	2.3870	44	108.756

Mean, 108.9815, ± .0369

Wildenstein,* in 1853, also made barium chromate the basis of his researches. A known weight of pure barium chloride was precipitated by a neutral alkaline chromate, and the precipitate allowed to settle until the supernatant liquid was perfectly clear. The barium chromate was then collected on a filter, washed with hot water, dried, gently ignited, and weighed. Here again arises the objection that the precipitate may have retained traces of alkaline salts, and again we find deduced an atomic weight which is too high. One hundred parts BaCrO₄ correspond to BaCl₂ as follows:

81.87	81.57
81.So	81.75
81.61	81.66
81.78	81.83
81.52	81.66
81.84	81.80
81.85	81.66
81.70	81.85
81.68	81.57
81.54	81.83
81.66	81.71
81.55	81.63
81.81	81.56
81.86	81.58
81.54	81.67
81.68	81.84

Mean, 81.702, ± .014

Next in order we have to consider two papers by Kessler, who employed a peculiar volumetric method entirely his own. In brief, he compared the oxidizing power of potassium anhydrochromate with that of the chlorate, and from

^{*} Journ. für Prakt. Chem., 59, 27.

his observations deduced the ratio between the molecular weights of the two salts.

In his earlier paper* the mode of procedure was about as follows: The two salts, weighed out in quantities having approximate chemical equivalency, were placed in two small flasks, and to each was added 100 cc. of a ferrous chloride solution and 30 cc. hydrochloric acid. The ferrous chloride was added in trifling excess, and, when action ceased, the amount unoxidized was determined by titration with a standard solution of anhydrochromate. As in each case the quantity of ferrous chloride was the same, it became easy to deduce from the data thus obtained the ratio in question. I have reduced all of his somewhat complicated figures to a simple common standard, and give below the amount of chromate equivalent to 100 of chlorate:

```
120.118
120.371
120.138
120.096
120.241
120.181
Mean, 120.191, ± .028
```

In his later papert Kessler substituted arsenic trioxide for the iron solution. In one series of experiments the quantity of anhydrochromate needed to oxidize 100 parts of the arsenic trioxide was determined, and in another the latter substance was similarly compared with the chlorate. The subjoined columns give the quantity of each salt proportional to 100 of As₂O₃:

$K_2Cr_2O_7$.	KClO ₃ .
98.95	41.156
98.94	41.116
99.17	41.200
98.98	41.255
99.08	41.201
99.15	41.086
	41.199
Mean, 99.045, ± .028	41.224

^{*} Poggend. Annal., 95, 208. 1855. † Poggend. Annal., 113, 137. 1861.

From the data given in the earlier paper, if we use our recent values for chlorine, potassium, and oxygen,

$$K_2Cr_3O_7 = 293.937, \pm .086$$

And from the later, $= 294.159, \pm .119$
General mean, $= 294.013, \pm .0697$

Finally, we come to the determinations published by Siewert,* whose work does not seem to have attracted general attention. He, reviewing Berlin's work, found that upon reducing silver chromate with hydrochloric acid and alcohol, the chromic chloride solution always retained traces of silver chloride dissolved in it. These could be precipitated by dilution with water; but, in Berlin's process, they naturally came down with the chromium hydroxide, making the weight of the latter too high. Hence too large a value for the atomic weight of chromium. In order to find a more correct value Siewert resorted to the analysis of sublimed, violet, chromic chloride. This salt he fused with sodium carbonate and a little nitre, treated the fused mass with water, and precipitated from the resulting solution the chlorine by silver nitrate in presence of nitric acid. The weight of the silver chloride thus obtained, estimated after the usual manner, gave means for calculating the atomic weight of chromium. His figures, reduced to a common standard, give, as proportional to 100 parts of chloride of silver, the quantities of chromic chloride stated in the third of the subjoined columns:

^{*} Zeitschrift Gesammt. Wissenschaften, 17, 530. 1861.

.2367 gm	n. Cr ₂ Cl ₄	gave .6396 gr	m. AgCl.	37.007
.2946	46	.7994	44	36.853
.2593	"	.7039	44	36.838
-4935 ·	"	1.3395	**	36.842
.5850	"	1.5884	66	36.830
.6511	"	1.76681	44	36.852
.5503	"	1.49391	44	36.836

Mean, 36.865, ± .0158

The first of these figures varies so widely from the others that we are justified in rejecting it; in which case the mean becomes $36.842, \pm .0031$.

Siewert also made two analyses of silver anhydrochromate by the following process. The salt, dried at 120°, was dissolved in nitric acid. The silver was then thrown down by hydrochloric acid, and, in the filtrate, chromium hydroxide was precipitated by ammonia. Reduced to a uniform standard, we find from his results, corresponding to 100 parts of AgCl, Ag₂Cr₂O₇, as in the last column:

Giving Berlin's single estimation equal weight with one of these, and combining, we get a general mean of 150.816, \pm .074.

Siewert's percentages of Cr₂O₃ obtained from Ag₂Cr₂O₇, are as follows, calculated from the above weighings.

Combining, as before, with Berlin's single result, giving the latter equal weight with one of these, we have a general mean of 35.236, $\pm .0335$.

For the ratio between silver chloride and chromic oxide, Siewert's two analyses of the anhydrochromate come out as follows. For 100 parts of AgCl we have of Cr₂O₃:

This figure, reduced to the standard of Berlin's work on the monochromate, becomes 26.525, \pm .034. Berlin's mean was 26.682, \pm .0076. The two means, combined, give a general mean of 26.676, \pm .074.

We may now consider the ratios before us, which are as follows:

(I.) Percentage Cr₂O₂ from Ag₂CrO₄, 23.014, ± .011
 (2.) Percentage Cr₂O₃ from Ag₂Cr₂O₇, 35.236, ± .0335
 (3.) AgCl: Ag₄CrO₄:: 100: 115.956, ± .023
 (4.) AgCl: Ag₄Cr₂O₇:: 100: 150.816, ± .074
 (5.) AgCl: Cr₂O₃:: 100: 26.676, ± .0074
 (6.) Percentage Cr₂O₃ in chromic sulphate, 39.1946, ± .0280
 (7.) Percentage Cr₂O₃ in ammonia chrome alum, 16.143, ± .0125
 (8.) BaSO₄: BaCrO₄:: 100: 108.9815, ± .0369
 (9.) BaCrO₄: BaCl₂:: 100: 81.702, ± .014
 (10.) Molecular weight of K₂Cr₂O₇, 294.013, ± .0697
 (11.) AgCl: CrCl₂:: 100: 36.842, ± .0031

From these ratios we can at once deduce five values for the molecular weight of Cr,O₃, as follows:

For barium chromate we get two values:

Finally, from these intermediate data we derive six values for the atomic weight of chromium:

				•
From	BaCrO ₄ C	r =	53.200,	± .064
"	Cr ₂ O ₃	· =	52.482,	810. 土
46	Ag ₂ CrO ₄ "	· ==	52.536,	± .074
6 6	Ag ₂ Cr ₂ O ₇	' =	52.188,	± .109
46	K,Cr,O, "	=	52.116,	± .078
4	CrCl3			
	General mean	' =	52.453,	± .015
	Or, if O = 16	' ==	52.574	

On account of the wide discrepancies between different data, and of the known constant errors vitiating some of the series of experiments, the foregoing general mean can have but little real value. In fact, a careful consideration of all the work represented in it will show that the most accurate estimate of the atomic weight of chromium must be deduced from the experiments of either Berlin, Kessler, or Siewert. Berlin's figures, taken by themselves, and combined, give, if the single analysis of silver anhydrochromate be assigned equal weight with a single analysis in the monochromate series, Cr = 52.389, $\pm .019$; or, if O = 16, Cr = 52.511.

Siewert's results, both for chromic chloride and the silver anhydrochromate, properly combined, give Cr = 52.009, \pm .025. If O = 16, this value becomes Cr = 52.129. In brief, the atomic weight of chromium may be nearly 52.5, or it may be 52. Only a revision of all the experiments could enable us to decide positively between these values. But as Siewert has pointed out probable sources of error in Berlin's work, I am inclined to give preference to the lower value.

MANGANESE.

Rejecting the early experiments of J. Davy and of Arfvedson, the first determinations of the atomic weight of manganese which we encounter are those of Turner* and of Berzelius.† Both of these chemists used the same method.

^{*} Trans. Roy. Soc. Edin., 11, 143. 1831.

[†] Lehrbuch, 5th Ed., 3, 1224.

The chloride of manganese was fused in a current of dry hydrochloric acid, and subsequently precipitated with a solution of silver. From the subjoined weighings I calculate the ratio given in the third column between MnCl, and 100 parts of AgCl:

```
4.20775 grm. MnCl<sub>2</sub> = 9.575 grm. AgCl. 43.945 } Berzelius. 3.063 " = 6.96912 " 43.950 } Berzelius. 12.47 grains MnCl<sub>2</sub> = 28.42 grains AgCl. 43.878—Turner.

Mean, 43.924, ± .015
```

Hence the molecular weight of MnCl, is 125.662, \pm .045.

Many years later Dumas* also made the chloride of manganese the starting point of some atomic weight determinations. The salt was fused in a current of hydrochloric acid, and afterwards titrated with a standard solution of silver in the usual way. 100 parts of Ag are equivalent to the quantities of MnCl, given in the third column:

```
3.3672 grm. MnCl<sub>2</sub> = 5.774 grm. Ag.
                                                  58.317
3.0872
                                                  58.326
                       5.293
2.9671
              "
                       5.0875
                                 "
                                                  58.321
I.I244
              "
                       1.928
                                                  58.320
                       2.251
1.3134
                                                  58.321
```

Mean, 58.321, ± .001

Hence MnCl₂ = 125.594, \pm .011. This, combined with Berzelius and Turner's figures, gives MnCl₂ = 125.598, \pm .011. And Mn = 54.858, \pm .031.

An entirely different method of investigation was followed by v. Hauer,† who, as in the case of cadmium, ignited the sulphate in a stream of sulphuretted hydrogen, and determined the quantity of sulphide thus formed. I subjoin his weighings, and also the percentage of MnS in MnSO₄ as calculated from them:

^{*} Ann. Chem. Pharm., 113, 25. 1860. † Journ. für Prakt. Chem., 72, 360. 1857.

```
57.660 per cent.
4.0626 grm. MnSO4 gave 2.3425 grm. MnS.
              66
                         2.8442
                                               57.613
4.9367
              "
                         3.0192
                                    "
                                                         "
                                               57.649
5.2372
              66
                                    66
                                               57.600
                                                         "
                         4.0347
7.0047
              "
                                    66
                         2.8297
4.9175
                                               57.543
                                    "
                         2.7955
                                               57.585
4.8546
4.9978
                         2.8799
                                               57.625
4.6737
                         2.6934
                                               57.629
                         2.7197
                                               57.572
4.7240
```

Mean, 57.608, ± .008

Hence Mn = 54.785, $\pm .031$.

This method of v. Hauer's, which seemed to give good results with cadmium, is, according to Schneider,* inapplicable to manganese; for the reason that the sulphide of the latter metal is liable to be contaminated with traces of oxysulphide. Such an impurity would bring the atomic weight out too high. The results of two different processes, one carried out by himself and the other in his laboratory by Rawack, are given by Schneider in this paper.

Rawack reduced manganoso-manganic oxide to manganous oxide by ignition in a stream of hydrogen, and weighed the water thus formed. From his weighings I get the values in the third column, which represent the Mn₃O₄ equivalent to one gramme of water:

```
4.149 grm. Mn<sub>2</sub>O<sub>4</sub> gave 0.330 grm. H<sub>2</sub>O.
                                                    12.5727
4.649
                             .370
                                                    12.5643
                                         "
6.8865
                             .5485
                                                    12.5552
                "
                                        "
7.356
                             .5855
                                                    12.5636
                 "
                                         "
                             .7135
                                                    12.5361
8.9445
                **
11.584
                             .9225
                                                    12.5572
```

Mean, 12.5582, ± .0034

Hence $Mn = 53.911, \pm .026$.

Here the most obvious source of error lies in the possible loss of water. Such a loss, however, would increase the apparent atomic weight of manganese; but we see that the value found is much lower than that obtained either by Dumas or v. Hauer.

^{*} Poggend. Annal., 107, 605.

Schneider himself effected the combustion of manganous oxalate with oxide of copper. The salt was not absolutely dry, so that it was necessary to collect both water and carbon dioxide. Then, upon deducting the weight of water from that of the original material, the weight of anhydrous oxalate was easily ascertained. Subtracting from this the CO₂, we get the weight of Mn. If we put CO₂ = 100, the quantities of manganese equivalent to it will be found in the last column:

```
1.5075 grm. oxalate gave .306 grm. H<sub>2</sub>O and .7445 grm. CO<sub>2</sub>. 61.3835
2.253
                          -4555
                                    66
                                             1.1135
                                                                61.4291
               "
                                     66
3.1935
                         .652
                                             1.5745
                                                                61.4163
               46
                                     **
                         1.028
                                             2.507
                                                                61.3482
5.073
                                                        Mean, 61.3943, \pm .0122
```

Hence $Mn = 53.904, \pm .014$.

This result agrees beautifully with the value calculated from Rawack's experiments.

Now to combine the four independent values which we have thus far obtained:

The considerations already cited, however, go to show that this general mean must be slightly affected by some plus constant error. It is probable, therefore, that a more correct figure will result from rejecting the first and second values in the above combination, and taking the data furnished by Rawack and Schneider alone. Combining their figures, we get as follows. Mn = 53.906, \pm .012. Or, if O = 16, Mn = 54.029.

Since the foregoing calculations were made Dewar and Scott* have reported the following experiments. From the

^{*} Nature, Sept. 15, 1881, p. 470.

complete analysis of silver permanganate, putting Ag = 108 and O = 16, they find in three estimations Mn = 55.51, 54.04, and 54.45. From the analysis of pure MnO_2 , made from the nitrate, Mn = 53.3 to 53.6. Up to the date of writing a detailed account of the methods employed has not been published.

IRON.

The atomic weight of iron has been determined almost exclusively from the composition of ferric oxide. Beyond this there are only a few comparatively unimportant experiments by Dumas relative to ferrous and ferric chlorides.

Most of the earlier data relative to the percentage of metal and oxygen in ferric oxide we may reject at once, as set aside by later investigations. Among this no longer valuable material there is a series of experiments by Berzelius, another by Döbereiner, and a third by Capitaine. The work done by Stromeyer and by Wackenroder was probably good, but I am unable to find its details. The former found 30.15 per cent. of oxygen in the oxide under consideration, while Wackenroder obtained figures ranging from a minimum of 30.01 to a maximum of 30.38 per cent.*

In 1844 Berzelius \dagger published two determinations of the ratio in question. He oxidized iron by means of nitric acid, and weighed the oxide thus formed. He thus found that when O = 100 Fe = 350.27 and 350.369.

Hence the following percentages of Fe in Fe,O,.

70.018 70.022 Mean, 70.020, ± .0013

About the same time Svanberg and Norlin; published

^{*} For additional details concerning these earlier papers I must refer to Oudemans' monograph, pp. 140, 141.

[†] Ann. Chem. Pharm., 50, 432. Berz. Jahresb., 25, 43.

[†] Berzelius' Jahresbericht, 25, 42.

two elaborate series of experiments; one relating to the synthesis of ferric oxide, the other to its reduction. In the first set pure piano-forte wire was oxidized by nitric acid, and the amount of oxide thus formed was determined. The results were as follows:

1.5257 grm	ı. Fe gave	2.1803 g	rm. Fe ₂ O ₃ .	69.977 pe	er cent. Fe.
2.4051	"	3.4390	44	69.936	46
2.3212	"	3.3194	"	69.928	66
2.32175	"	3.3183	£6	69.968	66
2.2772	"	3.2550	66	69.960	64
2.4782	"	3.5418	"	69.970	66
2.3582	46	3.3720	"	69.935	44

Mean, 69.9534, ± .0050

In the second series ferric oxide was reduced by ignition in a current of hydrogen, yielding the subjoined percentages of metal:

```
2.98353 grm. Fe<sub>2</sub>O<sub>3</sub> gave 2.08915 grm. Fe.
                                               70.025 per cent.
2.41515
                         1.6910
                                               70.015
                "
                          2.09455
                                               70.014
2.99175
                "
                                      "
3.5783
                          2.505925
                                               70.030
4.1922
                          2.9375
                                               70.072
                                               70.056
3.1015
                          2.17275
2.6886
                          1.88305
                                               70.036
```

Mean, 70.0354, ± .0055

It is evident that one or both of these series must be vitiated by constant errors, and that these probably arise from impurities in the materials employed. Impurities in the wire taken for the oxidation series could hardly have been altogether avoided, and in the reduction series it is possible that weighable traces of hydrogen may have been retained by the iron. At all events it is probable that the errors of both series are in contrary directions, and, therefore, in some measure compensatory.

In 1844 there was also published an important paper by Erdmann and Marchand.* These chemists prepared ferric oxide by the ignition of pure ferrous oxalate, and submitted

^{*} Journ. für Prakt. Chem., 33, 1.

133

it to reduction in a stream of hydrogen. Two sets of results were obtained with two different samples of ferrous oxalate, prepared by two different methods. For present purposes, however, it is not necessary to discuss these sets separately. The percentages of iron in Fe_2O_3 come out as follows:

```
70.013
69.962
69.979
70.030
69.977
70.044
70.015
70.055
```

Mean, 70.0094, ± .0080

In 1850 Maumené's* results appeared. He dissolved pure iron wire in aqua regia, precipitated with ammonia, filtered off the precipitate, washed thoroughly, ignited, and weighed, after the usual methods of quantitative analysis. The percentages of Fe in Fe₂O₃ are given in the third column:

1.482 grm. Fe gave 2.117 grm. Fe ₂ O ₃ .			70.005 per cent.		
1.452	**	2.074	66	70.010	44
1.3585	"	1.941	46	69.990	44
1.420	46	2.0285	66	70.002	44
1.492	"	2.1315	**	69.998	"
1.554	"	2.220	"	70.000	"

Mean, 70.0008, ± .0019

Two more results, obtained by Rivot† through the reduction of ferric oxide in hydrogen, remain to be noticed. The percentages are:

We have thus before us six series of results, which we may now combine.

^{*} Compt. Rend., Oct. 17, 1850. † Ann. Chem. Pharm., 78, 214. 1851.

Berzelius	70.020,	± .0013
Erdmann and Marchand	•	_
Svanberg and Norlin, Oxyd	69.9534,	± .0050
" Reduc	70.0354,	±.0055
Maumené	70.0008,	e100. ±
Rivot	69.33,	± .013
General mean	70.0075.	0100. ±

From this we get Fe = 55.891, $\pm .012$; or, if O = 16, this becomes 56.0195.

Dumas'* results, obtained from the chlorides of iron, are of so little weight that they might safely be omitted from our present discussion. For the sake of completeness, however, we will include them.

Pure ferrous chloride, ignited in a stream of hydrochloric acid gas, was dissolved in water and titrated with a silver solution in the usual way. One hundred parts of silver are equivalent to the amounts of FeCl, given in the third column:

Ferric chloride, titrated in the same way, gave these results:

These give us two additional values for Fe, as follows:

Combining these with the value deduced from the composition of Fe₂O₃, Fe = 55.891, \pm .012, we get this general mean, Fe = 55.913, \pm .012. If O = 16, this becomes Fe = 56.042.

^{*} Ann. Chem. Pharm., 113, 26. 1860.

COPPER.

The atomic weight of copper has been chiefly determined from the composition of the black oxide and the anhydrous sulphate. In dealing with the first named compound all experimenters have agreed in reducing it with a current of hydrogen, and weighing the metal thus set free.

The earliest experiments of any value were those of Berzelius,* whose results were as follows:

```
7.68075 grm. CuO lost 1.55 grm. O. 79.820 per cent. Cu in CuO. 9.6115 " 1.939 " 79.826 " " Mean, 79.823, ± .002
```

Erdmann and Marchand,† who come next in chronological order, corrected their results for weighing in air. Their weighings, thus corrected, give us the subjoined percentages of metal in CuO:

```
63.8962 grm. CuO gave 51.0391 grm. Cu. 79.878 per cent.
65.1590 " 52.0363 " 79.860 "
60.2878 " 48.1540 " 79.874 "
46.2700 " 36.9449 " 79.846 "

Mean, 79.8645, ± .0038
```

Still later we find a few analyses by Millon and Commaille.‡ These chemists not only reduced the oxide by hydrogen, but they also weighed, in addition to the metallic copper, the water formed in the experiments. In three determinations the results were as follows:

```
6.7145 grm. CuO gave 5.3565 grm. Cu and 1.5325 grm. H<sub>2</sub>O. 79.775 per cent.
3.3945 " 2.7085 " .7680 " 79.791 "
2.2240 grm. Cu. 79.770 "

Mean, 79.7787, ± .0043
```

For the third of these analyses the water estimation was not made, but for the other two it yielded results which, in

^{*} Poggend. Annal., 8, 177.

[†] Journ. für Prakt. Chem., 31, 389. 1844.

[‡] Fresenius' Zeitschrift, 2, 475. 1863.

the mean, would make the atomic weight of copper 63.087, \pm .222. This figure has so high a probable error that we need not consider it further.

The results obtained by Dumas* are wholly unavailable. Indeed, he does not even publish them in detail. He merely says that he reduced copper oxide, and also effected the synthesis of the subsulphide, but without getting figures which were wholly concordant. He puts Cu = 63.5.

Latest of all, and probably the best also, we have the determinations by Hampe.† First, he attempted to estimate the atomic weight of copper by the quantity of silver which the pure metal could precipitate from its solutions. This attempt failed to give satisfactory results, and he fell back upon the old method of reducing the oxide. From ten to twenty grammes of material were taken in each experiment, and the weights were reduced to a vacuum standard:

```
20.3260 grm. CuO gave· 16.2279 grm. Cu. 79.838 per cent.
20.68851 " 16.51669 " 79.835 "
10.10793 " 8.06926 " 79.831 "

Mean, 79.8347, ± .0013
```

Hampe also determined the quantity of copper in the anhydrous sulphate, CuSO₄. From 40 to 45 grammes of the salt were taken at a time, the metal was thrown down by electrolysis, and the weights were all corrected. I subjoin the results:

```
40.40300 grm. CuSO<sub>4</sub> gave 16.04958 grm. Cu. 39.724 per cent.
44.64280 " 17.73466 " 39.726 "

Mean, 39.725, ± .0007
```

We now have four series of experiments upon copper oxide, as follows:

```
      Berzelius
      79.823, ± .0020

      Erdmann and Marchand
      79.8645, ± .0038

      Millon and Commaille
      79.7787, ± .0043

      Hampe
      79.8347, ± .0013

      General mean
      79.830, ± .0010
```

^{*} Ann. d. Chim. et Phys., (3,) 55, 129.

[†] Fresenius' Zeitschrift, 13, 352.

For copper we have-

```
From composition of CuO....Cu = 63.181, ± .036

" CuSO<sub>4</sub>, (Hampe) ..... " = 63.171, ± .012

General mean ..... " = 63.173, ± .011
```

If O = 16, then Cu becomes = 63.318.

The close agreement between the two independent values for Cu is certainly very striking. It will be seen that Hampe's two estimates upon the sulphate carry (perhaps accidentally) much greater weight than all the experiments upon the oxide. This might seem like giving them undue credit, were it not for the fact of the remarkable concordance of the results above referred to. Either estimate for Cu would be valid without the other.

MOLYBDENUM.

If we leave out of account the inaccurate determination made by Berzelius,* we shall find that the data for the atomic weight of molybdenum lead to two independent estimates of its value; one near 92, the other near 96. The earlier results found by Berlin and by Svanberg and Struve lead to the lower number; the more recent work of Debray, Dumas, and Lothar Meyer sustains the higher. The latter value is the more probable, although both may be vitiated by constant errors in opposite directions.

The earliest investigation which we need especially to consider is that of Svanberg and Struve.† These chemists tried a variety of different methods, but finally based their conclusions upon the two following: first, molybdenum trioxide was fused with potassium carbonate, and the carbon dioxide which was expelled was estimated; secondly, molybdenum disulphide was converted into the trioxide by

^{*} Poggend. Annal., 8, 1. 1826.

[†] Journ. für Prakt. Chem., 44, 301. 1848.

roasting, and the ratio between the weights of the two substances was determined.

By the first method it was found that 100 parts of MoO, will expel the following quantities of CO_2 :

The carbon dioxide was determined simply from the loss of weight when the weighed quantities of trioxide and carbonate were fused together. It is plain that if, under these circumstances, a little of the trioxide should be volatilized, the total loss of weight would be slightly increased. A constant error of this kind would tend to bring out the atomic weight of molybdenum too low.

By the second method, the conversion by roasting of MoS, into MoO_s, Svanberg and Struve obtained these results. Two samples of artificial disulphide were taken, A and B, and yielded for each hundred parts the following of trioxide:

Three other experiments in series B gave divergent results, and, although published, are rejected by the authors themselves. Hence it is not necessary to cite them in this discussion. We again encounter in these figures the same source of constant error which apparently vitiates the preceding series, namely, the possible volatilization of the trioxide. Here, also, such an error would tend to reduce the atomic weight of molybdenum.

Upon discussing the data given in the foregoing para-

graphs we get somewhat noticeable results. From the carbon dioxide series, Mo = 91.711, $\pm .113$, a figure having no unusual interest. From the other series, if S = 31.987 and O = 15.9633, we get Mo = 92.979, $\pm .354$; but if we take S = 32 and O = 16, then Mo becomes = 92.133. In this case the higher values for oxygen and sulphur lead to a lower number for molybdenum. In the carbonate series the assumption of 12 and 16 for C and O, respectively, makes Mo = 92.033. In other words, if we assume the ordinary even numbers for C, O, and S, C synberg and C struve's two methods yield more nearly concordant results than when the revised values for these elements are taken.

Berlin,* a little later than Svanberg and Struve, determined the atomic weight of molybdenum by igniting a molybdate of ammonium and weighing the residual MoO₃. Here, again, a loss of the latter by volatilization may (and probably does) lead to too low a result. The salt used was $(NH_4)_4 Mo_5 O_{17}$. 3 $H_2 O$, and in it these percentages of MoO₃ were found:

81.598 81.612 81.558 81.555

Mean, 81.581, ± .0095

Hence Mo = 91.9817, $\pm .0776$; a result agreeing quite well with those of Svanberg and Struve.

Until 1859 the value 92 was generally accepted on the basis of the foregoing researches, but in this year Dumas† published some figures tending to sustain a higher number. He prepared molybdenum trioxide by roasting the disulphide, and then reduced it to metal by ignition in hydrogen. At the beginning the hydrogen was allowed to act at a comparatively low temperature, in order to avoid volatilization of trioxide; but at the end of the operation the heat

^{*} Journ. für Prakt. Chem., 49, 444. 1850.

[†] Ann. Chem. Pharm., 105, 84, and 113, 23.

was raised sufficiently to insure a complete reduction. From the weighings I calculate the percentages of metal in MoO₂:

.448 gm	n. MoO _a ga	ve .299 g	rm. Mo.	66.741 p	er cent.
.484	"	.323	66	66.736	64
.484	**	.322	66	66.529	66
.498	66	.332	44	66.667	66
-559	66	-373	44	66 .726	44
.388	66	.258	"	66.495	44
				Mean, 66.649,	± .030

In 1868 the same method was employed by Debray.* His trioxide was purified by sublimation in a platinum tube. His percentages are as follows:

5.514 grm	. MoO ₂ gave	3.667	grm.	Mo.		66.503 p	er cent.
7.910	44	5.265	"			66.561	66
9.031	44	6.015	44			66.604	44
					Mean,	66. \$56,	± .020

This mean, combined with that of Dumas', gives a general mean of 66.585, $\pm .017$.

Hence Mo = 95.429, $\pm .057$.

Debray also made two experiments upon the precipitation of molybdenum trioxide in ammoniacal solution by nitrate of silver. In his results, as published, there is curious discrepancy, which, I have no doubt, is due to typographical error. These results I am, therefore, compelled to leave out of consideration. They could not, however, exert a very profound influence upon the final discussion.

The most recent investigation upon the atomic weight of molybdenum is the discussion by Lothar Meyer† of the experimental results obtained by Liechti and Kemp‡ in their analyses of the chlorides. Of these compounds there are four: MoCl₂, MoCl₂, MoCl₄, and MoCl₅. The chlorine in each was estimated as silver chloride, and the molybdenum as disulphide. From these analyses Meyer deduces three

^{*} Compt. Rend., 66, 734.

[†] Ann. Chem. Pharm., 169, 365. 1873.

[†] Ann. Chem. Pharm., 169, 344.

sets of ratios, namely: between MoCl_n and n AgCl; between MoCl_n and MoS₂, and between MoS₂ and n AgCl. We will use only the first and last of these; the probable error of the atomic weight deduced from the second being relatively so high as to make the value connected with it comparatively unimportant. The analyses of the trichloride, being discordant, are here rejected.

By reducing the weighings published by Liechti and Kemp* to a common standard we get the following percentage results. In MoCl, the subjoined quantities of the original substance and of MoS, correspond to 100 parts of AgCl:

MoCl ₂ .	MoS_2 .
58.299	55.762
58. 194	55-591
58.524	56.065
	
Mean, 58,330, + .066	Mean, 55.806, + .00

Hence MoCl₂ = 166.902, \pm .188, and MoS₂ = 159.652, \pm .268.

With the tetrachloride similarly calculated we get these figures, corresponding to 100 parts AgCl:

$$MoCl_4$$
. MoS_4 .

41.492 27.957

41.319

Mean, 41.4055, \pm .0583

Hence $MoCl_4 = 236.914$, $\pm .358$, and MoS_2 , if given the weight of a single experiment in the dichloride series, = 159.964, $\pm .627$.

.5810 grm. MoCl₅ gave .3414

66

66

66

1.5222

.6465

^{*}These are as follows:

.2666 grm. MoCl₁ gave .2550 grm. MoS₁ and .4573 grm. AgCl.
.1811 " .1730 " .3112 "

.2530 " .2422 " .4320 "

.4126 grm. MoCl₄ gave .2780 " .9944 "

.1923 " — " .4654 "

For the pentachloride the following quantities balance 100 of AgCl:

Hence MoCl₅ = 272.587, $\pm .271$, and MoS₂ = 159.914, $\pm .287$.

We have now the molecular weight of each chloride, and three values for that of the disulphide. Combining the latter we get a general mean, as follows:

With these data, in addition to those given by Dumas and by Debray, we get five estimates of the atomic weight of molybdenum:

Or, if O = 16, Mo = 95.747.

It will at once be seen that the most reliable results are those obtained by the reduction of molybdenum trioxide. Traces of oxychlorides may possibly have contaminated the chlorides and augmented their atomic weight. Our final figure, therefore, may be a trifle too high, but the early value, 92, is unquestionably very far too low.

Since the foregoing discussion was written a single experiment by Rammelsberg * has been brought to my netice.

^{*} Berlin Monatsbericht, 1877, 574.

Closely following Dumas' method, he reduced molybdenum trioxide to metal, finding in it 66.708 per cent. of the latter. This figure comes within the limits of variation of Dumas' experiments, and therefore gives them additional confirmation. Its introduction into the general mean, however, would exert too little influence upon the latter to justify the labor of recalculation.

TUNGSTEN.

The atomic weight of tungsten has been determined from analyses of the trioxide, the hexchloride, and the tungstates of iron, silver, and barium.

The composition of the trioxide has been the subject of many investigations. Malaguti* reduced this substance to the blue oxide, and from the difference between the weights of the two compounds obtained a result now known to be considerably too high. In general, however, the method of investigation has been to reduce WO₃ to W in a stream of hydrogen at a white heat, and afterwards to reoxidize the metal, thus getting from one sample of material two results for the percentage of tungsten. This method is unquestionably accurate, provided that the trioxide used be pure.

The first experiments which we need consider are, as usual, those of Berzelius.† 899 parts WO₃ gave, on reduction, 716 of metal. 676 of metal, reoxidized, gave 846 WO₃. Hence these percentages of W in WO₃:

79.644, by reduction. 79.905, by oxidation.

Mean, 79.7745, ± .0880

These figures are far too high, the error being undoubtedly due to the presence of alkaline impurity in the trioxide employed.

^{*} Journ. für Prakt. Chem., 8, 179. 1836.

[†] Poggend. Annal., 8, 1. 1826.

Next in order of time comes the work of Schneider,* who, with characteristic carefulness, took every precaution to get pure material. His percentages of tungsten are as follows:

```
Reduction Series.

79.336

79.254

79.312

79.326

79.350

Mean, 79.3156, ± .0112

Oxidation Series.

79.329

79.324

79.328

Mean, 79.327, ± .0010
```

Closely agreeing with these figures are those of Marchand.†
published in the following year:

Reduction Series.

```
79.307

79.302

Mean, 79.3045, .0017

Oxidation Series.

79.321

79.352

Mean, 79.3365, ± .0105
```

The figures obtained by v. Borch‡ agree in mean tolerably well with the foregoing. They are as follows:

```
Reduction Series.

79.310

79.212

79.289

79.313

79.225

79.290

79.302

Mean, 79.277, ± .0106
```

^{*} Journ. für Prakt. Chem., 50, 152. 1850. † Ann. Chem. Pharm., 77, 261. 1851.

[‡] Journ. für Prakt. Chem., 54, 254. 1851.

Oxidation Series. 79.359

79·359 79·339

Mean, 79.349, ± .0067

Dumas* gives only a reduction series, based upon trioxide obtained by the ignition of a pure ammonium tungsten. The reduction was effected in a porcelain boat, platinum being objectionable on account of the tendency of tungstate to alloy with it. Dumas publishes only weighings, from which I have calculated the percentages:

2.784	grm. WO ₃ gave	2.208	grm.	W.	79.310 per cent.
2.994	46	2.373	"		79.259 "
4.600	4	3.649	"		79.3 2 6 "
.985	"	.781	"		79.289 "
.917	64	.727	66		79.280 "
.917	44	.728	"		79.389 "
1.717	44	1.362	66		79.324 "
2.988	46	2.370	"		79.317 "

Mean, 79.312, ± .009

The data furnished by Bernoulli† differ widely from those just given. This chemist undoubtedly worked with impure material, the trioxide having a greenish tinge. Hence the results are too high. These are the percentages of W:

```
Reduction Series.

79.556

79.526

79.553

79.558

79.549

78.736

Mean, 79.413, ± .091

Oxidation Series.

79.558

79.656

79.555

79.554
```

Mean, 79.581, ± .017

^{*} Ann. Chem. Pharm., 113, 23. 1860. † Poggend. Annal., 111, 573. 1860.

Two reduction experiments by Persoz* give the following results:

Finally, we have the work done by Roscoe.† This chemist used a porcelain boat and tube, and made six weighings, after successive reductions and oxidations, with the same sample of 7.884 grammes of trioxide. These weighings give me the following five percentages, which, for the sake of uniformity with foregoing series, I have classified under the usual, separate headings:

```
Reduction Series.

79.196
79.285
79.308

Mean, 79.263, ± .023

Oxidation Series.
79.230
79.299

Mean, 79.2645, ± .0233
```

There are still other experiments by Riche,‡ which I have not been able to get in detail. They cannot be of any value, however, for they give to tungsten an atomic weight of about ten units too low. We may therefore neglect this series, and go on to combine the others:

^{. *} Zeit. Anal. Chem., 3, 260. 1864. † Ann. Chem. Pharm., 162, 368. 1872.

[†] Journ. für Prakt. Chem., 69, 10. 1857.

TUNGSTEN.

Dumas	79.312,	± .009
Bernouilli, Reduction		
" Oxidation	79.581,	土 .017
Persoz	79.314,	± .007
Roscoe, Reduction	79.263,	± .023
" Oxidation	79.2645,	± .0233
General mean	79.3215,	±.00085

The rejection of the figures given by Berzelius and by Bernoulli exerts an unimportant influence upon the final result. There is, therefore, no practical objection to retaining them in the discussion.

In 1861 Scheibler* deduced the atomic weight of tungsten from analyses of barium metatungstate, BaO.4 WO₁.9 H₂O. In four experiments he estimated the barium as sulphate, getting closely concordant results, which were, however, very far too low. These, therefore, are rejected. But from the percentage of water in the salt a very good result was attained. The percentages of water are as follows:

13.053 13.054 13.045 13.010 13.022 Mean, 13.0368, ± .0060

The work of Zettnow,† published in 1867, was somewhat more complicated than any of the foregoing researches. He prepared the pure tungstates of silver and of iron, and from their composition determined the atomic weight of tungsten.

In the case of the iron salt the method of working was this: The pure, artificial FeWO, was fused with sodium carbonate, the resulting sodium tungstate was extracted by water, and the thoroughly washed, residual ferric oxide was dissolved in hydrochloric acid. This solution was then reduced by zinc, and titrated for iron with potassium permanganate. Corrections were applied for the drop in excess of

^{*} Journ. für Prakt. Chem., 83, 324.

[†] Poggend. Annal.. 130, 30.

permanganate needed to produce distinct reddening, and for the iron contained in the zinc. 11.956 grammes of the latter metal contained iron corresponding to 0.6 cc. of the standard solution. The permanganate was standardized by comparison with pure ammonium-ferrous sulphate, Am, Fe(SO₄)₂. 6 H,O, so that, in point of fact, Zettnow establishes directly only the ratio between that salt and the From Zettnow's four experiments in ferrous tungstate. standardizing I find that 1 cc. of his solution corresponds to 0.0365457 grammes of the double sulphate, with a probable error of $\pm .0000012$.

Three sets of titrations were made. In the first a quantity of ferrous tungstate was treated according to the process given above; the iron solution was diluted to 500 cc., and four titrations made upon 100 cc. at a time. The second set was like the first, except that three titrations were made with 100 cc. each, and a fourth upon 150 cc. In the third set the iron solution was diluted to 300 cc., and only two titrations upon 100 cc. each were made. In sets one and two thirty grammes of zinc were used for the reduction of each, while in number three but twenty grammes were taken. Zettnow's figures, as given by him, are quite complicated; therefore I have reduced them to a common standard. After applying all corrections the following quantities of tungstate, in grammes, correspond to 1 cc. of permanganate solution:

Mean, .0283549, ± .0000115

With the silver tungstate, Ag, WO, Zettnow employed two methods. In two experiments the substance was decomposed by nitric acid, and the silver thus taken into solution was titrated with standard sodium chloride. In three others the tungstate was treated directly with common salt, and the residual silver chloride collected and weighed. Here again, on account of some complexity in Zettnow's figures, I am compelled to reduce his data to a common standard. To 100 parts of AgCl the following quantities of Ag2WO4 correspond:

By First Method.

161.665

161.603

Mean, 161.634, ± .021

By Second Method.

161.687

161.651

161.613

Mean, 161.650, ± .014

General mean from both series, 161.645, ± .012

Finally, we have two analyses by Roscoe of tungsten hexchloride, published in the same paper with his results upon the trioxide. In one experiment the chlorine was determined as AgCl; in the other the chloride was reduced by hydrogen, and the residual tungsten estimated. By bringing both results into one form of expression we have for the percentage of chlorine in WCl_s:*

$$\begin{array}{c} 53.588 \\ 53.632 \\ \hline \\ \text{Mean, } 53.610, \pm .015 \end{array}$$

We have now five ratios from which to calculate the atomic weight of tungsten:

- (1.) Percentage of W in WO₃, 79.3215, ± .00085
- (2.) Percentage of H₂O in BaO.4WO₃.9H₂O, 13.0368, ± .0060

19.5700 grm. WCl₆ gave 42.4127 grm. AgCl. 10.4326 "4.8374" tungsten.

^{*} The actual figures are as follows:

```
(3.) Am<sub>2</sub>Fe(SO<sub>4</sub>)<sub>2</sub>.6H<sub>2</sub>O : FeWO<sub>4</sub> : : .0365457, \pm .0000012 : .0283549, \pm .0000115
```

- (4.) AgCl: Ag2WO4:: 100: 161.645, ± .012
- (5.) Percentage of Cl in WCl₂, 53.610, \pm .015

From these we get five values for tungsten, as follows:

From	(1)
44	(2) = 183.532, \pm .156
"	(3) = 183.923, \pm .120
**	(4) = 183.248, \pm .069
44	(5) = 183.639, \pm .109
	General mean " = 183.610, ± .032
	Or, if $O = 16$, then " = 184.032

URANIUM.

It is not the purpose of the present investigation to examine at all systematically such questions as are involved in the discussion whether the atomic weight of uranium is 120 or 240. For convenience we may use the formulæ based upon the smaller number, and, if eventually the larger value proves to be correct, it will be easy to double the figures which we obtain. Suffice it to say here, that the specific heat of the green oxide, according to Donath,* agrees best with the formula U_sO₄ and the lower atomic weight. the other hand, the value 240 fits best into such schemes as that given by Mendelejeff in his paper on the periodic law. An accurate determination of the specific heat of the metal itself is much needed, for the material with which Regnault worked was of uncertain quality; furthermore, the vapor density of some volatile uranium compounds ought to be ascertained.† Until some such data have been rigidly

^{*} Ber. d. Deutsch. Chem. Gesell., 12, 742. 1879.

[†] The value of 240 for uranium is strongly sustained by the recent experiments of Zimmermann upon the vapor density of the tetrachlorid and tetrabromid. For UBr₄ the vapor density is 19.46, while theory (U = 240) requires 19.36. For UCl₄ the v. d. 13.33 was found. Theory, 13.21. (Ber. der Deutsch. Chem. Gesell., 14, s. 1934. 1881.)

established the controversy over the two rival values can hardly be satisfactorily settled.

The earlier attempts to determine the atomic weight of uranium were all vitiated by the erroneous supposition that the uranous oxide was really the metal. The supposition, of course, does not affect the weighings and analytical data which were obtained, although these, from their discordance with each other and with later and better results, have now only a historical value.

For present purposes the determinations made by Berzelius,* by Arfvedson,† and by Marchand,‡ may be left quite out of account. Berzelius employed various methods, while the others relied upon estimating the percentage of oxygen lost upon the reduction of U₈O₄ to UO. Rammelsberg's || results also, although very suggestive, need no full discussion. He analyzed the green chloride, UCl,; effected the synthesis of uranyl sulphate from uranous oxide; determined the amount of residue left upon the ignition of the sodio and bario-uranic acetates; estimated the quantity of magnesium uranate formed from a known weight of UO, and attempted also to fix the ratio between the green and the black oxides. His figures vary so widely that they could count for little in the establishing of any general mean; and, moreover, they lead to estimates of the atomic weight which are mostly below the true value. For instance, twelve lots of U.O. from several different sources were reduced to UO by heating in hydrogen. The percentages of loss varied from 3.83 to 4.67, the mean being 4.121. These figures give values for the atomic weight of uranium ranging from 92.66 to 117.65, or, in mean, 107.50. Such discordance is due partly to impurity in some of the material studied, and illustrates the difficulties inherent in the problem to be solved. Some of the uranoso-uranic oxide was prepared by

^{*} Schweigg. Journ., 22, 336. 1818. Poggend. Annal., 1, 359. 1825.

[†] Poggend. Annal., 1, 245. Berz. Jahr., 3, 120. 1822.

[‡] Journ. für Prakt. Chem., 23, 497. 1841.

[|] Poggend. Annal., 55, 318, 1842; 56, 125, 1842; 59, 9, 1843; 66, 91, 1845. Journ. für Prakt. Chem., 29, 324.

calcining the oxalate, and retained an admixture of carbon. Many such points were worked up by Rammelsberg with much care, so that his papers should be scrupulously studied by any chemist who contemplates a redetermination of the atomic weight of uranium.

In 1841 and 1842 Peligot published certain papers* showing that the atomic weight of uranium must be somewhere near 120. A few years later the same chemist published fuller data concerning the constant in question, but in the time intervening between his earlier and his final researches other determinations were made by Ebelmen and by Wertheim. These investigations we may properly discuss in chronological order. For present purposes the early work of Peligot may be dismissed as only preliminary in character. It showed that what had been previously regarded as metallic uranium was in reality an oxide, but gave figures for the atomic weight of the metal which were merely approximations.

Ebelmen's \dagger determinations of the atomic weight of uranium were based upon analyses of uranic oxalate. This salt was dried at 100° , and then, in weighed amount, ignited in hydrogen. The residual uranous oxide was weighed, and in some cases converted into $U_{\bullet}O_{\bullet}$ by heating in oxygen. The following weights are reduced to a vacuum standard:

10. 1644	grm. oxalate gave	7.2939	grm. UO.		
12.9985	66	9.3312	"	Gain on oxidation, .	3685
11.8007	46	8.4690	44		3275
9.9923	"	7.1731	44	" .	2812
11.0887	44	7.9610	44	" ,	3105
10.0830	46	7.2389	"		
6.7940	44	4.8766	**		
16.0594	**	11.5290	44	"	4531

Reducing these figures to percentages, we may present the results in two columns. Column A gives the percentages of UO in the oxalate, while B represents the amount of U_3O_4 formed from 100 parts of UO:

^{*} Compt. Rend., 12, 735. 1841. Ann. Chim. Phys., (3,) 55. 1842.

[†] Journ. für Prakt. Chem., 27, 385. 1842.

A.	В.
71.924	
71.787	103.949
71.767	103.867
71.621	103.920
71.794	103.900
71.793	
71.778	
71.790	103.930
Mean, 71.782, \pm .019	Mean, 103.913, ± .009
column A, the molecular w	eight of UO = 134.523, ± .1

From column A, the molecular weight of UO = 134.523, ± .102
" B, " = 135.985, ± .326

Wertheim's * experiments were even simpler in character than those of Ebelmen. Sodio-uranic acetate, carefully dried at 200°, was ignited, leaving the following percentages of sodium uranate:

Hence the molecular weight of Na₂U₄O₇ = 634.865, \pm .191. And U = 119.282, \pm .048.

The final results of Peligot's † investigations appeared in 1846. Both the oxalate and the acetate of uranium were studied and subjected to combustion analysis. The oxalate was scrupulously purified by repeated crystallizations, and thirteen analyses, representing different fractions, were made. Seven of these gave imperfect results, due to incomplete purification of the material; six only, from the later crystallizations, need to be considered. In these the uranium

^{*} Journ. für Prakt. Chem., 29, 209. 1843.

[†] Compt. Rend., 22, 487.

was weighed as U_sO_4 , and the carbon as CO_2 . From the ratio between the CO_2 and U_sO_4 the atomic weight of uranium may be calculated without involving any error due to traces of moisture possibly present in the oxalate. I subjoin Peligot's weighings, and give, in the third column, the U_sO_4 proportional to 100 parts of CO_2 :

CO ₂ .	$U_{3}O_{4}$.	Ratio.
1.456 grm.	4.649 grm.	319.299
1.369 "	4.412 "	322.279
2.209 "	7.084 "	320.688
1.019 "	3.279 "	321.786
1.069 "	3.447 "	322.461
1.052 "	3.389 "	322.148

Mean, 321.443, ± .338

Hence $U_3O_4 = 423.342, \pm .451$.

From the acetate, $C_2H_3(UO)O_2.H_2O$, the following percentages of U_3O_4 were obtained:

5.061	grm. acetate	gave 3.354	grm. U ₃ O ₄ .	66.2715	per cent.
4.601	66	3.057	"	66.4421	66
1.869	• 44	1.238	44	66.2386	64
3.817	44	2.541	44	66.5706	"
10.182	"	6.757	"	66.3622	44
4.393	**	2.920	44	66.4694	44
2.868	**	1.897	"	66.1437	66
			3.5		

Mean, 66.3569, $\pm .038$

The acetate also yielded the subjoined percentages of carbon and of water. Assuming that the figures for carbon were calculated from known weights of dioxide, with C=12 and O=16, I have added a third column, in which the carbon percentages are converted into percentages of CO_1 :

H_2O .	Ċ.	CO ₂ .
21.60	11.27	41.323
21.16	11.30	41.433
21.10	11.30	41.433
21.20	11.10	40.700
		
Mean, 21.265, ± .187	11.24	41.222, ± .092

From all of these figures we may calculate the molecular weight of the uranic acetate as follows:

```
From percentage of U_3O_4.....C_2H_3(UO)O_2.H_3O=212.629, \pm .242

" CO_2..... " =212.999, \pm .476

" H_2O..... " =211.184, \pm 1.863

General mean..... " =212.685, \pm .214
```

We have now before us the molecular weights of four uranium compounds, giving us four values for U:

The four values for uranium combine as follows:

Or, if O = 16, U = 119.515, or 239.030.

Considering Peligot's figures by themselves, and combining values 3 and 4, we have U = 119.849, $\pm .123$; or, if 0 = 16, U = 120.125, or 240.250.

It is plain that the atomic weight of uranium needs to be scrupulously revised. The foregoing figures are by no means satisfactory. Chemically considered, it is probable that Peligot's work is the best, and that his results should be given preference. His figures from the oxalate and the acetate tally well with each other, whereas Ebelmen's two sets of results vary widely. From the percentage of UO yielded by the oxalate, Ebelmen's figures give a low value for U. From his oxidation of UO to U₃O₄ we get a value nearly two units higher. Peligot, in his work with the oxalate, found it, even after three or four crystallizations, to be contaminated with oxalic acid, and rejected the figures obtained from impure material. Probably Ebelmen's low values are due to the same impurity.

ALUMINUM.

The atomic weight of aluminum has been determined by Berzelius, Mather, Tissier, Dumas, Isnard, Terreil, and Mallet. The early calculations of Davy and of Thomson we may properly disregard.

Berzelius'* determination rests upon a single experiment. He ignited 10 grammes of dry aluminum sulphate, $Al_2(SO_4)_3$, and obtained 2.9934 grammes of Al_2O_3 as residue. Hence, if S = 31.987 and O = 15.9633, Al = 27.243.

In 1835† Mather published a single analysis of aluminum chloride, from which he sought to fix the atomic weight of the metal. 0.646 grm. of Al₂Cl₆ gave him 2.056 of AgCl and 0.2975 of Al₂O₃. These figures give worthless values for Al, and are included here only for the sake of completeness. From the ratio between AgCl and Al₂Cl₆, Al = 28.925.

Tissier's \ddagger determination, also resting on a single experiment, appeared in 1858. Metallic aluminum, containing .135 per cent. of sodium, was dissolved in hydrochloric acid. The solution was evaporated with nitric acid to expel all chlorine, and the residue was strongly ignited until only alumina remained. 1.935 grm. of Al gave 3.645 grm of Al₂O₃. If we correct for the trace of sodium in the aluminum, we have Al = 27.073.

Essentially the same method of determination was adopted by Isnard, who, although not next in chronological order, may fittingly be mentioned here. He found that 9 grm. of aluminum gave 27 grm. of Al. O3. Hence Al = 26.938.

In 1858 Dumas, in connection with his celebrated revision of the atomic weights, made seven experiments with aluminum chloride. The material was prepared in quantity,

^{*} Poggend. Annal., 8, 177.

[†] Silliman's Amer. Journ., 27, 241.

¹ Compt. Rend., 46, 1105.

^{||} Compt. Rend., 66, 508. 1868.

[§] Ann. Chim. Phys., (3,) 55, 151. Ann. Chem. Pharm., 113, 26.

sublimed over iron filings, and finally resublimed from metallic aluminum. Each sample used was collected in a small glass tube, after sublimation from aluminum in a a stream of dry hydrogen, and hermetically enclosed. Having been weighed in the tube, it was dissolved in water, and the quantity of silver necessary for precipitating the chlorine was determined. Reducing to a common standard, his weighings give the quantities of Al₂Cl₆ stated in the third column, as proportional to 100 parts of silver:

1.8786 gr	m. Al ₂ Cl ₆	= 4.543 g	rm. Ag.	41.352
3.021	44	7.292	44	41.459-Bad.
2.399	"	5.802	"	41.348
1.922	44	4.6525	"	41.311
1.697	44	4.1015	44	41.375
4.3165	44	10.448	66	41.314
6.728	44	16.265	46	41.365

In the second experiment the Al_2Cl_6 contained traces of iron. Rejecting this experiment the remaining six give a mean of $41.344, \pm .007$. Hence $Al = 27.441, \pm .082$.

In consequence of these figures of Dumas, the atomic weight of aluminum has generally of late years been put at 27.5, and the lower results deduced from the work of other investigators have been disregarded.

In 1879 Terreil* published a new determination of the atomic weight under consideration, based upon a direct comparison of the metal with hydrogen. Metallic aluminum, contained in a tube of hard glass, was heated strongly in a current of dry hydrochloric acid. Hydrogen was set free, and was collected over a strong solution of caustic potash. 0.410 grm. of aluminum thus were found equivalent to 508.2 cc., or .0455 grm. of hydrogen. Hence Al = 27.033.

About a year after Terreil's determination appeared the lower value for aluminum was thoroughly confirmed by J. W. Mallet.† After giving a full resumé of the work done by others, exclusive of Isnard, the author describes his own experiments, which may be summarized as follows:

^{*} Bulletin de la Soc. Chimique, 31, 153.

[†] Phil. Trans., 1880, p. 1003.

Four methods of determination were employed, each one simple and direct, and at the same time independent of the others. First, pure ammonia alum was calcined, and the residue of aluminum oxide was estimated. Second, aluminum bromide was titrated with a standard solution of silver. Third, metallic aluminum was attacked by caustic soda, and the hydrogen evolved was measured. Fourth, hydrogen was set free by aluminum, and weighed as water. Every weight was carefully verified, the verification being based upon the direct comparison, by J. E. Hilgard, of a kilogramme weight with the standard kilogramme at Washington. The specific gravity of each piece was determined, and also of all materials and vessels used in the weighings. During each weighing both barometer and thermometer were observed, so that every result represents a real weight in vacuo.

The ammonium alum used in the first series of experiments was specially prepared, and was absolutely free from ascertainable impurities. The salt was found, however, to lose traces of water at ordinary temperatures; a circumstance which tended towards a slight elevation of the apparent atomic weight of aluminum as calculated from the weighings. Two sets of experiments were made with the alum; one upon a sample air-dried for two hours at 21°-25°, the other upon material dried for twenty-four hours at 19°-26°. These sets, marked A and B respectively, differ slightly; B being the less trustworthy of the two, judged from a chemical standpoint. Mathematically it is the better of the two. Calcination was effected with a great variety of precautions, concerning which the original memoir must be consulted. To Mallet's weighings I append the percentages of Al₂O₃ deduced from them:

		Series A.			
8.2144 grm.	of the alum g	ave .9258 gr	m. Al ₂ O ₃ .	11.270 p	er cent.
14.0378	"	1.5825	66	11.273	44
5.6201	66	.6337	44	11.275	44
11.2227	44	1.2657	44	11.278	44
10.8435	**	1.2216	46	11.266	46

Mean, 11.2724, ± .0014

Series B.

12.1023 gra	n. of the alum gave	1.3660	grın. Al ₂ O ₂ .	11.287 p	er cent.
10.4544	"	1.1796	"	11.283	"
6.7962	"	.7670	44	11.286	"
8.5601	46	.9654	**	11.278	44
4.8992	44	.5528	**	11.283	44
				11.2824.	

Combined, these series give a general mean of 11.2793, $\pm .0008$. Hence Al = 27.075, $\pm .011$.

The aluminum bromide used in the second series of experiments was prepared by the direct action of bromine upon the metal. The product was repeatedly distilled, the earlier portions of each distillate being rejected, until a constant boiling point of 263.°3 at 747 mm. pressure was noted. The last distillation was effected in an atmosphere of pure nitrogen, in order to avoid the possible formation of oxide or oxy-bromide of aluminum; and the distillate was collected in three portions, which proved to be sensibly identical. The individual samples of bromide were collected in thin glass tubes, which were hermetically sealed after nearly filling. For the titration pure silver was prepared, and after fusion upon charcoal it was heated in a Sprengel vacnum in order to eliminate occluded gases. This silver was dissolved in specially purified nitric acid, the latter but very slightly in excess. The altiminum bromide, weighed in the sealed tube, was dissolved in water, precautions being taken to avoid any loss by splashing or fuming which might result from the violence of the action. To the solution thus obtained the silver solution was added, the silver being something less than a decigramme in deficiency. The remaining amount of silver needed to complete the precipitation of the bromine was added from a burette, in the form of a standard solution containing one milligramme of metal to each cubic centimetre. The final results were as follows, the figures in the third column representing the quantities of bromide proportional to 100 parts of silver. Series A is from the first portion of the last distillate of Al, Br; series

B from the second portion, and series C from the third portion:

		Series 2	<i>A</i> .	
6.0024 gr	m. Al ₂ Br	= 7.2793 g	rm. Ag.	82.458
8.6492	44	10.4897	66	82.454
3.1808	"	3.8573	44	82.462
		Series I	B .	
6.9617	44	8.4429	"	82.456
11.2041	44	13.5897	44	82.445
3.7621	66	4.5624	66	82.459
5.2842	66	6.4085	"	82.456
9.7338	46	11.8047	"	82.457
		Series (c.	
9.3515	"	11.3424	66	82.447
4,4426	**	5.3877	"	82.458
5.2750	**	6.3975	"	82.454
				<u> </u>
				Mean. 82.455. + .001

Hence Al = 27.046, $\pm .061$.

The high probable error of this result is due to the high probable error of the atomic weight of bromine.

The experiments to determine the amount of hydrogen evolved by the action of caustic soda upon metallic aluminum were conducted with pure metal, specially prepared, and with caustic soda made from sodium. The soda solution was so strong as to scarcely lose a perceptible amount of water by the passage through it of a dry gas at ordinary temperature. As the details of the experiments are somewhat complex, the original memoir must be consulted for them. The following results were obtained, the weight of the hydrogen being calculated from the volume, by Regnault's data corrected for the latitude and elevation of the University of Virginia:

Weight of Al.	Vol. of H.	Wt. of H.	At. Wt.
.3697 grm.	458.8 c. c.	.04106 grm.	27.012
.3769 "	467.9 "	.04187 "	27.005
.3620 "	449. I "	.04019 "	27.022
·7579 "	941.5 "	.08425 "	26.998
.7314 "	907.9"	.08125 "	27.006
.7541 "	936.4 "	.08380 "	26.996
		Me	an, 27.005, ± .0032

The closing series of experiments was made with larger quantities of aluminum than were used in the foregoing The hydrogen, evolved by the action of the caustic alkali, was dried by passing it through two drying tubes containing pumice stone and sulphuric acid, and two others containing asbestos and phosphorus pentoxide. Thence it passed through a combustion tube containing copper oxide heated to redness. A stream of dry nitrogen was employed to sweep the last traces of hydrogen into the combustion tube, and dry air was afterwards passed through the entire apparatus to reoxidize the surface of reduced copper, and to prevent the retention of occluded hydrogen. The water formed by the oxidation of the hydrogen was collected in three drying tubes. The results obtained were as follows. The third column gives the amount of water formed from 10 grammes of aluminum:

```
2.1704 grm. Al gave 2.1661 grm. H_2O. 9.9802
2.9355 " 2.9292 " 9.9785
5.2632 " 5.2562 " 9.9867
Mean, 9.9818, \pm .0017
Hence Al = 26.998, \pm .007.
```

In combining the various determinations of the atomic weight of aluminum into one general mean, we must arbitrarily assign weight to the single experiments of Berzelius, Isnard, Tissier, and Terreil. This may fairly be done by giving to each the probable error, and therefore the weight, of a single observation in Dumas' series. Mather's work may be ignored altogether:

```
Tissier = 27.096, \pm .201
    Isnard...... = 26.938, \pm .201
    Dumas ..... " = 27.441, \pm .082
    Terreil ..... " = 27.033, \pm .201
 44
    Mallet's alum experiments, " = 27.075, ± .011
         Al_2Br_6 " = 27.046, ± .061
 46
 ü
                "
                    " = 27.005, ± .003
         Н
         H.O-
                    " = 26.998, \pm .007
      General mean ..... " = 27.0092, ± .0028
```

If O = 16, Al = 27.075. Taking Mallet's work alone, Al = 27.0089, $\pm .0028$.

Evidently all the data except Mallet's might be rejected without affecting sensibly the final result. Dumas' work is clearly vitiated by constant errors, but the determinations by Isnard, Tissier, and Terreil may be regarded as having some confirmative value.

GOLD.

The only determinations of the atomic weight of gold which are worthy of consideration are those of Berzelius and of Levol.

The earliest method adopted by Berzelius* was that of precipitating a solution of gold chloride by means of a weighed quantity of metallic mercury. The weight of gold thus thrown down gave the ratio between the atomic weights of the two metals. In the single experiment which Berzelius publishes, 142.9 parts of Hg precipitated 93.55 of Au. Hence, using the value for mercury given in a preceding chapter, 199.712, Au = 196.113.

In a later investigation[†] Berzelius resorted to the analysis of potassio-auric chloride, 2KCl.AuCl₃. Weighed quantities of this salt were ignited in hydrogen; the resulting gold and potassium chloride were separated by means of water, and both were collected and estimated. The loss of weight upon ignition was, of course, chlorine. As the salt could not be perfectly dried without loss of chlorine, the atomic weight under investigation must be determined by the ratio between the KCl and the Au. If we reduce to a common standard, and compare with 100 parts of KCl, the equivalent amounts of gold will be those which I give in the last of the subjoined columns:

^{*} Poggend. Annal., 8, 177.

[†] Lehrbuch, 5 Aufl., 3, 1212.

```
4.1445 grm. K2AuCl3 gave .8185 grm. KCl and 2.159 grm. Au. 263.775
                   .44425 " 1.172 "
2.2495
                                                   263.815
                                    2.67225 "
           64
                              66
                                                   263.600
5.1300
                   1.01375
           "
                              "
                                    1.77725 "
                   .674
                                                   263.687
3.4130
                              "
4.19975
                    .8295
                                     2.188
                                                   263.773
```

Mean, 263.730, \pm .026

Hence Au = 196.186, $\pm .101$.

Still a third series of experiments by Berzelius* may be included here. In order to establish the atomic weight of phosphorus he employed that substance to precipitate gold from a solution of gold chloride in excess. Between the weight of phosphorus taken and the weight of gold obtained it was easy to fix a ratio. Since the atomic weight of phosphorus has been better established by other methods, we may properly reverse this ratio and apply it to our discussion of gold. 100 parts of P precipitate the quantities of Au given in the third column:

Hence Au = 195.303, $\pm .589$.

Levol's † estimation of the atomic weight under consideration can hardly have much value. A weighed quantity of gold was converted in a flask into AuCl₃. This was reduced by a stream of sulphur dioxide, and the resulting sulphuric acid was determined as BaSO₄. One gramme of gold gave 1.782 grm. BaSO₄. Hence Au = 195.794.

If we give this single experiment and Berzelius' single result with mercury each equal weight with one analysis in the potassio-auric chloride series, and include respectively the probable errors appertaining to Hg and to BaSO₄, we may combine all the data as follows:

^{*} Lehrbuch, 5 Aufl., 3, 1188.

[†] Ann. d. Chim. et d. Phys., (3,) 30, 355. 1850.

Or, if O = 16, Au = 196.606.

As gold is a metal which can be readily applied to the determination of the atomic weights of other elements, an experimental revision of its atomic weight is very desirable.

NICKEL AND COBALT.

On account of the close similarity of these metals to each other, their atomic weights, approximately if not actually identical, have received of late years much attention.

The first determinations, and the only ones up to 1852, were made by Rothhoff;* each with but a single experiment. For nickel 188 parts of the monoxide were dissolved in hydrochloric acid; the solution was evaporated to dryness, the residue was dissolved in water, and precipitated by silver nitrate. 718.2 parts of silver chloride were thus formed; whence Ni = 58.925. The same process was applied also to cobalt, 269.2 parts of the oxide being found equivalent to 1029.9 of AgCl. Hence Co = 58.817. These values are so nearly equal that their differences were naturally ascribable to experimental errors. They are, however, entitled to no special weight at present, since it cannot be certain from any evidence recorded that the oxide of either metal was absolutely free from traces of the other.

In 1852 Erdmann and Marchand published some results, but without details, concerning the atomic weight of nickel. They reduced the oxide by heating in a current of

^{*} Cited by Berzelius. Poggend. Annal., 8, 184. 1826.

[†] Journ. für Prakt. Chem., 55, 202. 1852.

hydrogen, and obtained values ranging from 58.2 to 58.6, when O = 16. Their results were not very concordant, and the lowest was probably the best.

In 1856, incidentally to other work, Deville* found that 100 parts of pure metallic nickel yielded 262 of sulphate; whence Ni = 59.15.

To none of the foregoing estimations can any importance now be attached. The modern discussion of the atomic weights under consideration began with the researches of Schneider in 1857. This chemist examined the oxalates of both metals, determining carbon by the combustion of the salts with copper oxide in a stream of dry air. The carbon dioxide thus formed was collected as usual in a potash bulb, which, in weighing, was counterpoised by a similar bulb, so as to eliminate errors due to the hygroscopic character of the glass. The metal in each oxalate was estimated, first by ignition in a stream of dry air, followed by intense heating in hydrogen. Pure nickel or cobalt was left behind in good condition for weighing. Four analyses of each oxalate were made, with the results given below. The nickel salt contained three molecules of water, and the cobalt salt two molecules:

			NiC ₂ O ₄ .3H ₂ O.		
1.1945	grm. gave	.528	grm. CO ₂ .	44.203 I	er cent.
2.5555	46	1.12625	44	44.072	46
3.199	46	1.408	**	44.014	**
5.020	44	2.214	46	44.104	**
				Mean, 44.098,	士.027

The following percentages of nickel were found in this salt:

^{*} Ann. Chim. Phys., (3,) 46, 182. 1856.

[†] Poggend. Annal., 101, 387. 1857.

$$CoC_2O_4$$
.2 H_2O .

1.6355	grin. gave	.781	grm. CO ₂ .	47.753 per cent.
1.107	"	.5295	46	47.832 "
2.309	"	1.101	"	47.683 "
3.007	"	1.435	44	47.722 "

Mean, 47.7475, ± .0213

The following were the percentages found for cobalt:

In a later paper* Schneider also gives some results obtained with a nickel oxalate containing but two molecules of water. This gave him 47.605 per cent. of CO₂, and the following percentages of nickel:

The conclusion at which Schneider arrived was, that the atomic weights of cobalt and nickel are not identical, being about 60 and 58 respectively. The percentages given above will be discussed at the end of this chapter in connection with all the other data relative to the constants in question.

The next chemist to take up the discussion of these atomic weights was Marignac, in 1857.† His original paper is not accessible to me, and I am therefore obliged to give only such features of it as I can get from abstracts and reviews. He worked with the chlorides and sulphates of nickel and cobalt, using apparently common gravimetric methods. The sulphates, taken as anhydrous, were first ignited to expel SO₂+ O, after which the residues were heated with weighed amounts of lead silicate. The increase in weight

^{*} Poggend. Annal., 107, 616.

[†] Jahresbericht, 1857, 225. Bibl. Univ. de Genève, (nouv. s.,) 1, 373.

was CoO or NiO respectively. The anhydrous chlorides were prepared from the hydrated salts by ignition in dry chlorine or hydrochloric acid. With cobalt, the monohydrated chloride, dried at 100° , was also employed. For nickel he gives the following values, referred probably to 0 = 16, S = 32, Ag = 108, Cl = 35.5:

To cobalt these values are assigned:

That is, contrary to Schneider's view, the two atomic weights are approximately the same. The values for nickel, however, run a little lower than those for cobalt; a fact which is probably not without significance. Marignac criticizes Schneider's earlier paper, holding that the nickel oxalate may have contained some free oxalic acid, and that the cobalt salt was possibly contaminated with carbonate or with basic compounds. In his later papers Schneider rejects these suggestions as unfounded, and in turn criticizes Marignac. The purity of anhydrous NiSO, is not easy to guarantee, and, according to Schneider, the anhydrous chlorides of cobalt and nickel are liable to be contaminated with oxides. This is the case even when the chlorides are heated in chlorine, unless the gas is carefully freed from all traces of air and moisture.

Dumas'* determinations of the two atomic weights were made with the chlorides of nickel and cobalt. The pure metals were dissolved in aqua regia, the solutions were repeatedly evaporated to dryness, and the residual chlorides were ignited in dry hydrochloric acid gas. The last two estimations in the nickel series were made upon NiCl, formed by heating the spongy metal in pure chlorine. In the third column I give the NiCl, or CoCl, equivalent to 100 parts of silver:

^{*} Ann. Chem. Pharm., 113, 25. 1860.

.9123 gr	m. NiCl ₂	= 1.515	grm. Ag.	60.218
2.295	14	3.8115	"	60.212
3.290	44	5.464	66	60.212
1.830	44	3.041	46	60.178
3.001	u	4.987	"	60.176
				Mean, 60.1992, ± .0062
2.352 gm	a. CoCl ₂ :	= 3.9035 g	rm. Ag.	60.254
4.210	"	6.990	"	60.229
3.592	46	5.960	44	60.268
2.492	• •	4.1405	**	60.186
4.2295	**	7.0255	"	60.202
				Mean, 60.2278, ± .011

These results give values for Co and Ni differing by less than a tenth of a unit; here, as elsewhere, the figure for Ni being a trifle the lower.

In 1863* the idea that nickel and cobalt have equal

atomic weights was strengthened by the researches of Russell. He found that the black oxide of cobalt, by intense heating in an atmosphere of carbon dioxide, became converted into a brown monoxide of constant composition. The ordinary oxide of nickel, on the other hand, was shown to be convertible into a definite monoxide by simple heating over the blast lamp. The pure oxides of the two metals, thus obtained, were reduced by ignition in hydrogen, and their exact composition thus ascertained. Several samples of each oxide were taken, yielding the following percentages of metal:

```
NiO.
             78.597
                      1st sample.
             78.584
             78.608
             78.581
                      2d sample.
             78.589
            78.583
             78.616
                      3d sample.
             78.590
             78.588
             78.590
             78.594
                      4th sample.
             78.597
             78.588
Mean of all, 78.593, \pm .0018
```

^{*} Journ. Chem. Soc., (2,) 1, 51.

```
78.591
78.588
78.550
78.598
78.614
78.603
78.591
78.591
78.588
78.592
78.598
78.598
78.598
78.598
78.598
78.595
78.598
78.595
78.598
78.595
78.596
78.596
```

Mean of all, 78.592, $\pm .0023$

These percentages are practically identical, and lead to essentially the same mean value for each atomic weight.

In a later paper Russell* confirmed the foregoing results by a different process. He dissolved metallic nickel and cobalt in hydrochloric acid and measured the hydrogen evolved. Thus the ratio between the metal and the ultimate standard was fixed without the intervention of any other element. About two-tenths of a gramme of metal, or less, was taken in each experiment. 100 parts by weight of Co or Ni give the following weights of H, calculated from the volume of the latter:

Ni.		Co.	
3.420)	3-395)
3.418		3.398	Ist sample.
3.416		3.397	l set emmpre.
3.417	st sample.	3.398 .	J
3.412		3.403)
3.415		3.401	2d sample.
3.416	j	3.401)

^{*} Journ. Chem. Soc., (2,) 7, 494. 1869.

Mean of all, 3.411, ± .001

A glance at the tabulated discussion which closes this chapter will show that these figures agree well with each other, and well with those found from the analyses of the oxides. The probable errors assigned in the hydrogen series may be a little too low, since they ought to be modified by the probable error of the weight of a unit volume of hydrogen. So insignificant a correction may, however, be neglected.

Some time after the publication of Russell's first paper, but before the appearance of his second, some other investigations were made known. Of these the first was by Sommaruga,* whose results, obtained by novel methods, closely confirmed those of Schneider and antagonized those of Dumas, Marignac, and Russell. The atomic weight of nickel Sommaruga deduced from analyses of the nickel potassium sulphate, K₂Ni(SO₄)₂.6H₂O, which, dried at 100°, has a perfectly definite composition. In this salt the sulphuric acid was determined in the usual way as barium sulphate, a process to which there are obvious objections. In the third column are given the quantities of the nickel salt proportional to 100 parts of BaSO₄:

0.9798	grm. gav	re 1.0462 g	rm. BaSO ₄ .	93.653
1.0537	46	1.1251	64	93.654
1.0802	66	1.1535	"	93.645
1.1865	44	1.2669	44	93.654
3.2100	44	3.4277	"	93.649
3.2124	"	3.4303	44	93.648

^{*} Sitzungsb. Wien Akad., 54, 2 Abth., 50. 1866.

For cobalt Sommaruga used the purpureo cobalt chloride of Gibbs and Genth. This salt, dried at 110°, is anhydrous and stable. Heated hotter, CoCl, remains. The latter, ignited in hydrogen, yields metallic cobalt. In every experiment the preliminary heating must be carried on cautiously until ammoniacal fumes no longer appear:

.6656 g	rm. gav	e .1588 g	rm. Co.	23.858 pe	er cent.
1.0918	44	.2600	44	23.814	"
.9058	66	.2160	"	23.846	"
1.5895	44	.3785	44	23.813	"
2.9167	**	.6957	"	23.847	"
1.8390	"	.4378	44	23.806	"
2.5010	"	.5968	46	23.808	"

Mean, 23.827, ± .006

Further along this series will be combined with a similar one by Lee. It may here be said that Sommaruga's paper was quickly followed by a critical essay from Schneider,* endorsing the former's work, and objecting to the results of Russell.

In 1867 still another new process for the estimation of these atomic weights was put forward by Winkler,† who determined the amount of gold which pure metallic nickel and cobalt could precipitate from a neutral solution of sodioauric chloride. Experimentally, the method seems to be quite accurate; practically, it involves a knowledge of the defectively ascertained atomic weight of gold. In order to obtain pure cobalt Winkler prepared purpureo-cobalt chloride, which, having been four or five times recrystallized. was ignited in hydrogen. His nickel was repeatedly purified by precipitation with sodium hypochlorite. From material thus obtained pure nickel chloride was prepared, which, after sublimation in dry chlorine, was also reduced by hydrogen. 100 parts of gold are precipitated by the quantities of nickel and cobalt given in the third columns respectively. In the cobalt series I include one experiment

^{*} Poggend. Annal., 130, 310.

[†] Zeit. Anal. Chem., 6, 18. 1867.

by Weselsky which was published by him in a paper presently to be cited:

.4367		44	.9666	46	45.179
.5189		44	1.1457	**	45.291
.6002		"	1.3286	44	45.175
				Mean,	45.209, ± .019
. 5890	grm. coba	lt precipitated	1 1.3045	grm. gold.	45.151
		44	.6981	44	45.080
.3147					
.3147 .5829		44	1.2913	66	45.141
•		"	1.2913	66	45.141 45.182
. 5829				**	

Weselsky's paper,* already cited, relates only to cobalt. He ignited the cobalticyanides of ammonium and of phenylammonium in hydrogen, and from the determinations of cobalt thus made deduced its atomic weight. His results are as follows:

```
.7575 grm. (NH<sub>4</sub>)<sub>6</sub>Co<sub>2</sub>Cy<sub>19</sub> gave .166 grm. Co. 21.914 per cent.
                                            .113
                                                                 21.972
   .5143
                                                         Mean, 21.943, ± .029
.8529 grm. (C<sub>8</sub>H<sub>8</sub>N)<sub>6</sub>Co<sub>2</sub>Cy<sub>12</sub> gave .1010 grm. Co. 11.842 per cent.
.6112
                                           .0723
                                                                 11.829
                                           .0850
.7140
                                                                 11.905
.9420
                                           .1120
                                                                 11.890
                                                        Mean, 11.8665, ± .0124
```

Finally, we come to the work done by Leet in the laboratory of Wolcott Gibbs. Like Weselsky, Lee ignited certain cobalticyanides and nickelocyanides in hydrogen and determined the residual metal. The double cyanides chosen were those of strychnia and brucia; salts of very high molecular weight, in which the percentages of metal are relatively low. A series of experiments with purpureo-cobalt

^{*} Ber. d. Deutsch. Chem. Gesell., 2, 592. 1868. † Am. Journ. Sci. and Arts, (3,) 2, 44. 1871.

chloride was also carried out. In order to avoid admixture of carbon in the metallic residues, the salts were first ignited in air, and then in oxygen. Reduction by hydrogen followed. The salts were in each case covered by a porous septum of earthenware, through which the hydrogen diffused, and which served to prevent the mechanical carrying away of solid particles; furthermore, heat was applied from above. The results attained were very satisfactory, and assign to nickel and cobalt atomic weights varying from each other by about a unit; Ni being nearly 58, and Co about 59. The exact figures will appear later. The cobalt results agree remarkably well with those of Weselsky. The following are the percentages of metal found:

```
In brescia nickelocyanide, Ni<sub>3</sub>Cy<sub>12</sub>(C<sub>23</sub>H<sub>26</sub>N<sub>2</sub>O<sub>4</sub>)<sub>6</sub>H<sub>6</sub>.10H<sub>2</sub>O.
                                        5.724
                                       5.729
                                        5.750
                                        5.733
                                        5.712
                                       5.729
                              Mean, 5.7295, ± .0034
In strychnia nickelocyanide, Ni2Cy12(C11H12N2O2)6: H6.8H2O.
                                       6.607
                                       6.613
                                       6.589
                                       6.607
                                       6.561
                                       6.595
                              Mean, 6.595, \pm .005
 In bruscia cobalticyanide, Co, Cy, (C, H, N, O, )6. H6. 20 H, O.
                                       3.759
                                       3.720
                                       3.739
                                       3.748
                                       3.747
                                       3.749
                              Mean, 3.7437, \pm .0036
```

```
In strychnia cobalticyanide, Co_2Cy_{12}(C_{21}H_{22}N_2O_2)_6. H_6.8H_2O.
                             4.583
                             4.596
                             4.554
                             4.564
                             4.577
                             4.549
                     Mean, 4.5705, ± .005
         In purpureo-cobalt chloride, Co2(NH2)10Cla.
                            23.575
                            23.587
                            23.586
                            23.579
                            23.569
                            23.581
                     Mean, 23.5795, ± .0019
```

The last series may be combined with Sommaruga's, thus:

```
Sommaruga _____ 23.827, ± .006
Lee _____ 23.5795, ± .0019
General mean ____ 23.6045, ± .0018
```

In discussing the atomic weights of nickel and cobalt, we may ignore the work of Rothhoff, Erdmann and Marchand and Deville. That of Marignac must also be omitted, for want of sufficient data. For nickel we have the following ratios. The probable error assigned in No. 4, is that of a single experiment in No. 2:

```
(1.) Per cent. of Ni in
                            NiC_2O_4.3H_2O_7, 29.084, \pm .006
            44
                                  44
                                            44.098, ± .027
 (2.)
                    CO<sub>2</sub> from
                    Ni in NiC_2O_4.2H_2O_7, 31.4076, \pm .0026
 (3.)
 (4.)
                    CO, from
                                             47.605, ± .053
            "
                    Ni in NiO, 78.593, ± .0018
 (5.)
                      " brucia nickelocyanide, 5.7295, ± .0034

" strychnia " 6.595, ± .005
 (6.)
 (7.)
 (8.) Ag: NiCl<sub>2</sub>:: 100: 60.1992, \pm .0062
 (9.) Ni : H :: 100 : 3.411, ± .001
(10.) Au: Ni:: 100: 45.209, ± .019
(11.) BaSO<sub>4</sub>: K_2Ni(SO_4)_2.6H_2O::100:93.6505, \pm .001
```

Since the proportion of water in the oxalates is not an absolutely certain quantity, the data concerning such salts

are best handled by employing the ratios between the carbon dioxide and the metal. Accordingly ratios (1) and (2) give a single value for Ni, and ratios (3) and (4) another. In all, we have nine values for the atomic weight in question:

If O = 16, Ni = 58.682.

In the foregoing result it will be seen that the two sets of figures due to Russell receive very great weight. This is because the one set is referred directly to hydrogen, without the intervention of the probable error of any other element; while the second set involves only the atomic weight of oxygen, of which the probable error is small. As regards accuracy of methods, however, and certainty concerning the purity of material, Russell's work is no better than Schneider's, and probably inferior to Lee's. Now values one to five in the above table represent the tolerably concordant results of Schneider, Lee, and Sommaruga. They, combined by themselves, give a general mean of Ni = 57.928, $\pm .0215$; or, if O = 16, of Ni = 58.062. This value, taking everything into account, I cannot but regard as more likely to prove correct than the larger mean deduced from all the At all events, the atomic weight of nickel needs further careful investigation.

For cobalt these ratios are available:

```
(1.) Per cent. of Co in CoC<sub>2</sub>O<sub>4</sub>.2H<sub>2</sub>O, 32.5555, ± .0149

(2.) " CO<sub>2</sub> from " 47.7475, ± .0213

(3.) " Co in CoO, 78.592, ± .0023

(4.) " purpureo-cobalt chloride, 23.6045, ± .0018

(5.) " " phenylammonium cobalticyanide, 11.8665, ± .0124

(6.) " ammonium " 21.943, ± .029
```

```
(7.) Per cent. of Co in brucia cobalticyanide, 3.7437, ± .0036
(8.) " " strychnia " 4.5705, ± .005
(9.) Ag: CoCl<sub>2</sub>:: 100: 60.2278, ± .011
(10.) Co: H:: 100: 3.4017, ± .0009
(11.) Au: Co:: 100: 45.151, ± .025
```

Hence we have ten values for Co, as follows:

If O = 16, $C_0 = 59.023$.

SELENIUM.

The atomic weight of this element was first determined by Berzelius,* who, saturating 100 parts of selenium with chlorine, found that 179 of chloride were produced. Further on these figures will be combined with similar results by Dumas.

We may omit, as unimportant for present purposes, the analyses of alkaline selenates made by Mitscherlich and Nitzsch,† and pass on to the experiments published by Sacc; in 1847. This chemist resorted to a variety of methods, some of which gave good results, while others were unsatisfactory. First, he sought to establish the exact composition of SeO₂, both by synthesis and by analysis. The former plan, according to which he oxidized pure selenium by

^{*} Poggend. Annal., 8, 1. 1826.

[†] Poggend. Annal., 9, 623. 1827.

¹ Ann. d. Chim. et d. Phys., (3,) 21, 119.

nitric acid, gave poor results; better figures were obtained upon reducing SeO, with ammonium bisulphite and hydrochloric acid, and determining the percentage of selenium set free:

```
.6800 grm. SeO<sub>2</sub> gave .4828 grm. Se. 71.000 per cent. 3.5227 " 2.5047 " 71.102 " 4.4870 " 3.1930 " 71.161 " Mean, 71.088, ± .032
```

In a similar manner Sacc also reduced barium selenite, and weighed the resulting mixture of barium sulphate and free selenium. This process gave discordant results, and a better method was found in calcining BaSeO₃ with sulphuric acid, and estimating the resulting quantity of BaSO₄. In the third column I give the amounts of BaSO₄ equivalent to 100 of BaSeO₃:

```
.5573 grm. BaSeO<sub>3</sub> gave .4929 grm. BaSO<sub>4</sub>. 88.444
.9942 " .8797 " 88.383
.2351 " .2080 " 88.473
.9747 " .8621 " 88.448
```

Mean, 88.437, ± .013

Still other experiments were made with the selenites of silver and lead; but the figures were subject to such errors that they need no further discussion here.

A few years after Sacc's work was published, Erdmann and Marchand made with their usual care a series of experiments upon the atomic weight under consideration.* They alalyzed pure mercuric selenide, which had been repeatedly sublimed and was well crystallized. Their method of manipulation has already been described in the chapter upon mercury. These percentages of Hg in HgSe were found:

```
71.726
71.731
71.741
————
Mean, 71.7327, ± .003
```

^{*} Journ. für Prakt. Chem., 55, 202. 1852.

The next determinations were made by Dumas,* who returned to the original method of Berzelius. Pure selenium was converted by dry chlorine into SeCl₄, and from the gain in weight the ratio between Se and Cl was easily deducible. I include Berzelius' single experiment, which I have already cited, and give in a third column the quantity of chlorine absorbed by 100 parts of selenium:

1.709	grm. Se absorb	3.049	grm. Cl.	178.409
1.810	46	3.219	46	177.845
1.679	66	3.003	"	178.856
1.498	**	2.688	46	179.439
1.944	44	3.468	44	178.395
1.887	46	3.382	**	179.226
1.935	64	3.452		178.398
				179.000—Berzelius.
				
	•			Mean, 178.696, ± .125

The question may here be properly asked, whether it would be possible thus to form SeCl₄ and be certain of its absolute purity? A trace of oxychloride, if simultaneously formed, would increase the apparent atomic weight of selenium. In point of fact, this method gives a higher value for Se than any of the other processes which have been adopted, and that value has the largest probable error of any one in the entire series. A glance at the table which summarizes the discussion at the end of this chapter will render this point sufficiently clear.

Latest of all, we come to the determinations made by Ekman and Pettersson.* They tried various methods of investigation, and finally decided upon the two following:

First. Pure silver selenite, Ag₂SeO₃ was ignited, leaving behind metallic silver in the subjoined percentages:

^{*} Ann. Chem. Pharm., 113, 32. 1860.

[†] Ber. d. Deutsch. Chem. Gesell., 9, 1210. 1876. Published in detail by the society at Upsala.

Second. A warm aqueous solution of selenious acid was mixed with HCl, and reduced by a current of SO₂. The reduced Se was collected upon a glass filter, dried, and weighed. Percentages of Se in SeO₂:

This series, combined with that of Sacc, 71.088, $\pm .032$, gives a general mean of 71.1907, $\pm .0016$.

There are now five series of figures from which to deduce the atomic weight of selenium:

```
    (1.) Per cent. of Se in SeO<sub>2</sub>, 71.1907, ± .0016
    (2.) BaSeO<sub>3</sub>: BaSO<sub>4</sub>:: 100: 88.437, ± .013
    (3.) Per cent. of Hg in HgSe, 71.7327, ± .003
    (4.) Se: SeCl<sub>4</sub>:: 100: 178.696, ± .125
    (5.) Per cent. of Ag in Ag<sub>2</sub>SeO<sub>2</sub>, 62.957, ± .005
```

From these we get the following values for selenium:

If O = 16, Se = 78.978.

TELLURIUM.

Particular interest attaches to the atomic weight of tellurium, on account of the speculations of Mendelejeff. According to the "periodic law" of that chemist, tellurium should lie between antimony and iodine, having an atomic weight greater than 120, and less than 127. Theoretically, Mendelejeff assigns it a value of Te = 125; but all the published determinations lead to a mean number higher than would be admissible under the aforesaid "periodic law." Whether theory or experiment is at fault remains to be discovered.

The first, and for many years the only, determinations of the constant in question, were made by Berzelius.* By means of nitric acid he oxidized tellurium to the dioxide, and from the increase in weight deduced a value for the metal. He published only his final results; from which, if O = 100, Te = 802.121. The three separate experiments give Te = 801.74, 801.786, and 802.838; whence we can calculate the following percentages of metal in the dioxide:

The next determinations were made by von Hauer,† who resorted to the analysis of the well crystallized double salt TeBr₄.2KBr. In this compound the bromine was estimated as silver bromide, the values assumed for Ag and Br being respectively 108.1 and 80. Recalculating, with our newer atomic weights for the above named elements, we get from v. Hauer's analyses, for 100 parts of the salt, the quantities of AgBr which are put in the third column:

^{*} Poggend. Annal., 28, 395. 1833 † Sitzungsb. Wien Akad., 25, 142.

```
2.000 grm. K<sub>2</sub>TeBr<sub>4</sub> gave 69.946 per cent. Br. 164.460
                66
                           69.8443
                                                    164.221
                46
                                         66
                            69.9113
2.934
                                                    164.379
                44
                            70.0163
                                         46
                                                    164.626
3.697
1.000
                            69.901
                                                    164.355
```

Mean, 164.408, \pm .045

From Berzelius' series we may calculate Te = 128.045, and from v. Hauer's Te = 127.419. Dumas,* by a method for which he gives absolutely no particulars, found Te = 129.

In 1879, with direct reference to Mendelejeff's speculations, the subject of the atomic weight of tellurium was taken up by Wills.† The methods of both Berzelius and von Hauer were employed, with various rigid precautions in the way of testing balance and weights, and to ensure purity of material. In the first series of experiments tellurium was oxidized by nitric acid to form TeO₂. The results gave figures ranging from Te = 126.31 to 129.34:

```
2.21613 grm. Te gave 2.77612 grm. TeO<sub>2</sub>. 79.828 per cent. Te.
             66
1.45313
                      1.81542
                                           80.044
             46
                                   66
                                                       "
2.67093
                      3.33838
                                           80.007
             44
                                   64
                                                       44
4.77828
                      5.95748
                                           80.207
                                                       "
                                           79.989
2.65029
                      3.31331
                                    Mean, 80.015, ± .041
```

In the second series tellurium was oxidized by aqua regia to TeO_2 ; with results varying from Te = 127.77 to 128.00:

```
2.85011 grm. Te gave 3.56158 grm. TeO<sub>2</sub>. 80.024 per cent. Te.
                       3.86897
3.09673
              66
                                    "
                                             80.040
5.09365
              "
                       6.36612
                                     "
                                             80.012
                                                         44
                       4.08064
                                                         46
3.26604
                                             80.037
                                      Mean, 80.028, \pm .004
```

Combining these series with that due to Berzelius, we have the following general mean:

^{*} Ann. d. Chim. et d. Phys., (3,) 55, 129. 1859. † Journ. Chem. Society, Oct., 1879, p. 704.

```
      Berzelius
      80.042, ± .005

      Wills, 1st series
      80.015, ± .041

      " 2d "
      80.028, ± .004

      General mean
      80.035, ± .003
```

Hence Te = 127.986, $\pm .035$.

By von Hauer's process, the analysis of TeBr₄.2KBr, Will's figures give results ranging from Te = 126.07 to 127.61. Reduced to a common standard, 100 parts of the salt yield the quantities of AgBr given in the third column:

```
1.70673 grm. K. TeBr. gave 2.80499 grm. AgBr.
                                                  164.349
                **
1.75225
                           2.88072
                                                  164.398
                **
                                        "
2.06938
                                                  164.657
                           3.40739
                           5.43228
                "
                                        "
3.29794
                                                  164.717
                66
2.46545
                           4.05742
                                                  164.571
                                           Mean, 164.538, \pm .048
```

Combined with von Hauer's mean, 164.408, \pm .045, this gives a general mean of 164.468, \pm .033. Hence Te = 127.170, \pm .173.

The two independent values for Te combine thus:

If O = 16, Te = 128.254.

A careful consideration of the foregoing figures, and of the experimental methods by which they were obtained, will show that they are not absolutely conclusive with regard to the place of tellurium under the periodic law. The atomic weight of iodine, calculated in a previous chapter, is 126.557. Wills' values for Te, rejecting his first series as relatively unimportant, range from 126.07 to 128.00; that is, some of them fall below the atomic weight of iodine, although none descend quite to the 125 assumed by Mendelejeff.

In considering the experimental methods, reference may properly be made to the controversy regarding the atomic weight of antimony. It will be seen that Dexter, estimating the latter constant by the conversion of the metal into Sb₂O₄, obtained a value approximately of Sb = 122. Dumas, working with SbCl₂, obtained a similar value. Schneider and Cooke, on the other hand, have established an atomic weight for antimony near 120, and Cooke in particular has traced out the constant errors which lurked unsuspected in the work of Dumas and Dexter. Now in some physical respects tellurium and antimony are quite similar. As constant errors vitiated the recently accepted values for Sb, so they may also effect our estimates for Te. The oxidation of Te by nitric acid resembles in minor particulars that of Sb. The analysis of K₂TeBr₆, gives a low value for Te, and yet the material may have contained traces of oxybromides, the presence of which would render even that lower value too high. A careful revision of the atomic weight of tellurium is still necessary.

VANADIUM.

Roscoe's determination of the atomic weight of vanadium is the only one having any present value. The results obtained by Berzelius* and by Czudnowicz † are unquestionably too high; the error being probably due to the presence of phosphoric acid in the vanadic acid employed. This particular impurity, as Roscoe has shown, prevents the complete reduction of V₂O₅ to V₂O₅ by means of hydrogen. All vanadium ores contain small quantities of phosphorus, which can only be detected with ammonium molybdate; a reaction unknown in Berzelius' time. Furthermore, the complete purification of vanadic acid from all traces of phosphoric acid is a matter of great difficulty, and probably never was accomplished until Roscoe undertook his researches.

In his determination of the atomic weight, Roscoet

^{*} Poggend. Annal., 22, 14. 1831.

[†] Poggend. Annal., 120, 17. 1863.

[‡] Journ. Chem. Soc., 6, pp. 330 and 344. 1868.

studied two compounds of vanadium; namely, the pentoxide, V₂O₅ and the oxychloride, VOCl₃. The pentoxide, absolutely pure, was reduced to V₂O₅ by heating in hydrogen, with the following results:

```
7.7397 grm. V<sub>2</sub>O<sub>8</sub> gave 6.3827 grm. V<sub>2</sub>O<sub>8</sub>. 17.533 per cent. of loss. 6.5819 " 5.4296 " 17.507 " 5.1895 " 4.2819 " 17.489 " 5.0450 " 4.1614 " 17.515 " 5.4296 grm. V<sub>2</sub>O<sub>8</sub>, reoxidized, gave 6.5814 grm. V<sub>3</sub>O<sub>8</sub>. 17.501 per cent. difference.
```

Mean, 17.509, ± .005

Hence $V = 51.264, \pm .025$.

Upon the oxychloride, VOCl₈, two series of experiments were made, one volumetric, the other gravimetric. In the volumetric series the compound was titrated with solutions containing known weights of silver, which had been purified according to the methods recommended by Stas. Roscoe publishes his weighings, and gives percentages deduced from them; his figures, reduced to a common standard, make the quantities of VOCl₈ given in the third column proportional to 100 parts of silver. He was assisted by two analysts:

		Analyst 1	1 .	
2.4322 gm	n. VOCl _s	= 4.5525 8	grm. Ag.	53.425
4.6840	46	8.7505	"	53.528
4.2188	"	7.8807	44	53-533
3.9490	"	7.3799	44	53.510
.9243	44	1.7267	**	53.5 3 0
1.4330	44	2.6769	44	53·53²
		Analyst I	В.	
2.8530	44	5.2853	44	53.980
2.1252	"	3.9535	"	53.755
1.4248		2.6642	44	53-479
			Mean,	53.586, ± .03

The gravimetric series, of course, fixes the ratio between VOCl₃ and AgCl. If we put the latter at 100 parts, the proportion of VOCl₃ comes out as given in the third column:

Analyst A.

1.8521	grm. VOCl	gave 4.5932 g	rm. AgCl.	40.323	
.7013	"	1.7303	44	40.531	
.7486	66	1.8467	**	40.537	
1.4408	86	3.5719	46	40.337	
·9453	**	2.3399	46	40.399	
1.6183	44	4.0282	44	40.174	
		Analyst	<i>B</i> .		
2.1936	44	5.4039	44	40.391	
2.5054	**	6.2118	"	40.333	
			M	ean, 40.378, ± .0	28

These two series give us two values for the molecular weight of VOCl₃:

From the volumetric series ...
$$VOCl_8 = 173.096, \pm .126$$

" gravimetric " ... " = 173.276, $\pm .141$
General mean ... " = 173.177, $\pm .094$

Hence $V = 51.104, \pm .104$.

Combining the two values for V we get the following result:

Or, if O = 16, V = 51.373.

ARSENIC.

For the determination of the atomic weight of arsenic two compounds have been studied; the chloride and the trioxide. The bromide may also be considered, since it was analyzed by Wallace in order to establish the atomic weight of bromine. His series, in the light of more recent knowledge, may properly be inverted, and applied to the determination of arsenic.

In 1826, Berzelius* heated arsenic trioxide with sulphur

^{*} Poggend. Annal., 8, 1.

in such a way that only SO₂ could escape. 2.203 grammes of As₂O₃, thus treated, gave a loss of 1.069 of SO₂. Hence As = 74.840. This is a close estimation; but, being drawn from a single experiment, has so little weight that it need not be included in our final general mean.

In 1845 Pelouze* applied his method of titration with known quantities of pure silver to the analysis of the trichloride of arsenic, AsCl₃. Using the old Berzelian atomic weights, and putting Ag = 1349.01, and Cl = 443.2, he found in three experiments for As the values 937.9, 937.1, and 937.4. Hence 100 parts of silver balance the following quantities of AsCl₃:

Later, the same method was employed by Dumas,† whose weighings, reduced to the foregoing standard, give the following results:

4.298 gm	m. AsCl,	= 7.673 g	rm. Ag.	Ratio, 56.015
5-535	"	9.880	"	56.022
7.660	66	13.686	"	55.970
4.680	44	8.358	**	55-993
				Mean, \$6,000, + .00

The two series of Pelouze and Dumas, combined, give a general mean of 56.014, \pm .0035, as the amount of AsCl, equivalent to 100 parts of silver. Hence As = 74.829, \pm .048, a value closely agreeing with that deduced from the single experiment of Berzelius.

The same process of titration with silver was applied by Wallace‡ to the analysis of arsenic tribromide, AsBr₃. This compound was repeatedly distilled to ensure purity, and was well crystallized. His weighings show that the quanti-

^{*} Compt. Rend., 20, 1047. † Ann. Chim. Phys., (3,) 55, 174. 1859.

[†] Philosophical Magazine, (4,) 18, 279.

ties of bromide given in the third column are proportional to 100 parts of silver:

```
8.3246 grm. AsBr_8 = 8.58 grm. Ag. 97.023
4.4368 " 4.573 " 97.022
5.098 " 5.257 " 96.970
Mean, 97.005, \pm .012
```

Hence As = 74.046, $\pm .058$. Why this value should be so much lower than that from the chloride is unexplained.

The volumetric work done by Kessler,* for the purpose of establishing the atomic weights of chromium and of arsenic, has already been described in the chromium chapter. In that investigation the amount of potassium dichromate required to oxidize 100 parts of As₂O₃ to As₂O₅ was determined, and compared with the quantity of potassium chlorate necessary to produce the same effect. From the molecular weight of KClO₃, that of K₂Cr₂O₇ was then calculable.

From the same figures, the molecular weights of KClO₃ and of K₂Cr₂O₇ being both known, that of As₂O₃ may be easily determined. The quantities of the other compounds proportional to 100 parts of As₂O₃ are as follows:

$K_2Cr_2O_7$	KClO ₃ .
98.95	41.156
98.94	41.116
99.17	41.200
98.98	41.255
99.08	41.201
99.15	41.086
	41.199
Mean, 99.045, ± .028	41.224
•	41.161
	41.193
	41.149
	41.126
	Mean, 41.172, ± .009

^{*} Poggend. Annal., 95, 204. 1855. Also 113, 134. 1861.

Another series with the bichromate gave the following figures:

```
99.08

99.06

99.10

98.97

98.97

Mean, 99.036, ± .019

Mean of previous series, 99.045, ± .028

General mean, 99.039, ± .016
```

Other defective series are given to illustrate the partial oxidation of the As_2O_3 by action of air. The foregoing figures give us two distinct values for the molecular weight of As_2O_3 . In calculating from the bichromate results the value for chromium deduced from Siewert's determinations will be used, viz., $Cr = 52.009, \pm .025$.

Hence As = 75.002, $\pm .018$.

The general mean for As comes out as follows:

If O = 16, then As becomes = 75.090.

ANTIMONY.

After some earlier, unsatisfactory determinations, Berzelius,* in 1826, published his final estimation of the atomic weight of antimony. He oxidized the metal by means of nitric acid, and found that 100 parts of antimony gave 124.8 of Sb_2O_4 . Hence, if O=16, Sb=129.03. The

^{*} Poggend. Annal., 8, 1.

value 129 remained in general acceptance until 1855, when Kessler,* by special volumetric methods, showed that it was certainly much too high. Kessler's results will be considered more fully further along, in connection with a later paper; for present purposes a brief statement of his earlier conclusions will suffice. Antimony, and various compounds of antimony, were oxidized partly by potassium anhydrochromate and partly by potassium chlorate; and from the amounts of oxidizing agent required, the atomic weight in question was deduced:

The figures given are those calculated by Kessler himself. A recalculation with our newer atomic weights for O, K, Cl, Cr, S, and C, would yield slightly lower values. It will be seen that five of the estimates agree closely, while one diverges widely from the others. It will be shown hereafter that the concordant values are all vitiated by constant errors, and that the exceptional figure is after all the best.

Shortly after the appearance of Kessler's first paper, Schneider † published some results obtained by the reduction of antimony sulphide in hydrogen. The material chosen was a very pure stibnite from Arnsberg, of which the gangue was only quartz. This was corrected for, and corrections were also applied for traces of undecomposed sulphide carried off mechanically by the gas stream, and for traces of sulphur retained by the reduced antimony. The latter sulphur was estimated as barium sulphate. From 3.2 to 10.6 grammes of material were taken in each experiment. The final corrected percentages of S in Sb₂S₃ were as follows:

^{*} Poggend. Annal., 95, 215.

[†] Poggend. Annal., 98, 293. 1856. Preliminary note in Bd. 97.

```
28.559
28.557
28.501
28.554
28.532
28.485
28.492
28.481

Mean, 28.520, ± .008
```

Hence, if S = 32, Sb = 120.3.

Immediately after the appearance of Schneider's memoir, Rose * published the result of a single analysis of antimony trichloride, previously made under his supervision by Weber. This analysis, if Cl = 35.5, makes Sb = 120.7, a value of no great weight, but in a measure confirmatory of that obtained by Schneider.

The next research upon the atomic weight of antimony was that of Dexter,† published in 1857. This chemist, having tried to determine the amount of gold precipitable by a known weight of antimony, and having obtained discordant results, finally resorted to the original method of Berzelius. Antimony, purified with extreme care, was oxidized by nitric acid, and the gain in weight was determined. From 1.5 to 3.3 grammes of metal were used in each experiment. The reduction of the weights to a vacuum standard was neglected as being superfluous. From the data obtained, we get the following percentages of Sb in Sb₂O₄:

79.268 79.272 79.255 79.266 79.253 79.271 79.264 79.260 79.286

^{*} Poggend. Annal., 98, 455. 1856. † Poggend. Annal., 100, 563.

79.274 79.232 79.395 79.379

Mean, 79.283, ± .009

Hence, if O = 16, Sb = 122.46.

The determinations of Dumas* were published in 1859. This chemist sought to fix the ratio between silver and antimonicus chloride, and obtained results for the atomic weight of antimony quite near to those of Dexter. The SbCl_a was prepared by the action of dry chlorine upon pure antimony; it was distilled several times over antimony powder, and it seemed to be perfectly pure. Known weights of this preparation were added to solutions of tartaric acid in water, and the silver chloride was precipitated without previous removal of the antimony. Here, as Cooke has since shown, is a possible source of error, for under such circumstances the crystalline argento-antimonious tartrate may also be thrown down and contaminate the chloride of silver. But be that as it may; Dumas' weighings, reduced to a common standard, give as proportional to 100 parts of silver, the quantities of SbCl, which are stated in the third of the subjoined columns:

1.876 gr	m. SbCl _s	== 2.660 g	rm. Ag.	70.526
4.336	46	6. 148	"	70.527
5.06 5	44	7.175	"	70.592
3-475	**	4.930	"	70.487
3.767	44	5.350	66	70.411
5.910	46	8.393	44	70.416
4.828	44	6.836	**	70.626

Mean, 70.512, ± .021

Hence, if Ag = 108, and Cl = 355, Sb = 122.

In 1861 Kessler's second paper† relative to the atomic weight of antimony appeared. Kessler's methods were somewhat complicated, and for full details the original memoirs must be consulted. A standard solution of potassium anhydrochromate was prepared, containing 6.1466

^{*} Ann. Chim. Phys., (3,) 55, 175.

[†] Poggend. Annal., 113, 145.

grammes to the litre. With this, solutions containing known quantities of antimony or of antimony compounds were titrated, the end reaction being adjusted with a standard solution of ferrous chloride. In some cases the titration was preceded by the addition of a definite weight of potassium chlorate, insufficient for complete oxidation; the anhydrochromate then served to finish the reaction. The object in view was to determine the amount of oxidizing agent, and therefore of oxygen, necessary for the conversion of known quantities of antimonious into antimonic compounds.

In the later paper Kessler refers to his earlier work, and shows that the values then found for antimony were all too high, except in the case of the series made with tartar emetic. That series he merely states, and subsequently ignores, evidently believing it to be unworthy of further consideration. For the remaining series he points out the sources of error. These need not be rediscussed here, as the discussion would have no value for present purposes; suffice it to say that in the series representing the oxidation of Sb_2O_3 with anhydrochromate and chlorate, the material used was found to be impure. Upon estimating the impurity and correcting for it, the earlier value of Sb = 123.80 becomes Sb = 122.36, according to Kessler's calculations.

In the paper now under consideration four series of results are given. The first represents experiments made upon a pure antimony trioxide which had been sublimed, and which consisted of shining colorless needles. This was dissolved, together with some potassium chlorate, in hydrochloric acid, and titrated with anhydrochromate solution. Six experiments were made, but Kessler rejects the first and second as untrustworthy. The data for the others are as follows:

Sb_2O_3 .	KClO ₃ .	K ₂ Cr ₂ O ₇ sol. in cc.
1.7888 grm.	.4527 grm.	19.2 cc.
1.6523 "	.4506 "	3.9 "
3.2998 "	.8806 "	16.5 "
1.3438 "	.3492 "	10.2 "

From these figures Kessler deduces Sb = 122.16.

These data, reduced to a common standard, give the following quantities of oxygen needed to oxidize 100 parts of Sb₂O₃ to Sb₂O₅. Each cubic centimetre of the K₂Cr₂O₇ solution corresponds to one milligramme of O:

In the second series of experiments pure antimony was dissolved in hydrochloric acid with the aid of an unweighed quantity of potassium chlorate. The solution, containing both antimonious and antimonic compounds, was then reduced entirely to the antimonious condition by means of stannous chloride. The excess of the latter was corrected with a strong hydrochloric acid solution of mercuric chloride, then, after diluting and filtering, a weighed quantity of potassium chlorate was added, and the titration with anhydrochromate was performed as usual. Calculated as above, the percentages of oxygen given in the last column correspond to 100 parts of antimony:

Sb.	KClO ₃ .	K2Cr2O, sol. cc.	Per cent. O.
1.636 grm.	0.5000 grm.	18.3	13.088
3.0825• "	0.9500 "	30.2	13.050
4.5652 "	1.4106 "	45-5	13.098

Mean, 13.079, ± .0096

This series gave Kessler Sb = 122.34.

The third and fourth series of experiments were made with pure antimony trichloride, SbCl₃, prepared by the action of mercuric chloride upon metallic antimony. This preparation, in the third series, was dissolved in hydrochloric acid, and titrated. In one experiment solid K₂Cr₂O₇ in weighed amount was added before titration: in the other two estimations KClO₃ was taken as usual. If, according to Siewert's work, we take Cr = 52.009, the percentages of oxygen in the last column correspond to 100 parts of SbCl₃:

```
Per cent. O.

1.8576 grm. SbCl<sub>3</sub> needed .5967 grm. K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub> and 33.4 cc. sol. 7.0338

1.9118 " .3019 " KClO<sub>3</sub> " 16.2 " 7.0321

4.1235 " .6801 " " " 23.2 " 7.0222

Mean, 7.0294, ± .0024
```

The fourth set of experiments was gravimetric. The solution of SbCl₃, mixed with tartaric acid, was first precipitated by hydrogen sulphide, in order to remove the antimony. The excess of H₂S was corrected by copper sulphate, and then the chlorine was estimated as silver chloride in the ordinary manner. 100 parts of AgCl correspond to the amounts of SbCl₃ given in the third column.

1.8662	grm. SbCl ₃ gave	3.483 grm.	AgCl.	53.580
1.6832	66	3.141	64	53.588
2.7437	44	5.1115	66	53.677
2.6798	66	5.0025	**	53.569
5.047	44	9.411	• •	53.629
3.8975	44	7.2585	• 6	53.696

Mean, 53.623, ± .015

The volumetric series with SbCl₃ gave Kessler values for Sb ranging from 121.16 to 121.47. The gravimetric series, on the other hand, yielded results from Sb = 124.12 to 124.67. This discrepancy Kessler rightly attributes to the presence of oxygen in the chloride; and, ingeniously correcting for this error, he deduces from both sets combined, the value of Sb = 122.37.

The several mean results for antimony agree so fairly with each other, and with the estimates obtained by Dexter and Dumas, that we cannot wonder that Kessler felt satisfied of their general correctness, and of the inaccuracy of the figures published by Schneider. Still, the old series of data obtained by the titration of tartar emetic with anhydrochromate contained no evident errors, and was not accounted for. This series,* if we reduce all of Kessler's figures to a single common standard, give a ratio between K₂Cr₂O₃ and C₄H₄KSbO₇. H₂O. 100 parts of the former will oxidize of the latter:

^{*} Poggend. Annal., 95, 217.

336.64 338.01 336.83 337.93 338.59 335.79

Mean, 337.30, \pm .29

From this, if $K_2Cr_2O_7 = 294.64$, Sb = 119.8.

The newer atomic weights found in the previous chapters of this work will be applied to the discussion of all these series further along. It may, however, be properly noted at this point, that the probable errors assigned to the percentages of oxygen in three of Kessler's series are too low. These percentages are calculated from the quantities of KClO₃ involved in the several reactions, and their probable errors should be increased with reference to the probable error of the molecular weight of that salt. The necessary calculations would be more laborious than the importance of the figures would warrant, and, accordingly, in computing the final general mean for antimony, Kessler's figures will receive somewhat higher weight than they are legitimately entitled to.

Naturally, the concordant results of Dexter, Kessler, and Dumas led to the general acceptance of the value of 122 for antimony as against the lower figure 120 of Schneider. Still, in 1871, Unger* published the results of a single analysis of Schlippe's salt, Na₃SbS₄.9H₂O. This analysis gave Sb = 119.76, if S = 32 and Na = 23, but no great weight could be attached to the determination. It served, nevertheless, to show that the controversy over the atomic weight of antimony was not finally settled.

More than ten years after the appearance of Kessler's second paper the subject of the atomic weight of antimony was again taken up, this time by Professor Cooke. His results appeared in the autumn of 1877,† and were conclusive in favor of the lower value, approximately 120. For full

^{*} Archiv. der Pharmacie, 197, 194. Quoted by Cooke.

[†] Proceedings American Academy, v. 13.

details the original memoir must be consulted; only a few of the leading points can be cited here.

Schneider analyzed a sulphide of antimony which was already formed. Cooke, reversing the method, effected the synthesis of this compound. Known weights of pure antimony were dissolved in hydrochloric acid containing a little nitric acid. In this solution weighed balls of antimony were boiled until the liquid became colorless; subsequently the weight of metal lost by the balls was ascertained. To the solution, which now contained only antimonious compounds, tartaric acid was added, and then, with a supersaturated aqueous sulphhydric acid, antimony trisulphide was precipitated. The precipitate was collected by an ingenious process of reverse filtration, converted into the black modification by drying at 210°, and weighed. After weighing, the Sb, S, was dissolved in hydrochloric acid, leaving a carbonaceous residue unacted upon. This was carefully estimated and corrected for. About two grammes of antimony were taken in each experiment and thirteen syntheses were performed. In two of these, however, the antimony trisulphide was weighed only in the red modification, and the results were uncorrected by conversion into the black variety and estimation of the carbonaceous residue. In fact, every such conversion and correction was preceded by a weighing of the red modification of the Sb.S.. The mean result of these weighings, if S = 32, gave Sb = 119.994. The mean result of the corrected syntheses gave Sb = 120.295. In these eleven experiments the following percentages of S in Sb₂S₃ were established:

```
28.57

28.60

28.57

28.43

28.42

28.53

28.50

28.49

28.58

28.50

28.51

Mean, 28.5182, ± .0120
```

These results, confirmatory of the work of Schneider, were presented to the American Academy in 1876. Still, before publication, Cooke thought it best to repeat the work of Dumas, in order to detect the cause of the old discrepancy between the values Sb = 120 and Sb = 122. Accordingly, various samples of antimony trichloride were taken, and purified by repeated distillations. The final distillate was further subjected to several recrystallizations from the fused state; or, in one case, from a saturated solution in bisulphide of carbon. The portions analyzed were dissolved in concentrated aqueous tartaric acid, and precipitated by silver nitrate, many precautions being observed. The silver chloride was collected by reverse filtration, and dried at temperatures from 110° to 120°. In one experiment the antimony was first removed by H,S. Seventeen experiments were made, giving, if Ag = 108 and Cl = 35.5, a mean value of Sb = 121.94. If we reduce to a common standard, Cooke's analyses give, as proportional to 100 parts of AgCl, the quantities of SbCl_a stated in the third column:

1.5974	grm. SbCl ₃	gave 3.0124	grm. AgCl.	53.028
1.2533	"	2.3620	**	53.061
.8876	**	1.6754	44	52.978
.8336	**	1.5674	44	53.184
.5326	66	1.0021	46	53.148
.7270	**	1.3691	**	53.101
1.2679	66	2.3883	44	53.088
1.9422	44	3.6646	66	52.999
1.7702	••	3.3384	**	53.025
2.5030	**	4.7184	"	53.048
2.1450	44	4.0410	46	53.081
1.7697	**	3.3281	fe .	53.175
2.3435	44	4.4157	46	53.072
1.3686	44	2.5813	• 44	53.020
1.8638	44	3.5146	**	53.030
2.0300	66	3.8282	46	53.028
2.4450	44	4.6086	• •	53.053

Mean, 53.066, ± .0096

This mean may be combined with that of Kessler's series, as follows:

Kessler		
General mean	53.2311.	+ .008

The results thus obtained with SbCl, confirmed Dumas' determination of the atomic weight of antimony as remarkably as the syntheses of Sb,S, had sustained the work of Schneider. Evidently, in one or the other series a constant error must be hidden, and much time was spent by Cooke in searching for it. It was eventually found that the chloride of antimony invariably contained traces of oxychloride, an impurity which tended to increase the apparent atomic weight of the metal under consideration. If was also found, in the course of the investigation, that hydrochloric acid solutions of antimonious compounds oxidize in the air during boiling as rapidly as ferrous compounds; a fact which explains the high values for antimony found by Kessler.

In order to render "assurance doubly sure," Professor Cooke also undertook the analysis of the bromide and the iodide of antimony. The bromide, SbBr₃, was prepared by adding the finely powdered metal to a solution of bromine in carbon disulphide. It was purified by repeated distillation over pulverized antimony, and by several recrystallizations from bisulphide of carbon. The bromine determinations resembled those of chlorine, and gave, if Ag = 108 and Br = 80, a mean value for antimony of Sb = 120. Reduced to a common standard, the fifteen analyses give the subjoined quantities of SbBr₃ proportional to 100 parts of silver bromide:

1.8621	grm. SbBr, ga	ve 2.9216 gr	m. AgBr.	63.736
.9856	"	1.5422	41	63.909
1.8650	44	2.9268	**	63.721
1.5330	"	2.4030	44	63.795
1.3689	44	2.1445	44	63.833
1.2124	44	1.8991	44	63.841
.9417	"	1.4749	**	63.848
2.5404	**	3.9755	"	63.901
1.5269	44	2.3905	46	63.874
1.8604	66	2.9180	44	68.756
1.7298	46	2.7083	**	63.870

3.2838	grm. SbBr ₃ gave	5.1398	grm. AgBr.	63.890
2.3589	44	3.6959	44	63.825
1.3323	¥	2.0863	66	63.859
2.6974	44	4.2285	44	63.791

Mean, 63.830, $\pm .008$

The iodide of antimony was prepared like the bromide, and analyzed in the same way. At first, discordant results were obtained, due to the presence of oxyiodide in the iodide studied. The impurity, however, was removed by subliming the iodide in an atmosphere of dry carbon dioxide. With this purer material, seven estimations of iodine were made, giving, if Ag = 108 and I = 127, a value for antimony of Sb = 120. Reduced to a uniform standard, Cooke's weighings give the following quantities of SbI_3 proportional to 100 parts of silver iodide:

1.1877 g	rm. SbI _a gave	1.6727	grm. AgI.	71.005
.4610	44	.6497	46	70.956
3.2527	**	4.5716	**	71.150
1.8068	66	2.5389	44	71.165
1.5970	**	2.2456	66	71.117
2.3201	66	3.2645	44	71.071
.3496	".	.4927	"•	70.956

Mean, 71.060, \pm .023

Although Cooke's work was practically conclusive, as between the rival values for antimony, his results were severely criticized by Kessler,* who, evidently, had read Cooke's paper in a very careless way. On the other hand, Schneider published in Poggendorff's Annalen a friendly review of the new determinations, which so splendidly vindicated his own accuracy. In reply to Kessler, Cooke undertook still another series of experiments with antimony bromide,† and obtained absolute confirmation of his previous results. To a solution of antimony bromide was added a solution containing a known weight of silver not quite sufficient to precipitate all the bromine. The excess

^{*} Berichte d. Deutsch. Chem. Gesell., 12, 1044. 1879.

[†] Amer. Journ. Sci. and Arts, May, 1880. Berichte, 13, 951.

of the latter was estimated by titration with a normal silver solution. Five analyses gave values for antimony ranging from 119.98 to 120.02, when Ag = 108 and Br = 80. Reduced to a common standard, the weights obtained gave the amounts of SbBr₂ stated in the third column as proportional to 100 parts of silver:

```
2.5032 grm. SbBr<sub>3</sub> = 2.2528 grm. Ag. III.115

2.0567 " 1.8509 " III.119

2.6512 " 2.3860 " III.115

3.3053 " 2.9749 " III.106

2.7495 " 2.4745 " III.113

Mean, III.114, ± .0014
```

Schneider,* also, in order to more fully answer Kessler's objections, repeated his work upon the Arnsberg stibnite. This he reduced in hydrogen as before, correcting scrupulously for impurities. The following percentages of sulphur were found:

These figures confirm his old results, and may be fairly combined with them and with the percentages found by Cooke, as follows:

```
Schneider, early series 28.520, ± .008

" late " 28.541, ± .0024

Cooke 28.5182, ± .0120

General mean 28.5385, ± .0023
```

We have now before us the following ratios, good and bad, from which to calculate the atomic weight of antimony. The single results obtained by Weber and by Unger, being unimportant, are not included:

```
(1.) Percentage of S in Sh_2S_p, 28.5385, \pm .0023

(2.) "Sb in Sh_2O_4, 79.283, \pm .009

(3.) O needed to oxidize 100 parts ShCl_3, 7.0294, \pm .0024

(4.) O "Sh_2O_3, 10.953, \pm .0075

(5.) O "Sb, 13.079, \pm .0096
```

The r determi ably as Schneid error m in searc ride of an imp weight in the solution ing boi explair In o Cooke : iodide . adding in carl tion ov tions fi tions r and B Reduc the sul

> I I I

1

of silv

2

1

^{*} Journ. für Prakt. Chem., (2,) 22, 131.

```
(6.) K<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>: tartar emetic :: 100: 337.30, ± .29
(7.) Ag: SbCl<sub>8</sub>:: 100: 70.512, ± .021
(8.) AgCl: SbCl<sub>8</sub>:: 100: 53.2311, ± .008
(9.) Ag: SbBr<sub>8</sub>:: 100: 111.114, ± .0014
(10.) AgBr: SbBr<sub>8</sub>:: 100: 63.830, ± .008
(11.) AgI: SbI<sub>8</sub>:: 100: 71.060, ± .023
```

Three of these ratios give estimates for the molecular veight of antimony trichloride, and two give corresponding values for the bromide. These values may be combined, as follows: First, for the chloride we have—

Hence Sb = 122.115, $\pm .055$.

For the bromide we get:

: 15

Hence Sb = 119.625, $\pm .063$.

From all the data eight values for Sb may be deduced. These fall into two groups; the one near the number 120, the other not far from 122. In making the calculation the atomic weights found in previous chapters are applied; the value selected for chromium being that deduced from Siewert's experiments:

Although the means of the four lower values and of the four higher values are thus shown to be approximately

equal in weight, we know from Cooke's experiments that the larger mean is vitiated by serious constant errors. Only in value 5, the result calculated from Dexter's experiments, has the constant error not been pointed out. Cooke considers it probable, however, that the Sb, O_4 involved in this work contained traces of some lower oxide, which, if present, would render the atomic weight of antimony apparently too high. Chemically considered, the preponderance of evidence is strongly in favor of values 1 to 3, deduced from the experiments of Schneider and of Cooke. These give a general mean of Sb = 119.955, \pm .036; or, if O = 16, this becomes Sb = 120.231.

This we may accept as most nearly the true result, and reject the data of Dexter, Dumas, and Kessler altogether.

Since this chapter was written, Pfeifer has compared the amount of antimony thrown down electrolytically, with the quantity of silver deposited by the same current in the same time. From rather meagre data he concludes that the atomic weight of antimony, thus determined, may be 121. Additional investigation is promised. The figures thus far published would weigh little as against Cooke's experiments. (Ann. Chem. Pharm., 209, 161. 1881.)

BISMUTH.

Early in the century the combining weight of bismuth was approximately fixed through the experiments of Lagerhjelm.* Effecting the direct union of bismuth and sulphur, he found that ten parts of the metal yield the following quantities of trisulphide:

^{*} Annals of Philosophy, 4, 358. 1814. Results adopted by Berzelius.

203

Hence B=215 in round numbers, a value now known to be much too high. Lagerhjelm also oxidized bismuth with nitric acid, and, after ignition, weighed the trioxide thus formed. Ten parts of metal gave the following quantities of Bi_2O_3 :

Hence, if O = 16, Bi = 211.85, a figure still too high.

In 1851 the subject of the atomic weight of bismuth was taken up by Schneider,* who, like Lagerhjelm, studied the oxidation of the metal with nitric acid. The work was executed with a variety of experimental refinements, by means of which every error due to possible loss of material was carefully avoided. For full details the original paper must be consulted; there is only room in these pages for the actual results, as follows. The figures represent the percentages of Bi in Bi₂O₃:

```
89.652
89.682
89.644
89.634
89.656
89.666
89.655
```

Mean, 89.6552, ± .0034

Hence Bi = 207.523, $\pm .082$; or, if O = 16, Bi = 208.001.

Finally, we come to the results obtained by Dumas.† Bismuth trichloride was prepared by the action of dry chlorine upon bismuth, and repeatedly rectified by distillation over bismuth powder. The product was weighed in a closed tube, dissolved in water, and precipitated with sodium carbonate. In the filtrate, after strongly acidulating.

^{*} Poggend. Annal., 82, 303. 1851. † Ann. de Chim. et de Phys., (3,) 55, 176. 1859.

with nitric acid, the chlorine was precipitated by a known amount of silver. The figures in the third column show the quantities of BiCl₂ proportional to 100 parts of silver:

3.506 gm	n. BiCl _s :	= 3.545 g	rm. Ag.	98.900
1.149	"	1.168	"	98.373
1.5965	46	1.629	66	98.005
2.1767	"	2.225	"	97.829
3.081	"	3.144	44	97.996
2.4158	"	2.470	66	97.806
1.7107	44	1.752	66	97.643
3.523	44	3.6055	" .	97.712
5.241	66	5.361	44	97.762

Mean, 98.003, ± .090

Hence Bi = 210.464, $\pm .294$.

The first three of the foregoing series of experiments were made with slightly discolored material, and may therefore be rejected. The remaining six percentages give a mean of 97.791; whence Bi = 209.78; or, if O = 16, Bi = 210.26.

As between the unaccordant results of Schneider and of Dumas, those of the former chemist are probably nearest correct. His method of determination was the more reliable, and the details which he gives concerning his manipulations afford strong presumptions of accuracy. Doubtless the bismuth trichloride used by Dumas, contained, like the corresponding antimony compounds, traces of oxychloride. We may fairly assume, for all practical purposes, that the atomic weight of bismuth cannot be far from 208.

TIN.

Stannic oxide and stannic chloride are the compounds which have been studied in estimating the atomic weight of tin.

The composition of stannic oxide has been fixed in two ways; by synthesis from the metal, and by reduction in hydrogen. For the first method we may consider the work of Berzelius, Mulder and Vlaanderen, and Dumas.

TIN. 205

Berzelius* oxidized 100 parts of tin by nitric acid, and found that 127.2 parts of SnO₂ were formed.

The work done by Mulder and Vlaanderen † was done in connection with a long investigation into the composition of Banca tin, which was found to be almost absolutely pure. For the atomic weight determinations, however, really pure tin was taken, prepared from pure tin oxide. This metal was oxidized by nitric acid, with the following results. 100 parts of tin gave of SnO₂:

Dumas toxidized pure tin by nitric acid in a flask of glass. The resulting SnO₂ was strongly ignited, first in the flask, and afterwards in platinum. His weighings, reduced to the foregoing standard, give for dioxide from 100 parts of tin the amounts stated in the third column:

In an investigation later than that previously cited, Vlaanderen \parallel found that when tin was oxidized in glass or porcelain vessels, and the resulting oxide ignited in them, traces of nitric acid were retained. When, on the other hand, the oxide was strongly heated in platinum, the latter was perceptibly attacked, so much so as to render the results uncertain. He therefore, in order to fix the atomic weight of tin, reduced the oxide by heating it in a porcelain boat in a stream of hydrogen. Two experiments gave Sn = 118.08, and Sn = 118.24. These, when O = 16, become, if reduced to the above common standard,

^{*} Poggend. Annal., 8, 177. † Journ. für Prakt. Chem., 49, 35. 1849.

[†] Ann. Chem. Pharm., 113, 26.

^{||} Jahresbericht, 1858, 183.

We have now four series of results showing the quantity of SnO, formed from 100 parts of tin. To Berzelius' single value may be assigned the probable error of a single experiment in Mulder and Vlaanderen's series:

Berzelius	127.200, \pm .041—Oxidation.
Mulder and Vlaanderen	127.517, ± .029- "
Dumas	127.105, ± .024— "
Vlaanderen	127.082, ± .012—Reduction.
	
General mean	$127.143, \pm .0098$

Dumas, in the paper previously quoted, also gives the results of some experiments with stannic chloride, SnCl₄. This was titrated with a solution containing a known weight of silver. From the weighings given, 100 parts of silver correspond to the quantities of SnCl₄ named in the third column:

All these data properly combined give us the following values for the atomic weight of tin:

If O = 16, this becomes Sn = 117.968.

TITANIUM.

The earliest determinations of the atomic weight of titanium are due to Heinrich Rose.* In his first investigation he studied the conversion of titanium sulphide into titanic acid, and obtained erroneous results; later, in 1829, he published his analyses of the chloride.† This compound was purified by repeated rectifications over mercury and over potassium, and was weighed in bulbs of thin glass. These were broken under water in tightly stoppered flasks; the titanic acid was precipitated by ammonia, and the chlorine was estimated as silver chloride. The following results were obtained. In a fourth column I give the TiO, in percentages referred to TiCl₄ as 100; and in a fifth column the quantity of TiCl₄ proportional to 100 parts of AgCl:

TiCl4.	TiO2.	AgCl.	Per cent. TiOz.	AgCl Ratio.
.885 grm.	.379 grm.	2.661 grm.	42.825	33.258
2.6365 "	1.120 "	7.954 "	42.481	33.147
1.7157 "	.732 "	5.172 "	42.665	33.173
3.0455 "	1.322 "	9.198"	43.423	33.100
2.4403 "	1.056 "	7.372 "	43.273	33.102
		M	ean, 42.933, ± .12	1 33.156, ± .019

If we directly compare the AgCl with the TiO, we shall find 100 parts of the former proportional to the following quantities of the latter:

From all these figures we can get three values for Ti, thus:

[#] Gilbert's Annalen, 1823, 67 and 129.

⁺ Poggend. Annal., 15, 145. Berz. Lehrbuch, 3, 1210.

These results will be discussed further along in connection with others.

Shortly after the appearance of Rose's paper, Mosander* published some figures giving the percentages of oxygen in titanium dioxide, from which a value for the atomic weight of titanium was deduced. Although no details are furnished as to experimental methods, and no actual weighings are given, I cite his percentages for whatever they may be worth:

40.814 40.825 40.610 40.180 40.107 40.050 40.780 40.660 39.830 Mean, 40.428

These figures give values for Ti ranging from 46.277 to 48.231; or, in mean, Ti = 47.045. They are not, however, sufficiently explicit to deserve any further consideration. It will be noticed that the highest value nearly coincides with Rose's lowest.

In 1847 Isidor Pierre made public a series of important determinations.† Titanium chloride, free from silicon and from iron, was prepared by the action of chlorine upon a mixture of carbon with pure, artificial, titanic acid. This chloride was weighed in sealed tubes, these were broken under water, and the resulting hydrochloric acid was titrated with a standard solution of silver after the method

^{*} Berz. Jahresbericht, 10, 108. 1831.

[†] Ann. de Chim. et Phys., (3,) 20, 257.

of Pelouze. I subjoin Pierre's weighings, and add, in a third column, the ratio of TiCl₄ to 100 parts of silver:

TiCl ₄ .	Ag.	Ratio.
.8215 grm.	1.84523 grm.	44.520
.7740 "	1.73909 "	44.506
·7775 "	1.74613 "	44.527
.7160 "	1.61219 "	44.412
.8085 "	1.82344 "	44.339
.6325 "	1.42230 "	44.470
.8155 "	1.83705 "	44.392
.8165 "	1.83899 "	44.399
.8065 "	1.81965 "	44.322

Mean, 44.432, ± .0173

It will be seen that the first three of these results agree well with each other and are much higher than the remaining six. The last four experiments were made purposely with tubes which had been previously opened, in order to determine the cause of the discrepancy. According to Pierre, the opening of a tube of titanium chloride admits a trace of atmospheric moisture. This causes a deposit of titanic acid near the mouth of the tube, and liberates hydrochloric acid. The latter gas being heavy, a part of it falls back into the tube, so that the remaining chloride is richer in chlorine and poorer in titanium than it should be. Hence, upon titration, too low figures for the atomic weight of titanium are obtained. Pierre accordingly rejects all but the first three of the above estimations:

From all of Pierre's......Ti = 49.889,
$$\pm$$
 .096 " the first three......" = 50.259, \pm .063

The memoir of Pierre upon the atomic weight of titanium was soon followed by a paper from Demoly,* who obtained much higher results. He also started out from titanic chloride, which was prepared from rutile. The latter substance was found to contain 1.8 per cent. of silica; whence Demoly inferred that the TiCl₄ investigated by Rose and by Pierre

^{*} Ann. Chem. Pharm., 72, 214. 1849. Berz. Jahresb., 30, 58.

might have been contaminated with SiCl, an impurity which would lower the value deduced for the atomic weight under consideration. Accordingly, in order to eliminate all such possible impurities, this process was resorted to: the chloride, after rectification over mercury and potassium, was acted upon by dry ammonia, whereupon the compound TiCl. 4NH, was deposited as a white powder. This was ignited in dry ammonia gas, and the residue, by means of chlorine, was reconverted into titanic chloride, which was again repeatedly rectified over mercury, potassium, and potassium amalgam. The product boiled steadily at 135°. This chloride, after weighing in a glass bulb, was decomposed by water, the titanic acid was precipitated by ammonia, and the chlorine was estimated in the filtrate as silver chloride. Three analyses were performed, yielding the following results. I give the actual weighings:

```
1.470 grm. TiCl<sub>4</sub> gave 4.241 grm. AgCl and .565 grm. TiO<sub>2</sub>.
2.330 " 6.752 " .801 "
2.880 " 8.330 " 1.088 "
```

The ".801" in the last column is certainly a misprint for .901. Assuming this correction, the results may be given in three ratios, thus:

Per cent. TiO2 from TiCl4.	TiCl, : 100 AgCl.	TiO2: 100 AgCl.
38.435	34.662	13.322
38.669	34.508	13.344
37.778	34-574	13.061
Mean, 38.294, ± .180	34.581, ± .030	13.242, ± .061

These three ratios give three widely divergent values for the atomic weight of titaniun;

The value assumed by Demoly is 56; who employs but one ratio and ignores practically the others.

Upon comparing Demoly's figures with those obtained by Rose, certain points of similarity are plainly to be noted. Both sets of results were reached by essentially the same method; and in both the discordance between the percentages of titanic acid and of silver chloride is glaring. This discordance can rationally be accounted for by assuming that the titanic chloride was in neither case absolutely what it purported to be; that, in brief, it must have contained impurities; such for example as hydrochloric acid, as shown in the experiments of Pierre, or possibly traces of oxychlorides. Considerations of this kind also throw doubt upon the results attained by Pierre, for he neglected the direct estimation of the titanic acid altogether, thus leaving us without means for correctly judging as to the character of his material. In fact, not one of the determinations of the atomic weight of titanium can be regarded as trustworthy. All depend upon the chloride, and the volatile chlorides of metals are as a class especially liable to contaminations of a kind most difficult to recognize. Possibly a series of good determinations might be based upon analyses of some of the titanofluorides. I subjoin a combination of the foregoing mean values, feeling that such a general average is a little better than any one set of determinations taken singly:

```
From Rose's analyses......Ti = 48.710, ± .105

" Pierre's " ......" = 49.889, ± .096

" Demoly's " = 52.191, ± .153

General mean....." = 49.846, ± .064

Or, if O = 16, Ti = 49.961.
```

This mean agrees with the average of all of Pierre's experiments.

ZIRCONIUM.

The atomic weight of zirconium has been determined by Berzelius, by Hermann, and by Marignac. Berzelius* ignited the neutral sulphate, and thus ascertained the ratio in it between the ZrO₂ and the SO₃. Putting SO₃ at 100, he gives the following proportional quantities of ZrO₂:

```
75.84
75.92
75.80
75.74
75.97
75.85
Mean, 75.853, ± .023
```

Hence Zr = 89.255, $\pm .039$; or, if O = 16, then Zr = 89.461.

Hermann's† estimate of the atomic weight of zirconium was based upon analyses of the chloride, concerning which he gives no details or weighings. From sublimed zirconium chloride he finds Zr = 831.8, when O = 100; and from two lots of the basic chloride $2ZrOCl_2.9H_2O$, Zr = 835.65 and 851.40 respectively. The mean of all three is 839.62; whence, with modern formulæ and O = 15.9633, Zr becomes = 89.354.

Marignac's results! were obtained by analyzing the double fluoride of zirconium and potassium. His weights are as follows:

```
1.000 grm. gave .431 grm. ZrO, and .613 grm. K2SO4.
         "
                .864
                           "
                                   1.232
                           "
                                             "
         "
                .282
 .654
                                    .399
                           "
5.000
         "
               2.169
                                   3.078
```

These figures give us three ratios. A, the ZrO, from 100

^{*} Poggend. Annal., 4, 126. 1825.

[†] Journ. für Prakt. Chem., 31, 77. Berz. Jahresb., 25, 147.

[‡] Ann. Chim. Phys., (3,) 60, 270. 1860.

parts of salt; B, the K, SO₄ from 100 parts of salt; and C, the ZrO₂ proportional to 100 parts of K, SO₄:

В.	C.
61.300	70.310
61.600	70.130
61.000	70.677
· 61.560	70.468
61.365, ± .094	70.396 , ± .079
Zr = 89.7 " = 91.4 " = 90.4	
	61.300 61.600 61.000 61.560 61.365, ± .094

General mean...... $= 90.328, \pm .113$

Or, if O = 16, Zr = 90.536.

Combining with Berzelius' work we get this result:

Or, if O = 16, Zr = 89.573.

These figures need little criticism. They show conclusively that the atomic weight of zirconium ought to be redetermined. Probably the method employed by Berzelius was the best with respect to manipulation, while on the other hand it is likely that Marignac worked with purer material. Hermann's experiments could hardly have yielded certain results, since the zirconium chloride might so easily become contaminated with traces of moisture and thence of oxygen.

THORIUM.

The atomic weight of thorium has been determined from analyses of the sulphate, oxalate, formate, and acetate, with widely varying results. The earliest figures are due to Berzelius,* who worked with the sulphate, and with the double sulphate of potassium and thorium. The thoria was precipitated by ammonia, and the sulphuric acid was estimated as BaSO₄. The sulphate gave the following ratios in two experiments. The third column represents the weight of ThO₂ proportional to 100 parts of BaSO₄:

The double potassium sulphate gave .265 grm. ThO₂, .156 grm. SO₃, and .3435 K₂SO₄. The SO₃, with the Berzelian atomic weights, represents .4537 grm. BaSO₄. Hence 100 BaSO₄ is equivalent to 58.408 ThO₃. This figure, combined with the two previous values for the same ratio, give a mean of 58.026, \pm .214. Hence ThO₂ = 269.940, \pm .997.

From the ratio between the K_2SO_4 and the ThO₂ in the double sulphate, ThO₂ = 268.284.

In 1861 new determinations were published by Chydenius,† whose memoir is accessible to me only in an abstract; which gives results without details. Thoria is regarded as a monoxide, ThO, and the old equivalents (O = 8) are used. The following values are assigned for the molecular weight of ThO, as found from analyses of several salts:

From Sulphate.	From K. Th. Sulphate.
66.33	67.02
67.13	
67.75	
68. 03	
Mean, 67	7.252, ± .201

^{*} Poggend. Annal., 16, 398. 1829. Lehrbuch, 3, 1224.

[†] Kemisk undersökning af Thorjord och Thorsalter. Helsingfors, 1861. An academic dissertation.

[†] Poggend. Annal., 119, 55. 1863.

Fro	m Acetate.	From Formate.	From Oxalate.
	67.31	68.06	65.87 \ Two results
	66.59	67.89	65.95 by Berlin.
	67.27	68.94	65.75
	67.06		65.13
	68.40	Mean, 68.297, ± .219	66.54
			65.85
Mean,	67.326, ± .201		
		Me	ean, 65.85, ± .123

We may fairly assume that these figures were calculated with O = 8, C = 6, and S = 16. Correcting by the values for these elements which have been found in previous chapters, ThO₂ becomes as follows:

The single result from the double potassium sulphate is included with the column from the ordinary sulphate, and the influence of the atomic weight of potassium is ignored.

Chydenius was soon followed by Marc Delafontaine, whose researches appeared in 1863.* This chemist especially studied thorium sulphate; partly in its most hydrous form, partly as thrown down by boiling. In Th(SO₄)₂.9H₂O, the following percentages of ThO₂ were found:

```
45.08

44.90

45.06

45.21

45.06

Mean, 45.062, ± .0332
```

Hence ThO, = 263.637, $\pm .256$.

The lower hydrate, $2\text{Th}(SO_4)_2.9\text{H}_2\text{O}$, was more thoroughly investigated. The thoria was estimated in two ways; first, (A,) by precipitation as oxalate and subsequent ignition; second, (B,) by direct calcination. These percentages of ThO₂ were found:

^{*} Arch. des Sci. Phys. et Nat., (2,) 18, 343.

Mean, 52.511, ± .047

Hence ThO₂ = 266.025, $\pm .363$.

In three experiments with this lower hydrate the sulphuric acid was also estimated, being thrown down as barium sulphate after removal of the thoria:

```
1.2425 grm. gave .400 SO<sub>2</sub>. (1.1656 grm. BaSO<sub>4</sub>.)
1.138 " .366 " (1.0665 " )
.734 " .2306 " (.6720 " )
```

The figures in parenthesis are reproduced by myself from Delafontaine's results, he having calculated his analyses with O=100, S=200, and Ba=857. These data may be reduced to a common standard, so as to represent the quantity of $2\text{Th}(SO_4)_2.9\text{H}_2O$ equivalent to 100 parts of $BaSO_4$. We then have the following results:

```
106.597
106.704
109.226
Mean, 107.509, ± .585
```

Hence ThO₂ = 259.555, ± 2.725 .

Delafontaine seems himself to have calculated from the ratio between the percentages of SO₃ and ThO₂; whence, with our revised values for S, Ba, and O, ThO₂ = 262.643.

Delafontaine's work was soon confirmed by Hermann,

^{*} Journ. für Prakt. Chem., 93, 114.

who published a single analysis of the lower hydrated sulphate, as follows:

Hence, from the ratio between SO₃ and ThO₂, ThO₂ = 263.030. Probably the SO₃ percentage was loss upon calcination.

The latest, and probably also the best determinations, are those of Cleve,* whose results, obtained from both the sulphate and the oxalate of thorium, agree admirably. The anhydrous sulphate, calcined, gave the subjoined percentages of thoria:

Hence ThO₂ = 265.380, $\pm .123$.

The oxalate was subjected to a combustion analysis, whereby both thoria and carbonic acid could be estimated. From the direct percentages of these constituents no accurate value can be deduced, there having undoubtedly been moisture in the material studied. From the ratio between CO₂ and ThO₂, however, good results are attainable. This ratio I put in a fourth column, making the thoria proportional to 100 parts of carbon dioxide:

Oxalate.	$Th O_2$.	CO ₂ .	Ratio.
1.7135 grm.	1.0189 grm.	.6736 grm.	151.262
1.3800 "	.8210 "	·5433 "	151.114
1.1850 "	.7030 "	.4650 "	151.183
1.0755 "	.6398 "	.4240 "	150.896
•			n, 151.114, ± .053
Hence ThO	$_{2}=265.357, \pm$	∟ .104.	

^{*} K. Svenska Vet. Akad. Handlinger. Bd. 2, No. 6. 1874.

There are now before us twelve estimates for the molecular weight of thoria. Two of these represent single experiments, and have no probable error attached to them; namely, the one due to Hermann, and the one deduced from Berzelius' K₂SO₄: ThO₂ ratio. A third value, from Delafontaine's sulphuric acid estimations, has so high a probable error that it could be rejected without influencing the general mean. These three values might all be excluded without serious objection; but it is perhaps better to arbitrarily assign them equal weight, average them together, and give their mean the same probable error as that which attaches to Berzelius' BaSO₄: ThO₂ series. This mean is indicated as "A" in the following combination:

```
Berzelius ____ " = 269.940, ± .997
Chydenius—Sulphate ..... " = 268.584, ± .803
       Acetate _____ "
                           = 268.735, \pm .805
       Formate ..... "
                           = 272.586, \pm .877
                           = 262.804, \pm .493
       Oxalate _____ "
Delafontaine—Higher hydrate ..... "
                           = 263.637, \pm .256
                           = 266.025, \pm .363
        Lower " ..... "
Cleve—Sulphate _____ "
                           = 265.380, \pm .123
    Oxalate ...... "
                           = 265.357, \pm .104
    General mean = 265.341, \pm .072
```

Hence Th = 233.414, \pm .0725; or, if O = 16, Th = 233.951.

These values vary from those derived from Cleve's experiments alone only in the second decimal.

GALLIUM.

Gallium has been so recently discovered, and obtained in such small quantities, that its atomic weight has not as yet been determined with much precision. The following data were fixed by the discoverer, Lecoq de Boisbaudran:*

^{*} Journ. Chem. Soc., 1878, p. 646.

3.1044 grammes gallium ammonium alum, upon ignition, left .5885 grm. Ga_2O_3 .

Hence Ga = 68.071. If O = 16, Ga = 68.233.

.4481 grammes gallium, converted into nitrate and ignited, gave .6024 grm. Ga₂O₃.

• Hence Ga = 69.538. If O = 16, Ga = 69.693.

These values, assigned equal weight, give these means:

If O = 15.9633, Ga = 68.854. If O = 16, Ga = 68.963.

In brief, for all practical purposes, 69 may be assumed as the atomic weight of gallium.

INDIUM.

Reich and Richter, the discoverers of indium, were also the first to determine its atomic weight.* They dissolved weighed quantities of the metal in nitric acid, precipitated the solution with ammonia, ignited the precipitate, and ascertained its weight. Two experiments were made, as follows:

.5135 grm. indium gave .6243 grm. In₂O₃.
.699 " .8515 "

Hence, in mean, In = 110.61, if O = 16; a value known now to be too low.

An unweighed quantity of fresh, moist indium sulphide was also dissolved in nitric acid, yielding, on precipitation,

.2105 grm. In₂O₃ and .542 grm. BaSO₄.

Hence, with $BaSO_4 = 233$, In = 111.544; also too low.

Soon after the publication of Reich and Richter's paper the subject was taken up by Winkler.† He dissolved indium in nitric acid, evaporated to dryness, ignited the residue, and weighed the oxide thus obtained.

^{*} Journ. für Prakt. Chem., 92, 484.

[†] Journ. für Prakt. Chem., 94, 8.

```
.5574 grm. In gave .6817 grm. In<sub>2</sub>O<sub>2</sub>.
.6661 " .8144 "
.5011 " .6126 "
```

Hence, in mean, if O = 16, In = 107.76; a result even lower than the values already cited.

In a later paper by Winkler* better results were obtained. Two methods were employed. First, metallic indium was placed in a solution of pure, neutral, sodio-auric chloride, and the amount of gold precipitated was weighed. I give the weighings and, in a third column, the amount of indium proportional to 100 parts of gold:

In.	Au.	Ratio.
.4471 grm.	.8205 grm.	57.782
.8445 "	1.4596 "	57.858
		Mean, 57.820, ± .026

Hence, if $Au = 196.155, \pm .095$, $In = 113.417, \pm .074$.

Winkler also repeated his earlier process, converting indium into oxide by solution in nitric acid and ignition of the residue. An additional experiment, the third as given below, was made after the method of Reich and Richter. The third column gives the percentage of In in In, O₂:

```
      1.124 grm. In gave 1.3616 grm. In O3.
      Per cent., 82.550

      1.015 " 1.2291 " " 82.581

      .6376 " .7725 " " 82.537
```

These figures were confirmed by a single experiment of Bunsen's,† published simultaneously with the specific heat determinations which showed that the oxide of indium was In₂O₃, and not InO as had been previously supposed:

```
1.0592 grm. In gave 1.2825 grm. In<sub>4</sub>O<sub>3</sub>. Per cent. In, 82.589
```

For convenience we may add this figure in with Winkler's series, which gives us a mean percentage of In in In_2O_3 of 82.564, \pm .0082. Hence, if O=15.9633, \pm .0035, In=113.385, \pm .060.

^{*} Journ. für Prakt. Chem., 102, 282.

[†] Poggend. Annal., 141, 28.

Combining results, we have the following general mean:

CERIUM.

Although cerium was discovered almost at the beginning of the present century, its atomic weight was not properly determined until after the discovery of lanthanum and didymium by Mosander. In 1842 the investigation was undertaken by Beringer,* who employed several methods. His cerium salts, however, were all rose-colored, and therefore were not wholly free from didymium; and his results are further affected by a negligence on his part to fully describe his analytical processes.

First, a neutral solution of cerium chloride was prepared by dissolving the carbonate in hydrochloric acid. This gave weights of ceroso-ceric oxide and silver chloride as follows. The third column shows the amount of CeO, proportional to 100 parts of AgCl:

CeOz.	AgCl.	Ratio.
.5755 grm.	1.419 grm.	40.557
.6715 "	1.6595 "	40.464
1.1300 "	2.786 "	40.560
.5366 "	1.3316 "	40.297
	ı	——— Mean, 40.469, ± .0415

The analysis of the dry cerium sulphate gave results as follows. In a fourth column I show the amount of CeO, proportional to 100 parts of BaSO₄:

^{*} Ann. Chem. Pharm., 42, 134.

Sulphate.	CeO ₂ .	$BaSO_4$.	Ratio.
1.379 grm.	.8495 grm.	1.711 grm.	49.649
1.276 "	.7875 "	1.580 "	49.836
1.246 "	.7690 "	1.543 "	49.838
1.553 "	·959 5 "	1.921 "	49.948

Mean, 49.819, ± .042

Beringer also gives a single analysis of the formate and the results of one conversion of the sulphide into oxide. The figures are, however, not valuable enough to cite.

The foregoing data involve one variation from Beringer's paper. Where I put CeO, as found he puts Ce,O₃. The latter is plainly inadmissible, although the atomic weights calculated from it agree curiously well with some other determinations. For instance, in the chloride series, the assumption of Ce,O₃ as the formula of the oxide obtained, gives Ce = 137.749, while CeO₂ makes Ce = 141.636. The former agrees with the results of Wolf, Wing, and others quite fairly; the latter is near the value obtained by Bührig. Obviously, the presence of didymium in the salts analyzed should tend to raise rather than to lower the apparent atomic weight of cerium.

Shortly after Beringer, Hermann* published the results of one experiment. 23.532 grm. of anhydrous cerium sulphate gave 29.160 grm. of BaSO₄. Hence 100 parts of the sulphate correspond to 123.926 of BaSO₄.

In 1848 similar figures were published by Marignac,[†] who found the following amounts of BaSO₄ proportional to 100 of dry cerium sulphate:

```
122.68
122.00
122.51
Mean, 122.40, ± .138
```

If we give Hermann's single result the weight of one experiment in this series, and combine, we get a mean value of 123.019, ± .113.

^{*} Journ. für Prakt. Chem., 30, 185. 1843.

[†] Arch. des Sciences Phys. et Nat., (1,) 8, 273. 1848.

Still another method was employed by Marignac. A definite mixture was made of solutions of cerium sulphate and barium chloride. To this were added, volumetrically, solutions of each salt successively, until equilibrium was attained. The figures published give maxima and minima for the BaCl, proportional to each lot of Ce,(SO₄)₈. In another column, using the mean value for BaCl, in each case, I put the ratio between 100 parts of this salt and the equivalent quantity of sulphate. The latter compound was several times recrystallized:

	Ce2	$(SO_4)_8$.		BaCl ₂ .	A	Patio.
First crys	tallizat	ion 11.0	11 grm.	11.990 12.050	grm. 9	.606
44	44	13.1	94 "	14.365 — 14.425	" 91	.657
Second	66	13.9	61 "	15.225 — 15.285	" 9	.518
44	44	12.6	27 "	13.761 — 13.821	" 91	.559
66	64	11.9	15 "	12.970 — 13.030	" 91	.654
Third	46	14.8	388 "	16.223 — 16.283	" 9	.602
44	44	14.1	13 "	15.383 — 15.423	" 9	1.755
Fourth	46	13.1	II "	14.270 14.330	" 91	.685
"	44	13.9	70 "	15.223 — 15.283	" 9	. 588
					_	

Mean, 91.625, ± .016

Omitting the valueless experiments of Kjerulf,* we come next to the figures published by Bunsen and Jegel† in 1858. From the air dried sulphate of cerium the metal was precipitated as oxalate, which, ignited, gave CeO₂. In the filtrate from the oxalate the sulphuric acid was estimated as BaSO₄:

```
1.5726 grm. sulphate gave .7899 grm. CeO<sub>2</sub> and 1.6185 grm. BaSO<sub>4</sub>.
1.6967 " * .8504 " 1.7500 "
```

Hence, for 100 parts BaSO₄, the CeO₂ is as follows:

$$48.804$$

$$48.575$$
Mean, 48.689 , $\pm .077$

One experiment was also made upon the oxalate:

.3530 grm. oxalate gave .1913 CeO2 and .0506 H2O.

Hence, in the dry salt, we have 63.261 per cent. of CeO2.

^{*} Ann. Chem. Pharm., 87, 12. † Ann. Chem. Pharm., 105, 45.

In each sample of CeO, the excess of oxygen over true Ce,O, was estimated by an iodometric titration; but the data thus obtained need not be further considered.

In two papers by Rammelsberg* data are given for the atomic weight of cerium, as follows. In the earlier paper cerium sulphate is analyzed, the cerium being thrown down by caustic potash, and the acid precipitated from the filtrate as barium sulphate:

```
.413 grm. Ce<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub> gave .244 grm. CeO<sub>2</sub> and .513 grm BaSO<sub>4</sub>.
```

Hence 100 BaSO₄ = 47.563 CeO₂, a value which may be combined with others, thus; this figure being assigned a weight equal to one experiment in Bunsen's series:

Beringer	49.819, ± .042
Bunsen and Jegel	$48.689, \pm .077$
Rammelsberg	47.563, ± .108
General mean	40.360. ± .035

It should be noted here that this mean is somewhat arbitrary, since Bunsen and Rammelsberg's cerium salts were undoubtedly freer from didymium than the material studied by Beringer.

In his later paper Rammelsberg gives these figures concerning cerium oxalate. 100 parts gave 10.43 of carbon and 21.73 of water. Hence the dry salt should yield 48.862 per cent. of CO,, whence Ce = 137.83.

In all of the foregoing experiments the ceroso-ceric oxide was somewhat colored, the tint ranging from one shade to another of light brown according to the amount of didymium present. Still, at the best, a faint color remained, which was supposed to be characteristic of the oxide itself. In 1868, however, some experiments of Dr. C. Wolf† were post-humously made public, which went to show that pure ceroso-ceric oxide is white, and that all samples previously studied were contaminated with some other earth, not necessarily didymium but possibly a new substance, the removal of

^{*} Poggend. Annal., 55, 65; 108, 44.

[†] Amer. Journ. Science and Arts, (2,) 46, 53.

which tended to lower the apparent atomic weight of cerium very perceptibly.

Cerium sulphate was recrystallized at least ten times. Even after twenty recrystallizations it still showed spectroscopic traces of didymium. The water contained in each sample of the salt was cautiously estimated, and the cerium was thrown down by boiling concentrated solutions of oxalic acid. The resulting oxalate was ignited with great care. I deduce from the weighings the percentage of CeO₂ given by the anhydrous sulphate:

Sulphate.	Water.	CeO2. Per	r cent. CeO ₂ .
1.4542 grm. 1.4104 " 1.35027 "	.19419 grm. .1898 " .1820 "	.76305 grm. ·7377 " .70665 "	60.559 60.437 60.487
	•	Mean,	60.494, ± .02

After the foregoing experiments the sulphate was further purified by solution in nitric acid and pouring into a large quantity of boiling water. The precipitate was converted into sulphate and analyzed as before:

Sulphate.	Water.	C602.	Per	r cent. Ce C) ₁ .
1.4327 grm. 1.5056 " 1.44045 "	.2733 grm. .2775 " .2710 "	.69925 g .7405 .7052	rm. "	60.311 60.296 60.300	
	,	-,-,-	Mean.	60.302. =	± .00₄

From another purification the following weights were obtained:

```
1.4684 grm. .1880 grm. .7717 grm. 60.270 per cent.
```

A last purification gave a still lower percentage:

```
1.3756 grm. .1832 grm. .7186 grm. 60.265 per cent.
```

The last oxide was perfectly white, and was spectroscopically free from didymium. In each case the CeO, was titrated iodometrically for its excess of oxygen. It will be noticed that in the successive series of determinations the percentage of CeO, steadily and strikingly diminishes, to an extent for which no ordinary impurity of didymium can

account. The death of Dr. Wolf interrupted the investigation, the results of which were edited and published by Professor F. A. Genth.

The experiments of Wolf seem to have hitherto escaped general notice, except from Wing, who has partially verified them.* This chemist, incidentally to other researches, purified some eerium sulphate after the method of Wolf, and made two similar analyses of it, as follows:

Sulphale.	Water.	CeO2.	Per cent. CeO ₂ .
1.2885 grm. 1.4090 "	.1707 grm. .1857 "	.6732 grm. .7372 "	60.225 60.263
		Me	an, 60.244, ± .012

The ceroso-ceric oxide in this case was perfectly white. The cerium oxalate which yielded it was precipitated boiling by a boiling concentrated solution of oxalic acid. The precipitate stood twenty-four hours before filtering.

We may now combine the results of Wolf and of Wing, as follows. The two concordant experiments of Wolf's series three and four may be united, giving a mean of 60.267, $\pm .001$:

Wolf,	1st series	60.494, ± .024
**	2d "3d and 4th series	60.302, ± .004
44	3d and 4th series	60.267, ± .001
Wing.		$60.244, \pm .012$
	General mean	60.271, ± .001

This mean, the percentage of CeO, in the anhydrous sulphate, gives Ce = 137.724; or, if O = 16, Ce = 138.039. This varies widely from the ordinarily accepted value as determined by Buehrig.

In 1875 Buehrig's † paper upon the atomic weight of cerium was issued. He first studied the sulphate, which, after eight crystallizations, still retained traces of free sulphuric acid. He found furthermore that the salt obstinately retained traces of water, which could not be wholly expelled by heat without partial decomposition of the material.

^{*} Amer. Journ. Sci. and Arts, (2,) 49, 358. 1870.

[†] Journ. für Prakt. Chem., 120, 222.

These sources of error probably affect all the previously cited series of experiments; although, in the case of Wolf's work, it is doubtful whether they could have influenced the atomic weight of cerium by more than one or two tenths of Buehrig also found, as Marignac had earlier shown, that upon precipitation of cerium sulphate with barium chloride the barium sulphate invariably carried down traces of cerium. Furthermore, the ceroso-ceric oxide from the filtrate always contained barium. For these reasons the sulphate was abandoned, and the atomic weight determinations of Buehrig were made with air-dried oxalate. This salt was placed in a series of platinum boats in a combustion tube behind copper oxide. It was then burned in a stream of pure, dry oxygen, and the carbonic acid and water were collected after the usual method. Ten experiments were made; in all of them the above named products were estimated, and in five analyses the resulting cerosoceric oxide was also weighed. By deducting the water found from the weight of the air-dried oxalate, the weight of the anhydrous oxalate is obtained, and the percentages of its constituents are easily determined. In weighing, the articles weighed were always counterpoised with similar materials. The following weights were found:

Oxala	te.	Water.		CO2.		CeO2	
9.8541	grm.	2.1987 §	grm.	3.6942	grm.		
9.5368	**	2.1269	"	3.5752	44		
9.2956	"	2.0735	"	3.4845	44		
10.0495	46	2.2364	"	3.7704	66		
10.8249	"	2.4145	44	4.0586	"		
9.3679	66	2.0907	•6	3.5118	44	4.6150	grm.
9.7646	44	2.1769	"	3.6616	44	4.8133	"
9.9026	66	2.2073	"	3.7139	**	4.8824	"
9.9376	66	2.2170	48	3.7251	"	4.8971	"
9.5324	46	2.1267	"	3.5735	44	4.6974	"

These figures give us the following percentages for CO, and CeO, in the anhydrous oxalate:

CO ₂ .	CeO2.
48.256	
48.249	
48.248	

Mean, 48,2546, + .001	63.4316. + .0032
48.253	63.430
48.249	63.429
48.262	63.446
48.257	63.436
48.258	63.417
48.257	
48.257	
CeO ₃ .	CO _s .

From percentage CO₂ Ce = 141.228, ± .025 " CeO₂ " = 141.141, ± .020

Obviously the single oxalate experiments of Jegel and of Rammelsberg would exert no appreciable influence upon these mean results. They may therefore be ignored.

In combining all of these data in one general mean, we may begin as usual by tabulating our ratios:

(1.) BaSO₄: Ce₂(SO₄)₃:: 100: 123.019, ± .113 (2.) BaSO₄: CeO₂:: 100: 49.360, ± .035 (3.) BaCl₂: Ce₃(SO₄)₃:: 100: 91.625, ± .016 (4.) AgCl: CeO₃:: 100: 40.469, ± .0415 (5.) Percentage CeO₂ from anhydrous sulphate, 60.271, ± .001 (6.) " " oxalate, 63.4316, ± .0032 (7.) " CO₃ " 48.2546, ± .001

These ratios give us four values for the molecular weight of CeO₂ and two values for Ce₂(SO₄)₃:

Hence we have three independent values for the atomic weight of cerium, as follows:

From molecular weight of CeO₂.......Ce = 139.563, ± .024

" Ce₂(SO₄)₃....." = 141.281, ± .083

From ratio (7,) CO₂ in oxalate......" = 141.228, ± .025

General mean....." = 140.424, ± .017

Or, if O = 16, Ce = 140.747.

Buehrig's results alone, both sets combined, give Ce = 141.198, $\pm .020$; or, if O = 16, Ce = 141.523.

Wolf and Wing's figures alone make Ce = 137.724; or, if O = 16, Ce = 138.039.

The latter result is subject to the errors pointed out by Buehrig as involved in the use of cerium sulphate; but the ceroso-ceric oxide obtained in the analyses was pure white. Buehrig's ceroso-ceric oxide, on the other hand, was yellow. In neither case was didymium present. All things considered, therefore, it is probable that the lower result is too low and the higher result too high. How near the general mean of all may be to the truth we have no evidence to show. It is clear that new determinations are needed, made with material yielding white ceroso-ceric oxide, and with avoidance of the sources of error which Buehrig pointed out.

LANTHANUM.

Leaving out of account the work of Mosander, and the valueless experiments of Choubine, we may consider the estimates of the atomic weight of lanthanum which are due to Hermann, Rammelsberg, Marignac, Czudnowicz, Holzmann, Zschiesche, Erk, and Cleve.

From Rammelsberg* we have but one analysis. .700 grm. of lanthanum sulphate gave .883 grm. of barium sulphate. Hence 100 parts of BaSO₄ are equivalent to 79.276 of La₂(SO₄)₈.

^{*} Poggend. Annal., 55, 65.

Marignac,* working also with the sulphate of lanthanum, employed two methods. First, the salt in solution was mixed with a slight excess of barium chloride. The resulting barium sulphate was filtered off and weighed; but, as it contained some occluded lanthanum compounds, its weight was too high. In the filtrate the excess of barium was estimated, also as sulphate. This last weight of sulphate, deducted from the total sulphate which the whole amount of barium chloride could form, gave the sulphate actually proportional to the lanthanum compound. The following weights are given:

$La_2(SO_4)_3$.	BaCl ₂ .	ist BaSO4.	2d BaSO.
4.346 grm.	4.758 grm.	5.364 grm.	.115 grm.
4.733 "	5.178 "	5.848 "	.147 "

Hence we have the following quantities of La₂(SO₄)₃ proportional to 100 parts of BaSO₄. Column A is deduced from the first BaSO₄ and column B from the second, after the manner above described:

	A.	В.
	81.022	· 83.281
	80.934	83.662
Mean,	80.978, ± .030	Mean, 83.471, ± .128
From	A	La = 138.776
66	B	" = 147.474

A agrees best with other determinations, although, theoretically, it is not so good as B.

Marignac's second method, described in the same paper with the foregoing experiments, consisted in mixing solutions of La₂(SO₄)₃ with solutions of BaCl₂, titrating one with the other until equilibrium was established. The method has already been described under cerium. The weighings give maxima and minima for BaCl₂. In another column I give La₂(SO₄)₃ proportional to 100 parts of BaCl₂, mean weights being taken for the latter:

^{*} Archives des Sci Phys. et Naturelles, (1,) 11, 29. 1849.

$La_2(SO_4)_3$.	BaCl ₂ .	Ratio.
11.644 grm.	12.765 — 12.825 grm.	91.004
12.035 "	13.195 — 13.265 "	90.968
10.690 "	11.669 — 11.749 "	91.297
12.750 "	13.920 — 14.000 "	91.332
10.757 "	11.734 — 11.814 "	91.362
12.672 "	13.813 — 13.893 "	91.475
9.246 "	10.080 — 10.160 "	91.364
10.292 "	11.204 — 11.264 "	91.615
10.192 "	11.111 — 11.171 "	91.482

Mean, 91.322, \pm .048

Hence La = 140.484.

Although not next in chronological order, some still more recent work of Marignac's * may properly be considered here. The salt studied was the sulphate of lanthanum, purified by repeated crystallizations. In two experiments the salt was calcined, and the residual oxide weighed; in two others the lanthanum was precipitated as oxalate, and converted into oxide by ignition. The following percentages are given for La₂O₃:

The atomic weight determinations of Holzmann † were made by analyses of the sulphate and iodate of lanthanum, and the double nitrate of magnesium and lanthanum. In the sulphate experiments the lanthanum was first thrown down as oxalate, which, on ignition, yielded oxide. The sulphuric acid was precipitated as BaSO₄ in the filtrate.

$La_2(SO_4)_8$.	La_2O_3 .	BaSO₄.
.9663 grm.	.5157 grm.	1.1093 grm.
.6226 "	·3323 "	.7123 "
.8669 "	.4626 "	.9869 "

^{*} Ann. de Chim. et de Phys., (4,) 30, 68. 1873. † Journ. für Prakt. Chem., 75, 321. 1858.

These results are best used by taking the ratio between the BaSO₄, put at 100, and the La₂O₃. The figures are then as follows:

$$46.489$$

$$46.652$$

$$46.873$$
Mean, 46.671 , $\pm .075$

In the analyses of the iodate the lanthanum was thrown down as oxalate, as before. The iodic acid was also estimated volumetrically, but the figures are hardly available for present discussion. The following percentages of La₂O₃ were found:

The formula of this salt is La₂(IO₈)₆.3H₂O.

The double nitrate, La₂(NO₃)₅.3Mg(NO₃)₂.24H₂O, gave the following analytical data:

Salt.	H_2O .	MgO.	La_2O_3 .
.5327 grm.	.1569 grm.	.0417 grm.	.1131 grm.
.5931 "	.1734 "	.0467 "	.1262 "
.5662 "	.1647 "	.0442 "	.1197 "
·3757 "		.0297 "	.0813 ''
.3263 "		.0256 "	.0693 "

These weighings give the subjoined percentages of La,O,:

These data of Holzmann give values for the molecular weight of La₂O₃ as follows:

Czudnowicz* based his determination of the atomic weight of lanthanum upon one analysis of the air-dried sulphate. The salt contained 22.741 per cent. of water.

.598 grm. gave .272 grm. La₂O₃ and .586 grm. BaSO₄.

The La₂O₃ was found by precipitation as oxalate and ignition. The BaSO₄ was thrown down from the filtrate. Reduced to the standards already adopted these data give for the percentage of La₂O₃ in the anhydrous sulphate the figure 58.668. 79.117 parts of the salt are proportional to 100 parts of BaSO₄.

Hermann† studied both the sulphate and the carbonate of lanthanum. From the anhydrous sulphate, by precipitation as oxalate and ignition, the following percentages of La,O₃ were obtained:

The carbonate, dried at 100°, gave the following percentages:

Reckoning from the ratio between CO, and La₂O₃ the molecular weight of the latter becomes 325.896.

Zschiesche's texperiments consist of six analyses of lanthanum sulphate, which salt was dehydrated at 230°, and afterwards calcined. I subjoin his percentages, and in a fourth column deduce from them the percentage of La₂O₃ in the anhydrous salt:

H_2O .	SO ₃ .	La_2O_3 .	La ₂ O ₃ in anhydrous salt.
22.629	33.470	43.909	56.745
22.562	33.306	44.132	56.9 64
22.730	33.200	44.070	57.034

^{*} Journ. für Prakt. Chem., 80, 33. 1860

[†] Journ. für Prakt. Chem., 82, 396. 1861.

[†] Journ. für Prakt. Chem., 104, 174.

H_2O .	SO ₃ .	La_2O_3 .	La203 in anhydrous salt.
22.570	33-333	44.090	56.947
22.610	33.160	44.240	57.150
22.630	33.051	44.310	5 7.27 7
			Mean, 57.021, ± .051

Erk * found that .474 grm. of La₂(SO₄)₃, by precipitation as oxalate and ignition, gave .2705 grm. of La₂O₃, or 57.068 per cent. .7045 grm. of the sulphate also gave .8815 grm. of BaSO₄. Hence 100 parts of BaSO₄ are equivalent to 79.921 of La₂(SO₄)₃.

Last of all, and probably best of all, we come to the determinations of Cleve.† Strongly calcined La₂O₃, spectroscopically pure, was dissolved in nitric acid, and then, by evaporation with sulphuric acid, converted into sulphate:

```
      1.9215 grm. La<sub>2</sub>O<sub>3</sub> gave
      3.3365 grm. sulphate.
      57.590 per cent.

      2.0570
      "
      3.5705
      "
      57.611
      "

      1.6980
      "
      2.9445
      "
      57.667
      "

      2.0840
      "
      3.6170
      "
      57.617
      "

      1.9565
      "
      3.3960
      "
      57.612
      "
```

Mean, 57.619, ± .0085

From the last column, which indicates the percentage of La₂O₃ in La(SO₄)₃, we get, if SO₃ = 80, La = 139.15.

We may now combine the similar means into general means, and deduce a value for the atomic weight of lanthanum. For the percentage of oxide in sulphate we have six estimates, as follows. The single experiments of Czudnowicz and of Erk are assigned the probable error and weight of a single experiment in Hermann's series:

Czudnowicz	58.668,	土 .027
Erk	57.068,	土 .027
Hermann	57.654,	土 .016
Zschiesche	57.021,	± .051
Marignac	57.5475,	土 .0115
Cleve	57.619,	± .0085
General mean	57.620,	± .0059

^{*} Jenaisches' Zeitschrift, 6, 306. 1871.

[†] K. Svenska Vet. Akad. Handlingar, Bd. 2, No. 7. 1874.

For the quantity of La₂(SO₄)₃ proportional to 100 parts of BaSO₄, we have five experiments, which may be given equal weight and averaged together:

Marignac	81.022
"	80.934
Rammelsberg	79.276
Czudnowicz	79.117
Erk	79.921
Mean,	80.054, ± .270

In all, there are seven ratios from which to calculate:

- (1.) Percentage of La₂O₃ in La₂(SO₄)₃, 57.620, \pm .0059
- (2.) BaCl₂: La₂(SO₄)₃:: 100: 91.322, \pm .048—Marignac.
- (3.) BaSO₄: La₂(SO₄)₃:: 100: 80.054, \pm .270
- (4.) BaSO₄: La₂O₃:: 100: 46.671, ± .075—Holzmann.
- (5.) Percentage of La₂O₃ in iodate, 23.447, ± .0216—Holzmann.
- (6.) " magnesian nitrate, 21.3056, ± .058—Holzmann.
- (7.) " carbonate, 68.47—Hermann.

These ratios give five values for the molecular weight of lanthanum oxide, and two for that of the sulphate:

From (2).....La₂(SO₄)₈ = 568.488,
$$\pm$$
 .320
" (3)..... " = 558.624, \pm 1.888
General mean, " = 568.212, \pm .316

Hence La = 140.346, $\pm .160$.

Here the value derived from ratio (7) is given the weight of a single experiment in ratio (1.) Hence La = 138.460, ± .031.

Combining the two values for La, we get this final result:

From La₂O₃La = 138.460,
$$\pm$$
 .031

" La₂(SO₄)₃ " = 140.346, \pm .160

General mean " = 138.526, \pm .030

Or, if
$$O = 16$$
, La = 138.844.

Since this value is a little under and Cleve's a little over 139, the latter figure may fairly be used in all calculations involving a knowledge of the atomic weight of lanthanum.

DIDYMIUM.

The atomic weight of didymium has been determined by Marignac, Hermann, Zschiesche, Erk, and Cleve. Mosander's early experiments we may leave out of account.

Marignac * mixed a solution of the sulphate with a slight excess of barium chloride, filtered, weighed the precipitate, and estimated the excess of barium in the filtrate by the ordinary method. The first precipitate always contained didymium, and therefore weighed too much. By deducting the weight of the second precipitate, representing the excess of the barium chloride, from the weight of barium sulphate theoretically formable, the weight of the latter proportional to the quantity of didymium salt taken was found:

$Di_2(SO_4)_3$.	$BaCl_2$.	rst BaSO4.	≥d BaSO₄.
3.633 grm.	3.902 grm.	4.412 grm.	.084 grm.
3.862 "	4.227 "	4.679 "	.075 "
3.330 "	3.552 "	4.027 "	.088 "
1.386 "	1.477 "	1.681 "	.014 "

These figures give us a ratio between the sulphates of didymium and barium which we may express as follows. Column A gives the Di₂(SO₄)₃ proportional to 100 parts of BaSO₄, as calculated from the first precipitate of the latter. Column B gives a similar ratio calculated with the second BaSO₄ precipitate, this being deduced from the total BaSO₄ which the chloride used could form:

Mean, 82.455, ± .052	
	Mean, 84.320, ± .414
82.247—Erk.	
82.451	84.425
82.692	85.545
82.539	82.626
82.344	84.685
A.	ь.

^{*} Arch. des Sci. Phys. et Naturelles, (1,) 11, 29. 1849.

To A I have added a single result of Erk's, to be described further along. It will be seen that although A is theoretically defective, its figures are much more concordant than those in B. In fact, the latter would almost vanish for the final general mean for the atomic weight of didymium:

In a later paper* Marignac adopts two other methods for establishing the atomic weight of didymium. The carefully dehydrated sulphate was taken, the didymium was precipitated as oxalate, and the latter, ignited, yielded oxide. The following percentages of oxide were found:

The chloride of didymium was also studied. As the anhydrous salt could not be obtained in an absolutely definite state, Marignae prepared neutral solutions of it and determined the ratio between didymium oxide and silver chloride. The latter compound was first precipitated in the usual way, and filtered off; the excess of silver in the filtrate was removed by hydrochloric acid, and after that the didymium was thrown down as oxalate and weighed as oxide. The subjoined weights of AgCl and Di₂O₃ were found. In a third column I give the ratio between the two compounds, putting AgCl at 100:

AgCl.	Di_2O_3 .	Ratio.
10.058 grm.	3.946 grm.	39.232
5.029 "	1.960 "	38.974
5.844 "	2.276 "	38.946

Mean, 39.051, ± .061

Hence Di = 143.637, $\pm .263$.

^{*} Ann. d. Chim. et d. Phys., (3,) 38, 148. 1853.

Hermann's* determination of the atomic weight of didymium rests on a single experiment with the sulphate. By precipitation as oxalate and subsequent ignition, he found that this salt yielded 58.14 per cent. of Di₂O₃.

Zschiesche† also analyzed didymium sulphate, which he dehydrated at 230°, and afterwards converted into oxide by calcination. I give his percentages, and also, in a fourth column, the percentage of oxide from the *anhydrous* sulphate as deduced from his figures:

H_2O .	SO ₈ .	Di_2O_3 .	Di ₂ O ₂ in anhyd. salt.
23.19	32.97	43.83	57.070
23.03	32.39	44.58	57.919
23.00	32.56	44.95	58.006
23.547	31.938	44.515	58. 225
22.550	32.870	44.570	57-554

The salt used in the first experiment probably contained lanthanum. Rejecting this, the mean of the figures remaining in the fourth column is 57.926, $\pm .094$. Hence Di = 141.007.

Erk,‡ to whom reference has already been made, estimated didymium in the sulphate by precipitation as oxalate and calcination to oxide:

$Di_2(SO_4)_8$ Di_2O_3 .		Per cent. Di ₂ O ₃ .
.556 grm.	.323 grm.	58.094
.674 "	.3915 "	58.087

Hermann's single result for this percentage, 58.14, agrees more nearly with Erk's series than with any other. It may therefore be averaged in with Erk's two experiments, giving a mean of 58.107, \pm .0112. Erk also obtained from .7065 grm. of sulphate .859 grm. BaSO₄. This experiment has already been averaged with Marignac's earlier results.

The latest determinations of the atomic weight of didymium were published by Cleve | in 1874. Strongly calcined

^{*} Journ. für Prakt. Chem., 82, 367. 1861.

⁺ Journ. für Prakt. Chem., 107, 74.

[†] Jenaisches' Zeitschrift, 6, 306. 1871.

^{||} K. Svenska Vet. Akad. Handlingar, Bd. 2, No. 8. These figures were kindly transcribed for me by Professor Delafontaine of Chicago, as I had not access to a copy of the original memoir.

didymium oxide was dissolved in nitric acid, the solution was evaporated with sulphuric acid, and the weight of the resulting sulphate was ascertained. I subjoin the weighings and the percentage of Di₂O₃ in Di₂(SO₄)₃:

Di_2O_3 .	$Di_2(SO_4)_2$.	Per cent. Di ₂ O ₃ .
2.257 grm:	3.844 grm.	58.715
1.086 "	1.8485 "	58.750
1.1525 "	1.9615 "	58.756
1.3635 " .	2.319 "	58.797
1.9655 "	3-3435 "	58.786
1.528 "	2.599 "	58.792
		Mean, 58.766, ± .0087

Hence Di = 146.804. If $SO_3 = 80$, Di = 147.021.

This determination is undoubtedly the best of all, and might properly be accepted to the exclusion of the others. Still, it is worth while to combine all the figures into one general mean. For the percentage of Di₂O₃ in Di₂(SO₄)₃ we have the following data:

General mean	58.451, ± .0059
Cleve	58.766, ± .0087
Zschiesche	$57.926, \pm .094$
Erk and Hermann	$58.107, \pm .0112$
Marignac	

For the atomic weight of didymium we have now three independent values:

```
From per cent. Di<sub>2</sub>O<sub>3</sub> in Di<sub>2</sub>(SO<sub>4</sub>)<sub>2</sub>......Di = 144.604, ± .031

" Marignac's chloride analyses......." = 143.637, ± .263

" Marignac's and Erk's BaSO<sub>4</sub> ratio..." = 143.929, ± .189

General mean..........." = 144.573, ± .0306
```

If O = 16, Di = 144.906.

THE YTTRIUM GROUP.

The atomic weights of the metals in this group can only be said to have been determined approximately. Not only do great difficulties attend the purification of the material used for study and the separation of the earths from each other, but there have been and still are grave doubts as to the actual nature of some of the latter. The figures for scandium, yttrium, and ytterbium seem to be tolerably good; those for decipium, philippium, thulium, erbium, and terbium are little more than estimates; for samarium we have no data whatever. All the atomic weights in this group are based upon analyses or syntheses of sulphates; and from analogy to the cerium metals all of these elements are regarded as forming sesquioxides.

SCANDIUM.

Cleve,* who was the first to make accurate experiments on the atomic weight of this metal, obtained the following data. 1.451 grm. of sulphate, ignited, gave .5293 grm. of Sc₂O₃. .4479 grm. of Sc₂O₃, converted into sulphate, yielded 1.2255 grm. of the latter, which, upon ignition, gave .4479 grm. of Sc₂O₃. Hence, for the percentage of Sc₂O₃ in Sc₂(SO₄), we have:

Hence, if $SO_1 = 80$, Sc = 45.044.

Later and better results are those of Nilson,† who converted scandium oxide into the sulphate. I give in a third column the percentage of oxide in sulphate:

^{*} Compt. Rend., 89, 419. † Compt. Rend., 91, 118.

```
.3379 grm. Sc<sub>2</sub>O<sub>3</sub> gave .9343 grm. Sc<sub>2</sub>(SO<sub>4</sub>)<sub>3</sub>. 36.166 per cent. 3015 " .8330 " 36.194 " .2998 " .8257 " 36.187 " .3192 " .8823 " 36.178 "
```

Mean, 36.181, ± .004

Hence Sc = 43.980, $\pm .015$; or, if O = 16, then Sc = 44.081. If SO₃ = 80, then Sc = 44.032. These values are doubtless very nearly correct.

YTTRIUM.

For yttrium we need consider only the determinations of Popp, Delafontaine, Bahr and Bunsen, and Cleve.

Popp* evidently worked with material not wholly free from earths of higher molecular weight than yttria. The yttrium sulphate was dehydrated at 200°; the sulphuric acid was then estimated as barium sulphate; and after the excess of barium in the filtrate had been removed, the yttrium was thrown down as oxalate, and ignited to yield oxide. The following are the weights given by Popp:

Sulphate.	$BaSO_4$.	Yt_2O_3 .	H_2O .
1.1805 gm.	1.3145 grm.	.4742 grm.	.255 grm.
1.4295 "	1.593 "	·5745 "	.308 "
.8455 "	.9407 "	.3392 "	. 1825 "
1.045 "	1.1635 "	.4195 "	.2258 "

Eliminating water, these figures give us for the percentages of Yt₂O₃ in Yt₂(SO₄)₃ the values in column A. In column B I put the quantities of Yt₂O₃ proportional to 100 parts of BaSO₄:

A.	В
51.237	36.075
51.226	36.064
51.161	36.058
51.209	36.055
Mean, 51.208, ± .011	Mean, 36.063, ± .003

From B, Yt = 101.880. The values in A will be combined with similar data from other experimenters.

^{*} Ann. Chem. Pharm., 131, 179.

In 1865 Delafontaine* published some results obtained from yttrium sulphate, the yttrium being thrown down as oxalate and weighed as oxide. In the fourth column I give the percentages of Yt₂O₃ reckoned from the anhydrous sulphate:

Sulphate.	Yt_2O_3 .	H_2O .	Per cent. Yt2O3.
.9545 grm.	.371 grm.	.216 grm.	50.237
2.485 "	.9585 "	.565 "	49.922
2.153 "	.827 "	· 4935 "	49.834
		M	 ean, 49.998, ± .081

In another paper † Delafontaine gives the following percentages of Yt₂O₃ in dry sulphate. The mode of estimation was the same as before:

Bahr and Bunsen,‡ and likewise Cleve, adopted the method of converting dry yttrium oxide into anhydrous sulphate, and noting the gain in weight. Bahr and Bunsen give us the two following results. I add the usual percentage column:

Yt_2O_3 .	$Yt_2(SO_4)_3$.	Per cent. Yt2O3.
.7266 grm.	1.4737 grm.	49.304
.7856 "	1.5956 "	49.235
•		
		Mean, 49.2695, ± .0233

Cleve's || results are published in a joint memoir by Cleve and Hoeglund, and are as follows:

^{*} Ann. Chem. Pharm., 134, 108.

[†] Arch. des Sci. Phys. et Nat., (2,) 25, 119. 1866.

[†] Ann. Chem. Pharm., 137, 21. 1866.

[|] K. Svenska Vet. Akad. Handlingar, Bd. 1, No. 8.

Yt, O,.	$Yt_2(SO_4)_3$.	Per cent. Yt, O.
1.4060 grm.	2.8925 grm.	48.608
1.0930 "	2.2515 "	48.545
1.4540 "	2.9895 "	48.637
1.3285 "	2.7320 "	48.627
2.3500 "	4.8330 "	48.624
2.5780 "	5.3055 "	48.591

Mean, 48.605, $\pm .0096$

This series is unquestionably the best of all. From it, if $SO_3 = 80$, Yt = 89.485.

Combining all these data we have the subjoined general mean for the percentage of Yt_2O_3 :

From the general mean of all, Yt = 97.616. From the mean after excluding Popp's work, Yt = 89.816, \pm .067; or, if O = 16, Yt = 90.023.

YTTERBIUM.

For ytterbium we have one very good set of determinations by Nilson.* The oxide was converted into the sulphate after the usual manner:

$Yb_{2}O_{3}$.	$Yb_2(SO_4)_3$.	Per cent. Yb ₂ O ₃ .
1.0063 grm.	1.6186 grm.	62.171
1.0139 "	1.6314 "	62.149
.8509 "	1.3690 "	62.155
·7371 "	1.1861 "	62.145
1.0005 "	1.6099 "	62.147
.8090 "	1.3022 "	62.126
1.0059 "	1.6189 "	62.134
		Mean, 62.147, ± .0036

^{*} Compt. Rend., 91, 56. 1880.

Hence Yb = 172.761, \pm .038. If O = 16, then Yb = 173.158. If SO₃ = 80, Yb = 173.016. The true number cannot be far from 173.

ERBIUM.

Since the earth which was formerly regarded as the oxide of this metal is now known to be a mixture of two or three different oxides, the older determinations of its molecular weight have little more than historical interest. Nevertleless the work done by several investigators may properly be cited, since it sheds some light upon certain important problems.

First, Delafontaine's * early investigations may be considered. A sulphate, regarded as erbium sulphate, gave the following data. An oxalate was thrown down from it, which, upon ignition, gave oxide. The percentages in the fourth column refer to the anhydrous sulphate. In the last experiment water was not estimated, and I assume for its water the mean percentage of the four preceding experiments:

Sulphate.	Er_2O_3 .	H_2O .	Per cent. Er_2O_3 .
.827 grm.	.353 grm.	.177 grm.	54.308
1.0485 "	·4475 "	.226 "	54.407
.8 03 "	.3415 "	.171 "	54.035
1.232 "	.523 "	.264 "	54.028
1.1505 "	· 4 95 "		54.760

Mean, 54.308, ± .0915

Bahr and Bunsen† give a series of results, representing successive purifications of the earth which was studied. The final result, obtained by the conversion of oxide into sulphate, was as follows:

.7870 grm. oxide gave 1.2765 grm. sulphate. 61.653 per cent. oxide.

Hoeglund,‡ following the method of Bahr and Bunsen, secured these results:

^{*} Ann. Chem. Pharm., 134, 108. 1865.

[†] Ann. Chem. Pharm., 137, 21. 1866.

[‡] K. Svenska Vet. Akad. Handlingar, Bd. 1, No. 6.

Er_3O_3 .	$Er_2(SO_4)_3$.	Per cent. Er203.	
1.8760 grm.	3.0360 grm.	61.792	
1.7990 "	2.9100 "	61.821	
2.8410 "	4.5935 "	61.848	
1.2850 "	2.0775 "	61.853	
1.1300 "	1.827 "	61.850	
.8475 "	1.370 "	61.861	
		-	

Mean, 61.8375, ± .0063

Humpidge and Burney* give data as follows:

```
1.9596 grm. \text{Er}_2(\text{SO}_4)_3 gave 1.2147 grm. \text{Er}_2\text{O}_3. 61.987 per cent.
1.9011 " 1.1781 " 61.965 " \frac{61.965}{61.976}, \pm.0074
```

Combining all four series we get the subjoined general mean for the percentage of oxide in sulphate. Bahr and Bunsen's single experiment is given the probable error of one experiment in Hoeglund's series:

Delafontaine	54.308,	\pm	.0915
Bahr and Bunsen	61.653,	±	.0178
Hoeglund	61.8375,	±	.0063
Humpidge and Burney			
¶ General mean			
Rejecting the first	61.880,	\pm	.0046

From the mean of all, Er = 170.379, $\pm .082$; or, if O = 16, Er = 170.770. From Bahr and Bunsen's determination, Er = 168.683; and from Humpidge and Burney's highest, Er = 171.428.

The foregoing data were all published before the composite nature of the supposed erbia was fully recognized. It will be seen, however, that three sets of results were fairly comparable, while Delafontaine evidently studied an earth widely different from that investigated by the others. Since the discovery of ytterbium, some light has been thrown on the matter. The old erbia is a mixture of at least three earths, to one of which, a rose-colored body, the name erbia is now restricted. For the atomic weight of the true erbium

^{*} Journ. Chem. Society, Feb., 1879, p. 116.

Cleve* gives three values, but without data concerning weighings or methods. Doubtless the oxide was converted into sulphate, and the calculations were made with SO₃ = 80:

166.00 166.21 166.25 ———— Mean, 166.153

With SO_s = 79.874, this becomes 165.891, and if only 0 = 16, 166.273. These figures are undoubtedly the nearest yet reached to the true value. According to Thalén,† who reasons from spectroscopic evidence, the erbium of Hoeglund was largely ytterbium.

TERBIUM, SAMARIUM, PHILIPPIUM, DECIPIUM, THULIUM, HOLMIUM, AND SORET'S EARTH X.

Concerning these substances, real or alleged, the data are exceedingly vague. For phillippium Delafontaine gives an atomic weight approximating to 123 or 125, and in the same memoir decipium is put at 171. It seems probable that philippium may be identical with Cleve's holmium and the metal of Soret's earth X, while decipium comes near Cleve's thulium, for which the discoverer gives a value of about 170.7.|| If decipium and thulium are identical, or if either proves to be erbium or ytterbium contaminated with the other, then we shall have a triad of metals with atomic weights ranging from Er = 166 to Yb = 173, strikingly parallel with lanthanum, cerium, and didymium. If we take the natural arrangement of the elements as tabulated after Mendelejeff's plan, somewhat modified in Roscoe and Schorlemmer's "Treatise on Chemistry, we find that such a triad should exist, and, furthermore, that another similar

^{*} Compt. Rend., 91,-382.

[†] Poggend. Beiblätter, 5, 122. 1881.

[‡] Arch. des Sci. Phys. et Nat., Mars, 1880.

^{||} Compt. Rend., 91, 329. 1880.

[&]amp; Vol. 2, Part 2, p. 507.

group ought to lie between indium and tin. The latter triad should have atomic weights ranging from 114 to 117; and here possibly, or else forming a triad with yttrium, the other metals of this group may lie.

COLUMBIUM.*

The atomic weight of this metal has been determined by Rose, Hermann, Blomstrand, and Marignac. Rose† analyzed a compound which he supposed to be chloride, but which, according to Rammelsberg,‡ must have been nearly pure oxychloride. If it was chloride, then the widely varying results give approximately Cb = 122; if it was oxychloride, the value becomes nearly 94. If it was chloride, it was doubtless contaminated with tantalum compounds.

Hermann's || results seem to have no present value, and as for Blomstrand's, I am not able to get at a copy of his original memoir. The results of the latter chemist are thus summed up in Becker's "Digest." Three chlorine estimations in the pentachloride give, in mean, Cb = 96.67. Eleven weighings of columbic acid from the same compound make Cb = 96.16. Other experiments on sodium columbate lead Blomstrand to regard 95 as the most probable value.

Marignac¶ made about twenty analyses of the potassium fluoxycolumbate, CbOF₃.2KF.H₂O. 100 parts of this salt give the following percentages:

^{*} This name has priority over the more generally accepted "niobium," and therefore deserves preference.

[†] Poggend. Annal., 104, 439. 1858.

[†] Poggend. Annal., 136, 353. 1869.

^{||} Journ. für Prakt. Chem., 68, 73. 1856.

Acta Univ. Lund, 1864.

[¶] Archives des Sci. Phys. et Nat., (2,) 23, 258. 1865.

From the mean percentage of Cb_2O_5 , Cb = 93.217. If O = 16, this becomes 93.431.

From the mean between the extremes given for K_2SO_4 , Cb = 93.812. If O = 16, this becomes 94.027.

As Deville and Troost's* results for the vapor density of the chloride and oxychloride agree fairly well with Cb = 94, we may adopt this value as approximately correct.

TANTALUM.

The results obtained for the atomic weight of this metal by Berzelius,† Rose,‡ and Hermann || may be fairly left out of account as valueless. These chemists could not have worked with pure preparations, and their data are sufficiently summed up in Becker's "Digest."

Marignac § made four analyses of a pure potassium fluotantalate, and four more experiments upon the ammonium salt. The potassium compound, K₂TaF₇ was treated with sulphuric acid, and the mixture was then evaporated to dryness. The potassium sulphate was then dissolved out by water, while the residue was ignited and weighed as Ta₂O₄. 100 parts of the salt gave the following quantities of Ta₂O₅ and K₂SO₄:

Ta_2O_5 .	K ₂ SO ₄ .
56.50	44.37
56.75	44.35
56.55	44.22
56.56	44.24
Mean, 56.59, ± .037	Mean, 44.295, ± .026

^{*} Comptes Rend., 56, 891. 1863.

[†] Poggend. Annal., 4, 14. 1825. Lehrbuch, 3, 1209.

[†] Poggend. Annal., 99, 80. 1856.

^{||} Journ. für Prakt. Chem., 70, 193. 1857.

² Archives des Sci. Phys. et Nat., 26, 89, serie 2. 1866.

From these figures, 100 parts of K₂SO₄ correspond to the subjoined quantities of Ta₂O₅:

127.338 127.960 128.178 127.848 Mean, 127.831, ± .120

The ammonium salt, (NH₄), TaF₇, ignited with sulphuric acid, gave these percentages of Ta₂O₅. The figures are corrected for a trace of K₂SO₄ which was always present:

63.08 63.24 63.27 63.42 Mean, 63.25, ± .047

Hence we have four values for Ta:

Or, if O = 16, Ta = 182.562.

If we assume K = 39, O = 16, F = 19, S = 32, and N = 14; the percentage of K_2SO_4 from K_2TaF_7 gives Ta = 181.912; and the analyses of the ammonium salt make Ta = 182.020. Evidently, 182 is not far from the true value.

PLATINUM.

For this metal we have to consider only experiments by Berzelius, by Andrews, and by Seubert. In an early paper Berzelius* reduced platinous chloride, and found it to contain 73.3 per cent. of platinum. Hence, Pt = 194.204, a

^{*} Poggend. Annal., 8, 177. 1826.

value very near that obtained most recently by Seubert. In his later investigations, Berzelius* studied the potassium chloroplatinate, K₂PtCl₆. 6.981 parts of this salt, ignited in hydrogen, lost 2.024 of chlorine. The residue consisted of 2.822 platinum, and 2.135 potassium chloride. From these data we may calculate the atomic weight of platinum in four ways:

The last of these values is undoubtedly the most reliable, since it involves no errors due to the possible presence of moisture in the salt analyzed. If O=16, the value becomes Pt=197.104.

The work done by Andrews † is even less satisfactory than the foregoing, for the reason that its full details seem never to have been published. Andrews dried potassium chloroplatinate at 105°, and then decomposed it by means of zinc and water. The excess of zinc having been dissolved by treatment with acetic and nitric acids, the platinum was collected upon a filter and weighed, while the chlorine in the filtrate was estimated by Pelouze's method. Three determinations gave as follows for the atomic weight of platinum:

If we assume that these values were calculated with K = 39 and Cl = 35.5, the mean, corrected by our later figures for these elements, becomes Pt = 197.382. If O = 16, this becomes Pt = 197.836. Unfortunately, Andrews does not, in his brief note upon the subject, indicate the manner by which his calculations were made.

^{*} Poggend. Annal., 13, 468. 1828.

[†] British Association Report, 1852. Chem. Gazette, 10, 380.

Latest of all we have to consider the experiments of Seubert.* This chemist prepared very pure chloroplatinates of ammonium and potassium, and from their composition deduced the atomic weight of the metal under consideration. The ammonium salt, (NH₄), PtCl₆ was analyzed by heating in a stream of hydrogen, expelling the excess of that gas by a current of carbon dioxide, and weighing the residual metal. In three experiments the hydrochloric acid formed during such a reduction was collected in an absorption apparatus, and estimated by precipitation as silver chloride. Three series of results are given for the percentage of platinum in this salt, together with another single result which may be considered alone. Here are the figures:

Series I.	Series II.	Series III.
43.957	43.871	43.990
43.948	43.876	43.986
43.960	43.872	44.001
43.946	43.881	44.020
43.963	43.875	43.994
43.961	43.879	43.996
		44.004
Mean, 43.956 , $\pm .002$	Mean, 43.876, \pm .001	44.026
		43.998
		-
		Mean, 44.001, ± .003

These series represent three preparations. The additional single experiment above referred to was made with material belonging to series II, but recrystallized from water. This salt gave 43.955 per cent. of platinum, a figure to which we may assign the probable error of one experiment in the first series. Combining, we get the subjoined general mean percentage of Pt in (NH₄)₂PtCl₆:

Series	I	43.956, ± .002
66	II	43.876, \pm .001
66	III	44.001, ± .003
Extra	experiment	43.955, \pm .004
	General mean	43.907, ± .0009

^{*} Ber. der Deutsch. Chem. Gesell., 14, 865. 1881.

Hence Pt = 194.314, \pm .078. If N = 14, and Cl = 35.5, then Pt = 194.906. Calculating with Stas' values for N and Cl, Seubert gets from the four results combined above, the following figures for Pt, respectively: 194.685, 194.039, 195.034, 194.665.

For the chlorine estimations in the ammonium salt the subjoined weighings are given:

Salt.	Pt.	AgCl.
2.7054 grm.	1.1871 grm.	5.2226 grm.
2.2748 "	.9958 "	4.3758 "
3.0822 "	1.3561 "	5.9496 "

Hence 100 parts of AgCl correspond to the following quantities of salt:

$$51.802
51.986
51.805
Mean, 51.864, \pm .041$$

Hence, calculating directly from the ratio between 6AgCl and $(NH_4)_2$ PtCl₆, Pt = 196.871, \pm .363.

Seubert himself reckons the percentage of chlorine from the weight of silver chloride, and then calculates the ratio between Cl_6 and Pt. He thus finds, with Stas' value for Cl, Pt = 195.330.

The potassium salt, K, PtCl₆, was also analyzed by ignition in hydrogen, treatment with water, and weighing both the platinum and the potassium chloride. These percentages were found:

Pt.	KCI.
40.119	30.706
40.120	30.728
40.076	30.698
40.070	30.66 6
40.107	30.700
40.120	30.627
40.114	30.710
40.130	30.621
Mean 40.107, \pm .005	Mean, $30.682_{1} \pm .009$

From the first column_____Pt = 194.370, ± .068
" second " = 194.645, ± .213

If K = 39, and Cl = 35.5, the first column gives Pt = 194.933. Seubert, from the percentage of platinum, gets Pt = 194.392; and from the ratio 2KCl: Pt he finds Pt = 194.494.

As with the ammonium salt, three experiments were made upon the potassium compound to determine the amount of chlorine lost upon reduction in hydrogen. I cite the weighings, and add in a fourth column the quantity of K₂PtCl₆ proportional to 100 parts of AgCl. This AgCl represents but four atoms of the chlorine:

Salt.	Pt.	AgCl.	Ratio.
6.7771 grm.	2.7158 grm.	7.9725 grm.	85.006
3.5834 "	1.4372 "	4.2270 "	84.774
4-4139 "	1.7713 "	5.2144 "	84.648
		Mean.	

Hence Pt = 195.002, $\pm .415$. If K = 39, Ag = 108, and Cl = 35.5, then Pt = 194.955. Seubert, calculating the percentage of chlorine and thence the ratio Cl₄: Pt, gets Pt = 194.631.

Combining all the values we have the following result for the atomic weight of platinum:

```
1. From per cent. Pt in (NH_4)_2PtCl<sub>6</sub> .....Pt = 194.314, ± .078

2. " 6AgCl: (NH_4)_2PtCl<sub>6</sub> ratio......" = 196.871, ± .363

3. " per cent. Pt in K_2PtCl<sub>6</sub> ......" = 194.370, ± .068

4. " " KCl " ......" = 194.645, ± .213

5. " 4AgCl: K_1PtCl<sub>6</sub> ratio....." = 195.002, ± .415

General mean ....." = 194.415, ± .049
```

Or, if O = 16, Pt = 194.867.

Seubert, taking the arithmetical mean of his eight values, gets Pt = 194.620. He regards, however, those results as best which are dependent upon the percentage of platinum in the ammonium salt, and upon the complete analysis of the potassium compound. These give him a mean of Pt = 194.461, which, if corrected by reduction to a vacuum standard, becomes Pt = 194.34.

In will be noticed that three of the ratios, calculated with

K = 39, N = 14, Ag = 108, and Cl = 35.5, give nearly Pt = 195, namely:

194.906 194.933 194.955

The general mean of all, if O = 16, gives Pt = 194.867. Hence, for all practical calculations, the value 195 may be safely employed.

OSMIUM.

The atomic weight of this metal has been determined by Berzelius and by Fremy.

Berzelius* analyzed potassium osmichloride, igniting it in hydrogen like the corresponding platinum salt. 1.3165 grammes lost .3805 of chlorine, and the residue consisted of .401 grm. of potassium chloride, with .535 grm. of osmium. Calculating only from the ratio between the Os and the KCl, we have, Os = 198.494; or, if O = 16, Os = 198.951.

Fremy's determination \uparrow is based upon the composition of osmium tetroxide. No details as to weighings or methods are given; barely the final result is stated. This, if O = 15.9633, is Os = 199.190. If O = 16, Os = 199.648.

Berzelius' work is evidently entitled to preference, although neither determination is in any sense equal to the present requirements of chemical science. The values given are doubtless several units too high.

IRIDIUM.

The only early determination of the atomic weight of iridium was made by Berzelius,‡ who analyzed potassium iridichloride by the same method employed with the platinum and the osmium salts. The result found from a single

^{*} Poggend. Annal., 13, 530. 1828.

[†] Compt. Rend., 19, 468. Journ. für Prakt. Chem., 33, 410. 1844.

[‡] Poggend. Annal., 13, 435. 1828.

255

analysis was not far from Ir = 196.7. This is now known to be too high. I have not, therefore, thought it worth while to recalculate Berzelius' figures, but give his estimation as it is stated in Roscoe and Schorlemmer's "Treatise on Chemistry."

In 1878 the matter was taken up by Seubert,* who had at his disposal 150 grammes of pure iridium. From this he prepared the iridichlorides of ammonium and potassium, (NH₄), IrCl₆ and K, IrCl₆, which salts were made the basis of his determinations. The potassium salt was dried by gentle heating in a stream of dry chlorine.

Upon ignition of the ammonium salt in hydrogen, metallic iridium was left behind in white coherent laminæ. The percentages of metal found in seven estimations were as follows:

The potassium salt was also analyzed by decomposition in hydrogen with special precautions. In the residue the iridium and the potassium chloride were separated after the usual method, and both were estimated. Eight analyses gave the following results, expressed in percentages:

Ir.	₂KCl.	Cl ₄ .
39.881	30.829	29.290
39.890	30.842	29.277
39.86 8	30.813	29.300
39.876	30.835	29.289 ·
39.877	30.825	29.287
39.879	30.811	29 .310
39:882	30.814	29.285
39.883	30.792	29.288
Mean, 39.880, ± .0015	$\phantom{00000000000000000000000000000000000$	29.291, ± .0024

^{*} Ber. d. Deutsch. Chem. Gesell., 11, 1767.

From these data several values for the atomic weight of iridium may be calculated:

If O = 16, this becomes Ir = 193.145.

In the potassium salt, instead of calculating from the percentages directly, we may reckon upon the ratios between Ir and Cl₄, and between Ir and 2KCl:

Or, if O = 16, Ir = 192.982.

Again, we may combine this mean with the value derived from the ammonium iridichloride, and so estimate the relative importance of the latter:

If O = 16, this becomes Ir = 193.094.

We may assume, then, from all the facts before us, that if O = 16, the atomic weight of iridium varies from the even number 193 only within the limits of experimental error.

PALLADIUM.

The atomic weight of palladium has been studied by Berzelius and by Quintus Icilius. In an early paper Berzelius* found that 100 parts of the metal united with 28.15 of sulphur. Hence Pd = 113.63, a result which is unquestionably far too high.

^{*} Poggend. Annal., 8, 177. 1826.

In a later paper* Berzelius published two analyses of potassium palladiochloride, K₂PdCl₄. The salt was decomposed by ignition in hydrogen, as was the case with the double chlorides of potassium with platinum, osmium, and iridium. Reducing his results to percentages, we get the following composition for the substance in question:

Pd.	≥KCl.	Cl₂.	
32.726	46.044	21.229	
32.655	45.741	21.604	
Mean, 32.690	45.892	21.416	

From these percentages, calculating directly, very discordant results are obtained:

Obviously, the only way to get satisfactory figures is to calculate from the ratio between the Pd and 2KCl. Doing this, we get, Pd = 105.737; or, if O = 16, Pd = 105.981.

This last value varies so slightly from the even number 106 that the latter may be safely used for all purposes of chemical calculation.

The determination made by Quintus Icilius* need be given only for the sake of completeness. He ignited potassium palladichloride in hydrogen, and found the following amounts of residue. His weights are here recalculated into percentages:

64.708 64.965 64.781 Mean, 64.818

From this mean, Pd = 111.879. Upon looking at the values deduced from Berzelius' figures, it will be seen that

^{*} Poggend. Annal., 13, 454. 1828.

^{† &}quot;Die Atomgewichte vom Pd, K, Cl, Ag, C, und H, nach der Methode der kleinsten Quadrate berechnet." Inaug. Diss. Göttingen, 1847. Contains no other original analyses.

the highest, 110.796, is calculated from the chlorine lost upon igniting the palladiochloride. The same kind of error which vitiates that result probably affects also these data drawn from the palladiochloride.

RHODIUM.

Berzelius* determined the atomic weight of this metal by the analysis of sodium and potassium rhodiochlorides, Na,RhCl₅, and K,RhCl₅. The latter salt was dried by heating in chlorine. The compounds were analyzed by reduction in hydrogen, after the usual manner. Reduced to percentages the analyses come out as follows:

•	In Na ₈ RhCl ₈ .	
Rħ.	зNaCl.	Cl ₃ .
26.959	45.853	27.189
27.229	45.301	27.470
		27.616
Mean, 27.094	45-577	
		27.425
	In K2RhCl3.	
Rh.	2KCl.	CI ₃ .
28.989	41.450	29.561

From the analyses of the sodium salt we get the following values for Rh:

These are discordant figures, and indicate some doubt as to purity of material. The last value is fairly good, however, and is confirmed by results from the potassium compound:

^{*} Poggend. Annal., 13, 435. 1828.

If O = 16, this becomes Rh = 104.285.

RUTHENIUM.

The atomic weight of this metal has been determined only by Claus.* Although he employed several methods, the only results worthy of present notice come from the analysis of potassium rutheniochloride, K,RuCl. The salt was dried by heating to 200° in chlorine gas, but even then retained a trace of water. The percentage results of analysis are as follows:

Ru.	≥KCl.	CI ₈ .
28.96	40.80	30.24
28.48	41.39	30.22
28.91	41.08	30.04
Mean, 28.78	41.09	30.17

Reckoning directly from the percentages we get the following discordant values for Ru:

From perce	entage of	metalRu	=	103.016
46	"	KCl"	=	107.190
"	"	Cl ₈ "	=	96.854

Obviously, the best result is to be obtained from the ratio between Ru and 2KCl. This gives Ru = 104.217; or, if O = 16, Ru = 104.457. But little weight can be attached to this determination.

^{*} Journ. für Prakt. Chem., 34, 435. 1845.

	•	
	•	
		-
	`	

APPENDIX.

ON DUMAS' CORRECTION AND PROUT'S HYPOTHESIS.

In the year 1815 Prout put forth his famous hypothesis that the atomic weights of all the elements were multiples of that of hydrogen. His views were adopted by many chemists, but opposed by others; among them Berzelius and Turner; and down to the present day "Prout's Law" has been the subject of earnest controversy. Of course the fact was early recognized that in its original form the hypothesis could not stand, and accordingly it was modified by Dumas in such manner that half and quarter multiples of the atomic weight of hydrogen were considered as well as the whole numbers.

But of late years Prout's hypothesis, even with its elastic modification, has been in disfavor. Only a few chemists still clung to it as the representative of a veritable law. The researches of Stas were especially directed towards ascertaining its truth or falsity; and his results, as well as those obtained by Marignac, were such as to lead most chemists to the belief that it had been forever overthrown. The atomic weights determined by Stas agreed neither with whole, half, nor quarter multiples of that of hydrogen, and the variations seemed to be wholly outside the range of recognizable experimental errors.

In 1878, however, a probable source of error in some of Stas' researches was pointed out by Dumas.* Many of Stas' ratios had involved the use of pure metallic silver, which had been fused under a cover of borax containing a little

nitre. Such silver Dumas heated to redness in a Sprengel vacuum, and found that it gave up weighable quantities of oxygen, which had been absorbed by the metal when in the melted state. In one experiment a kilogramme of silver gave 82 milligrammes of occluded gas, and in three other cases 226,140, and 249 milligrammes respectively were found. In other words, the silver which had been considered pure by Stas and others, was really not pure, and a correction became necessary in nearly all series of atomic weight determinations.

The amount of this correction, which I think may hereafter be appropriately designated as "Dumas' correction," will naturally vary in different cases, and in no particular case can we tell, without actual examination of the silver employed, exactly how great it should be. We may, however, assume that all the metallic silver heretofore used in establishing atomic weight ratios was subject to it; and, reckoning from the largest error indicated in the experiments of Dumas, namely, 249 milligrammes of oxygen in the kilogramme of metal, we may ascertain its tendency with reference to Prout's law.

In the chapter upon the atomic weights of silver, chlorine, bromine, iodine, potassium, sodium, and sulphur, twenty ratios are given, of which nine are subject to Dumas' correction. Applying it as suggested above, we get the following results. The values previously found and given in the chapter just quoted, we may designate as uncorrected. For convenience in future reference I assume that O = 16:

	Uncorrected.	Corrected.	Difference
Silver	107.923	107.896	027
Chlorine	35.451	35.478	+ .027
Bromine	79.951	79.978	+ .027
Iodine	126.848	126.875	+ .027
Potassium	39.109	39.083	026
Sodium	23.051	23.024	027
Sulphur	32.058	32.058	

The result of the correction, it will be seen, is generally favorable to Prout's hypothesis. Of the seven elements

under consideration, one has its atomic weight unaffected, one is rendered less in accord with the hypothesis, and five approximate more closely than before to even multiples or multiples half of hydrogen.

In the later chapters of this work the effect of Dumas' correction is generally less striking. One general statement, however, may be made concerning it. Whenever the atomic weight of a metal is calculated from the ratio between its haloid salts and metallic silver, the total effect of Dumas' correction, including the above corrections for the halogens themselves, will be to lower the final result. This point will be further considered presently. Only chlorine, bromine, and iodine have their atomic weights raised by the correction.

In view of Dumas' correction the question naturally arises as to how far other metals, used in atomic weight researches, may occlude gaseous impurities. For example, when the atomic weight of oxygen is fixed by the synthesis of water over copper oxide, may not the copper occlude appreciable quantities of the hydrogen in which it cools? If it does, then the apparent weight of metallic copper would be too high, and the atomic weight of oxygen would come out too low. Such an error might possibly account for the difference between 16 and 15.9633 in the atomic weight of oxygen, and it would also increase the atomic weight of copper as determined by the same process. At all events, every metal of which the atomic weight has been determined by the reduction of its compounds in hydrogen, ought to be scrupulously investigated with reference to the possible occlusion of gaseous impurities. With all of these metals the effect of such impurities would be to render the apparent atomic weights decidedly too high.

Although every series of atomic weight determinations must be considered by itself, and weighed on its own merits, it may not be out of place for me just here to point out two general sources of error in addition to the one we have been considering. First, every value after oxygen, with one or two partial exceptions, involves whatever error may attach

to the atomic weight of oxygen. If the latter be 16, instead of 15.9633, this error in some instances becomes multiplied to a large fraction of a unit, as the subjoined example will show.

Other similar errors are repeated continually. The value assigned to any element is necessarily affected by whatever errors may attach to the atomic weights of those other elements through whose medium it is compared with the standard, hydrogen. Thus, the atomic weight of carbon depends upon that of oxygen; calcium depends upon both carbon and oxygen; and fluorine, as determined from calcium fluoride, involves the foregoing elements, together with sulphur, silver, and chlorine. Since, however, some atomic weights are affected by plus errors and others by minus errors, there is a fortunate tendency to compensation of errors in cases like that of fluorine, and, in reality, better results are obtained than considerations such as these would lead us to look for.

Another general source of error is to be found in the fact that some of the weighings involved in our discussions had been reduced to absolute standards, while others were merely uncorrected weighings in air. The errors thus introduced into the work are doubtless small, but still they ought not to be absolutely ignored.

Now, having considered the larger classes of errors, we may properly pass on to a comparison of our atomic weights with reference to Prout's hypothesis. In order to facilitate work, I have tabulated the figures in two columns, one giving atomic weights referred to hydrogen as unity, the other based upon the standard of oxygen as exactly sixteen. Such imperfectly known elements as decipium, philippium, samarium, terbium, and thulium are not included.

APPENDIX.

TABLE OF ATOMIC WEIGHTS.

	IABLE OF A	OMIC WI	
	H = 1.	O = 16.	Remarks.
Aluminum	27.009, ± .003	27.075	Cashah and Sahmidada data
Antimony	119.955, ± .036	120.231	Cooke's and Schneider's data.
Arsenic Barium	74.918, \pm .016	75.090 137.007	
Bismuth	$207.523, \pm .082$	208.001	From Schneider's data.
Boron	10.941, ± .023	10.966	l rom bonned b ann.
Bromine	79.768, ± .019	79.951	
Cadmium	111.835, \pm .024	112.092	İ
Cæsium	$132.583, \pm .024$	132.918	
Calcium	39.990, ± .010	40.082	4
Carbon	11.9736, \pm .0028	12.0011	
Cerium	140.424, ± .017	140.747	Buehrig's data give 141.523. (O = 16.)
Chlorine	$35.370, \pm .014$	35.451	P 6: 11.
Chromium	52.009, ± .025	52.129	From Siewert's data.
Cobalt Columbium	58.887, ± .008	59.023	From one ratio only.
Copper	93.812 63.173, ± .011	94.027 63.318	From one ratio only.
Didymium	144.573, ± .031	144.906	Cleve's data give 147.021.
Erbium	165 800	166.273	$(SO_8 = 80.)$ From Cleve's data only.
Fluorine	165.891 18.984, ± .0065	19.027	From Cieve's data omy.
Gallium'	68.854	68.963	Imperfectly determined.
Glucinum	9.085, ± .0055	9.106	Nilson and Pettersson's data.
Gold	196.155, ± .095	196.606	
Hydrogen	1.0000	1.0023	
Indium	113.398, ± .047	113.659	
Iodine	126.557, ± .022	126.848	
Iridium	$192.651, \pm .033$	193.094	Seubert's data.
Iron	$55.913, \pm .012$	56.042	
Lanthanum	138.526, ± .030	138.844	
Lead Lithium	206.471, ± .021	206.946 7.0235	
Magnesium	7.0073, ± .007 23.959, ± .005	24.014	Marchand and Scheerer's data.
Manganese	53.906, ± .012	54.029	Schneider and Rawack's data.
Mercury	199.712, ± .042	200.171	
Molybdenum	95.527, ± .051	95.747	
Nickel	57.928, ± .022	58.062	Schneider, Sommaruga, and Lee.
Nitrogen	14.0210, ± .0035	14.029	
Osmium	198.494	198.951	Very doubtful.
Oxygen	15.9633, ± .0035	16.000	
Palladium	105.737	105.981	Badly determined.
Phosphorus	$30.958, \pm .007$	31.029	Combando dasa
Platinum	194.415, ± .049	194.867	Seubert's data.
Potassium Rhodium	39.019, ± .012	39.109 104.285	Badly determined.
Rubidium	104.055 85.251, ± .018	85.529	Dadry determined.
Ruthenium	104.217	104.457	Badly determined.
Scandium	43.980, ± .015	44.081	
Selenium	78.797, ± .011	78.978	

TABLE OF ATOMIC WEIGHTS-CONTINUED.

	H == 1.	O = 16.	Remarks.
Silicon Silver Sodium Strontium Sulphur Tantalum Tellurium Thallium Thorium Tin Titanium Tungsten Uranium Vanadium Ytterbium Yttrium Zinc Zirconium	28.195, ± .066 107.675, ± .0096 22.998, ± .011 87.374, ± .032 31.984, ± .012 182.144, ± .466 127.960, ± .034 203.715, ± .073 117.698, ± .040 49.846, ± .064 183.610, ± .032 238.482, ± .082 51.256, ± .024 172.761, ± .038 89.816, ± .067 64.9045, ± .019 89.367, ± .037	28.260 107.923 23.051 87.575 32.058 182.562 128.254 204.183 233.951 117.968 49.961 184.032 239.030 51.373 173.158 90.023 65.054	Very badly determined. Imperfectly determined. Crookes' data. Imperfectly determined. If SO ₁ = 80, Yb = 173.016. Doubtful. Axel Erdmann's data. Doubtful.

At the close of his admirable paper on the atomic weight of aluminum Mallet makes substantially the following argument in favor of Prout's hypothesis. Citing the atomic weights of eighteen elements which he considers well determined, he shows that ten of them have values falling within one-tenth of a unit of whole numbers. Now, what is the mathematical probability that this close approximation to conformity with Prout's law, in ten cases out of eighteen, is purely accidental, as those chemists who reject the hypothesis seem to hold? Working this problem out, Mallet finds the probability in favor of mere coincidence to be in the ratio of 1: 1097.8, and hence he concludes that Prout's views are still worthy of respectful consideration.

Applying Mallet's reasoning to the table of atomic weights now before us, we find that in the first column, when H=1, twenty-five elements out of sixty-six have values falling within the limits of one-tenth of a unit variation from whole numbers. But many of the figures which fall without this limit involve the variation of oxygen multiplied many times over. We must therefore study the second column, which assumes that the atomic weight of oxygen is exactly six-

teen. Here we have forty elements falling within the limit of variation assigned by Mallet, and twenty-six falling without. The variations we may properly study in some detail.

Taking first the elements whose atomic weights vary from even multiples of unity by less than a tenth of a unit, we have to consider the following: aluminum, arsenic, barium, bismuth, boron, bromine, cadmium, cæsium, calcium, carbon, cobalt, columbium, didymium, fluorine, gallium, hydrogen, iridium, iron, lead, lithium, magnesium, manganese, nickel nitrogen, osmium, oxygen, palladium, phosphorus, scandium, selenium, silver, sodium, sulphur, thorium, tin, titanium, tungsten, uranium, yttrium, and zinc. Of these, aluminum, arsenic, barium, bismuth, cadmium, calcium, carbon, cobalt, columbium, fluorine, hydrogen, iridium, iron, lithium, magnesium, manganese, nickel, nitrogen, phosphorus, scandium, sodium, sulphur, tungsten, uranium, yttrium, and zinc have plus variations, while boron, bromine, cæsium, didymium, gallium, lead, osmium, palladium, selenium, silver, thorium, tin, and titanium fall slightly under the units to which they approximate. Oxygen, as the standard of comparison, of course shows here no variation, its possible error having been transferred to hydrogen.

Of the foregoing elements it will be seen that twenty-six have plus variations from whole numbers, while thirteen are minus. Among the latter, boron, gallium, osmium, palladium, thorium, and titanium have been but roughly determined. Bromine, by Dumas' correction, has its variation diminished. In the cases of lead, cæsium, selenium, and tin, the cause of variation, supposing one to exist, remains to be determined. The value for osmium is undoubtedly several units too high, so that its agreement with Prout's law may be considered purely accidental. As for didymium, the figure assigned is the mean of all determinations; whereas Cleve's data, calculated with SO₃ = 80, make Di = 147.021, a variation which, like most of the others, is far within the limits of ordinary experimental error. In the

case of silver it has already been shown that Dumas' correction is unfavorable to it considered in its bearings upon Prout's law. Silver is the only element among those having minus variations which could carry very much weight against the hypothesis.

Among the elements whose variations are plus, columbium, uranium, and yttrium have been poorly determined. Yttrium especially may be considered doubtful. weights of aluminum, arsenic, barium, cadmium, lithium, phosphorus, and sodium involve Dumas' correction to a greater or less extent, and will be lowered by its application, that is, brought nearer to whole numbers. For aluminum, certain other causes for variation were pointed out in the chapter upon that metal; and it may be noted that the direct ratio between it and hydrogen gives Al = 27.998. ± .007. Here the variation is less than the probable error. For calcium, and consequently for fluorine also, sources of plus error were indicated in the discussion of their respective atomic weights, and reiteration here is unnecessary. iridium, iron, nickel, and tungsten all involve such errors as may arise from the possible occlusion of hydrogen by the metals after reduction from their compounds. For scandium, the atomic weight, calculated with $SO_{3} = 80$, becomes 44.032, a variation much within the limits of experimental error. For carbon and bismuth the variations are insignifi-In short, in the majority of instances the errors may be diminished by corrections which are in all probability needed, and which can be easily pointed out. The more carefully we scrutinize the data the more probable Prout's hypothesis appears.

Among the twenty-six elements whose atomic weights are removed by more than a tenth of a unit from whole numbers, chlorine, rubidium, and strontium have values nearly half multiples of that of hydrogen, and in each case Dumas' correction will make the approximation still closer. Erbium, gold, indium, lanthanum, rhodium, ruthenium, silicon, and zirconium may be dismissed from consideration as too imperfectly determined to carry much weight in the present

discussion. For chromium, copper, molybdenum, and vanadium I have no criticisms to offer; but the remaining elements may be considered individually.

The value assigned to antimony, 120.231, is the general mean of Cooke's and Schneider's work upon the bromide, iodide, and sulphide. If Ag = 108, Br = 80, and I = 127, Cooke's data for the bromide and iodide give the following values for Sb, all of which fall within a tenth of a unit of the whole number 120:

Early bromid	le serie	s	Sb	=	119.901
Late	**		"	==	120.009
Iodide series			"	=	119.973

In the case of cerium, the value assigned in the table is the general mean of all reputable determinations. But it is subject to doubt on account of the facts observed by Wolf and by Wing, whose ceroso-ceric oxide was white, while that of all other observers was yellowish. Wolf's and Wing's data, calculated with O = 16, give Ce = 138.039. Cerium, then, is not an established exception to Prout's law.

Glucinum and ytterbium have their atomic weights calculated from analyses of the sulphates. But if Prout's law is true, $SO_3 = 80$. Calculated with this figure, we have Gl = 9.096 and Yb = 173.016. Both elements thus fall within reasonable limits of variation from the hypothetical values.

Iodine is one of the most important seeming exceptions. If we assume Ag = 108, and calculate the atomic weight of iodine only from the direct ratio between iodine and silver, we have, with Dumas' correction applied, I = 126.966; that is, it comes within one-tenth of a unit of the whole number 127.

The atomic weight of mercury depends upon analyses of the chloride, oxide, and sulphide. Of these three compounds the purity of the chloride is most easily assured. Calculated from its composition, with Cl = 35.5, Hg = 199.971. With so high an atomic weight small errors are easily multiplied.

For the atomic weight of platinum Scubert's data give five values, ranging both above and below the round number 195. Calculated with integer values for the other elements, three of these figures fall very close to 195, as follows:

From per cent. Pt in
$$(NH_4)_2$$
PtCl₆----Pt = 194.906
" K_2 PtCl₆------ " = 194.933
From chlorine estimation in K_2 PtCl₆-- " = 194.955

Potassium is the most serious exception of all. But if O=16 and Dumas' correction be applied, the general mean from all the available data becomes K=39.083. That is, potassium falls within the limit of 0.1 variation.

The atomic weight assigned to tantalum is the mean of four values. Two of these, recalculated with integers, come out as follows:

For tellurium I need only call attention to the discrepancies between the several sets of determinations made by Wills. A reference to the chapter on tellurium will show that his figures give results ranging from Te = 126.07 to Te = 129.34. The mean value is therefore too much subject to doubt to carry weight as an exception.

As for thallium, the last case to be considered, I have already shown that Crookes' data, recalculated with integer values for N and O, give Tl = 204.008. That is, instead of an exception, we have here an admirable instance in support of Prout's hypothesis.

Enough has been said in this brief resumé to show that none of the seeming exceptions to Prout's law are inexplicable. Some of them, indeed, carefully investigated, support it strongly. In short, admitting half multiples as legitimate, it is more probable that the few apparent exceptions are due to undetected constant errors, than that the great number of close agreements should be merely accidental. I began this recalculation of the atomic weights

with a strong prejudice against Prout's hypothesis, but the facts as they came before me have forced me to give it a very respectful consideration. All chemists must at least admit that the strife over it is not yet ended, and that its opponents cannot thus far claim a perfect victory.

A .	Barium chloride and lanthanum sul-
	phate 230
Allen and Johnson 91	and silver 57
Allen and Pepys 6	and silver chloride 60
Aluminum 156	chromate 121, 122
bromide 159	metatungstate 147
chloride156	nitrate 61
oxide 156	selenite 177
sulphate156	silicofluoride 85
Ammonia alum 158	sulphate and barium chlo-
Ammonia chrome alum 120	ride 61
Ammonium chloride 40	and barium chromate_ 121
chloroplatinate 251	and barium fluoride 80
cobalticyanide 172	and barium nitrate 61
fluotantalate 249	and barium selenite 177
iridichloride 255	and cerium dioxide, 222, 224
molybdate 139	and didymium sulphate, 236
Anderson 76	and glucinum sulphate, 97
Andrews 57, 250	and gold 163
Antimony 188, 269	and indium oxide 219
bromide 198, 200	and lanthanum sul-
chloride, 190, 191, 193, 194, 197	phate 229
iodide 199	and magnesium sul-
oxide 188, 190	
sulphide 189, 196, 200	phate 100 and nickel potassium
Antimony compounds, oxidation	sulphate
of 189, 191, 192, 193, 194	and thallium sulphate, 93
Antimony and potassium tartrate_ 194	and thorium sulphate_ 214
Arago	and yttrium oxide 241
Arfvedson 87, 127, 151	uranate151
Arsenic 185	and uranyl acetate 151
bromide186	Beringer 221
chloride186	Berlin 118, 139
trioxide 123, 187	Bernoulli 145
Arsenious oxide and potassium an-	Beryllium. See Glucinum 96
hydrochromate 123, 187	Berzelius 1, 6, 10, 14, 19, 21, 27, 31,
Arsenious oxide and potassium	39, 60, 61, 68, 70, 72, 78, 82, 84,
chlorate 187	86,87,100,108,117,127,131,135,
Atomic weights, table of 265, 266	137, 143, 151, 156, 162, 176, 180,
Awdejew 97	183, 185, 188, 205, 212, 214, 248,
_	249, 254, 256, 258
В.	Biot and Arago 6, 39
i	Bismuth 202
Bahr 102, 242, 244	chloride 203
Bahr and Bunsen 242, 244	oxide 203
Balard 21	sulphide202
Barium 57	Blomstrand 247
chloride and barium chro-	Boisbaudran 218
mate 122	Borax 84
and barium sulphate 70	Borch 144
and cerium sulphate 223	Boron
18	•
10	273

Boron bromide 84	Cobaltiswanida of abanulamma
chloride 84	Cobalticyanide of phenylammo-
	nium 172
Brauner 96 Bromine 9, 21	of strychnia174
Brucia cobalticyanide 173	Columbic acid247
pickelowepide	Columbium247
nickelocyanide 173	oxychloride 247
Buehrig 226	pentachloride 247
Buff 6	Cooke 29, 112, 195, 198, 199
Bunsen 90, 91, 92, 220, 242, 244	Copper 135
Bunsen and Jegel 223	oxide
Burney 245	sulphate136
_	subsulphide 136
C. ,	Crookes 95
· ·	Czudnowicz 183, 233
Cadmium 111	
bromide II2	<u>_</u>
chloride 112	D.
oxalate 112	
sulphate	Davy 6, 78, 127
sulphide III	Debray 98, 140
Cæsium 91	Decinium 2.6
bitartrate 91	Decipium 246 Delafontaine 215, 242, 244
chloride 91	
Calcium 67	De Luca
carbonate 67	Demoly 209
and calcium sulphate_ 69	De Saussure 6
chloride70	Deville 84, 248 Deville and Troost 248
fluoride 78	
oxide and calcium sulphate, 70	Dewar and Scott130
sulphate 69, 70	Dexter190
Capitaine 131	Diamond 54
Carbon 50	Didymium 236
dioxide 54, 56	chloride 237
monoxide55	oxalate 237
Cavendish6	oxide 237
Cerium 221, 269	sulphate236
chloride21	Diehl 87
formate 222	Dœbereiner 72, 131
oxalate 223, 224, 227	Dulong and Berzelius 1, 6, 39 Dumas, 1, 2, 16, 27, 28, 30, 58, 64, 67
oxide 221, 224	Dumas, 1, 2, 10, 27, 28, 30, 58, 04, 07
sulphate, 221, 222, 223, 224, 225	70, 75, 79, 80, 83, 84, 85, 106, 112,
sulphide 223	128, 134, 136, 139, 145, 156, 167,
Chlorine 9	178, 181, 186, 191, 203, 205, 206,
Choubine 229	261
Chrome alum	Dumas and Boussingault 6, 39
Chromium 117	and Stas 54
acetate 118	Dumas' correction 261
chloride 124	
sulphate 120	-
Chydenius 214	E.
Clark2	
Claus 259	Ebelmen 152
Cleve 217, 234, 238, 240, 242, 246	Bkman and Petterson 178
Cleve and Hoeglund 242	Erbium 244
Cobalt164	oxide 244
chloride 166, 167	sulphate244
oxalate 166, 168	Erdmann (Axel) 108
sulphate 166	Erdmann and Marchand_ 1, 4, 28, 54.
Cobalticyanide of ammonium 172	68, 69, 115, 132, 135, 164, 177
of brucia 173	Erk234, 238

F.	Indium sulphide 219
_	Iodine9, 269
Faget12	and silver 25
Favre 109	Iridium 254
Ferric chloride 134	Iron 131
oxide	chlorides 134
Ferrous chloride134	oxide131
tungstate 147	tungstate 147
Fluorine 78	Isnard 156
Fluor spar 78 Fourcroy 6	J.
Fremy 254	J.
1 1cmy 254	Jacquelain 82, 101, 108, 118
G.	Jegel 223
-	Johnson and Allen 91
Gallium 218	Johnson and Ameni 91
ammonium alum	K.
nitrate 219	
oxide 219	Kemp 140
Gay-Lussac 108	Kessler 122, 187, 189, 191, 199
Genth 226	Kirwan
Gerhardt II	Kjerulf 223
Glucinum 96, 269	Klatzo
and ammonium oxalate 08	Kralovanzky 87
sulphate 97	
Gmelin	L.
Godeffroy 90, 91	
Gold	Lagerhjelm 202
and cobalt 172	Lamy 93
indium 220	Lanthanum 229
mercury 162	carbonate 233
nickel 172	iodate 232
phosphorus 163	magnesium nitrate 232
Graphite 54	oxalate231
	oxide 232
Ħ.	sulphate 229
• 0	Laurent
Hagen 87	Lavoisier 6, 39
Hampe	Lead 72
Hauer 111, 128, 180 Hebberling 93	chloride 74 chromate 118
Hermann, 87, 212, 216, 222, 233, 238,	fluoride 80
247, 248	
Hilgard 158	nitrate
Hoeglund 242, 244	and lead sulphate 74
Holmium246	oxide 72
Holzmann 231	and lead sulphate 74
Humpidge 96, 245	sulphate 72
Humpidge and Burney 245	and lead fluoride 80
Huntington 112	and lead nitrate 74
Hydrogen 1, 3, 6, 7	and lead oxide 74
and aluminum 157, 160	Le Conte 7, 40
cobalt 169	Lee 172
nickel 169	Lefort 121
	Lenssen 112
I.	Liebig 21, 51
_	Liebig and Redtenbacher 51
Iceland spar67	Liechti and Kemp 140
219	Lithium
oxide 219	carbonate

Lithium chloride 87, 89	Nickel oxalate 165, 166
nitrate 89	oxide 164, 168
sulphate 88	potassium sulphate 170
Loewig 21	sulphate 165, 166
Louyet 78, 79	Nilson 96, 99, 240, 244 and Pettersson 96, 99
Louyet 70, 79	Niobium. See Columbium 247
M.	Nitrogen
	Nitzsch 176
Macdonnell100	Nordenfeldt102
Magnesite103	Norlin 131
Magnesium 100	
carbonate 103	O .
chloride 106	
lanthanum nitrate 232	Osmium 254
oxalate [02]	Oxygen
oxide 101	P.
sulphate 100 uranate 151	••
Malaguti 143	Palladium 256
Mallet 87, 157, 267	sulphide 256
Manganese 127	Peligot 118, 151, 153
chloride 128	Pelouze, 11, 32, 40, 57, 64, 82, 85, 186
dioxide 131	Penny 10, 15, 32, 41, 45, 47
oxalate 130	Pepys6 Persoz146
sulphate 128	Persoz 146
sulphide128	Pettersson
Manganoso-manganic oxide 129	Phenylammonium cobalticyanide 172
Marchand, 1, 4, 28, 54, 68, 69, 103, 115, 132, 135, 144, 151, 164, 177	Philippium 246
Marchand and Scheerer 103	Phosphorus 82
Marignac, 9, 11, 12, 14, 15, 16, 18, 19,	and gold 163
22, 23, 25, 40, 42, 44, 57, 59, 60,	pentoxide82
62, 65, 74, 166, 212, 222, 230, 231,	trichloride 83
236, 237, 247, 248	Piccard 90
Mather 156	Pierre 208
Maumené 9, 12, 19, 52, 133	Platinum 249, 270
Mendelejeff's law 96, 150, 180, 246	dichloride 249
Mercury 114, 269 and gold 162	Popp 241 Potassium 9, 270
chloride116, 269	anhydrochromate 123, 187
oxide 115	and tartar emetic 104
selenide 177	aurichloride 162
sulphide 116	bromate 2I
Meyer, Lothar 96, 140	bromide 23
Millon 24, 116, 135	chlorate10
and Commaille 135	and arsenious oxide 187
Mitscherlich and Nitzsch	and potassium nitrate_ 47
Moberg 120 Molybdenum 137	chloride and gold 162 iridium 256
chlorides 140	osmium 254
sulphide 137	palladium 257
trioxide 137	platinum 250
Mosander 208, 229, 236	potassium nitrate_ 46
Mulder and Vlaanderen 205	rhodium 259
	ruthenium 259
N.	silver 18
Mishal	silver chloride 19
Nickel 164 chloride 166, 167	silver nitrate 44
· · · · · · · · · · · · · · · · · · ·	fluoride 70

Potassium fluoxycolumbate 247	; Scheibler 147
iodate24	Schiel
iridichloride 254, 255	Schlippe's salt
nitrate and potassium chlo-	Schneider, 129, 144, 165, 166, 171, 189.
rate 47	199, 200, 203
nitrate and potassium chlo-	Schrötter S2
ride 46	Sefström 114
osmichloride 254 palladichloride 257	Seguin 6 Selenium 176
palladiochloride 257	chloride 176, 178
perchlorate II	dioxide 176, 179
perchlorate 11 platinochloride 250, 252	Seubert 251, 255
rhodiochloride258	Siewert 124
rutheniochloride 259	Silicon 85
sulphate and potassium fluo-	chloride85
ride 70	dioxide 85
and potassium tantalo-	Silver 9
fluoride 248	and aluminum bromide 160
and thoria 214	chloride 157
and zirconia 213	ammonium chloride 40
tantalofluoride 248	antimony 202
tartrantimonite 194	bromide 200
zircofluoride 213 Prout's hypothesis 261	chloride 191 arsenious bromide 187
Purpureo-cobalt chloride 171, 174	chloride 186
1 asparco-coom: cmorac 1/1, 1/4	barium chloride 57
Q.	bismuth chloride 204
₹.	cadmium bromide 113
Quintus Icilius 257	chloride 112
Z	calcium chloride 70
12	cobalt chloride 168
R.	ferric chloride 134
	ferric chloride 134 ferrous chloride 134
Rammelsberg, 142, 151, 224, 229, 247	ferric chloride 134 ferrous chloride 134 lead chloride 74
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51	ferric chloride 134 ferrous chloride 134 lead chloride 74 lithium chloride 89
Rammelsberg, 142, 151, 224, 229, 247 Rawack	ferric chloride 134 ferrous chloride 134 lead chloride 74 lithium chloride 89 magnesium chloride 106
Rammelsberg, 142, 151, 224, 229, 247 Rawack	ferric chloride 134 ferrous chloride 134 lead chloride 74 lithium chloride 89 magnesium chloride 106 manganese chloride 128
Rammelsberg, 142, 151, 224, 229, 247 Rawack	ferric chloride 134 ferrous chloride 134 lead chloride 74 lithium chloride 89 magnesium chloride 106 manganese chloride 128 nickel chloride 168
Rammelsberg, 142, 151, 224, 229, 247 Rawack	ferric chloride 134 ferrous chloride 134 lead chloride 74 lithium chloride 89 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146	ferric chloride 134 ferrous chloride 134 lead chloride 74 lithium chloride 89 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219	ferric chloride 134 ferrous chloride 134 lead chloride 74 lithium chloride 89 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133	ferric chloride 134 ferrous chloride 134 lead chloride 74 lithium chloride 89 magnesium chloride 106 manganese chloride 128 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183	ferric chloride 134 ferrous chloride 74 lead chloride 74 lithium chloride 89 magnesium chloride 106 manganese chloride 168 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rosc 190, 207, 247, 248	ferric chloride 134 ferrous chloride 134 lead chloride 74 lithium chloride 89 magnesium chloride 106 manganese chloride 108 nickel chloride 108 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 25 silver nitrate 41 sodium chloride 32
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoff 164	ferric chloride 134 ferrous chloride 134 lead chloride 89 lithium chloride 89 magnesium chloride 106 manganese chloride 108 nickel chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 64
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoff 164 Rubidium 90	ferric chloride 134 ferrous chloride 134 lead chloride 74 lithium chloride 89 magnesium chloride 106 manganese chloride 168 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 85 silver nitrate 41 sodium chloride 32 strontium chloride 32 strontium chloride 266
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Rosce 146, 183 Rose 190, 207, 247, 248 Rothhoff 164 Rubidium 90 chloride 90 Russell 168, 160	ferric chloride 134 ferrous chloride 134 lead chloride 74 lithium chloride 89 magnesium chloride 106 manganese chloride 168 phosphorus trichloride, 83 potassium bromide 23
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Rosce 146, 183 Rose 190, 207, 247, 248 Rothhoff 164 Rubidium 90 chloride 90 Russell 168, 160	ferric chloride
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoff 164 Rubidium 90 chloride 90	ferric chloride 134 ferrous chloride 134 lead chloride 74 lithium chloride 89 magnesium chloride 106 manganese chloride 108 nickel chloride 108 phosphorus trichloride, 83 potassium bromide 23 chloride 18 iodide 25 silicon chloride 25 silver nitrate 41 sodium chloride 32 strontium chloride 206 tianium chloride 206 tianium chloride 206 vanadium oxychloride, 184 occlusion of oxygen by 262
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoff 164 Rubidium 90 Russell 168, 169 Ruthenium 259	ferric chloride
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Rosce 146, 183 Rose 190, 207, 247, 248 Rothhoff 164 Rubidium 90 chloride 90 Russell 168, 160	ferric chloride
Rammelsberg, 142, 151, 224, 229, 247 Rawack	ferric chloride
Rammelsberg, 142, 151, 224, 229, 247 Rawack 129 Redtenbacher 51 Regnault 6, 39, 150 Reich and Richter 219 Reynolds 96 Rhodium 258 Riche 146 Richter 219 Rivot 133 Roscoe 146, 183 Rose 190, 207, 247, 248 Rothhoff 164 Rubidium 90 chloride 90 Russell 168, 169 Ruthenium 259 Sacc 176	ferric chloride
Rammelsberg, 142, 151, 224, 229, 247 Rawack	ferric chloride
Rammelsberg, 142, 151, 224, 229, 247 Rawack	ferric chloride
Rammelsberg, 142, 151, 224, 229, 247 Rawack	ferric chloride
Rammelsberg, 142, 151, 224, 229, 247 Rawack	ferric chloride
Rammelsberg, 142, 151, 224, 229, 247 Rawack	ferric chloride

~ ''	11		0.31 3.1 . 41 42
Suver	chlorate	14	
	chloride	14	uranate 153
	and ammonium platin-		Sommaruga 170
	chloride	252	Soret's earth X 246
	and antimony chloride,		Stas, 13, 14, 16, 18, 21, 22, 23, 24, 25,
	194, 197		26, 28, 30, 32, 38, 40, 41, 42, 43,
	and barium chloride	60	44, 46, 48, 54, 55, 73, 75, 89
	and boron chloride	84	Stibnite 180 000
			Stibnite 189, 200 Stromeyer 64, 87, 111, 131
	and ceroso-ceric oxide,		Stromeyer 04, 67, 111, 131
	and chromic chloride_		Strontium 64
	and didymium oxide_		chloride and silver 64
	and lithium chloride	87	and strontium sulphate, 66
	and manganese chlo-		Struve 30, 31, 61, 137
	ride	128	Strychnia cobalticyanide 174
	and molybdenum chlo-		nickelocyanide 173
			Sulphur 9, 27
	rides	141	Surphur
	and potassium chlo-		Svanberg,31, 76, 102, 116, 131, 137,
	ride	19	and Nordenfeldt 102
	and potassium platin-		and Norlin 131
	chloride	253	and Struve 31, 137
	and rubidium chloride,		
	and silicon chloride	8 6	T.
	and silver bromide	21	=
	chromate		Tantalum 248, 270
		•	1antalum 240, 2/0
	iodide	27	oxide 248
	nitrate	43	Tartar emetic 194
	sulphide	31	Tellurium 180, 270
	tungstate	149	dioxide 180, 181
	and thallium chloride,		potassium bromide 180, 181
	93, 94		Terbium 246
	and titanium chloride,	207	Terreil157
	dioxide_ 207,		Thalén246
	and vanadium oxychlo-		11 11
	ride		chloride 93
	chromate	119	iodide 94
	iodate	24	nitrate 95
	iodide	25	sulphate 93
	and antimony iodide	100	Thomsen 8
	and silver chloride	27	Thomson 6, 39
	and thallium iodide	94	Thorium 214
	malate	52	acetate215
	nitrate	41	formate 215
	and potassium chloride,	44	oxalate 215, 217
	and silver chloride	43	oxide 214
	oxalate	53	sulphate 214, 215, 217
	permanganate	131	Thulium 246
	racemate	51	Tin 204
	selenite	178	dioxide 205
	sulphate	30	tetrachloride 206
		28	Tissier156
	sulphideshloride		
	and silver chloride	31	Titanium
	tartrate	51	ammoniochloride 210
	tungstate	149	chloride207, 209, 210
Sodio-	uranic acetate 151,	153	dioxide 207, 208
Cadina		21	sulphide 207
Some	n g		
Somu			Troost
Source	chlorate	32	Troost
Soutui	chloratechloride	32 32	Troost
Source	chloridecolumbate	32 32 247	Troost
	chloratechloride	32 32 247 70	Troost

σ.	Winkler 171, 219 Wolf 224
Unger 195	Wrede6
Uranium 150	
acetate 153	X .
oxalate	
oxides 151	X, Soret's earth 246
sulphate 151	
tetrabromide 150	Y.
tetrachloride 150, 151	
	Ytterbium 243, 269
▼.	oxide 243
• •	sulphate243
Vanadium 183	Yttrium 241
oxides 183	oxide 241
oxychloride 184	sulphate 241
Vauquelin6	• • • • • • • • • • • • • • • • • • • •
Vlaanderen 205	Z .
Viamucium	
W.	Zettnow 147
	Zimmermann 150
Wackenroder131	Zinc 108
Wallace 21, 186	oxalate100
Weber 190	oxide 108
Weeren 97	Zirconium 212
Wertheim 153	chloride 212
Weselsky 172	dioxide 212
Wildenstein122	potassium fluoride 212
Wills 181	sulphate212
Wing 226	Zschiesche 233, 238



SMITHSONIAN MISCELLANEOUS COLLECTIONS.

437

CHECK LIST

04

PUBLICATIONS

OF THE

SMITHSONIAN INSTITUTION,

DECEMBER, 1881.



WASHINGTON, D. C.: 1881.



CHECK LIST

01

PUBLICATIONS OF THE SMITHSONIAN INSTITUTION, To December, 1881.

Where no price is affixed the work cannot be furnished.

Publications marked * do not appear in the Contributions, Col ections, or Reports.

1	allas, G. M. Ienry, Jos.	Journal of Regents, Report of Organization Committee Digest of Act of Congress, Address at Laying Corner Stone, Exposition of Bequest, First Report of Secretary, First Report of the Institution, Second Report of Institution, Third Report of Institution, Programme of Organization, Correspondence, Squier & Davis,	8₹0.#	32 8 8 48 38 208 64 4	1846 1847 1847 1847 1848 1847 1848 1849	
C D D E H F G H I J K L M	•	Digest of Act of Congress, Address at Laying Corner Stone, Exposition of Bequest, First Report of Secretary, First Report of the Institution, Second Report of Institution, Third Report of Institution, Programme of Organization,	8vo.* 8vo.* 8vo.* 8vo.* 8vo.* 8vo.* 4vo.*	8 8 48 38 208 64 4	1847 1847 1848 1848 1847 1848 1849	
D D E H F G H I L	•	Address at Laying Corner Stone, Exposition of Bequest, First Report of Secretary, First Report of the Institution, Second Report of Institution, Third Report of Institution, Programme of Organization,	8vo # 8vo.# 8vo.# 8vo.# 8vo.# 4to.#	8 8 48 38 208 64	1847 1848 1847 1848 1849 1847	
E H F G H I J K L	•	Exposition of Bequest, First Report of Secretary, First Report of the Institution, Second Report of Institution, Third Report of Institution, Programme of Organization,	8vo.* 8vo.* 8vo.* 8vo.* 4to.*	8 48 38 208 64 4	1847 1848 1847 1848 1849	
F G H I J K L	Cenry, Jos.	First Report of Secretary, First Report of the Institution, Second Report of Institution, Third Report of Institution, Programme of Organization,	8vo.* 8vo.* 8vo.* 4to.*	48 38 208 64 4	1848 1847 1848 1849 1847	
G H I J K L		First Report of the Institution, Second Report of Institution, Third Report of Institution, Programme of Organization,	8vo.# 8vo.# 8vo.# 4to.#	38 208 64 4	1847 1848 1849 1847	
H I J K L		Second Report of Institution, Third Report of Institution, Programme of Organization,	8vo.# 8vo.# 4to.#	208 64 4	1848 1849 1847	
I J K L		Third Report of Institution, Programme of Organization,	8vo.# 4to.#	64	1849 1847	
J K L		Programme of Organization,	4to.#	4	1847	
K L M		'		_		
L		Correspondence, Squier & Davis,	8vo #	١.,	1045	
3 1			J, U	8	1847	
		First Report of Organization Committee,	8 v o.#	8	1846	
747		Reports of Institution up to Jan. 1849,	8₹0.#	72	1849	
14		Officers, Regents, Act, &c.,	8 v o.#	14	1846	
0		Act to establish Smithsonian Institution,	8 v o.#	8	1846	
P O	wen, R. D.	Hints on Public Architecture,	4to.#	140	1849	10.00
Q		Check List of Periodicals,	4to.#	28	1853	
1 8	Iquier, E. G. and Davis, E. H.	Ancient Monuments of Mississippi Valley,	8. C. 1,	346	1848	
2		Smithsonian Contributions to Knowledge	B. C. 1,	360	1848	

No.	Author.	Title.	Size On Series.	PAGES.	DATE.	PRICE.
3	Walker, S. C.	Researches, Planet Neptune	8. C. 11,	60	1849	
4	Walker, S. C.	Ephemeris of Neptune for 1848,	8. C. 11,	8	1848	
5	Walker, S. C.	Ephemeris of Neptune for 1849,	8. C. 11,	32	1849	
6	Walker, S. C.	Ephemeris of Neptune for 1850,	S. C. 11,	10	1850	
*	Walker, S. C.	Ephemeris of Neptune for 1851,	S. C. 11,	10	1850	
8	Downes, John	Occultations in 1848,	4to.#	12	1848	
9	Downes, John	Occultations in 1849,	4to.#	24	1848	
10	Downes, John	Occultations in 1850,	4to.#	26	1849	
11	Downes, John	Occultations in 1851,	8. C. 11,	26	1850	ı
12	Lieber, Francis	Vocal Sounds of L. Bridgeman,	S. C. 11,	32	1850	
13	Ellet, Charles	Physical Geography of U. S.	S. C. 11,	64	1850	
14	Gibbes, R. W.	Memoir on Mosasaurus,	8. C. 11,	14	1850	
15	Squier, E. G.	Aboriginal Monuments of N. Y.	8. C. 11,	188	1850	
16	Agassiz, Louis	Classification of Insects,	8. C. 11,	28	1850	
17	Hare, Robert	Explosiveness of Nitre,	8. C. 11,	20	1850	
18	Gould, Jr., B. A.	Discovery of Neptune,	8 vo. #	56	1850	
19	Guyot, A.	Directions for Meteorological Observations,	8 v o.#	40	1850	
20	Bailey, J. W.	Microscopic Examination of Soundings,	S. C. 11,	16	1851	
21		Annual Report of Smithsonian Institution for 1849	8 v o.	272	1850	
22	Gray, Asa	Plants Wrightians. Part I,	8. C. 111,	146	1852	
23	Bailey, J. W.	Microscopic Observations in S. Carolina, Georgia, and Florida,	S. C. 11,	48	1851	
24	Walker, S. C	Ephemeris of Neptune, 1852. Appendix I,	S. C. 111,	10	1853	
2 5	Jewett, Chas. C.	Public Libraries of United States,	8vo.	21 0	1851	•
26		Smithsonian Contributions to Knowledge,	S. C. 11,	564	- 1	
27	Booth, J. C. and Morfit, C.	Improvements in Chemical Arts,	M. C. 11,	216	1852	.!
28		Annual Report of Smithsonian Institution for 1850,	8 v o.	326	1851	
29	Downes, John	Occultations in 1852,	S. C. 111,	84	851	

No.	AUTHOR.	Title.	SIZE OR · SERIES.	PAGES.	DATE.	PRICE.
30	Girard, Charles	Fresh-Water Fishes of N. America	8. C. 111,	80	1851	
31	Suyet, A.	Meteorological Tables,		212	1852	
32	Harvey, Wm. H.	Marine Algae of North America. Part I,	8. C. 111,	152	1852	
33	Davis, Chas. H.	Law of Deposit of Flood Tide,	8. C. 111,	14	1852	.75
34	į	Directions for Collecting Speci- mens,	M. C. 11,	40	1859	free
3 5	Locke, John	Observations on Terrestrial Mag- netism,	8. C. 111,	30	1852	
36	Secchi, A.	Researches on Electrical Rheo- metry,	8. C. 111,	60	185 2	
37	Whittlesey, Ch.	Ancient Works in Ohio,	8. C. 111,	20	1851	
38		Smithsonian Contributions to Knowledge,	8. C. 111,	564	1852	
39		Smithsonian Contributions to Knowledge,	8. C. 1v,	426	1852	
40	Riggs, S. R.	Dakota Grammar and Dictionary,	8. C. 1 v ,	414	1852	
41	Leidy, Joseph	Extinct American Ox,	S. C. ▼,	20	1852	
42	Gray, Asa	Plante Wrightians. Part II,	S. C. v,	120	1853	
43	Harvey, Wm. H.	Marine Alges of North America. Part II,	8. C. v,	262	1853	
44	Leidy, Joseph	Flora and Fauna within Living Animals,	S. C. ▼,	68	1858	
45	Wyman, Jostries	Anatomy of Rana Pipiens,	S. C. ▼,	52	1853	
46	Torrey, John	Plants: Fremontianse,	8. C. vi,	24	1853	-
47	Jewett, Chas. C.	Construction of Catalogues of Libraries,	8 v o.#	108	1853	-50
48	Girard, Charles	Bibliographia Americana Ilis- torico Naturalis,	8 vo. #	64	1852	
49	Baird, S. F. and Girard C.	Catalogue of Serpents,	M. C. 11,	188	1853	1.00
50	Stimpson, Wm.	Marine Invertebrata of Gr. Manan	S. C. v1,	68	1853	1.50
51		Annual Report of Smithsonian Institution for 1851,	8 v o.	104	1852	
52	Coffin, Jas. H.	Winds of the Northern Hemisphere,	S. C. v1,	200	1853	
53	Stanley, J. M.	Catalogue of Portraits of Indians,	M. C. 11,	76	1852	
54	Downes, John	Occultations in 1853,	8. C. vi,	86	1853	

No.	AUTHOR.	Title.	Size or Series	PAGES	DATE.	PRICE.
 55	· · · · · · · · · · · · · · · · · · ·	Smithsonian Contributions to Knowledge,	S. C. v,	538	1853	
56	† !	Smithsonian Contributions to Knowledge,	S. C. v1,	484	1854	
57	!	Annual Report of Smithsonian Institution for 1852,	8 ▼ 0.	96	1853	
58	Leidy, Joseph	Ancient Fauna of Nebraska,	8. C. vi,	120	1853	'
59	Chappelsmith, J.	Tornado in Indiana,	8. C. v11,	32	1855	.25
60	Torrey, John	Batis Maritima,	S. C. VI,		1853	
61	Torrey, John	Darlingtonia Californica,	S. C. v1,	8	1853	1
62	Melsheimer, F.E.	Catalogue of Coleoptera,	8vo.#	190	1853	2.06
63	Bailey, J. W.	New Species of Microscopic Organisms,	8. C. v11,	16	1854	
64		List of Foreign Correspondents of Smithsonian Institution,	8vo.#	16	1856	i
65		Registry of Period. Phenomena,	folio,#	4	1854)
66		Annular Eclipse, May 26, 1854	8 vo. •	14	1854	
67		Annual Report of Smithsonian Institution for 1853,	8vo.	310	1854	
68	Mitchell, B. R. & Turner, W. W.	Vocabulary of Jargon of Oregon,	8vo.#	22	1853	
69		List of American Correspondents of Smithsonian Institution,	8 v o.#	16	1853	
70	Lapham, I. A.	Antiquities of Wisconsin,	S. C. v 11,	108	1855	
71	Haven, S. F.	Archmology of the United States,	S. C. v111,	172	1856	
72	Leidy, Joseph	Extinct Sloth Tribe of N. America,	S. C. v11,	70	1855	
73		Publications of Societies in Smith- sonian Library,	S. C. VII,	40	1855	
74		Catalogue of Smithsonian Publications,	M. C. v,	52	1862	
75		Annual Report of Smithsonian Institution for 1854,	8 v o.	464	1855	I
76		Smithsonian Contributions to Knowledge,	S. C. v 11,	262	1855	;
77		Annual Report of Smithsonian Institution for 1855,	8vo.	440	1856	!
78		Smithsonian Contributions to Knowledge,	8. C. v 111,	566	1856	ı

No.	Author.	Title.	Siza or Saries.	PAGES.	DATE.	PRICE.
79	Runkle, John D.	Tables for Planetary Motion,	S. C. 11,	64	1856	1.00
80	Alvord, Benj.	Tangencies of Circles and Spheres,	S. C. v111,	16	1856	1.00
81	Olmsted, D.	Secular Period of Aurora Borealis	S. C. VIII,	52	1856	
82	Jones, Joseph	Investigation on A. Vertebrata,	S. C. v111,	150	1856	1.50
83	Meech, L. W.	Relative Intensity of Heat and Light of the Sun,	S. C. 1x,	58	1856	
84	Force, Peter	Auroral Phenomena in North Latitudes,	8. C. v111,	122	1856	
85		Publications of Societies in Smith- onian Library. Part II,	S. C. VIII,	3 8	1856	
86	Mayer, Brants	Mexican History and Archæology	8. C. 1x,	36	1856	
87	Coffin, Jas. H.	Psychrometrical Tables,	M. C. 1,	20	1856	.25
88	Gibbs, W. and Genth, F. A.	Ammonia Cobalt Bases,	S. C. 1x,	72	1856	1.00
89	Brewer, Th. M.	North American Oology. Part I,	8. C. x1,	140	1857	5.00
90	Hitchcock, E.	Illustrations of Surface Geology,	8. C. 1x,	164	1857	
91		Annual Report of Smithsonian Institution for 1856,	8 v o.	468	1857	
92		Smithsonian Contributions to Knowledge,	S. C. 1x,	482	1857	
93		Meteorological Observations for 1855,	8vo.#	118	1857	
94	Runkle, John D.	Asteroid Supplement to New Tables for $b \frac{(i)}{s}$,	S. C. 1x,	72	1857	.50
95	Harvey, Wm. H.	Marine Algee of North America. Part III,	S. C. x,	142	1858	
96	Harvey, Wm. H.	Marine Algæ of North America. 3 parts complete,	4to.	568	1858	
97	Kane, E. K.	Magnetic Observations in the Arctic Sens,	S. C. x,	72	1859	1.00
98	Bowen, T. J.	Yoruba Grammar and Dictionary,	S. C. x,	232	1858	
99		Smithsonian Contributions to Knowledge,	S. C. x,	462	1858	
100	Gillis, J. M.	Eclipse of the Sun, Sept. 7, 1858,	8. C. x1,	22	1859	.50
101	Hill Thos.	Map of Solar Eclipse, Mar. 15, '58,	8 v o. *	8	1858	free
102	Osten Sacken, R.	Catalogue of Dipters of North America,	M. C. 111,	112	1858	

No.	Author.	Title.	SIZE OR SERIES.	PAGES.	DATE.	PRICE.
103	Caswell, A.	Meteorological Observations, Pro- vidence, R. I.,	S. C. x11,	188	1860	2.00
104	Kane, E. K.	Meteorological Observations in Arctic Seas,	8. C. x1,	120	1859	
105	Baird, S. F.	Catalogue of North American Mammals,	4to.#	22	1857	.50
106	Baird, S. F.	Catalogue of North American Birds,	4to.#	42	1858	.50
107		Annual Report of Smithsonian Institution for 1857,	8 v o.	438) 1858	1
108	Baird, S. F.	Catalogue of N. American Birds,	M. C. 11,	24	1859	.25
109		Annual Report of Smithsonian Institution for 1858,	8₹0.	448	1859	1
110		Annual Report of Smithsonian Institution for 1859,	`8 v o.	450	1860	i I
111	•	Smithsonian Contributions to Knowledge,	S. C. x1,	50	1859	'i
112		Smithsonian Contributions to Knowledge,	8. C. x11,	54	1860	
113	Bache, A. D.	Magnetic and Meteorological Observations at Girard Coll. Pt. I,	8. C. x11,	2	1859	.25
114	Sonntag, A.	Terrestrial Magnetism in Mexico,	S. C. x1,	9:	2 1859	1.25
115		Report on Invention of Electro- Magnetic Telegraph,	М. С. п,	4	1861	free
116	Rhees, Wm. J.	List of Public Libraries, &c.	8 v o.#	84	1859	1
117		Catalogue of Publications, &c., in Smithsonian Library,	M. C. 111,	284	1859	2.00
118	Morris, John G.	Catalogue of Lepidopters of North America,	M. C. 111,	76	1860	1.00
119	Whittlesey, Ch.	Fluctuations of Level in N. A. Lakes,	8. C. x11,	2 8	1860	1.00
120	Hildreth, S. P. and Wood, J.	Meteorological Observations at Marietta, O.,	S. C. xvi,	52	1867	1.00
121	Bache, A. D.	Magnetic and Meteorological Observations at Girard Coll. Pt. II,	8. C. x111,	28	1862	.25
122		Smithsonian Miscellaneous Collections,	M. C. 1,	738	1862	
123	 	Smithsonian Miscellaneous Collections,	M. C. 11,	715	1862	
124		Smithsonian Miscellaneous Collections,	M. C. 111,	772	1862	

No.	AUTHOR.	Title.	SIZE OR SERIES.	PAGES.	DATE.	PRICE.
125		Smithsonian Miscellaneous Collections,	M. C. 1v,	762	1862	•
126	Le Conte, John L.	Colcoptora of Kansas and New Mexico,	S. C. x1,	64	1859	1.25
127	Loomis, E.	Storms in Europe and America, Dec. 1836,	S. C. xı,	2 8	1860	1.25
128	Lea, Carpenter,	Check List of Shells in N. America	M. C. 11,	52	1860	. 2 5
129	Kane, E. K.	Astronomical Observations in the Arctic Sens,	S. C. x11,	56	1860	1.00
130	Kane, E. K.	Tidal Observations in the Arctic Seas,	S. C. x 111,	90	1860	1.50
131	Smith, N. D.	Meteorological Observations in Arkansas from 1840 to 1859,	S. C. x11,	96	1860	1.25
132	Bache, A. D.	Magnetic and Meteorological Observations at Girard Coll. Pt.III	S. C. x 111,	16	1862	.25
133	Morris, John G.	Synopsis of Lepidoptera of North America. Part I,	M. C. 1v,	386	1862	2.00
134	Hagen, H.	Synopsis of Neuropters of North America,	M. C. 1 v ,	368	1861	
135	Mitchell, S. W.	Venom of the Rattlesnake,	S. C. X11,	156	1860	
136	Le Conte, John L.	Classification of Colcoptera of North America. Part I,	M. C. 111,	312	1862	
137		Circular to Officers of Hudson's Bay Co.,	M.C. VIII,	6	1860	
138	Morgan, L. H.	Circular as to Degrees of Relationship,	M. C. 11,	34	1860	
139		Collecting Nests and Eggs of North American Birds,	M. C. 11,	22	1860	free
140	Le Conte, John L.	List of Coleoptera of North America. Part I,	M. C. v1,	82	1866	.75
141	Loew, H. and Osten Sacken	Monographs of Diptera. Part I,	M. C. vi,	24 6	1862	1.50
142	Binney, W. G.	Bibliography of North American Conchology. Part I,	M. C. v,	658	1863	8.00
143	Binney, W. G.	Land and Fresh-Water Shells of North America. Part II,	M. C. v11,	172	1865	1. 2 5
144	Binney, W. G.	Land and Fresh-Water Shells of North America. Part III,	M. C. v11,	128	1865	 1.00
145	Prime, Temple	Monograph of American Corbiculads,	M. C. v11,	99	1865	.75

No.	Author.	Title.	or or Series.	PAGES.	DATE.	PRICE.
146	M'Clintock, Sir F. L.	Meteorological Observations in the Arctic Seas,	S. C. x 111,	164	1862	1.50
147		Annual Report of Smithsonian Institution for 1860,	8 v o.	448	1861	
148		Directions for Meteorological Observations,	M. C. 1,	72	1860	
149		Annual Report of Smithsonian Institution for 1861,	8 v o.	464	1862	
150	:	Annual Report of Smithsonian Institution for 1862,	8vo.	446	1863	
151	1	Smithsonian Contributions to Knowledge,	8. C. x111,	558	1863	
152	Carpenter, P. P.	Lectures on Mollusca,	8vo.*	140	1861	
153	Guyot, A.	Tables, Meteorological and Physical,	M. C. 1,	63 8	1859	
154		List of Foreign Correspondents of Smithsonian Institution,	M. C. v,	56	1862	
155	Whittlesey, Ch.	Ancient Mining on Lake Superior	S. C. xiii,	34	1863	
156	Egleston, T.	Catalogue of Minerals,	M. C. v11,	56	1863	.5
157	Henry, Jos. and Coffin, J. H.	Results of Meteorological Observations from 1854 to 1859,	4to.#	1270	1661	
158	:	Smithsonian Miscellaneous Collections,	M. C. v,	774	1864	
159	Mitchell, S. W. & Morehouse, G. R.	Anatomy and Physiology of Respiration in Chelonia,	S. C. XIII,	50	1863	1.0
160	Gibbs, G.	Instructions for Ethnology and Philology,	M. C. v11,	56	1863	
161	Gibbs, G.	Dictionary of the Chinook Jargon	M. C. v11,	60	1863	.5
162	Bache, A. D.	Magnetic and Meteorological Obs. at Girard Coll. Pt. IV, V,& VI,	8. C. x 111,	78	1862	1.0
163	i	Circular on History of Grass- hoppers,	M. C. 11,	4	1860	fre
164	1	Smithsonian Museum Miscellanea	M.C. v111,	88	1962	.5
165	Allen, H.	Monograph of the Bats of North	M. C. v11,	1	1864	
166	Bache, A. D.	Magnetic Survey of Pennsylvania	S. C. XIII,	88	1863	' 1.0
167	Le Conte, Jno. L.	New Species of North America Coleoptera. Part I,	M. C. v1,	180	1866	j.0
168		Circular Relative to Birds from Middle and South America,	 M.C. viii,	١.	1863	fre

No.	Author.	Title.	SIZE OR SERIES.	PAGES.	DATE.	PRICE.
169		Smithsonian Miscellaneous Collections,	M. C. v1,	888	1867	
170		Comparative Vocabulary,	4to.#	2 0	1863	free
171	Loew, H.	Monograph of the Diptera of North America. Part II,	M. C. v1,	372	1864	
172	Meek, F. B. and Hayden, F. V.	Palmontology of the Upper Missouri. Part I,	S. C. xiv,	168	1865	
173	Dean, John	Gray Substance of the Medulla Oblongata,	S. C. xvi,	80	1864	2.50
174	Binney, W. G.	Bibliography of North American Conchology. Part II,	M. C. 1x,	302	1864	2.00
175	Bache, A. D.	Mag. and Met. Observ. at Girard Coll. Parts VII, VIII, & IX,	S. C. x1v,	72	1864	1.00
176		Circular, Collecting North American Shells,	м. с. п,	4	1860	free
177	Meek, F. B.	Check List of Invertebrate Fos- sils of North America,	M. C. v11,	42	1864	.25
178		Circular to Entomologists,	M.C. v111,	2	1860	free
179		Catalogue of Publications of So- cieties,	M. C. 1x,	596	1866	
180	Draper, H.	Construction of a Silvered Glass Telescope,	S. C. xiv,	60	1864	1.00
181	Baird, S. F.	Review of American Birds in Smithsonian Museum. Part I,	М. С. хп,	484	1866	2.00
182		Results of Meteorological Observations from 1854-1859. Vol. II,	4to.#	546	1864	2.50
183	Meek, F. B.	Check List of Invertebrate Fos- sils of North America,	M. C. VII,	34	1864	.25
184		Smithsonian Contributions to Knowledge,	S. C. xiv,	490	1865	
185		List of Birds in Mexico, &c.,	8vo.*	8	1863	free
186	Bache, A. D.	Mng. and Met. Observ. at Girard College. Parts X, XI, & XII,	S. C. xiv,	42	1865	.50
187		Annual Report of Smithsonian Institution for 1863,	8 v o.	420	1864	
188		Annual Report of Smithsonian Institution for 1864,	8vo.	450	1865	
189	Scudder, S. H.	Catalogue of Orthoptera of North America,	M C. viii,	110	1868	1.00
190	Henry, Joseph	Queries Relative to Tornadoes,	M. C. x,	4	1865	free

No.	Author.	TITLE.	Size or Series.	PAGES	DATE.	PRICE.
191		Smithsonian Miscellaneous Collections,	M. C. v11,	878	1867	
192	Leidy, Joseph	Cretaceous Reptiles of the U. S.,	S. C. xiv,	142	1865	
193		Duplicate Shells from Expedition of Capt. Wilkes,	8 v o.#	4	1865	free
194	Binney, W. G & Bland, T.	Land and Fresh-Water Shells of North America. Part I,	M.C. v111,	328	1869	
195	Bache, A. D.	Girard College Observations. Complete. Parts I to XII,	s. c.	262	1865	
196	Hayes, I. I.	Physical Observations in the Arctic Seas,	S. C. xv,	286	1867	
197	Whittlesey, Ch.	Glacial Drift of Northwestern States,	S. C. xv,	3 8	1866	
198	Kane, E. K.	Physical Observations in the Arctic Seas. Complete,	8. C.	340	1860	
199	Newcomb, S.	Orbit of Neptune,	S. C. xv,	116	1866	
200	Conrad, T. A.	Check List of the Invertebrate Fossils of North America,	M. C. v11,	46	1866	.2
20 1	Stimpson, Wm.	Hydrobiinss and Allied Forms,	M. C. v11,	64	1865	.5
202	Pumpelly, R.	Geological Researches in China, Mongolia, &c.	S. C. xv,	173	1866	3.5
2 03		List of Works published by Smithsonian Institution,	M. C. vii.	12	1866	
204	Cleaveland, P.	Meteorological Observations, Brunswick, Me., 1807-1859,	S. C. xvi,	- 60	1867	1.0
2 05		Circular for Archmology and Eth- nology,	M.C. v111,	2	1867	free
206		Smithsonian Contributions to Knowledge,	8. C. xv,	620	1867	
207		Relative to Scientific Investiga- tions in Russian America,	M.C. v111,	10	1867	
208	Pickering, Chas.	Gliddon Mummy Case in Smithsonian Institution,	S. C. xv1,	6	1867	۱ ا
209		Annual Report of the Smithsonian Institution for 1865,	8 v o.	496	1866	
210		Arrangement of Families of Birds in Smithsonian Institution,	M.C. v111,	8	1866	
211		Smithsonian Contributions to Knowledge,	S. C. xv i,	498	1870	

_						
No.	AUTHOR.	TITLE.	Size OR Series.	PAGES.	DATE.	PRICE.
212		Smithsonian Miscellaneous Collections,	M.C. v111,	921	1869	
213		Smithsonian Miscellaneous Collections,	M. C. 1x,	918	1869	
214		Annual Report of Smithsonian Institution for 1866,	8 v o.	470	1867	
215		Annual Report of Smithsonian Institution for 1867,	8₹0.	506	1868	
216		List of Photograph Portraits of North American Indians,	M. C. xiv,	42	1867	
217	Hoek, M.	Meteoric Shower, 1867, Nov. 13,	8vo.#	4	1867	
218	Morgan, L. H.	Systems of Consanguinity and Affinity,	S. C. xv 11,	616	1869	
219	Osten Sacken, B.	Monograph of Diptera of North America. Part IV,	M .C. v111,	3 58	1869	2.00
2 20	Swan, Jas. G.	Indians of Cape Flattery,	S. C. xv1,	118	1870	
221	Coffin, James H.	Orbit, &c., of Meteoric Fire Ball, July 20, 1860,	S. C. xv1,	56	1869	1.00
222	Schott, Chas. A.	Tables of Rain and Snow in United States,	S.C.xviii,	178	1872	8.00
223	Gould, B. A.	On the Transatlantic Longitude,	S. C. xvi,	110	1869	1.00
224		Annual Report of Smithsonian Institution for 1868,	8 v o.#	473	1869	
225		List of Foreign Correspondents of Smithsonian Institution,	8 v o.#	56	1870	
226		List of Publications of Smithson- ian Institution,	8 v o.	34	1869	
227	Gill, Theod.	Families of Mollusks,	M. C. x,	65	1871	.25
228		Annual Report of Smithsonian Institution for 1869,	8vo.	43 0	1871	
229		Smithsonian Contributions to Knowledge,	S. C. xv11,	616	1871	
2 30	Gill, Theod.	List of Families of Mammals,	M. C. x1,	104	1872	.25
231	Baird, S. F.	Memoranda Food Fishes,	M. C. x,	8	1871	
232	Stockwell, J. N.	Secular Variations of Orbits of Planets,	8. C. xv iii,	220	1872	2.00
233	Perrel, Wm.	Converging Series, Ratio of Dia- meter, and Circum. of Circles,	S.C.xv111,	6	1871	.25
234	Baird, S. F.	Circular Relative to Food Fishes,	M. C. x,	14	1871	free

No.	Auteor.	Title.	SIZE OR SERIES.	PAGES.	DATR.	PRIOR.
235	Henry, Joseph	Circular Relative to Thunder- storms,	M. C. x,	,	1871	free
236	Henry, Joseph	Circular Relative to Heights,	M. C. x,	2	1871	free
2 37	Henry, Joseph	Circular Relative to Lightning- rods,	M. C. z,	4	1871	free
238	Rhees, Wm. J.	List of American Libraries, and Public Institutions,	M. C. x,	256	1872	free
239	Harkness, Wm.	Magnetic Observations on the Monadnock,	8. C. xvni,	226	1872	2.00
240	Barnard, J. G.	Problems of Rotary Motion,	8. C. xix,	56	1872	
241	Wood, H. C.	Fresh-Water Algæ of N. America,	8. C. x1x,	272	1872	7.50
242	Clark, H. J.	Lucernariss.	8. C. xx11,	138	1878	7.54
243		List of Foreign Correspondents of Smithsonian Institution,	M. C. x,	68	1872	
244		Annual Report of Smithsonian Institution for 1870,	870.	494	1871	
245		Check List of Smithsonian Publications to July, 1872,	M. C. z,	21	1872	
246		Smithsonian Contributions to Knowledge,	8. C. xvm,	646	1872	
247	Gill, Theod.	List of Families of Fishes,	M. C. x1,	96	1872	
248	Hilgard, E. W.	Geology of Lower Louisiana,	S. C. xx111,	38	1872	
249		Annual Report of Smithsonian Institution for 1871,	8 v o.	473	1873	
250		Smithsonian Miscellaneous Col- lections,	M. C. x,	***	1873	
251	Rau, Charles	Memoir of Von Martius,	R. 1869,		1871	
252	Carpenter, P. P.	American Mollusca,	M. C. x,	l	1872	
253	Tryon, G. W.	Mon. of Strepomatids. Land and Fresh Water Shells. Part IV,	M. C. xvi,	i i	1873	. 64
254	De Saussure, H.	Monograph of Hymenoptera,	M. C. x\∴,	1	1875	2.00
2 55	Clarke, F. W.	Specific Gravity Tables. Part I,	M. C. x11,	1	1873	
256	Loew, H.	Monograph Diptera. Part III,	M. C. x1,	l i	1873	201
257	Baird, S. F.	Systematic List of For. Corresp.	M. C. x,		1872	- 05
258	Watson, S.	Botanical Index,	M. C. xv,		1878	
259	Jones, Jos.	Antiquities of Tennessee,	S. C. xx11,	181	1876	3.04
260		Regulations of Smith. Inst'n,		42	1872	

No.	Auteor.	Title.	Size or Series.	PAGES.	DATE.	PRICE.
26 1	Packard, A. S.	Directions for Collecting and Pre- serving Insects,	M. C. x1,	60	1873	free
262	Newcomb, S.	Orbit of Uranus,	8. C. x1x,	296	1873	3.00
263		Astronomical Telegram Circular,	M. C. x11,	4	1873	
264	LeConte, J. L.	New Species Coleopters. Part II,	M. C. x1,	74	1873	
26 5	LeConte, J. L.	Classification Coleoptera. Part II,	M. C. x1,	72	1873	
266	Woodward, J. J.	Toner Lecture I. Cancerous Tumors,	M. C. xv,	44	1873	.25
267	Swan, J. G.	Haidah Indians,	S. C. xx1,	22	1874	1.00
26 8	Coffin, J. H.	Winds of the Globe,	S. C. xx,	781	1875	
269	Rabel, Simeon,	Sculptures of Santa Lucia Cosumal- whuspa in Guatemala.	8. C. xx11,	94	1878	
270	Osten Sacken, C. B.	Catalogue of Diptera of North Am.,	M. C. xvi,	324	1878	2.00
271		Annual Report of Smithsonian Institution for 1872,	8 v o.	456	1873	
272	 - 	Smithsonian Contributions,	S. C. x1x,	660	1874	
273		Miscellaneous Collections,	M. C. x1,	796	1874	
274		Miscellaneous Collections,	M. C. x11,	767	1874	
2 75	<u> </u>	Annual Report of Smithsonian Institution for 1873,	8 v o.	452	1874	
276	Clarke, F. W.	Specific Heat Tables. Part II,	M. C. xiv,	58	1876	.50
277	Schott, C. A.	Temperature Tables,	S. C. xxi,	360	1876	3.00
2 78		Check List of Smithsonian Publications,	м. с.	24	1874	
279	DaCosta, J. M.	Toner Lecture III. The Heart,	M. C. xv,	32	1874	. 25
2 80	Alexander, S.	Harmonies of Solar System,	S. C. xx1,	104	1875	1.00
281	Newcomb, S.	Planetary Motion,	S. C. xx1,	40	1874	
282	Wood, H. C.	Toner Lecture IV. Study of Fever,	M. C. xv,	50	1875	.25
283	Gill, Theod.	Catalogue of Fishes,	M. C. xiv,	56	1875	.25
284		Smithsonian Contributions,	S. C. xx,	794	1876	
285		Smithsonian Contributions,	S. C. xxi,	543	1876	
286		Annual Report of Smithsonian Institution for 1874,	8₹0.	416	1875	
287	Rau, Charles,	Archmological Collection, Nat. Museum,	8. C. xx11,	118	1876	2.00

No.	Author.	Title.	Siza or Saries.	PAGES.	DATE.	Paios.
288	Clarke, F. W.	Specific Gravity Tables. Supp. I,	M. C. xiv,	62	1876	,50
289	Clarke, F. W.	Tables, Expan. by Heat. Pt. III.	M. C. xiv,	58	1876	.50
290		List of Smithsonian Publications,	8vo. x111,	12	1876	
291	Brown-Séquard,	Toner Lecture II. The Brain,	M. C. xv,	26	1877	.25
2 92	Cope, E. D.	Batrachia. Bulletin National Museum, No. 1,	 M. C. x111,	106	1875	
29 3	Kidder, J. L., Coues, E.	Birds Kerguelen Island. Bulle- tin National Museum, No. 2,	М. С. хпі,	61	1875	
294	Kidder, J. L., and others,	Nat. Hist. Kerguelen Island. Bulletin Nat. Museum, No. 3,	 M. C. x1111,	122	 1876 	•
295	Lawrence, G. N.	Birds of Mexico. Bulletin Nat. Museum, No. 4,	M. C. x111,	56	1875	
296	Goode, G. B.	Fishes of Bermuda. Bull. Nat. Museum, No. 5,	M. C. x111,	82	 1876 	
297	Goode, G. B.	Classification of Animal Resources, etc. Bulletin Nat. Museum, No. 6,	M. C. xiii,	139	1876	
29 8		Annual Report of Smithsonian Institution for 1875,		422	1876	1
299		Annual Report of Smithsonian Institution for 1876,	8 v o.	488	1377	1
300	Keen, W. W.	Toner Lecture V. Continued Fevers,	M. C. xv,	72	 1877 	.2
301		Check List of Smithsonian Publications to July, 1877,	M. C. xiv,	72	1877	ı
B02	Adams, Wm.	Toner Lecture VI. Subcutaneous Surgery,	M. C. xv.	30	 1877 	.25
803	Streets, Thos. H.	Natural History of Hawaiian and Fanning Islands and Lower California. Bulletin National Museum, No. 7,	M. C. xiii,	172	1877	
304	Dall, Wm. H.	Index to names applied to Bra- chiopoda. Bulletin Nat. Mus., No. 8,	M. C. xiii,	88	1877	
305	Jordan, David S.	North Amer. Ichthyology, No. 1. Review of Rafinesque's N. Am. Fishes. Bulletin of Nat. Mus., No. 9,	M. C. xm,	53	1877	
306	Jordan, David S.	North Amer. Ichthyology, No. 2. Notes on Cottidae, &c. Bull. Nat. Mus., No. 10.	M. C. x111,	120	1877	
307	Baird, S. F.	Report on Centennial Exhibition of 1876,	R. 1876,	22	1877	

No.	Author.	Title.	SIZE OR SERIES.	PAGES.	DATE.	PRICE.
30 8 .	Jordan, D. S., and Brayton, A. W.	North Amer. Ichthyology, No. 3 Distribution of Fishes of S. C Ga., and Tenn. Bulletin Nat Museum, No. 12,	м. с.	237	1878	
309		List of Foreign Correspondents of the Smithsonian Institution to Jan. 1878,	M. C. xv,	120	1878	
310	Barnard, J. G.	Internal Structure of the Earth,	S. C. XXIII,	19	1877	.25
311	Holden, Edw. S.	Index Catalogue of Books Relat- ing to Nebulæ,	M. C. xiv,	126	1877	.50
312	,	Smithsonian Miscellaneous Coll. (Bull. N. M., 1-10.)	M. C. x111,	982	1878	
3 13	Eggers, Baron,	Flora of St. Croix and Virgin Islands. Bull. Nat. Mus., No. 13,	м. с.	186	1879	
314		Smithsonian Miscellaneous Collections.	M. C. x1 v ,	911	1878	
315		Smithsonian Miscellaneous Collections.	M. C. xv,	880	1878	
316		Circular in Reference to American Archæology,	M. C. xv,	15	1878	free
317	Elliot, D. G.	Classification and Synopsis of Trochilidse,	8. C. xxiii,	289	1879	3 .0 0
318	Dall, Wm. H.	Remains of Man from Caves in Alcutian Islands.	S. C. xx11,	44	1878	2.00
3 19	Baird, S. F.	Circular. Inquiries Relative to Crawfish and Crustacea,	M. C. xv,	8	1878	free
320	Baird, S. F.	Circular Kelating to Collections of Living Reptiles,	M. C. xv,	2	1878	free
32 1	Shakespeare, E. O.	Toner Lecture, VII. Inflamma- tion in Arteries atter Ligature, etc.	M. C. xvi,	74	1879	-25
322		Smithsonian Miscellaneous Collections.	M. C. xv1,	950	1880	
323		Annual Report of the Smithsonian Institution for 1877,	8 vo.	50 0	1878	
324		Circular relative to Scientific and Literary Exchanges,	M. C. MVI,	2	1879	free
325		Circular. Business Arrangements of the Smithsonian Institution,	M. C. xvi,	7	1879	free
326	Goode, G. Brown,	Catalogue of Collection of Animal Resources and Fisheries of U. S. Bulletin Nat. Museum, No. 14,		367	1879	

No.	Author.	TITLE.	SIZE OR SERIES.	PAGES.	DATE.	Putch.
327	Smithson, James,	Scientific writings of,	M. C. xx1,	166	1879	.75
32 8	Rhees, Wm. J.	Smithsonian Institution. Docu- ments Relative to its Origin and History,	M. C. x v n,	1027	1879	2.50
329	Rhoes, Wm. J.	Smithsonian Institution. Jour- nals of Board of Regents and Reports of Committees,	M.C.xviii,	851	1879	2 00
330	Rhees, Wm. J.	Smithson and his Bequest,	M. C. xxi,	76	1879	.75
3 31	Rau, Charles,	The Palenque Tablet,	8. C. xx11,	90	1879	2.00
332	1	Proceedings of the Nat. Museum for 1878. Vol I,	! ¡M. C. xix,	524	1879	1.00
333		Proceedings of the Nat. Museum for 1879. Vol. II,	M. C. xix,	504	1880	1.00
334	Elliot, D. G.	List of Described Species of Hum- ming-Birds,	M. C. xvi,	22	1879	.23
335	Rhees, Wm. J.	List of Principal American Libra- ries, Museums, Societies, etc.,		, 6	1879	free
336		Smithsonian Miscellaneous Collections,	M C. xvii,	1034	1880	
337	!	Smithsonian Miscellaneous Collections,	M.C. xviii,	851	1880	
33 8	Welling, J. C.	Life and Character of Joseph Henry,	M. C. xxi,	30	1880	
339	Taylor, Wm. B.	The Scientific Work of Joseph Henry,	M. C. xxi,	225	1880	
340	!	Smithsonian Contributions to Knowledge,	! 8. C. xx11,	544	1880	
\$4 1	i	Annual Report of Smithsonian Institution for 1878,	8 v o.	575	1879	
342	Kumlien, L.	Contributions to Natural History of Arctic America. Bulletin of the National Museum, No. 15,		179	1879	
343	Henry, Joseph,	Annual Reports of Sec. of S. I., 1865-77,	 R .	548	1880	free
344		Check List of Smithsonian Publications,	M. C. xvi,	16	1879	free
345		Annual Report of Smithsonian Institution for 1879,	¦ 8 ∀ 0.	631	1850	
846		Smithsonian Contributions to Knowledge,	! 8. C. xx111,		1881	•••
347		Cloud Charts,	•	2	1851	lica

No.	AUTHOR.	Title.	SIZE OR SERIES.	PAGES	DATE.	PRIOR.
348	Baird, S. F.	Fishes of New Jersey Coast,	R. 1854,	40	1855	
349	Waring, G. E.	Toner Lecture VIII. Sanitary Drainage of Washington.	M.C.	26	1880	.25
350		Map of Stars. (For Aurora obs.)	*	1	1856	free
351	Cooper, J. G.	Forests of North America,	R. 1858,	36	1859	
352	Whitney, W. D.	Lectures on Linguistics,	R. 1863,	22	1864	free
353	Schott, C. A.	Tables of Rain Fall, (2d edition).	s. c.	269	1881	3.00
354	Delaunay, M.	Essay on Velocity of Light,	R. 1864,	31	1864	free
355	Wetherill, C. M.	Ozone and Antozone,	R. 1864,	12	1864	
356		Memorial of Joseph Henry,	M. C. xxi,	532	1880	free
357	Wood, H. C.	Researches on Fever,	S. C. XXIII	2 63	1878	2.50
358	Becker, G. J.	Atomic Weights. Constants of Nature. Part IV,	M. C.	152	1880	1.00
359	ı	Planisphere of the Heavens,	*	1	1864	free
360	Descr, E.	Palafittes, or Lacustrian Con- structions,	R. 1865,	53	1865	
361	Baegert, J.	Aborigines of California,	R. 1863-4,	41	1865	
362	Rau, Charles,	Artificial Shell Deposits in New Jersey,	R. 1864,	6	1865	
363	Lewis, James,	Instructions for collecting Land and Fresh Water Shells,	R. 1866,	8	18 6 0	free
364	Lilljeborg, W.	Classification of Birds,	R 1865,	16	1866	free
36 5	Ross, Hardisty, Jones, Gibbs,	Tinneh or Chepewyan Indians,	R. 1866,	25	1856	
366	Edwards, A. M.	Directions for collecting Diato- macea,	.*	7	1867	free
367	Rothrock, J. T.	Flora of Alaska,	R. 1867,	33	1867	free
36 8	Rau, Charles,	Indian Pottery,	R. 1866,	11	1867	
369	Abbe, C.	Dorpat and Poulkova,	R. 1867,	23	1867	!
370	Rau, Charles,	Flint Implements in Illinois,	R. 1858,	9	1868	
371	Newton, H. A.	Metric Tables,	R. 1865,	23	1868	free
372	Rau, Charles,	Drilling in Stone without Metal,	R. 1868,	11	1869	
373	1	Meteorological Stations and Observers,	R. 1868,	42	1869	free
374	Schott, C. A.	Rain Charts for Summer, Winter, and Year,	8. C. xviii,	3	1870	free

No.	Author.	TITLE.	Size or Series.	PAGES. DATE.	Paics.
3 75	Taylor, W. B.	Origin and Nature of Force,	R. 1870,	19 1870	free
876	Hunt, T. Sterry	Chemistry of the Earth,	R. 1869,	26 187	free
377	Babinet, M.	Diamond and Precious Stones,	R. 1870,	33 187	free
378	Roehrig, F. L. O.	Dakota Language,	R. 1871,	19 187	free
379	Henry, Joseph,	Eulogy on Alex. D. Bache,	R. 1870,	28 187	free
380	Peabody, A. P.	Scientific Education of Mechanics,	R. 18 72 ,	13 187	3 free
381	Schott, C. A.	Temperature Chart of U. S. for Year, etc.,	•	1 187	3 free
382	Rau, Charles,	North American Stone Implements,	R. 1872,	16 187	3
8 83	Bransford, J. F.	Archmological Researches in Nicaragua,	s. c.	100 188	
384	Baird, S. F.	Circular for Shipping Fresh Fish,	M. C.	4 188	free
385	Rau, Charles,	Ancient Aboriginal Trade in North America,	R. 1872,	49 187	3
386	Brezina, A.	Crystallography,	R. 1872,	36 187	4 free
387	Schott, C. A.	Temperature Charts, Summer Winter, Year,	8. C. xxi,	3 187	4 free
388	Schott, C. A.	Temperature Chart—Year,	' # 	1	4 free
389	Henry, Joseph,	Investigation of Illuminants,	R. 1880,	25 188	1 free
890	Hilgard, J. E.	Tides and Tidal Action in Har- bors,	R. 1874,	22 187	
391		Act to Establish Smithsonian Institution,	M.C.xviii,	10 187	5 free
392	Romer, F. F.	Prehistoric Antiquities of Hun- gary,	R. 1876,	9 187	7
393	Gillman, Henry,	Mound-builders. Ancient Man in Michigan,	R. 1875,	13 187	
194	Abbott, C. C.	Stone-age in New Jersey,	R. 1875,	136 187	
39 5	Taylor, Wm. B.	Kinetic Theories of Gravitation,	R. 1876,	80 187	7 free
396	McParlin, T. A.	History and Climate of New Mexico,	R. 1877,	30 187	7 free
397	Mason, Otis T.	Latimer Collection of Antiqui- ties,	R. 1876,	23 187	7 free
39 8	Abbe C.	Meteorological Memoirs (Translations),	R. 1877,	104 187	8 free

No.	AUTHOR.	Title.	Size OR Series.	Ранв.	DATE.	PRICE.
399	Holmgren, F.	Color-blindness,	R. 1877,	72	1878	free
400	Jones, Chas. C.	Aboriginal Structures in Ga.,	R. 1877,	13	1878	free
401	Weismann, A.	Change of Mexican Axolotl to Amblystoma,	R. 1877,	29	1878	free
402	Rau, Charles,	Stock in - trade of Aboriginal Lapidary,	R. 1877,	9	1878	free
403	Rau, Charles,	Gold Ornament from Florida,	R. 1877,	6	187 8	free
404	Haldeman, S. S.	Polychrome Bead from Fiorida,	R. 1877,	6	1878	free
405	Taylor, Wm. B.	Henry and the Telegraph,	R. 1878,	103	1879	free
406	Henry, Joseph,	Researches in Sound,	R. 1878,	106	1879	free
407	Gray, Asa,	Memoir of Joseph Henry,	R. 1878,	35	1879	free
408	Baird, S. F.	Report of Secretary of Smith- sonian Institution for 1878,	R. 1878,	60	1879	free
409	Sherman, W. T. Parker, P. and Baird, S. F.	Report of Museum Building Commission and of the Archi tects for 1879,	R. 1879,	18	1880	free
410	Holden, E. S.	Reports of Astronomical Observatories for 1879,	R. 1879,	60	1880	free
411	Lautenbach, B. F.	Irritation of Polarised Nerve,	R. 1878,	59	1880	free
412	Cope, Edw. D	Zoological Position of Texas. Bull. Nat. Mus., No. 17.	M C.	51	1880	
413	Goode, G. B.	Berlin Fisherv Exhibit. Bull. Nat. Mus., No. 18.	M. C.	278	1880	
4 i 4	Schott, C. A.	Base-Chart of the U. S.	*	1	1880	.15
+15	Knight, Edw. H.	Savage Weapons,	R. 1879,	90	1880	free
4!8		Smithsonian Miscellaneous Collections,	M. C. xix,	1034	1880	
417	Mayer, Alfred M.	Henry as a Discoverer,	M. C. xxi,	36	1880	free
418	Baird, S. F.	Report of Secretary of Smith- sonian Institution for 1879,	R. 1879,	76	1880	free
419	Baird, S. F.	Report of Secretary of Smith- sonian Institution for 1880,	R. 1880,	88	1881	free
	Mason, Otis T.	Anthropological Investigations in 1879,	R. 1879,	30	1881	free
	Boehmer, G. H.	Index to Smithsonian Anthropological Articles,	R. 1879,	10	1881	free
422	Ridgway, Robt.	Nomenclature of N. American Birds Bull. Nat. Mus., No. 21,	M. C.	94	1881	.25

No.	Author.	TITLE.	Size OR Series.	PAGES.	DATR.	Paten.
42 3		Smithsonian Miscellaneous Collections,	M. C. xx,	846	1881	
424		Smithsonian Miscellaneous Collections,	M. C. xx:,	773	1881	
425		Proceedings of National Museum for 1880. Vol. III,	M. C.	594	1881	
426	Holden, E. S. and Hastings, C. S.	Synopsis of Herschel's Writings,	R. 1880,	118	1881	free
427	Holden, E. S.	Recent Progress in Astronomy,	R. 1880,	39	1 8 81	free
42 8	Hawes, G. W.	Recent Progress in Geology and Mineralogy,	R. 1880,	30	1881	free
129	Barker, G. F.	Recent Progress in Physics and Chemistry,	R. 1880,	65	1881	ſree
43 0	Farlow, W. G.	Recent Progress in Botany,	R. 1880,	19	1881	free
431	Gill, Theodore,	Recent Progress in Zoology,	R. 1880,	62	1881	free
432	Mason, Otis T.	Recent Progress in Anthro-	R. 1680,	51	1881	free
433	Mason, Otis T.	Visit to Luray Cavern, Va.	R. 1880,	12	1881	free
434	Sherman, W. T. Parker, P. and Baird, S. F.	Report of National Museum Building Commission and of the Architects for 1880,			1881	free
435	Loud, F. H.	Discussion of Snell's Barometric Observations,	R. 1880,	23	1881	free
436		List of Periodicals received by Smithsonian Institution,	R. 1880,	9	1881	free
437		Check List of Publications of Smithsonian Institution,	м. с.	22	1881	free
43 8	Holden E. S. and Boehmer, G. H.	Reports of Astronomical Observatories for 1880,	R. 1880,	128	1881	free

WORKS IN PREPARATION.

Monograph of Chitonide. By P. P. CARPENTER.

Bibliography of the Fishes of Western North America. By Theodore Glll. Bulletin of the National Museum, No. 11.

Synopsis of the Fishes of the United States. By David S. Jordan. Bulletin of the National Museum, No. 16.

An Index of Names used for Zoological Genera, comprising 70,000 titles. By Samuel H. Scudder. 8vo. Bulletin of the National Museum, No. 19.

The Writings of American Zoologists. Index Bibliography, No. I. Publications of Spencer Fullerton Baird. By G. Brown Goode. 8vo. Bulletin of the National Museum, No. 20.

Flora of the District of Columbia. By LESTER F. WARD. Bulletin of the National Museum, No. 22.

Collector's Manual of Marine Zoology. By RICHARD RATHBUN. Bulletin of the National Museum, No. 23.

Tables showing the amount of Precipitation of Rain and Snow for each Month and Year at upwards of 2000 stations in the United States. By Charles A. Schott and E. H. Courtenay.

Regulations for the Telegraphic Announcement of Astronomical Discoveries. By S. F. BAIRD.

The Constants of Nature. Part V. A Revision of the Atomic Weights. By Frank W. Clarke.

List and Index of Publications of the Smithsonian Institution from 1846 to 1881. By Wm. J. Rhees.

List of Foreign Correspondents. By GEO. H. BOEHMER.

• • . • • ·



• •

SMITHSONIAN MISCELLANEOUS COLLECTIONS.

CATALOGUE OF PUBLICATIONS

OF THE

SMITHSONIAN INSTITUTION,

(1846 - 1882,)

WITH AN

ALPHABETICAL INDEX OF ARTICLES

IN THE

SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE, MISCELLANEOUS COLLECTIONS, ANNUAL REPORTS, BULLETINS AND PROCEEDINGS
OF THE U. S. NATIONAL MUSEUM, AND REPORT
OF THE BUREAU OF ETHNOLOGY.

BY WILLIAM J. RHEES, CHIEF CLERK OF THE INSTITUTION.

WASHINGTON: SMITHSONIAN INSTITUTION. 1882.

JUDD & DETWEILER, PRINTERS, WASHINGTON, D. C.

CONTENTS.

									Page.
1.	Smithsonian Contributions			•	•	•	•		7
2.	Miscellaneous Collections		٠.	•	•	•	•	•	VI
3.	Annual Reports								VΠ
4.	Bulletins of the U.S. National	Μι	ıseum				•		VIII
5.	Proceedings of the U.S. Nation	nal	Muse	um				•	IX
6.	Report of Bureau of Ethnology	7			•				IX
7.	Copyright	•	•						IX
8.	Use of illustrations								x
9.	Size of editions								x
10.	Distribution of publications			-	_				ХI
11.	Rules of distribution .			Ī			Ĭ	•	XII
12.	Form of application for publication	Itio	ns .		•	·	·		XIII
13.	Price list of publications .	-010		•	•	•	•	•	XIV
	of Publications, A to Q	•	•	•	•	•	•	•	1
		•	1 45	· ′Q	•	•	•	• •	3
	of Publications, Regular Sessified List of Separate Pu				•	•	•	•	91
ULA	Anatomy, Physiology, Medici				•	•	•	•	91 92
	Anthropology	цо,	anu c	urgei	y	•	•	•	92
	Architecture	•	:	•	:	•	•	•	94
	Astronomy			•	•	•	•		94
	Bibliography								95
	Biography								95
	Biology								96
	Botany		•		• .		•		96
	Chemistry and Technology		•				•		97
	Electricity and Magnetism			•		•			97
	Geology		•	•					98
	Mathematics			•	•				98
	Meteorology				•			•	99
	Microscopy		•			•			100
	Mineralogy				•				100
	Miscellaneous		•	•	•				100
	Palæontology		•	•				•	102
	Philology								102
							(1	тт)	

	•									Page.
CLASSIFIED LIST	of Sepai	RATE I	UBL	ICATI(ONS-	-Conti	nued.			
Physical Ge	ography					•				103
Physics						•		•		103
Terrestrial I	Physics	•		•		•				104
Zoölogy—B	irds .	•		•						104
\mathbf{F}	ishes .					•				105
Ir	nsects .					•		•		105
M	ammals					•		•		106
M	[ollusks		٠.			•				106
R	adiates							•		107
R	eptiles		·					•		107
SMITHBONIAN CON	TRIBUTIO	NS TO	Kno	WLEI	GE-	•				
Table showi							, &c.	•		108
MISCELLANEOUS C	•									
Table showi	ng date, r	umber	of p	ages,	illust	rations	, &c.	•	•	109
ANNUAL REPORTS	OF THE	Instit	rutic)N						
Table showi	ng date, r	umber	of p	ages,	illust	rations	, &c.			110
PUBLICATIONS OF	THE U.	S. Nat	TION.	L M	USEUI	и				
Table showi	ng titles,	number	r in S	Smith	oniar	series	, &c.			111
Bulletins of	•						•			
Table showi							. &c.			112
Bulletins of T	_		_	_				ماسس		
Table showi					•			Olum	103	- 112
Proceedings of	•		_	_			, u c.	•	•	112
		•					a .			110
Table showing	•		_	_			, αc.	•	•	113
PUBLICATIONS OF							_			
Table showing	_		_	_				•	•	113
Publications having	-						n seri	ies b	ut	
included in	the Annu	al Rep	orts (of the	Insti	itution		•		114
Publications not in	cluded in	regula	ur ser	ies of	"Co	ntribu	ions,"	" C	ol-	
lections," or	"Report	s "		•	٠.					116
List of papers publ	ished in	the Co	ntrib	oution	s to	Know	edge.	givi	ng	
authors, title								•		117
ALPHABETICAL IN							blicati	ions		121

PREFACE.

The present catalogue embraces all the articles published by the Smithsonian Institution from its organization in 1846 to the first of July, 1882, a period of thirty-six years.

At the beginning nothing was issued but pamphlets explanatory of the plan of the Institution and brief annual reports of the proceedings of the Board of Regents, indicated in the catalogue by the letters A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, Q. An elaborate work, (P in the catalogue), by ROBERT DALE OWEN, on public architecture, with special reference to the plans of the Smithsonian Institution, prepared on behalf of the Building Committee, was printed at the expense of the Institution in 1849, but did not form part of the regular series organized by the Secretary of the Institution, Prof. Henry.

1. SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE.

The series entitled "Smithsonian Contributions to Knowledge," in quarto form, was commenced in 1848 by the publication of Squier and Davis' Ancient Monuments of the Mississippi Valley. The following "Advertisement" of the first volume, prepared by Prof. Henry, has been inserted in every succeeding volume to indicate the character and design of the series:

"This volume is intended to form the first of a series of volumes, consisting of original memoirs on different branches of knowledge published at the expense and under the direction of the Smithsonian Institution. The publication of this series forms part of a general plan adopted for carrying into effect the benevolent intentions of James Smithson, Esq., of England. This gentleman left his property in trust to the United States of America to found at Washington an institution which should bear his own name, and have for its objects 'the increase and diffusion of knowledge among men.' This trust was accepted by the Government of the United States, and an act of Congress was passed August 10, 1846, constituting the President and the other principal executive officers of the General Government, the Chief Justice of the Supreme Court, the Mayor of Washington, and such other persons as they might elect honorary members, an establishment under the name of the 'Smithsonian Institution, for the increase and diffusion of knowledge among

men.' The members and honorary members of this establishment are to hold stated and special meetings for the supervision of the affairs of the Institution and for the advice and instruction of a Board of Regents, to whom the financial and other affairs are entrusted.

"The Board of Regents consists of three members ex-officio of the establishment, namely, the Vice-President of the United States, the Chief Justice of the Supreme Court, and the Mayor of Washington, together with twelve other members, three of whom are appointed by the Senate from its own body, three by the House of Representatives from its members, and six citizens appointed by a joint resolution of both houses. To this Board is given the power of electing a Secretary and other officers, for conducting the active operations of the Institution.

"To carry into effect the purposes of the testator, the plan of organization should evidently embrace two objects, one, the increase of knowledge by the addition of new truths to the existing stock; the other, the diffusion of knowledge thus increased among men. No restriction is made in favor of any kind of knowledge, and hence each branch is entitled to and should receive a share

of attention.

"The act of Congress, establishing the Institution, directs, as part of the plan of organization, the formation of a Library, a Museum, and a Gallery of Art, together with provisions for physical research and popular lectures, while it leaves to the Regents the power of adopting such other parts of an organization as they may deem best suited to promote the objects of the bequest.

bequest.

"After much deliberation, the Regents resolved to divide the annual income, thirty thousand nine hundred and fifty dollars, into two equal parts, one part to be devoted to the increase and diffusion of knowledge by means of original research and publications, the other half of the income to be applied in accordance with the requirements of the act of Congress to the gradual formation of a Library, a Museum, and a Gallery of Art."

(The Programme of Organization, adopted December 8, 1847, follows)

"In accordance with the rules adopted in the Programme of Organization, each memoir in this volume has been favorably reported on by a Commission appointed for its examination. It is however, impossible, in most cases, to verify the statements of an author; and, therefore, neither the Commission nor the Institution can be responsible for more than the general character of a memoir."

The total number of papers published in the 23 volumes of "Contributions" is 119, with an aggregate of 12,456 pages, 1,567 wood cuts, 523 plates, and 16 maps, each volume averaging 541½ pages.

2. MISCELLANEOUS COLLECTIONS.

In the year 1862, another series was instituted, entitled "Smithsonian Miscellaneous Collections" each volume of which has the following preface:

"The present series, entitled "Smithsonian Miscellaneous Collections," is intended to embrace all the publications issued directly by the Smithsonian

PREFACE. VII

Institution in octavo form; those in quarto constituting the "Smithsonian Contributions to Knowledge." The quarto series includes memoirs, embracing the records of extended original investigations and researches, resulting in what are believed to be new truths, and constituting positive additions to the sum of human knowledge. The octavo series is designed to contain reports on the present state of our knowledge of particular branches of science; instructions for collecting and digesting facts and materials for research; lists and synopses of species of the organic and inorganic world; museum catalogues; reports of explorations; aids to bibliographical investigations, etc.; generally prepared at the express request of the Institution and at its expense.

"The position of a work in one or the other of the two series will sometimes depend upon whether the required illustrations can be presented more

conveniently in the quarto or the octavo form.

"In the Smithsonian Contributions to Knowledge, as well as in the present series, each article is separately paged and indexed, and the actual date of its publication is that given on its special title page, and not that of the volume in which it is placed. In many cases works have been published and largely distributed years before their combination into volumes.

"While due care is taken on the part of the Smithsonian Institution to insure a proper standard of excellence in its publications, it will be readily understood that it cannot hold itself responsible for the facts and conclusions of the authors, as it is impossible in most cases to verify their statements."

The total number of papers published in the 23 volumes of "Miscellaneous Collections" is 122, each volume averaging 882½ pages, with an aggregate of 20,299 pages, 2,868 wood cuts, and 43 plates.

3. Annual Reports.

By the act of Congress organizing the Institution it was made the duty of the "Board of Regents to submit at each session a report of the operations, expenditures, and condition of the Institution." These Annual Reports form a third series of Smithsonian publications. They consist of the reports of the Secretary to the Board of Regents of the operations and condition of the Institution; the reports of committees of the Board; reports of lectures; extracts from correspondence; original or translated articles relating to the history and progress of science, etc.

The first report was submitted by the Board to the second session of the 29th Congress, 1847, and formed an octavo pamphlet of 38 pages. A similar report was presented annually thereafter, varying in size from 64 pages to 326, printed in pamphlet form with paper covers up to 1853, when Congress ordered the report to be bound in cloth. In the volume for that year the essential portion of the contents of the preceding seven reports was reprinted,

VIII PREFACE.

and this is now considered as the first of a set of Smithsonian Reports. The number of pages was limited between 1854 and 1876 to 400. In the latter year this restriction was removed, and since then the average number of pages has been 600.

The number of copies of these reports for general distribution ordered by Congress has been very variable, the largest being 7,500 in 1874 and 1875, and the smallest 150 in 1847. The number of copies granted the Institution each year is shown in the following table:

Number of extra copies furnished the Institution by Congre	s for	r distribution.
--	-------	-----------------

FOR THE YEAR.	No. of Copies.	FOR THE YEAR.	No. of Copies.	FOR THE YEAR.	No. of Copies.
1847	150	1859	2,000	1871	5,000
1848	1,000	1860	2,000	1872	5,000
1849	500	1861	2,000	1873	6,00 0
1850	1,000	1862	2,000	1874	7,500
1851	2,000	1863	2,000	1875	7,500
1852	2,000	1864	2,000	1876	6,500
1853	3,000	1865	2,000	1877	6,500
1854	2,500	1866	2,000	1878	6,500
1855	2,500	1867	2,000	1879	7,000
1856	2,500	1868	2,000	1880	7,000
1857	5,000	1869	3,000] 1	
1858	5,000	1870	5,000	11	

The total number of pages in the 35 volumes of Annual Reports is 14,419, average 412 pages; total number of woodcuts, 1,898.

4. Bulletins of the U.S. National Museum.

In the year 1875 a fourth series of publications (octavo) was commenced, entitled "Bulletins of the National Museum," intended to illustrate the collections of natural history and ethnology belonging to the United States, constituting the National Museum, of which the Smithsonian Institution is the custodian.

Twenty of these Bulletins have been published, with an aggregate of 3,103 pages, 45 plates, and 1 map.

5. Proceedings of the U.S. National Museum.

In imitation of the practice of those learned societies which publish periodically descriptions of new species, &c., in the form of proceedings of weekly or monthly meetings, and thus present to the world the discoveries connected with the establishment at the earliest practicable moment, it appeared to be very desirable that the National Museum should have some medium of prompt publication for announcing descriptions of specimens received, (many of which are new species,) as well as other interesting facts relative to natural history furnished by correspondents of the Institution. To meet this want a fifth series of publications, (octavo,) entitled "Proceedings of the National Museum," was commenced in 1880. They are printed in successive signatures as fast as material sufficient for 16 pages is prepared, and distributed at once to scientific societies and leading active working naturalists in this country and in Europe,* each signature having printed at the bottom of its first page the date of actual issue, for settling any questions as to priority of publication. Of this series four volumes have been published, comprising 2,221 pages, with 28 cuts and 19 plates.

6. REPORTS OF THE BUREAU OF ETHNOLOGY.

The sixth series of publications is the annual report (in Imperial octavo) of the Bureau of Ethnology, placed by Congress in charge of the Smithsonian Institution. The first volume of this series was issued in 1881, and consists of 638 pages, with 343 cuts, 54 plates, and 1 map.

The distribution of this volume to individuals is wholly by Members of Congress and the Director of the Bureau, Major J. W. Powell—the Institution having copies at its disposal only for the libraries on its regular list of distribution for its own full series.

7. COPYRIGHT.

No copyright has ever been secured on the publications of the Institution. They are left free to be used by compilers of books without any restrictions, except that full credit shall be given to the name of Smithson for any extracts which may be made from them.

^{*} Prof. Baird's report for 1880.

8. Use of Illustrations.

Copies of the wood cuts used by the Institution are granted to authors or publishers on payment of the actual cost of production of electrotypes, and promise to give proper reference to the article in which they originally appeared.

9. Size of Editions.

In the first experiments of the Smithsonian system of publication, the proper magnitude of the editions necessary to meet the immediate and future demand could not be accurately ascertained. The number of copies of the Contributions then fixed upon, has since been found inadequate, although it was larger than that usually issued by other institutions. The edition has, therefore, been augmented, until at the present time 1,000 copies of each article are set aside to be combined into volumes, and an extra number, varying with the probable demand, struck off for separate distribution, and for sale.

Each article is complete in itself, with separate paging, title, and index, and without any necessary relationship to others combined with it in the same volume.

Of the early volumes of Smithsonian Contributions, the edition, for reasons already explained, was less than of the succeeding ones, so that complete sets cannot now be furnished.

In the year 1862, the plan of stereotyping every article printed by the Institution was adopted, the plates being carefully preserved, thus making it practicable at any time to issue new editions except where expensive lithographic plates were used, a limited number, only, of impressions from these having been taken.

A number of the earlier articles in octavo were out of print before the commencement of the series of "Miscellaneous Collections," and consequently are not included in them.

The printing of the "Bulletins" and "Proceedings" is authorized by the DEPARTMENT OF THE INTERIOR and paid for out of its fund. An edition of 1,000 copies is published, of which one-half is distributed by the Department of the Interior and one-half by the Institution. As the pages are stereotyped, the cost of additional copies is alight; and for the purpose of making sure that a sufficient number of sets will be accessible forever to

PREFACE. XI

students in all parts of the world, it has been considered expedient to print 1,500 additional copies of each for incorporation in the Miscellaneous Collections.*

10. DISTRIBUTION OF PUBLICATIONS.

The distribution of the publications of the Institution is a matter which requires much care and judicious selection, the great object being to make known to the world the truths which may result from the expenditure of the Smithson fund. For this purpose the Contributions are so distributed as to be accessible to the greatest number of readers; that is, to large central libraries.

The volumes of Contributions are presented on the express condition that, while they are carefully preserved, they shall be accessible at all times to students and others who may desire to consult them, and be returned to the Institution in case the establishments to which they are presented at any time case to exist.

These works, it must be recollected, are not of a popular character, but require profound study to fully understand them; they are, however, of importance to the professional teacher and the popular expounder of science. They contain the materials from which general treatises on special subjects may be elaborated.†

Full sets of the publications cannot be given to all who apply for them, since this is impossible with the limited income of the Institution, and, indeed, if care be not exercised in the distribution, so large a portion of the income will be annually expended on the production of copies for distribution of what has already been published that nothing further can be done in the way of new publications. It must be recollected that every addition to the list of distribution not only involves the giving of the publications which have already been made, but also of those which are to be made hereafter.‡

The rules governing the distribution of the Smithsonian publications are appended. To enable institutions not coming within their provisos, as well as individuals, to procure copies of such as may be desired, a small number is set aside and sold by the Institution at a price which is intended merely to cover the actual cost of their publication.

^{*} Prof. Baird's report for 1880.

[†] Prof. Henry's report for 1876.

[‡] Prof. Henry's report for 1873.

XII PREFACE.

11. Rules for Distribution of the Publications of the Smithsonian Institution.

To Institutions.

The publications of the Smithsonian Institution are furnished:

- 1st. To learned societies of the first class, which present complete series of their publications to the Institution.
- 2d. To libraries of the first class, which give in exchange their catalogues and other publications; or an equivalent, from their duplicate volumes.
- 3d. To colleges of the first class, which furnish catalogues of their libraries and of their students, and all publications relative to their organization and history.
 - 4th. To public libraries containing 25,000 volumes.
- 5th. To smaller public libraries, where a large district would be otherwise unsupplied.
- 6th. Institutions devoted exclusively to the promotion of particular branches of knowledge may receive such Smithsonian publications as relate to their respective objects.

To Individuals.

The gratuitous distribution to individuals, of the publications of the Institution, is restricted:

- 1st. To those who are engaged in original research in the branch of science to which the book asked for pertains.
 - 2d. To those who require it in the business of instruction.
 - 3d. To donors to the museum or library of the Institution.

18

12. Form of Application for Publications.

Date,

To the Smithsonian Institution, Washington, D. C.	To	the	Smiths	onian	Institution,	Washington,	D.	C
---	----	-----	--------	-------	--------------	-------------	----	---

In behalf of the	,	we respectfully apply	
or the publications of the Smiths	onian Institution, on	condition that all vol-	
tmes received shall be carefully	preserved, be accessible	le to any person who	
may wish to consult them, and be	returned to the Smit	hsonian Institution in	
ase the establishment at any time	e ceases to exist.		
1. Name of Establishment			
2. Location—Town			
State			
3. When established			
4. Character	·····		
5. Buildings and property			
6. Permanent fund			
7. Annual income			
8. Volumes in library	•		
9. Number of persons having use of b	ooks		
10. Date of last catalogue of library	•		
11. What publications made			
(Send printed list if possible.)			
12. Names of Officers: President			
Secretary			
Librarian			
13. Addresses of principal scientific me	n connected with the Est	ablishment and subjects in	
which specially interested			
•			
I recommend the above application.			
	Member of Congress,		
	District	State	

13. PRICE LIST OF SMITHSONIAN PUBLICATIONS.

Where no price is given the work is out of print, and cannot be furnished. Of those marked "free" the edition is limited, and copies are only given to those specially interested in the subjects to which they pertain, who are collaborators of the Institution or contributors to its library or museum.

										_					
P	\$10 00	60	\$1 00	120	\$1.00	180	\$ 1 00	240		300	\$0 25	360	free.	420	free.
ī	₩TO OO	61	1 00	121	25	181	2 00	241	\$7 50	301	#U 20	361	1100.	421	free.
2		62	2 00	122	6 00	182	2 50	242	7 50	302	25	362	free.	422	\$15
3		63	ĩõõ	123	6 00	183	25	243	free.	303	60	363	free.	423	6 91
ĭ.		64	free.	124	0 00	184	12 00	244	1 00	304	50	364	free.	424	6 99
4		65		125	6 00	185	free.	245	free.	306	50	365	free.	425	2 (0
6		66		126	1 25	186	50	246	12 00	306	50	366	free.	426	free.
7		67		127	1 25	187	50	247	50	307		367	free.	427	free.
8		68		128	25	188	50	248		308	50	368		423	1.00
9		69		129	1 00	189	1 00	249		309	free.	369		429	tree
10		70	6 00	130	1 50	190	free.	250		310	25	370	free.	430	free.
11		71		131	1 25	191		251	free.	311	50	371	free.	431	free.
12	4 00	72	3 00	132	25	192	5 00	252		312		372	free.	432	Irre.
13	2 00	73	25	133	2 00	193	free.	253		313	50	373	free.	433	free.
14	1 00	74		134		194		254	2 00	314	6 00	374	free.	434	free
15	4 00	75		135		195		255		315	6 00	375	free.	435	free.
16		76	12 00	136		196		256	2 60	316	free.	376	free.	436	free. free
17	50	77		137		197	1 00	257		317	3 00	377	free.	437	
18 19		78 79	12 00	138	free.	198 199	2 00	256 259	2 00 l 3 00	318 319	2 00	378	free.	438 439	frer. 25
20	1 00	80	1 00	140	1ree. 75	200	2 UU 25	260	3 00	320	free. free.	379 380	free. free.	440	30
21	1 00	81		141	1 50	200	50	261	free.	321	25	381	free.	441	1 50
22	1 00	82	1 50	142	3 00	202	3 50	262	3 00	322	600	382	free.	442	50
23	2 00	83	1 25	143	1 25	203	0 00	263	0 00	323		383	\$2 50	443	1 (0
24		84	1 50	144	1 00	204	1 00	264		324	free.	384	free.	444	30
25	50	85	25	145	75	205	free.	265		325	free.	385	free.	445	ine
25 26		86		146	1 50	206	12 00	266	25	326	50	386	free.	446	free.
27	50	87	25	147		207	free.	267	1 00	327	75	387	free.	447	free.
28	1 00	88	1 00	148	free.	208	50	268		328	5 00	388	free.	446	free
29		89	5 00	149		209		269		329		389	free.	449	free.
29 30 81		90	5 00	150		210	25	270	2 00	830		390	free.	450	free
81		91		151	12 00	211	12 00	271		331	2 00	391	free.	451	free.
82 83		92	12 00	152		212	6 00	272	12 00	832	2 00	392	free.	452 453	free. free
84 84	75	93 94	50	153 154		213 214	6 00 1 00	273 274	6 00 6 00	833 334	2 00 25	393 394	free.	454	free.
35	free.	95	6 80	155		215	100	275	8 80	335	free.	395	free.	455	free.
36	1 00	96	20 00	156	50	216	25	276	80	336	6 00	396	free.	456	free
87	- ~ -	97	1 00	157	•	217		277	3 00	337	6 00	397	free.	457	free.
88		98	4 00	158		218		278	free.	338		398	free.	458	free.
89		99	12 00	159	1 00	219	2 00	270	25	339		399	free.	450	free.
40		100	50	160	1	220		280	1 00	340	12 00	400	free.	460	free.
41	1 00	101	free.	161	50	221	1 00	281	75	341	50	401	free.	461	free
42		102		162	1 00	222	3 00	282	25	342	50	402	free.	403	free.
48		103	2 00	163	free.	223	1 00	283	25	343	free.	403	free.	463	30
44		104		164	50	224		284	12 00	344	free.	404	free.	464	free
45		105	50	165		225		285	12 00	845	50	405	free.	465 466	free. free
46 47		106	50	166	1 00	226	free.	286	50	346	12 00	406	free.	467	3 (1)
48	50	107 108	25	167 168	1 00	227 228	25 50	287 288	2 00	347 348	free.	407 408		468	6 (1)
49	1 00	109	20	169	free. 6 00	228	12 00	289	50	349	25	409	free. free.	469	free
50	1 50	110		170	free.	230	25	200	free.	350	free.	410	free.	470	3.0
51	100	iii	12 00	171	MEG.	231	w	201	25	351	nec.	411	free.	471	free.
52	5 00	112	12 00	172		232	2 00	292	50	352	free.	412	50	472	free.
53	80	113	25	173	2 50	233	25	293	50	353	8 00	413	50	473	free.
54	50	114	1 26	174	2 00	234	free.	294	80	354	free.	414	15	474	free
88		115	free.	175	1 00	235	free.	295	50	3.55		415	free.	475	600
56 57		116		176	free.	236	free.	296	50	366	50	416	6 00	476	_
57	1 00	117	2 00	177	25	237	free.	297	50	357	2 50	417	free.	477	free
58	5 00	118	1 00	178	free.	238	free.	298	50	358	1 00	418	free.	478	:5
50	25	119	100	179	3 00	239	2 00	299		350	free.	419	free.		

LIST OF PUBLICATIONS

OF THE

SMITHSONIAN INSTITUTION.

Note.—A to Q indicate early publications not embraced in the regular series.

- A. Journal of Proceedings of the Regents of the Smithsonian Institution, at the city of Washington, beginning on the first Monday of September, 1846. 1846. 8vo., pp. 32.
- B. Report of the Organization Committee of the Smithsonian Institution, with the resolutions accompanying the same and adopted by the Board of Regents; also, the Will of the testator, the Act accepting the bequest, and the Act organizing the Institution. 1847. 8vo., pp. 32.
- C. Digest of the Act of Congress establishing the Smithsonian Institution.

 August 10, 1846. 8vo., pp. 8.
- D. Address delivered on occasion of laying the Corner Stone of the Smithsonian Institution, May 1, 1847. By George M. Dallas, Chancellor of the Institution. 1847. 8vo., pp. 8.
- E. Smithson's Bequest. Professor Henry's exposition before the New Jersey Historical Society, at its meeting in Princeton, on Wednesday, September 27. 1847. 8vo., pp. 8.
- F. First Report of the Secretary of the Smithsonian Institution to the Board of Regents; giving a Programme of Organization, and an account of the operations during the year. Presented December 8, 1847. 1848. 8vo., pp. 48.
- G. [First] Report from the Board of Regents, submitted to Congress, of the operations, expenditures, and condition of the Smithsonian Institution. Senate Doc. 211; 29th Congress, 2d Session. 1847. 8vo., pp. 38.
- H. Second Report of the Board of Regents of the Smithsonian Institution, to the Senate and House of Representatives, showing the operations, expenditures, and condition of the Institution during the year 1847. 30th Congress, 1st Session. Senate Miscellaneous No. 23. 1848. 8vo., pp. 208.

(1)

CLASSIFIED LIS	T OF SERAP.	ame T	ים וד	TC A TIT	\Wa	-Cont	hanni			Page.
	Geography	7115 1	UDL	ICAII	JAB	СОПС	mucu.			103
· Physics	осовтириј	•	•	•	•	•	•	•	•	103
	l Physics	•	•	•	•	•	•	•	•	104
Zoölogy—			•	•	•	•	•	•	•	104
22001083	Fishes .	•	•	•	•	•	•	•	•	105
	Insects .	•	•	•	•	•	•	•	•	105
	Mammals	•	•	•	•	•	•	•	•	106
	Mollusks	•	•	•	•	•	•	•	•	106
	Radiates	•	•	•	•	•	•	•	•	107
	Reptiles		•	•	•	•	•	•	•	107
Smithsonian C	•	· ·	V No		· ·	•	•	•	•	
	wing date, nu		-			- ration	18, &c.	•		103
Miscellaneous Table sho	Collection wing date, nu		of p	ages,	illust	ration	ıs, &c.	•	•	109
Annual Repor Table sho	ers of the l				illust	ration	18, & c.		•	110
Publications of	_	. Nat	TION	AL M	USEU	м—				111
Bulletins of Table sho	F THE U.S. wing date, nu						ıs, &c.		•	112
Bulletins of	THE U.S. N	ATIO	NAL	Muse	UM, c	ollect	ed in v	olum	es	
Table sho	wing date, nu	ımber	of p	ages,	illust	ration	s, &c.			112
Proceedings	-		-	-						
Table sho	wing date, nu	ımber	of p	ages,	illust	ration		•		113
Publications of Table sho	of the Burn wing date, nu						s, &c.			113
Publications hav	ing separate in the Annua							ies b	ut	114
Publications not	included in	regula						"C	· ol-	
lections,"	or "Reports	"		•	٠,	•	•			116
List of papers pu authors, t	ablished in the						vledge,	givir	ng	117
AT DEFADENTICAT							hlione	-	-	191

PREFACE.

The present catalogue embraces all the articles published by the Smithsonian Institution from its organization in 1846 to the first of July, 1882, a period of thirty-six years.

At the beginning nothing was issued but pamphlets explanatory of the plan of the Institution and brief annual reports of the proceedings of the Board of Regents, indicated in the catalogue by the letters A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, Q. An elaborate work, (P in the catalogue), by ROBERT DALE OWEN, on public architecture, with special reference to the plans of the Smithsonian Institution, prepared on behalf of the Building Committee, was printed at the expense of the Institution in 1849, but did not form part of the regular series organized by the Secretary of the Institution, Prof. Henry.

1. SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE.

The series entitled "Smithsonian Contributions to Knowledge," in quarto form, was commenced in 1848 by the publication of Squier and Davis' Ancient Monuments of the Mississippi Valley. The following "Advertisement" of the first volume, prepared by Prof. Henry, has been inserted in every succeeding volume to indicate the character and design of the series:

"This volume is intended to form the first of a series of volumes, consisting of original memoirs on different branches of knowledge published at the expense and under the direction of the Smithsonian Institution. The publication of this series forms part of a general plan adopted for carrying into effect the benevolent intentions of James Smithson, Esq., of England. This gentleman left his property in trust to the United States of America to found at Washington an institution which should bear his own name, and have for its objects 'the increase and diffusion of knowledge among men.' This trust was accepted by the Government of the United States, and an act of Congress was passed August 10, 1846, constituting the President and the other principal executive officers of the General Government, the Chief Justice of the Supreme Court, the Mayor of Washington, and such other persons as they might elect honorary members, an establishment under the name of the 'Smithsonian Institution, for the increase and diffusion of knowledge among

- other rivers. By Charles Ellet, Jr. 1850. 4to., pp. 64, 2 woodcuts, 1 plate. (S. C. 11.)
- 14. A Memoir on Mosasaurus, and the three Allied New Genera, Holcodus, Conosaurus, and Amphorosteus. By Robert W. Gibbes. November, 1850. 4to., pp. 14, 3 plates of 28 figures. (S. C. 11.)
- 15. Aboriginal Monuments of the State of New York. Comprising the results of original surveys and explorations; with an illustrative appendix. By E. G. SQUIER. 1850. 4to., pp. 188, 79 woodcuts. 14 plates of 33 figures. (S. C. 11.)
- 16. The Classification of Insects from Embryological Data. By Louis Agassiz, 1850. 4to., pp. 28, 8 woodcuts, one plate of 23 figures. (S. C. II.)
- 17. Memoir on the Explosiveness of Nitre, with a view to elucidate its agency in the tremendous explosion of July, 1845, in New York. By ROBERT HARE. 1850. 4to., pp. 20. (S. C. II.)
- 18. Report on the History of the Discovery of Neptune. By BENJAMIN APTHORP GOULD, Jr. 1850. 8vo., pp. 56.
- 19. Directions for Meteorological Observations, intended for the first class of observers. By Arnold Guyot. 1850. 8vo., pp. 40, 9 woodcuts.
- 20. Microscopical Examination of Soundings, made by the United States Coast Survey off the Atlantic coast of the United States. By J. W. Bailey. January, 1851. 4to., pp. 16 and 1 plate of 68 figures. (S. C. II.)
- Fourth Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1849. 31st Congress, 1st Session. Senate Miscellaneous No. 120, 8vo., pp. 64, with appendix of 207 pp. House of Representatives Miscellaneous No. 50. 1850. 8vo., pp. 272.

Report of Prof. J. HENRY, and Proceedings of the Board.

GRAY, ASA. Account of Lindheimer's, Fendler's and Wright's botanical explorations in New Mexico and California.

AGASSIZ, LOUIS. On the formation of a museum.

List of meteorological observers.

JEWETT, G. C. Report on library and catalogue system.

JEWETT, C. C. Report on public libraries of the United States.

22. Plantæ Wrightianæ Texano-Neo-Mexicanæ. By Asa Gray. Part I March, 1852. 4to., pp. 146, 10 plates of 127 figures. (S. C. 111.

An account of a collection of plants made by Charles Wright in Western Texas, New Mexico, and Sonora, in the years 1851 and 1852.

23. Microscopical Observations made in South Carolina, Georgia, and Florida. By J. W. Bailey. 1851. 4to., pp. 48, 3 plates of 83 figures. (S. C. II.)

- 24. Ephemeris of the Planet Neptune for the year 1852. By SEARS C. WALKER. 1853. 4to. pp. 10. (S. C. III.)
- Notices of Public Libraries in the United States of America. By Chas.
 Jewett. Printed by order of Congress as an appendix to the Fourth Annual Report of the Board of Regents of the Smithsonian Institution. 1851. 8vo., pp. 210.
- Smithsonian Contributions to Knowledge. Vol. II. 1851. 4to., pp. 572, 89 woodcuts, 24 plates.

WALKER, S. C. Researches relative to Neptune. No. 3.

LIEBER, F. Vocal sounds of Laura Bridgman. No. 12.

BAILEY, J. W. Microscopical soundings off Atlantic Coast. No. 20.

ELLET, C. Physical geography of the Mississippi Valley. No. 13.

GIBBES, R. W. Mosasaurus and three allied genera. No. 14.

AGASSIZ, L. Classification of insects from embryological data. No. 16.

HARE, R. Explosiveness of nitre. No. 17.

BAILEY, J. W. Microscopical observations in S. C., Ga., Fla. No. 23.

SQUIER, E. G. Aboriginal monuments of State of New York. No. 15.

WALKER, S. C. Ephemeris of Neptune for 1848. No. 4.

WALKER, S. C. Ephemeris of Neptune for 1846, '47, '48, '49. No. 5.

WALKER, S. C. Ephemeris of Neptune for 1850. No. 6.

WALKER, S. C. Ephemeris of Neptune for 1851. No. 7.

DOWNES, J. Occultations visible in the United States in 1851. No. 11.

- 27. On Recent Improvements in the Chemical Arts. By James C. Booth and Campbell Morfit. 1852. 8vo., pp. 216. (M. C. II.)
- Fifth Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1850. Special session, March, 1851. Senate Miscellaneous No. 1. 1851. 8vo., pp. 145. (Extra edition of 326 pp.)

CONTENTS.

Report of Prof. J. Henry, and Proceedings of the Board.

Jewett, C. C. General catalogue system for libraries.

Baird, S. F. Report on museum, and statistics of British Museum.

Memorial of the Regents to Congress, relative to the Smithson Fund.

Squier, E. G. Antiquities of Nicaragua.

Report of Commission on General Stereotype Catalogue of Pub. Libraries.

Culbertson, T. A. Expedition to the Mauvaises Terres and Upper Missouri.

PORTER, T. C. List of plants of Upper Missouri. HARRIS, E. List of birds and mammalia of Missouri river. CULBERTSON, T. A. Indian tribes of the Upper Missouri. JEWETT, C. C. Copyright books from 1846-1849.

29. Occultations visible in the United States during the year 1852. By JOHN DOWNES. 1851. 4to., pp. 34. (S. C. III.)

- 30. Contributions to the Natural History of the Fresh Water Fishes of North America. By Charles Girard. Part I.—A monograph of the Cottoids. December, 1851. 4to., pp. 80, 3 plates of 48 figures. (S. C. III.)
- 31. A Collection of Meteorological Tables, with other tables useful in Practical Meteorology. By Arnold Guyot. 1852. 8vo., pp. 212.
- 32. Nereis Boreali-Americana: or, Contributions to a History of the Marine Algæ of North America. By William Henry Harvey. Part I.—Melanospermeæ. January, 1852. 4to., pp. 152, 12 colored plates of 29 figures. (S. C. III.)
- 33. The Law of Deposit of the Flood Tide: its Dynamical Action and Office. By CHARLES HENRY DAVIS. 1852. 4to., pp. 14. (S. C. III.)
- 34. Directions for Collecting, Preserving, and Transporting Specimens of Natural History. March, 1859. 8vo., pp. 40, 6 woodcuts. (M. C. II.)
- 35. Observations on Terrestrial Magnetism. By John Locke. April. 1852. 4to., pp. 30. (S. C. III.)
- 36. Researches on Electrical Rheometry. By A. SECCHI. May, 1852. 4to., pp. 60, 3 plates of 15 figures. (S. C. III.)
- 37. Descriptions of Ancient Works in Ohio. By Chas. WHITTLESEY. 1851. 4to., pp. 20, 7 plates of 18 figures. (S. C. III.)
- Smithsonian Contributions to Knowledge. Vol. III. 1852. 4to.,
 pp. 562, and 35 plates.

LOCKE, J. Terrestrial magnetism. No. 35.

SECCHI, A. Electrical rheometry. No. 36.

GIRARD, C. Monograph of the cottoids. No. 30.

HARVEY, W. H. Marine algo of North America. Part I. No. 32.

GRAY, A. Planto Wrightiano Texano-Neo-Mexicano. Part I. No. 22.

DAVIS, C. H. Law of deposit of the flood tide. No. 33.

WHITTLESEY, C. Descriptions of ancient works in Ohio. No. 37.

WALKER, S. C. Ephemeris of the planet Neptune for 1852. No. 24.

DOWNES, J. Occultations visible in United States during 1852. No. 24.

39. Smithsonian Contributions to Knowledge. Vol. IV. 1852. 4to., pp. 426.

CONTENTS.

- Riggs, S. R. Dakota Grammar and Dictionary. No. 40.
- 40. Grammar and Dictionary of the Dakota Language. Collected by the members of the Dakota mission. Edited by S. R. RIGGS. 1852. 4to., pp. 414. (S. C. IV.)

- 41. Memoir on the Extinct Species of American Ox. By JOSEPH LEIDY. December, 1852. 4to., pp. 20, 4 plates of 15 figures. (S. C. v.)
- 42. Plantæ Wrightianæ Texano-Neo-Mexicanæ. By Asa Gray. Part II. February, 1853. 4to., pp. 120, 4 plates of 39 figures. (S. C. v.)
- Nereis Boreali-Americana; or, Contributions to a History of the Marine Algæ of North America. By W. H. HARVEY. Part II.—Rhodospermeæ. March, 1853. 4to., pp. 262, 24 plates, colored, of 64 figures. (S. C. v.)
- 44. A Flora and Fauna within Living Animals. By Joseph Leidy. April, 1853. 4to., pp. 68, 10 plates of 140 figures. (S. C. v.)
- 45. Anatomy of the Nervous System of Rana pipiens. By JEFFRIES WY-MAN. March, 1853. 4to., pp. 52, 4 woodcuts, 2 plates of 29 figures. (S. C. v.)
- Plantæ Frémontianæ: or, Descriptions of Plants collected by J. C. Frémont in California. By John Torrey. 1853. 4to., pp. 24, 10 plates of 89 figures. (S. C. vi.)
- On the Construction of Catalogues of Libraries, and their publication by means of separate stereotyped titles. With rules and examples. By Charles C. Jewett. 1852. 8vo., pp. 78. 1853. 8vo., pp. 108.
- 48. Bibliographia Americana Historico-Naturalis; or Bibliography of American Natural History for the year 1851. 1851. By CHARLES GIRARD. December, 1852. 8vo., pp. 64.
- Catalogue of North American Reptiles in the Museum of the Smithsonian Institution. By S. F. BAIRD and C. GIRARD. Part I.—Serpents. January, 1853. 8vo., pp. 188. (M. C. II.)
- Synopsis of the Marine Invertebrata of Grand Manan: or the region about the mouth of the Bay of Fundy, New Brunswick. By Wm. STIMPSON. March, 1853. 4to., pp. 68, 3 plates of 37 figures. (S. C. vi.)
- 51. Sixth Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1851. 32d Congress, 1st session, Senate Mis. No. 108. 1852. 8vo., pp. 104.

Report of Prof. J. Heney, and Proceedings of the Board.

Jewett, C. C. Smithsonian library and copyright system.

Baird, S. F. Natural history explorations in the United States in 1851.

FOREMAN, E. Meteorological system and correspondence.

Leidy, J. Report on fossils from Nebraska.

Turner, W. W. Indian philology.

- 51. Report for 1851—Continued.
 - Report of Committee of American Association for Promotion of Science, on a system of combined meteorological observations for North America.
- 52. Winds of the Northern Hemisphere. By James H. Coffin. November, 1853. 4to., pp. 200, 6 woodcuts, 13 plates of 238 figures. S. C. vi.)
- 53. Catalogue of Portraits of North American Indians, and Sketches of Scenery, etc., painted by J. M. STANLEY. Deposited with the Smithsonian Institution. December, 1852. 8vo., pp. 76. (M. C. II.)
- 54. Occultations of Planets and Stars by the Moon, during the year 1853. Ву Јони Downes. 1853. 4to., pp. 36. (S. C. vi.)
- 55. Smithsonian Contributions to Knowledge. Vol. V. 1853. 4to., pp. 538, 4 woodcuts, 45 plates.

CANTENTS

LEIDY, J. Flora and fauna within living animals. No. 44.

LEIDY, J. Extinct species of American ox. No. 41.

WYMAN, J. Anatomy of the nervous system of Rana pipiens. No. 45 HARVEY, W. H. Marine algos of North America. Part II. No. 43. Gray, A. Plantos Wrightianse. Part II. No. 42.

56. Smithsonian Contributions to Knowledge. Vol. VI. 1854. 4to., pp. 484, 9 woodcuts, 53 plates.

CONTENTS.

Torrey, J. Plantæ Frémontianæ. No. 46.

TORREY, J. Batis maritima. No. 60.

TORREY, J. Darlingtonia californica. No. 61.

STIMPSON, W. Marine invertebrata of Grand Manan. No. 50. COFFIN, J. H. Winds of the Northern Hemisphere. No. 52.

LEIDY, J. Ancient fauna of Nebraska. No. 58.

Downes, J. Occultations during the year 1858. No. 54.

57. Seventh Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1852. 32d Congress, 2d Session, Senate Mis. No. 53. 1853. 8vo., pp. 96.

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

JEWETT, C. C. Report on library and the Halliwell manuscripts.

BAIRD, S. F. Scientific explorations in America in 1852.

FOREMAN, E. Report on meteorological system.

- 58. The Ancient Fauna of Nebraska: or, a Description of Remains of Extinct Mammalia and Chelonia, from the Mauvaises Terres of Nebraska. By Joseph Leidy. June, 1853. 4to., pp. 126, 3 woodcuts. 25 plates of 126 figures. (S. C. VI.)
- 59. Account of a Tornado near New Harmony, Indiana, April 30, 1852.

with a map of the track, etc. By John Chappelsmith. April, 1855. 4to., pp. 12, 2 woodcuts, 1 map, 1 plate. (S. C. vii.)

- 60. Observations on the Batis maritima of Linnæus. By John Torrey. April, 1853. 4to., pp. 8, 1 plate of 21 figures. (S. C. vi.)
- 61. On the Darlingtonia Californica; a new pitcher-plant from Northern California. By John Torrey. April, 1858. 4to., pp. 8, 1 plate of 9 figures. (S. C. vi.)
- Catalogue of the Described Coleoptera of the United States. By F. E.
 Melsheimer. July, 1853. 8vo., pp. 190.
- 63. Notes on New Species and Localities of Microscopical Organisms. By J. W. Bailey, February, 1854. 4to., pp. 16, 7 woodcuts, 1 plate of 39 figures. (S. C. VII.)
- 64. List of Foreign Institutions in Correspondence with the Smithsonian Institution. 1856. 8vo., pp. 16.
- 65. Registry of Periodical Phenomena. Folio, pp. 4.
- 66. The Annular Eclipse of May 26, 1854. 8vo., pp. 14, 1 map.
- 67. Eighth Annual Report of the Board of Regents of the Smithsonian Institution for the year 1853. 33d Congress, 1st Session, Senate Doc. No. 73, pp. 269. House of Representatives, Mis. Doc. No. 97. 1854. 8vo., pp. 310.*

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

JEWETT, C. C. Report on library.

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

BLODGET, L. List of meteorological observers.

PEARCE, J. A. Report of Committee of Regents on distribution of Smithsonian income.

MEACHAM, J. Minority report of Committee on income.

SMITHSON, J. Will of

RUSH, R. Letter from, relative to James Smithson.

GILBERT, D., President of the Royal Society. Notice of Smithson.

SMITHSON, J. List of papers presented by, to the Royal Society.

SMITHSON, J. Contributions to the Annals of Philosophy.

Act of Congress accepting Bequest, July 1, 1836.

Act of Congress to establish the Smithsonian Institution, August 10, 1846.

BERBIEN, J. McP. Construction of the Act establishing the Smithsonian Institution.

HENRY, J. Address on the Smithsonian Institution.

^{*}This is the first of the series of annual reports published by Congress as a bound volume.

67. Report for 1853-Continued.

..

"

"

EVERETT, E.; SPARKS, J.; PIERCE, B.; LONGFELLOW, H. W.; GRAY, A. Report of American Academy of Arts and Sciences on Organization of Smithsonian Institution.

- First Report of the Secretary, Dec. 8, 1847. (Reprint.) HENRY, J. Second Report of the Secretary for 1848.
 - " Third Report of the Secretary for 1849. "
 - Fourth Report of the Secretary for 1850. "
 - Fifth Report of the Secretary for 1851. Sixth Report of the Secretary for 1852.
- RUSH MITCHELL, with additions by W. W. TURNER. April, 1853. 8vo., pp. 22. 69. List of Domestic Institutions in correspondence with the Smithsonian

68. Vocabulary of the Jargon or Trade Language of Oregon. By B.

- Institution. 1853. 8vo., pp. 16. 70. The Antiquities of Wisconsin, as Surveyed and Described, by I. A.
- LAPHAM. May, 1855. 4to., pp. 108, 65 woodcuts, one map, 54 plates of 112 figures. (S. I. vii.) 71. Archæology of the United States; or Sketches, Historical and Bibliographical, of the progress of information and opinion respecting Ver
- July, 1856. 4to., pp. 172. (S. C. VIII.) 72. A Memoir on the Extinct Sloth tribe of North America. By JOSEPH LEIDY. June, 1855. 4to., pp. 70, 16 plates of 139 figures. 🔌 C. vII.)

tiges of Antiquity in the United States. By SAMUEL F. HAVEN.

- 73. Publications of Learned Societies and Periodicals in the Library of the Smithsonian Institution. December 31, 1854. Part 1. 1855. 4to., pp. 40. (S. C. VII.)
- 74. Catalogue of Publications of the Smithsonian Institution. Corrected to June, 1862. 8vo., pp. 52. (M. C. v.)
- 75. Ninth Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1854. Senate Mis. Doc. No. 24, 33d Congress, 2d session. House of Representatives, Mis. Doc. No. 51. 1855. 8vo., pp. 464, 4 woodcuts.

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

BAIRD, S. F. Report on publications, exchanges, museum, and explorations in the years 1858 and 1854.

ALEXANDER, B. S. Report of architect.

List of meteorological stations and observers.

MARSH, G. P. Lecture on the camel.

BRAINARD, D. Lecture on nature and cure of bites of serpents and the wounds of poisoned arrows.

LOOMIS, E. Lecture on the zone of small planets between Man and Jupiter.

75. Report for 1854-Continued.

CHANNING, W. F. Lecture on the American fire alarm telegraph.

REED, H. Lectures on the Union.

RUSSELL, R.; HENRY, J. Lectures and notes on meteorology.

HARE, R. On John Wise's observation of a thunderstorm.

GIBBONS, H. Climate of San Francisco.

LOGAN, T. M. Meteorological observations at Sacramento, California. HATCH, F. W. Meteorological observations at Sacramento, California.

FROEBEL, J. Remarks contributing to the physical geography of the North American Continent.

STRANG, J. J. Natural history of Beaver Island, Michigan.

EOFF, J. Habits of the black bass of the Ohio.

HEAD, J. E. Natural history of the country about Fort Ripley, Minn.

PARVIN, J. B. Habits of the gopher of Illinois.

MANN, C. Habits of a species of salamander.

Hoy, P. R. On the amblystoma luridum, a salamander inhabiting Wis.

CARLETON, J. H. Diary of an excursion in New Mexico.

BAIRD, S. F. Fishes on the coast of New Jersey and Long Island.

JACKSON, C. T. Catalogue of rocks, minerals, and ores collected on geological survey in Michigan.

LOCKE, J. Catalogue of rocks, minerals, ores, and fossils.

FOSTER, J. W. Catalogue of rocks, minerals, etc.

WHITNEY, J. D. Catalogue of rocks, minerals, etc.

OWEN, D. D. Catalogue of geological specimens.

BERLANDIER, L. Catalogue of collection of historical and geographical manuscripts, maps, etc.

HENRY, J. Circular respecting new report on libraries.

HENRY, J. Circular respecting copyrights.

76. Smithsonian Contributions to Knowledge. Vol. VII. 1855. 4to., pp. 260, 74 woodcuts, 72 plates, two maps.

CONTENTS.

CHAPPELSMITH, J. Tornado near New Harmony, Indiana. No. 59.

BAILEY, J. W. New species and localities of microscopic organisms. No. 63.

LAPHAM, I. A. Antiquities of Wisconsin. No. 70.

LEIDY, J. Extinct sloth tribe of North America. No. 72.

Publications of societies and periodicals in Smithsonian Library. Part I. No. 78.

77. Tenth Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1855. 34th Congress, 1st session, Senate Mis. Doc. 73. House of Representatives Mis. Doc. 113. 8vo., pp. 440, 79 woodcuts.

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of meteorological observers.

Correspondence:

HAMILTON COLLEGE, Clinton, N. Y. Examination of Spencer's Telescope.

77. Report for 1855-Continued.

AMERICAN ACADEMY OF ARTS AND SCIENCES, Boston, Mass. Thanks to Smithsonian Institution for Exchanges.

STONE, WM. J. On plaster casts of antique and modern statues, etc. ILLINOIS STATE BOARD OF EDUCATION. Meteorological system for every State.

BUTLER, A. P. Report of Senate Judiciary Committee on the Management of the Smithsonian Institution, Feb. 6, 1855.

HARVEY, W. H. Lecture on marine algæ.

Morris, J. G. Lecture on natural history as applied to farming and gardening.

Morris, J. G. Lecture on insect instincts and transformations.

CHACE, G. I. Lecture on oxygen and its combinations.

SMITH, J. L. Lecture on meteoric stones.

SNELL, E. S. Lecture on planetary disturbances.

LOGAN, T. M. On the climate of California.

Morris, O. W.; HENRY, J. Quantity of rain at different heights.

GUYOT, A.; HENRY J. Directions for meteorological observations.

HENRY, J. Earthquake directions.

HENRY, J. Aurora directions.

GREEN, J. Account of a new barometer.

HENRY, J. Registration of periodical phenomena.

MASTERMAN, S. Observations on thunder and lightning.

LETTERMAN, J. Sketch of the Navajo Indians.

CLINGMAN, T. L. Topography of Black Mountain, North Carolina.

ROBINSON, E.; LUDEWIG, H. E.; SQUIER, E. G.; MURPHEY, H. C.:

Hodgson, W. B.; IRVING, W.; PRESCOTT, W. H.; SPARKS, J.:

BANCROFT, G.; HAWKS, F. L. Communications relative to publication of Spanish works on New Mexico, by Buckingham Smith.

MULLER, J. Report on recent progress in physics-Galvanism.

78. Smithsonian Contributions to Knowledge. Vol. VIII. 1856. 4to., pp. 564, 52 woodcuts, 9 plates.

CONTENTS.

HAVEN, S. F. Archæology of the United States. No. 71.

OLMSTED, D. Recent secular period of aurora borealis. No. 81.

ALVORD, B. Tangencies of circles and of spheres. No. 80.

JONES, J. Chemical and physiological investigations relative to vertebrata. No. 82.

FORCE, P. Auroral phenomena in higher northern latitudes. No. 84. Publications of societies and periodicals in Smithsonian Library. Part II. No. 85.

- 79. New Tables for determining the Values of the Coefficients in the Perturbative Function of Planetary Motion, which depend upon the ratio of the mean distances. By JOHN D. RUNKLE. November, 1856. 4to., pp. 64. (S. C. IX.)
- 80. The Tangencies of Circles and of Spheres. By BENJAMIN ALVORD. January, 1856. 4to., pp. 16, 25 woodcuts, 9 plates of 20 figures. (S. C. viii.)

- 81. On the recent Secular Period of the Aurora Borealis. By DENISON OLMSTED. May, 1856. 4to., pp. 52. (S. C. VIII.)
- 82. Investigations, Chemical and Physiological, relative to certain American Vertebrata. By JOSEPH JONES. July, 1856. 4to., pp. 150, 27 woodcuts. (S. C. VIII.)
- 83. On the Relative Intensity of the Heat and Light of the Sun upon Different Latitudes of the Earth. By L. W. MEECH. November, 1856. 4to., pp. 58, 5 woodcuts, 6 plates of 9 figures. (S. C. IX.)
- 84. Record of Auroral Phenomena observed in the higher Northern Latitudes. By Peter Force. July, 1856. 4to., pp. 122. (S. C. VIII.)
- 85. Publications of Learned Societies and Periodicals in the Library of the Smithsonian Institution, Part II. May, 1856. 4to., pp. 38. (S. C. VIII.)
- 86. Observations on Mexican History and Archæology, with a special notice of Zapotec Remains, as delineated in Mr. J. G. Sawkins's drawings of Mitla, etc. By Brantz Mayer. November, 1856. 4to., pp. 36, 17 woodcuts, 4 plates of 6 figures. (S. C. ix.)
- 87. Psychrometrical Table for Determining the Elastic Force of Aqueous Vapor, and the Relative Humidity of the Atmosphere from indications of the Wet and the Dry Bulb Thermometer, Fahrenheit. By James H. Coffin. 1856. 8vo., pp. 20. (M. C. I.)
- 88. Researches on the Ammonia-cobalt Bases. By Wolcott Gibbs and Frederick Aug. Genth. December, 1856. 4to., pp. 72, 21 woodcuts. (S. C. ix.)
- 89. North American Oölogy. By Thomas M. Brewer. Part I. Raptores and Fissirostres. 1857. 4to., pp. 140, 5 plates of 193 figures. (S. C. xi.)
- 90. Illustrations of Surface Geology. By Edward Hitchcock. April, 1857. 4to., pp. 164, 2 woodcuts, 12 plates of 89 figures. (S. C. ix.)
- 91. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1856. 34th Congress, 3d session, Senate, No. 54, House of Representatives, Mis. Doc. No. 55. 1857. 8vo., pp. 468, 69 woodcuts.

Report of Prof. J. HENRY, and Proceedings of the Board.

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of meteorological stations and observers.

KOHL, J. G. On a collection of the charts and maps of America.

REID, D. B. Architecture in relation to ventilation, warming, lighting, fire-proofing, acoustics, and the general preservation of health.

91. Report for 1856-Continued.

No. 88.

HENRY, J. Syllabus of a course of lectures on physics. HENRY, J. Acoustics applied to public buildings.

BAIRD, S. F. Directions for collecting, preserving, and transporting specimens of natural history.

GILL, T. Fishes of New York.

Guest, W. E. Ancient Indian remains near Prescott, Canada West.

SHARPLESS, T., and PATTERSON, R. Phonography.

Institutions in which phonography is taught.

geology of Trinidad. BABBAGE, C. On tables of the constants of nature and art.

HENRY, J. On the mode of testing building materials, and an account

WALL, G. P., and SAWKINS, J. G. Report on the survey of the economic

of the marble used in the extension of the United States Capitol.

SMALLWOOD, C. Description of the observatory at St. Martin, Isle Jes. Canada East.

MEECH, L. W. Relative intensity of the heat and light of the sun. MULLER, J. Report on recent progress in physics-electricity.

92. Smithsonian Contributions to Knowledge. Vol. IX. 1857. 4to., pp. 480, 45 woodcuts, 22 plates.

CONTENTS.

MEECH, L. W. Intensity of heat and light of sun upon different latter tudes. No. 88. HITCHCOCK, E. Illustrations of surface geology. No. 90.

MAYER, B. Mexican history and archæology, and Zapotec remains.

GIBBS, W. and GENTH, F. A. Researches on ammonia-cobalt bases.

RUNKLE, J. D. New tables, planetary motion. No. 79.

- RUNKLE, J. D. Asteroid supplement to new tables. No. 94.
- 93. Smithsonian Meteorological Observations for the year 1855. (Printed for examination by the observers.) 1857. 8vo., pp. 118.
- 94. Asteroid Supplement to New Tables for determining the Values of b(i) and its derivatives. By John D. Runkle. May, 1857. 4to. pp. 72. (S. C. 1x.)
- 95. Nereis Boreali-Americana: or, Contributions to the History of the Marine Alge of North America. By WILLIAM HENRY HARVEY. Part III.—Chlorospermeæ. March, 1858. 4to., pp. 142, 14 plates of 44 figures. (S. C. x.)
- 96. Nereis Boreali-Americana: or, Contributions to a History of the Marine Algæ of North America. By WILLIAM HENRY HARVEY. Three parts in one vol., with 50 plates. May, 1858. 4to., pp. 568. [No. 32, 43, 95.]
- 97. Magnetical Observations in the Arctic Seas. By ELISHA KENT KANE Made during the Second Grinnell Expedition in search of Sir John

- Franklin, in 1853-'54-'55, at Van Rensselaer Harbor, and other points on the west coast of Greenland. Reduced and discussed by Charles A. Schott. 1859. 4to., pp. 72, 1 woodcut, 2 plates. (S. C. x.)
- 98. Grammar and Dictionary of the Yoruba Language. With an introductory description of the country and people of Yoruba. By T. J. Bowen. June, 1858. 4to., pp. 232, 1 map. (S. C. x.)
- 99. Smithsonian Contributions to Knowledge. Vol. X. 1858. 4to., pp. 462, 1 woodcut, 16 plates, 1 map.

HARVEY, W. H. Marine algo. Part III. Chlorospermess. No. 95. KANE, E. K. Magnetical observations in the Arctic seas. No. 97. BOWEN, T. J. Grammar and dictionary of the Yoruba language. No. 98.

- 100. An Account of the Total Eclipse of the Sun on September 7, 1858, as observed near Olmos, Peru. By J. M. Gilliss. April, 1859.
 4to., pp. 22, 1 woodcut, 1 plate. (S. C. xr.)
- 101. Map of the Solar Eclipse of March 15, 1858. By Thomas Hill. January, 1858. 8vo., pp. 8, 1 plate.
- 102. Catalogue of the described Diptera of North America. By R. OSTEN SACKEN. January, 1858. 8vo., pp. 116. October, 1859. (M. C. III.)
- 103. Meteorological Observations made at Providence, Rhode Island, from December, 1831, to May, 1860. By A. Caswell. October, 1860. 4to., pp. 188. (S. C. XII.)
- 104. Meteorological Observations in the Arctic Seas. By E. K. KANE. Made during the Second Grinnell Expedition in search of Sir John Franklin, in 1853-'54-'55, at Van Rensselaer Harbor, and other points on the west coast of Greenland. Reduced and discussed by Charles A. Schott. November, 1859. 4to., pp. 120, 10 woodcuts. (S. C. XI.)
- 105. Catalogue of North American Mammals, chiefly in the Museum of the Smithsonian Institution. By Spencer F. Baird. July, 1857. 4to., pp. 22.
- 106. Catalogue of North American Birds, chiefly in the Museum of the Smithsonian Institution. By Spencer F. Baird. October, 1858. 4to., pp. 42.
- 107. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1857. 35th Congress, 1st Session, Senate Mis. Doc. No. 272, House of Representatives, No. 135. 1858. 8vo., pp. 438, 100 woodcuts.

107. Report for 1857-Continued.

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

BAIRD, S. F. Report on publications, exchanges, museum, and experiences.

List of meteorological stations and observers.

STANLEY, J. M. Report of Committee of Regents on gallery of Indian portraits.

CONTAXAKI, MISS ELIZA B. Present of ornamental album from Greece. Chase, S. P. On telegraph.

GALE, L. D. On telegraph.

HALL, J. On telegraph.

Mason, C. On telegraph.

HENRY, J. On telegraph.

HENRY, J. On telegraph, deposition of, Sept. 1849.

HENRY, J. Communication relative to a publication by Prof. Morse.

Felton, C. C. Report of special committee of Board of Regents on the communication of Prof. Henry relative to the electro-magneta-

telegraph.

HENRY, J. History of the electro-magnetic telegraph.

LECONTE, J. Lecture on coal.

ALEXANDER, S. Lecture on vastness of the visible creation.

FENDLER, A. Meteorology and ethnology, Colonia Tovar, Venezuela. S. A.

LOGAN, T. M. Meteorology of Sacramento, California.

DEWEY, C. On best hours to find mean temperatures.

WISSNER, J. Meteorology of the District of Columbia.

MASTERMAN, S. Observations on natural phonomena, shooting staraurora, etc.

Müller, J. Report on recent progress in physics. (Electricity, garvanism.)

- 108. Catalogue of North American Birds, chiefly in the Museum of the Smithsonian Institution. By Spencer F. Baird. 1859. 8vo. pp. 24. (M. C. II.)
- 108*. Same Title, (printed for labelling, with one side of each leaf blank
- 109. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1858. 35th Congress, 2d Session, Senate Mis. Doc. No. 49. House Rep., No. 57. 1859. 8vo., pp. 448, 48 woodcuts.

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

Correspondence:

Schleiden, R. Free freight between Germany and United States to Smithsonian Institution by the North German Lloyd.

SABINE, E. Or continuance of magnetic observations.

BAIRD, S. F. Report on publications, exchanges, museum, and explora-

List of meteorological stations and observers.

CASWELL, A. Lecture on astronomy.

CUVIER, M. Memoir of Priestley.

BAIRD, S. F. Instructions for collecting nests and eggs of American birds.

BAIRD, S. F. Instructions for collecting insects.

100. Report for 1858—Continued.

LECONTE, J. L. Instructions for collecting coleoptera.

CLEMENS, B. Instructions for collecting hymenoptera.

UHLER, P. R. Instructions for collecting orthoptera.

UHLER, P. R. Instructions for collecting hemiptera.

UHLER, P. R. Instructions for collecting neuroptera.

LOEW, H.; OSTEN SACKEN, B. Instructions for collecting diptera.

CLEMBNS, B. Instructions for collecting lepidoptera.

TAYLOR, A. S. Grasshoppers and locusts of America.

MOTSCHULSKY, V. On mean's of destroying the grasshopper.

MARTINS, C. Vegetable colonization of the British Isles of Shetland, Faroe, and Iceland.

DECANDOLLE, A. Causes which limit vegetable species towards the north in Europe and similar regions.

COOPER, J. G. Distribution of the forests and trees of North America, and catalogue of the native trees of the United States.

BLACKISTON; BLAND; WILLIS, J. R. Birds of Nova Scotia.

BLAND; WILLIS J. R. Birds of Bermuda.

DUPREZ, M. F. Atmospheric electricity.

MÜLLER, J. Recent progress in physics. (Galvanism.)

HENRY, J. Meteorological stations, cost of establishment of.

HODGINS, J. G. Meteorological stations of Upper Canada.

DUDLEY, T. Earthquake at New Madrid, Missouri.

NAILL, D. W. Dispersion of a cloud by an electrical discharge.

HARE, R. Method of forming small weights.

FRIEDLÄNDER, J. Plan of a Bibliography.

Lyon, S. S. Antiquities from Kentucky.

GARDINER, R. H. Barometer, rain and snow gauges.

GUEST, W. E. Snow gauge.

GARDINER, R. H. Opening and closing of Kennebec river, Maine.

CANUDAS, A. Earthquakes in Guatemala.

Humphreys, A. A. Method of ascertaining the amount of water in rivers.

110. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1859. 36th Congress, 1st Session, House of Representatives, Mis. Doc. No. 90. 1860. 8vo., pp. 450, 57 woodcuts.

CONTENTS.

Report of Prof. J. HENRY, and Proceedings of the Board.

BAIRD, S. F. Report of publications, exchanges, museum, and explora-

List of meteorological stations and observers.

PEARCE, J. A. Notice of Richard Rush.

FELTON, C. C. Notice of W. W. Turner.

FELTON, C. C. Notice of Washington Irving.

BACHE, A. D. Notice of James P. Espy.

BACHE, A. D. Notice of G. Würdemann.

HENRY, J. Notice of Parker Cleaveland.

Correspondence:

DUKE OF NORTHUMBERLAND. Presentation of books.

2 HENRY, J. Account of Priestley's lens.

110. Report for 1859—Continued.

CUNARD, E. Free freight to Smithsonian Institution, between United States and England.

LOGAN, W. E. Request for duplicate shells.

Ross, B. R. Observations in Hudson's Bay Territory.

JOHNSON, S. W. Lectures on agricultural chemistry.

CABPENTER, P. P. Lectures on the shells of the Gulf of California.

MÆDLER, M. Movement of the stars around a central point.

DE LA RIVE, A. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1858, to June, 1859.

RETZIUS, A. Present state of ethnology in relation to the form of the human skull.

FLOURENS, M. Memoir of Pyramus de Candolle.

AIRY, G. B. On the means which will be available for correcting the measures of the sun's distance in the next twenty-five years.

Powell, B. Report on the state of knowledge of radiant heat, made to the British Association in 1882, 1840, and 1854.

HILGARD, J. E. Description of the magnetic observatory at the Smithsonian Institution.

Poggendorff, J. C. On the use of the galvanometer as a measuring instrument.

MALLET, R. On observations of earthquake phenomena.

CASELLA, L. Description of meteorological instruments.

GREEN, J.; WÜRDEMANN, W. On filling barometer tubes.

Welsh, J. The construction of a standard barometer, and apparatus and processes employed in the verification of barometers at the Kew Observatory.

111. Smithsonian Contributions to Knowledge. Vol. XI. 1859. 4to, pp. 502, 20 woodcute, 23 plates.

CONTENTS.

Brewer, T. M. North American Oölogy. Part 1. Raptores and Fissirostres. No. 89.

GILLINS, J. M. Total eclipse of the sun, September 7, 1858, in Peru. No. 100.

Bache, A. D. Magnetic and meteorological observations. Part I. No. 113.

KANE, E. K. Meteorological observations in the Arctic seas. No. 194 LECONTE, J. L. Coleoptera of Kansas and Eastern New Mexico. No. 196

SONNIAG, A. Observations on terrestrial magnetism in Mexico. No. 114. Loomis, E. On certain storms in Europe and America, December, 1836. No. 127.

112. Smithsonian Contributions to Knowledge. Vol. XII. 1860. 4to., pp. 538, 15 woodcuts, 3 plates.

CONTENTS.

KANE, E. K. Astronomical observations in the Arctic seas. No. 129.
WHITTLESEY, C. Fluctuations of level in North American lakes. No. 119.

- 112. Contributions to Knowledge. Vol. XII-Continued.
 - CASWELL, A. Meteorological observations at Providence, Rhode Island, 28½ years. No. 103.
 - SMITH, N. D. Meteorological observations near Washington, Arkansas, 20 years. No. 181.
 - MITCHELL, S. W. Researches upon venom of the rattlesnake. No. 135.
- 113. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, in 1840-'41-'42-'43-'44-'45. Part I. Investigation of the eleven year period in the amplitude of the solar-diurnal variation and of the disturbances of the magnetic declination, By A. D. BACHE. November, 1859. 4to., pp. 22, 5 woodcuts. (S. C. XI.)
- 114. Observations on Terrestrial Magnetism in Mexico. Conducted under the direction of Baron Von Müller, with notes and illustrations of an examination of the volcano Popocatepetl and its vicinity. By August Sonntag. February, 1860. 4to., pp. 92, 4 woodcuts, one plate. (S. C. XI.)
- 115. Proceedings of the Board of Regents of the Smithsonian Institution, in relation to the Electro-Magnetic Telegraph. 1861. 8vo., pp. 40, 7 woodcuts. (M. C. II.)
- 116. List of Public Libraries, Institutions, and Societies in the United States and British Provinces of North America. By WILLIAM J. RHEES. 1859. 8vo., pp. 84.
- 117. Catalogue of Publications of Societies and of other Periodical Works in the Library of the Smithsonian Institution, July 1, 1858. Foreign Works. 1859. 8vo., pp. 264. (M. C. III.)
- 118. Catalogue of the Described Lepidoptera of North America. By John G. Morris. May, 1860. 8vo., pp. 76. (M. C. III.)
- 119. On Fluctuations of Level in the North American Lakes. By Charles Whittlesey. July, 1860. 4to., pp. 28, 2 plates of 4 figures. (S. C. XII.)
- 120. Results of Meteorological Observations made at Marietta, Ohio, between 1826 and 1859, inclusive. By S. P. HILDRETH. To which are added results of observations taken at Marietta, by Joseph Wood, between 1817 and 1823. Reduced and discussed by Charles A. Schott. September, 1867. 4to., pp. 52, 14 woodcuts. (S. C. xvi.)
- 121. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, in 1840-'41'-42'-43-'44-'45. By A. D. BACHE. Part II.—Investigation of the solar-diurnal variation in the magnetic declination and its annual inequality. June, 1862, 4to., pp. 28, 8 woodcuts. (S. C. XIII.)

122. Smithsonian Miscellaneous Collections. Vol. I. 1862. 8vo., pp. 738. 23 woodcuts.

CONTENTS.

Directions for meteorological observations. No. 148. COFFIN, J. H. Psychrometrical tables. No. 87. Guyor, A. Meteorological and physical tables. No. 153.

123. Smithsonian Miscellaneous Collections. Vol. II. 1862. 8vo., pp. 714, 33 woodcuts.

CONTENTS.

BOOTH, J. C.; MORFIT, C. Recent improvements in chemical arts. No. 27.

Proceedings of Board of Regents in relation to the electro-magnetic telegraph. No. 115.

STANLEY, J. M. Catalogue of portraits of North American Indians. No. 53.

BAIRD, S. F. Catalogue of North American birds. No. 108.

BAIRD, S. F.; GIRARD, C. Catalogue of North American reptiles. Serpents. No. 49.

Check-list shells North America. No. 128.

Directions for collecting specimens of natural history. No. 34.

HENRY, J. Circular to officers Hudson Bay Company. No. 137.

Instructions for collecting nests and eggs. No. 189.

North American grasshoppers. No. 163.

North American shells. No. 176.

Morgan, L. H. Circular respecting relationship. No. 138.

124. Smithsonian Miscellaneous Collections. Vol. III. 1862. 8vo.. pp. 776, 49 woodcuts.

CONTENTS.

OSTEN SACKEN, R. Catalogue diptera North America. No. 102.

MORRIS, J. G. Catalogue described lepidoptera North America. No. 115.

LE CONTE, J. L. Classification coleoptera. I. No. 136.

Catalogue publications of societies in Smithsonian library. No. 117.

125. Smithsonian Miscellaneous Collections. Vol. IV. 1862. 8vo., pp. 762, 30 woodcuts.

CONTENTS.

HAGEN, H. Synopsis of North American neuroptera. No. 134.

MORRIS, J. G. Synopsis of North American lepidoptera. No. 133

- 126. The Coleoptera of Kansas and Eastern New Mexico. By John L Le Conte. December, 1859. 4to., pp. 64, 2 plates of 33 figures. 1 map. (S. C. xi.)
- 127. On Certain Storms in Europe and America, December, 1836. By ELIAS LOOMIS. February, 1860. 4to., pp. 28, 13 plates. (S. C. XI.

- 128. Check-lists of the Shells of North America. By Isaac Lea, P. P. Carpenter, Wm. Stimpson, W. G. Binney, and Temple Prime. June, 1860. 8vo., pp. 52. (M. C. 11.)
- 129. Astronomical Observations in the Arctic Seas. By E. K. KANE.

 Made during the Second Grinnell Expedition in search of Sir John
 Franklin, in 1853, 1854, and 1855, at Van Rensselaer Harbor, and
 other points in the vicinity of the northwest coast of Greenland.
 Reduced and discussed by Charles A. Schott. May, 1860. 4to.,
 pp. 56, 3 woodcuts, 1 map. (S. C. XII.)
- 130. Tidal Observations in the Arctic Seas. By E. K. KANE. Made during the Second Grinnell Expedition in search of Sir John Franklin, in 1853, 1854, and 1855, at Van Rensselaer Harbor. Reduced and discussed by Charles A. Schott. October, 1860. 4to., pp. 90, 3 woodcuts, 4 plates. (S. C. XIII.)
- 131. Meteorological Observations made near Washington, Arkansas, from 1840 to 1859, inclusive. By NATHAN D. SMITH. October, 1860. 4to., pp. 96. (S. C. XII.)
- 132. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, in 1840-'41-'42-'43-'44-'45. By A. D. BACHE. Part III.—Investigation of the influence of the moon on the magnetic declination. June, 1862. 4to., pp. 16, 3 woodcuts. (S. C. XIII.)
- 133. Synopsis of the Described Lepidoptera of North America. By John G. Morris. Part I.—Diurnal and crepuscular lepidoptera. February, 1862. 8vo., pp. 386, 30 woodcuts. (M. C. IV.)
- 134. Synopsis of the Neuroptera of North America. With a list of the South American species. By Hermann Hagen. July, 1861. 8vo., pp. 368. (M. C. IV.)
- 135. Researches upon the Venom of the Rattlesnake. With an investigation of the anatomy and physiology of the organs concerned. By S. Weir Mitchell. December, 1860. 4to., pp. 156, 12 woodcuts. (S. C. XII.)
- 136. Classification of the Coleoptera of North America. By John L. Le
 CONTE. Part I. March, 1862. 8vo., pp. 312, 49 woodcuts. (M. C. III.)
- 137. Circular to Officers of the Hudson's Bay Company. 1860. 8vo., pp.
 6. (Μ. С. π; Μ. С. viii.)
- 138. Circular in reference to the Degrees of Relationship among different Nations. By Lewis H. Morgan. January, 1860. 8vo., pp. 34. (M. C. 11.)

- 139. Instructions in reference to Collecting Nests and Eggs of North American Birds. January, 1860. 8vo., pp. 22, 20 woodcuts. (M. C. 11.)
- 140. List of the Coleoptera of North America. Prepared for the Smithsonian Institution. By John L. Le Conte. Part I. April, 1866. 8vo., pp. 82. (M. C. vi.)
- 141. Monographs of the Diptera of North America. By H. LOEW. Part I. Edited, with additions, by R. Osten Sacken. April, 1862. 8vo., pp. 246, 15 woodcuts, 2 plates of 42 figures. (M. C. vi.)
- 142. Bibliography of North American Conchology, previous to the year 1860. By W. G. BINNEY. Part I.—American Authors. March, 1863. 8vo., pp. 658. (M. C. v.)
- 143. Land and Fresh-Water Shells of North America. By W. G. BINNEY.
 Part II.—Pulmonata Limnophila and Thalassophila. September,
 1865. 8vo., pp. 172, 372 woodcuts. (M. C. vII.)
- 144. Land and Fresh-Water Shells of North America. By W. G. BINNEY. Part III.—Ampullariidæ, Valvatidæ, Viviparidæ, fresh-water Rissoidæ, Cyclophoridæ, Truncatellidæ, fresh-water Neritidæ, Helicinidæ. September, 1865. 8vo., pp. 128, 253 woodcuts. (M. C. vil.)
- 145. Monograph of American Corbiculadæ, (recent and fossil.) By TEMPLE PRIME. December, 1865. 8vo., pp. 92, 86 woodcuts. (M. C. VII.)
- 146. Meteorological Observations in the Arctic Seas. By Francis Leppold McClintock. Made on board the Arctic searching yacht "Fox," in Baffin Bay and Prince Regent's Inlet, in 1857, 1858, and 1859. Reduced and discussed by Charles A. Schott. May, 1862. 4to., pp. 164, 15 woodcuts, 1 map. (S. C. XIIL)
- 147. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1860. 36th Congress, 2d Session. Senate Mis. Doc., No. 21, 1861. 8vo., pp. 448, 73 woodcuts.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of meteorological stations and observers.

List of meteorological material contributed to Smithsonian Institution.

147. Report for 1860-Continued.

Correspondence:

CITIZENS OF PHILADELPHIA. Memorial relative to Lowe's aeronautic voyages across the Atlantic.

LEPSIUS, R. Presentation of books on Egypt.

MÜLLER, F. On Smithsonian exchanges.

HINCKS, WM. On Smithsonian exchanges.

HENRY, J. Reply to memorial on Lowe's aeronautic voyage.

HENRY, J. On currents of the atmosphere and aerial navigation.

ROGERS, F. Lectures on roads and bridges.

CARPENTER, P. P. Lectures on mollusca, or shell fish, and their allies.

Morlot, A. General views on archeology.

The microscope.

NICKLÉS, M. J. Scientific Congress of Carlsruhe, 1858.

CUVIER, M. Memoir of René Just Hauy.

SABINE, E. Magnetic storms.

GARDINER, R. H. Disappearance of ice.

FENDLER, A. Temperature of St. Louis, Missouri.

DEWEY, C. Best hours for temperature observations.

HENRY, J. Description of Smithsonian anemometer.

NEWTON, A. Suggestions for saving parts of the skeleton of birds.

Vollum, E. P. On the wingless grasshopper of California.

Wurdeman, G. Specimens of flamingo, etc., from South Florida.

GESNER, W. Habits of pouched rat, or salamander of Georgia.

BARNARD, V. Birds of Chester county, Pennsylvania.

COOPER. J. G. Forests and trees of Florida and the Mexican Boundary.

- 148. Directions for Meteorological Observations and the Registry of Periodical Phenomena. 1860. 8vo., pp. 72, 23 woodcuts. (M. C. I.)
- 149. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1861. 37th Congress, 2d Session, House of Representatives Mis. Doc., No. 77. 1862. 8vo., pp. 464, 25 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on publications, exchanges, museum, and explora-

List of meteorological stations and observers.

List of meteorological material contributed to Smithsonian Institution.

KING, C. B. Catalogue of engravings presented to the Smithsonian Institution.

Correspondence:

FLADGATE, CLARKE, and FINCH. Smithson's residuary legacy.

KUNHARDT & Co. Free freight to Smithsonian Institution, between United States and Germany.

TROYON, F. Lacustrian habitations.

MORLOT, A. Afchæology.

HILL, A. J. Map of antiquities of United States.

ROYAL ACADEMY OF SCIENCE, MADRID. Exchanges.

149. Report for 1861—Continued.

MEXICAN Soc. OF GEOG. AND STATISTICS. Exchanges.

HAGEN, H. Neuroptera.

HAMILTON COLLEGE, CLINTON, N. Y. Exchange of specimens.

Codd, J. A. Acknowledgment for books.

SHEPARD, C. U. Arrangement of mineralogical collection.

University of Tobonto. Scientific co-operation.

ROYAL HORTICULTURAL SOCIETY, LONDON. Exchange of publica-

BETHUNE, C. J. Acknowledgment for books.

PEASE, W. H. Acknowledgment for shells, etc. Circular of the Institute of Rupert's Land.

CRUMMELL, A: Facts respecting Liberia College.

GIBBS, G.; AND OTHERS. Recommendation of Shea's Indian .:guistics.

SHEA, J. G. Account of library of Indian linguistics.

WOOLSEY, T. D. Eulogy on Cornelius C. Felton.

Cox, S. S. Eulogy on Stephen A. Douglas.

Rogers, F. Lecture on bridges.

ALEXANDER, S. Lecture on the relations of time and space.

HAYES, I. I. Lecture on Arctic explorations.

FLOURENS, M. Memoir of Geoffroy Saint Hilaire.

LAUGEL, A. The sun, its chemical analysis.

LEE, DR. Progress of astronomical photography.

LESPIAULT, PROF. Small planets between Mars and Jupiter.

DUFOUR, C.; KÄMTZ. Scintillation of the stars.

DAUBRÉE, M. Mctamorphism and the formation of crystalline rocks.

CRAIG, B. F. Nitrification.

HUNT, T. S. History of petroleum or rock-oil.

ALLEN, Z. Explosibility of coal oils.

Destructive effect of iron-rust.

TROYON, F. Lacustrian cities of Switzerland.

RUTIMEYER, A. Fauna of Middle Europe during the Stone Age.

TROYON, F. Report on ethnological collections of Museum at Lausanne.

TROYON, F. Archæological researches made at Concise.

PEALE, T. R. Ancient mound in St. Louis, Missourt.

GIBBS, G. Instructions for archæological investigations

HENRY, J. Circular, ancient mining in Lake Superior Copper Region

Suggestions relative to ethnological map of Morgan, L. H. America.

Cours, E.; Prentiss, S. S. List of birds of the District of Columbia PRIZE QUESTIONS:

Holland Society of Science, Harlem.

Batavian Society of Experimental Philosophy of Rotterdam.

Society of Arts and Sciences of Utrecht.

Royal Academy of Netherlands.

150. Annual Report of the Board of Regents of the Smithsonian Institu tion, for the year 1862. 37th Congress, 3d session, House of Repre sentatives Mis. Doc., No. 25. 1863. 8vo., pp. 446, 94 woodcuts.

150. Report for 1862—Continued.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of meteorological stations and observers.

List of meteorological material contributed to Smithsonian Institution. Correspondence:

STANTON, E. M. Deposit of Beaufort library.

CANBY, E. Deposit of Bishop Johns' library.

SIBLEY, H. Meteorological telegrams.

ENGLEMANN, G. Heights of mountains in Colorado.

GUYOT, A. On mountain measurements.

HUBBARD, J. S. Investigations of Biela's comet.

DABLINGTON, W. History of Chester county, Pa.

LYMAN, THEODORE. Ethnographical collections.

DE SCHLAGINTWEIT, H. Ethnographical collections.

DAA, L. K. Ethnological Museum of Norway.

DILLE, I. Antiquities in Missouri and Tennessee.

REID, A. Skulls and mummy from Patagonia.

GIBBS, G. Philological circular.

GIBBS, G. Ethnological map of United States.

GIBBS, G. Indian vocabularies.

WATKINS, E. A. Systems of relationship of Cree Indians.

FLACHENECKER, G. Indian languages.

PEASE, W. H. Natural history of Pacific islands.

WIRTZ, H. R. Herbarium captured in Tennessee.

METTENIUS, G. Acknowledgment for ferns.

DE SAUSSURE, H. Hymenoptera.

ROMERO, M. Explorations of John Xantus in Mexico.

BACHE, A. D. Eulogy on James A. Pearce.

BARNARD, F. A. P. Lectures on the undulatory theory of light.

WILSON, D. Lectures on physical ethnology.

Morlor, A. Lecture on the study of high antiquity.

LUBBOCK, J. North American archæology.

FLOURENS, M. Historical sketch of the Academy of Science of Paris.

FLOURENS, M. Memoir of Leopold von Buch.

FLOURENS, M. Memoir of Louis Jacques Thenard.

QUATREFAGES, M. Memoir of Isidore Geoffroy St. Hilaire.

PHIPSON, T. L. The catalytic force, or studies on the phenomena of contact.

HERSCHEL, J. On atoms.

LESLEY, J. P. On the classification of books.

RIED, A. Human remains from Patagonia.

PRIZE QUESTIONS:

London Institution of Civil Engineers.

Provincial Society of Arts and Sciences, Utrecht.

Royal Danish Society of Science.

Academy of Sciences of the Institute of Bologna.

151. Smithsonian Contributions to Knowledge. Vol. XIII. 1863. 4to., pp. 558; 80 woodcuts, 4 plates, 3 maps.

ONTENTS.

KANE, E. K. Tidal observations, Arctic seas. No. 180.

McClintock, F. L. Meteorological observations, Arctic seas. No. 146.

WHITTLESEY, C. Ancient mining on shores of Lake Superior. No. 155.

BACHE, A. D. Discussion, Girard College observations. Part II. No. 121.

BACHE, A. D. Discussion, Girard College observations. Part III. No. 182.

Bache, A. D. Discussion, Girard College observations. Parts IV, V. VI. No. 162.

BACHE, A. D. Magnetic survey of Pennsylvania, &c. No. 166.

MITCHELL, S. W.; Morehouse, G. R. Researches upon anatomy and physiology of chelonia. No. 159.

- 152. Lectures on Mollusca or "Shell-fish," and their Allies. Prepared for the Smithsonian Institution. By Philip P. Carpenter. 1861. 8vo., pp. 140.
- 153. Tables, Meteorological and Physical, Prepared for the Smithsonian Institution. By Arnold Guyot. Third edition, revised and enlarged. 1859. 8vo., pp. 638. (M. C. I.)

CONTENTS.

- I. Thermometrical tables, A.
- II. Hygrometrical tables, B.
- III. Barometrical tables, C.IV. Hypsometrical tables, D.
- T. M. tarabarical assistance, 2.
- V. Meteorological corrections, E. VI. Miscellaneous tables, F.
- 154. List of Foreign Correspondents of the Smithsonian Institution. Cor-
- rected to January, 1862. May, 1862. 8vo., pp. 56. (M. C. v.)
- 155. Ancient Mining on the Shores of Lake Superior. By CHARLES WHITTLESEY. April, 1863. 4to., pp. 34, 30 woodcuts, 1 map. S. C. XIII.)
- 156. Catalogue of Minerals, with their Formulas, etc. By T. Eglestos. June, 1863. 8vo., pp. 56. (M. C. vil.)
- 157. Results of Meteorological Observations made under the direction of the United States Patent Office and the Smithsonian Institution, from the year 1854 to 1859, inclusive, being a Report of the Commissioner of Patents made at the first session of the 36th Congress. Vol. I. 1861. 36th Congress, 1st session, Senate Ex. Doc. 4to., pp. 1270.
- 158. Smithsonian Miscellaneous Collections. Vol. V. 1864. 8vo., pp. 774.

158. Miscellaneous Collections. Vol. V—Continued. CONTENTS.

- BINNEY, W. G. Bibliography of North American conchology. No. 142. Catalogue of publications of the Smithsonian Institution to June, 1862. No. 74.
- List of foreign correspondents of the Smithsonian Institution to January, 1862. No. 154.
- 159. Researches upon the Anatomy and Physiology of Respiration in the Chelonia. By S. Weir Mitchell and George R. Morehouse. April, 1863. 4to., pp. 50, 10 woodcuts. (S. C. XIII.)
- 160. Instructions for Research Relative to the Ethnology and Philology of America. By George Gibbs. March, 1863. 8vo., pp. 56. (M. C. VII.)
- 161. A Dictionary of the Chinook Jargon or Trade Language of Oregon. By George Gibbs. March, 1863. 8vo., pp. 60. (M. C. vii.)
- 162. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, in 1840-'41-'42-'43-'44-'45. Second section, comprising Parts IV, V, and VI. Horizontal force. Investigation of the eleven (or ten) year period and of the disturbances of the horizontal component of the magnetic force, with an investigation of the solar diurnal variation, and of the annual inequality of the horizontal force, and of the lunar effect on the same. By A. D. BACHE. November, 1862. 4to., pp. 78, 11 woodcuts. (S. C. XIII.)
- 163. Circular in Reference to the History of North American Grasshoppers. January, 1860. 8vo., pp. 4. (M. C. II.)
- 164. Smithsonian Museum Miscellanea. 1862. 8vo., pp. 88. (M. C. VIII.)

CONTENTS.

- Abbreviations of names of States and Territories of North America, for labelling insects, shells, &c.
- A series of small figures, from 1-1648.
- A series of medium figures, from 1-2747.
- A series of large figures, from 1-2599.
- Blank check-list of specimens.
- No. 5. consists of columns of figures from 1 to 1,000, and of two series, 25 and 50, to the 8vo. column. All these are stereotyped and printed with from one to eight columns on each page, with blank spaces of greater or less extent, as may be required.
- 165. Monograph of the Bats of North America. By H. Allen. June, 1864. 8vo., pp. 110, 73 woodcuts. (M. C. vii.)

- 166. Records and Results of a Magnetic Survey of Pennsylvania and parts of Adjacent States, in 1840 and 1841, with some additional records and Results of 1834, 1835, 1843 and 1862, and a map. By A. D. BACHE. October, 1863. 4to., pp. 88, 1 map. (S. C. XIII.)
- 167. New Species of North American Coleoptera. By John L. Le Conte. Part I. March, 1863. April, 1866. 8vo., pp. 180. (M. C. vi.)
- 168. Circular Relative to Collections of Birds from Middle and South America. December, 1863. 8vo., pp. 2. (M. C. VIII.)
- 169. Smithsonian Miscellaneous Collections. Vol. VI. 1867. 8vo., pp. 888. 15 woodcuts, 7 plates of 106 figures.

LOEW, H. Monograph of diptera. Part I. No. 141.

LOEW, H. Monograph of diptera. Part II. No. 171.

LE CONTE, J. L. List of coleoptera of North America. No. 140.

- LE CONTE, J. L. New species of North American coleoptera. No. 107.
- 170. Comparative Vocabulary. May, 1863. 4to., pp. 20.
- 171. Monographs of the Diptera of North America. By H. Loew.
 Part II. Edited by R. Osten Sacken. January, 1864. 8vo.,
 pp. 372, 5 plates of 44 figures. (M. C. vi.)
- 172. Palsontology of the Upper Missouri; a Report upon Collections made principally by the Expeditions under command of Lieutenant G. K. Warren, United States Topographical Engineers, in 1855 and 1856. Invertebrates. By F. B. MEEK and F. V. HAYDEN. Part I. April, 1865. 4to., pp. 158, 48 woodcuts, 5 plates of 41 figures. S. C. XIV.)
- 173. The Gray Substance of the Medulla Oblongata and Trapezium. By JOHN DEAN. February, 1864. 4to., pp. 80, 5 woodcuts, 16 plates of 65 figures. (S. C. xvi.)
- 174. Bibliography of North American Conchology previous to the year 1860. By W. G. BINNEY. Part II. Foreign authors. June, 1864. 8vo., pp. 302. (M. C. IX.)
- 175. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, in 1840, 1841, 1842. 1843, 1844 and 1845. Third section, comprising Parts VII, VIII. and IX. Vertical force. Investigations of the eleven (or ten) year period and of the disturbances of the vertical component of the magnetic force, and appendix on the magnetic effect of the aurora borealis; with an investigation of the solar diurnal variation, and of the

- annual inequality of the vertical force; and of the lunar effect on the vertical force, the inclination, and total force. By A. D. BACHE. April, 1864. 4to., pp. 72, 14 woodcuts. (S. C. xiv.)
- 176. Circular in Reference to Collecting North American Shells. January, 1860. 8vo., pp. 4. (M. C. π.)
- 177. Check-list of the Invertebrate Fossils of North America. Cretaceous and Jurassic. By F. B. Meek. April, 1864. 8vo., pp. 42. (M. C. VII.)
- 178. Circular to Entomologists. 1860. 8vo., pp. 2. (M. C. VIII.)
- 179. Catalogue of Publications of Societies and of Periodical Works, belonging to the Smithsonian Institution, January 1, 1866. 1866. 8vo., pp. 596. (M. C. ix.)
- 180. On the Construction of a Silvered Glass Telescope, fifteen and a half inches in aperture, and its use in Celestial Photography. By Henry Draper. July, 1864. 4to., pp. 60, 53 woodcuts. (S. C. xiv.)
- 181. Review of American Birds in the Museum of the Smithsonian Institution. By S. F. BAIRD. Part I.—North and Middle America. June, 1864—June, 1866. 8vo., pp. 484, 80 woodcuts. (M. C. XII.)
- 182. Results of Meteorological Observations made under the Direction of the United States Patent Office and the Smithsonian Institution, from the year 1854 to 1859, inclusive, being a Report of the Commissioner of Patents made at the first session of the 36th Congress. Vol. II. Part I. 36th Congress, 1st session. Senate Ex. Doc. 1864. 4to., pp. 546.
- 183. Check-list of the Invertebrate Fossils of North America. Miocene. By F. B. Meek. November, 1864. 8vo., pp. 34. (M. C. vii.)
- 184. Smithsonian Contributions to Knowledge. Vol. XIV. 1865. 4to., pp. 490, 158 woodcuts, 25 plates.

- BACHE, A. D. Discussion Girard College observations. Parts VII, VIII, IX. No. 175.
- BACHE, A. D. Discussion Girard College observations. Parts X, XI, XII. No. 186.
- DRAPER, H. Construction of silvered glass telescope and its use in celestial photography. No. 180.
- MEEK, F. B.; HAYDEN, F. V. Palsontology of the Upper Missouri. No. 172.
- LEIDY, J. Cretaceous reptiles of the United States. No. 192.

- 185. List of the Described Birds of Mexico, Central America, and the West Indies not in the Collection of the Smithsonian Institution. January 1, 1863. 8vo., pp. 8.
- 186. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, 1840-'41-'42-'43-'4-Fourth section, comprising Parts X, XI and XII. Dip and total force; analysis of the disturbances of the dip and total force; discussion of the solar diurnal variation and annual inequality of the dip and total force; and discussion of the absolute dip, with the final values for declination, dip and force between 1841 and 1845. By A. D. BACHE. April, 1865. 4to., pp. 42, 8 woodcuts. (S. C. XIV.)
- 187. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1863. 38th Congress, 1st Session. House of Representatives, Mis. Doc. No. 83. 1864. 8vo., pp. 420, 56 woodcuts. CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report of publications, exchanges, museum, and explorations.

List of meteorological stations and observers.

Meteorological material contributed to Smithsonian Institution.

Correspondence:

GOULD, B. A. Discussion of Piazzi's astronomical observations. PACKARD, F. A. Project of an outline history of education in the United States.

CHAMBER OF COMMERCE OF BORDEAUX. Exchange of publications. AGRICULTURAL ASSOCIATION OF MILAN. Exchange of publications. IRWIN, B. J. D. Tucson meteorite.

SANTIAGO AINSA. Tucson meteorite.

Poole, H. Cone-in-cone.

HUNGARIAN NATIONAL MUSEUM. Acknowledgment for birds. UNIVERSITY OF CHRISTIANIA, NORWAY. Ethnological specimens presented.

IMP. GEOLOG. INSTITUTE, VIENNA. Tertiary fossils presented. British Museum. Electrotypes of engravings of shells granted. FISHER, J. G. Acknowledgment of perrennibranchiates.

HAIDINGER, W. Honorary medal to Von Martius.

HUDSON'S BAY Co. Kennicott's explorations.

WHITNEY, W. D. Lectures on the principles of linguistic science. DEBEAUMONT, ELIE. Memoir of C. F. Bautemps-Beaupré.

ALEXANDER, C. A. Origin and history of the Royal Society of London. WETHERILL, C. M. Modern theory of chemical types.

DE LA RIVE, A. Phenomena which accompany the propagation of elatricity in highly rarefled elastic fluids.

MARCET, PROF. Report on Society of Physics and Natural History. Geneva, from July, 1862, to June, 1863.

187. Report for 1863-Continued.

PLATEAU, J. Experimental and theoretical researches on the figures of equilibrium of a liquid mass withdrawn from the action of gravity. Part I.

History of discovery relative to magnetism.

GAUTIER, PROF. Researches relative to the nebulæ.

MERINO, St. M. Figure of the earth.

ARAGO, F. Æronautic voyages, etc.

GLAISHER, JAS. Account of balloon ascensions.

BARGERT, JAC. Aboriginal inhabitants of the California peninsula.

JONES, J. M. Kjækken-mædding in Nova Scotia.

Morlot, A. Abstract of Dr. Keller's report on lacustrian settlements.

RAU, C. Agricultural implements of the North American stone period.

TROWBRIDGE, D. Ancient fort and burial ground in Tompkins county, New York.

KELLEY, O. H. Ancient town in Minnesota.

FOSTER, J. W. Ancient relics in Missouri.

DANILSEN, A. F. Mound in East Tennessee..

Purple dyeing, ancient and modern.

PEALE, T. R. Method of preserving lepidoptera.

FIGANIERRE, M. Account of remarkable accumulation of bats.

Tables of English and French weights and measures.

Table for conversion of centigrade degrees to Fahrenheit's scale.

Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1864.
 38th Congress, 2d Session. House of Representatives, Mis. Doc. 1865.
 8vo., pp. 450, 50 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

WEBB, T. W. Account of Prof. Henry Draper's telescope.

LEIDY, Jos. Review of "Cretaceous reptiles of United States."

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of foreign institutions from which donations to the library have been received in exchange. 1860-64.

List of meteorological stations and observers.

Meteorological material contributed to Smithsonian Institution.

WALLACH, R.; HENRY, J. Report of Committee of Regents in relation to the fire, January, 1865.

FOURIER, Jos. Memoir of Delambre.

DELAUNAY, M. Essay on the velocity of light.

WETHERILL, C. M. Ozone and antozone.

Jamin, J. Vegetation and the atmosphere.

BECQUEREL, M. Preservation of copper and iron in salt water.

Prescrvation of wood.

Caoutchouc and gutta percha.

VON KAROLYI; CRAIG, B. F. Products of the combustion of gun-cotton and gun-powder.

PETTENKOFER, Max. Apparatus for testing the results of perspiration and respiration.

LAMONT, J. The solar eclipse of July 18, 1860.

188 Report for 1864—Continued.

DEPRADOS, M. LE BARON. Eclipse of the sun, April 25, 1865.

DUBY, M. Report on the transactions of the Society of Physics and Natural History of Geneva, 1861.

DECANDOLLE, A. Report on the transactions of the Society of Physics and Natural History, 1862.

TROYON, FRED. On the crania helvetica.

PLATEAU, J. Experimental and theoretical researches on the figures of equilibrium of a liquid mass withdrawn from the action of gravity, etc. Parts II, III, IV.

RAU, C. Artificial shell deposits in New Jersey.

GIBBS, G. The intermixture of races.

BARGERT, J. Aboriginal inhabitants of the California peninsula. Part II. MORLOT, A. The study of high antiquity in Europe.

PRIZE QUESTIONS:

Holland Society of Sciences at Harlem.

Imperial Society of Natural Sciences of Cherbourg.

Royal Prussian Academy of Sciences.

Imperial Academy of Sciences at Vienna.

EXPLORATIONS:

Scientific expedition to Mexico. A report addressed to the Emperor of France by the Minister of Public Instruction.

KIRBY, W. W. A journey to the Yukon, Russian America.

FEILNER, J. Explorations in Upper California in 1860. •

Hoy, P. R. Journal of an exploration of Western Missouri in 1854. Tables of English and French weights and measures.

Table of chemical equivalents of sixty-three elements.

- 189. Catalogue of the Orthoptera of North America described previous to 1867. By SAMUEL H. SCUDDER. October, 1868. 8vo., pp. 110. (M. C. VIII.)
- 190. Queries Relative to Tornadoes. By JOSEPH HENRY. 8vo., pp. 4, 1 woodcut. (M. C. x.)
- 191. Smithsonian Miscellaneous Collections. Vol. VII. 1867. 8vo., pp. 878, 813 woodcuts.

CONTENTS.

ALLEN, H. Monograph of bats of North America. No. 165.

BINNEY, W. G. Land and fresh-water shells of North America. Part II. No. 143.

BINNEY, W. G. Land and fresh-water shells of North America. Part III. No. 144.

STIMPSON, W. Hydrobiinæ. No. 201.

PRIME, T. Monograph of American corbiculadse. No. 145.

CONRAD, T. A. Check-list of fossils—socene and oligocene. No. 200.

MEEK, F. B. Check-list of fossils-miocene. No. 188.

MEEK, F. B. Check-list of fossils—cretaceous and jurassic. No. 177.

EGLESTON, T. Catalogue of minerals. No. 156.

GIBBS, G. Dictionary of Chinook jargon. No. 161.

GIBBS, G. Instructions for ethnological and philological research. No. 160.

List of works published by the Smithsonian Institution to January. 1894. No. 208.

- 192. Cretaceous Reptiles of the United States. By JOSEPH LEIDY. May, 1865. 4to., pp. 142, 35 woodcuts, 20 plates of 309 figures. (S. C. XIV.)
- 193. Duplicate Shells Collected by the United States Exploring Expedition under Captain C. Wilkes. 8vo., pp. 4.
- 194. Land and Fresh-water Shells of North America. Part I.—Pulmonata geophila. By W. G. Binney and T. Bland. February, 1869. 8vo., pp. 328, 723 woodcuts. (M. C. VIII.)
- 195. Discussion of the Magnetic and Meteorological Observations made at the Girard College Observatory, Philadelphia, in 1840-'41-'42-'43-'44-'45. Parts I to XII inclusive. (Nos. 113^{xI}, 121^{xIII}, 132^{xIII}, 162^{xIII}, 175^{xIV}, 186^{xIV}.) 4to., pp. 262, 49 woodcuts.
- 196. Physical Observations in the Arctic Seas. By Isaac I. Haves. Reduced and discussed by Charles A. Schott. June, 1867. 4to., pp. 286, 15 woodcuts, 3 maps, 3 plates. (S. C. xv.)
- 197. On the Fresh-water Glacial Drift of the Northwestern States. By CHARLES WHITTLESEY. May, 1866. 4to., pp. 38, 14 woodcuts, 1 map, 1 plate. (S. C. xv.)
- 198. Physical Observations in the Arctic Seas. By ELISHA KENT KANE. Made during the Second Grinnell Expedition in Search of Sir John Franklin, in 1853, 1854, and 1855, at Van Rensselaer Harbor and other points on the West Coast of Greenland. Reduced and discussed by Charles A. Schott. Part I.—Magnetism. II.—Meteorology. III.—Astronomy. IV.—Tides. (Nos. 97^x, 104^{xi}, 129^{xii}, 130^{xiii}.) 1859-'60. 4to., pp. 340, 17 woodcuts, 1 map, 6 plates.
- 199. An Investigation of the Orbit of Neptune, with General Tables of its Motion. By Simon Newcomb. January, 1866. 4to., pp. 116. (S. C. xv.)
- 200. Check-list of the Invertebrate Fossils of North America, Eocene and Oligocene. By T. A. CONRAD. May, 1866. 8vo., pp. 46. (M. C. VII.)
- 201. Researches upon the Hydrobiinæ and Allied Forms; chiefly made upon materials in the Museum of the Smithsonian Institution. By WILLIAM STIMPSON. August, 1865. 8vo., pp. 64, 32 woodcuts. (M. C. VII.)
- 202. Geological Researches in China, Mongolia, and Japan, during the years 1862 to 1865. By RAPHAEL PUMPELLY. August, 1866. 4to., pp. 173, 18 woodcuts, 3 plates of 10 figures, 6 plates of 15 maps. (S. C. xv.)

- 203. List of Works published by the Smithsonian Institution. January, 1866. 8vo., pp. 12. (M. C. vii.)
- 204. Results of Meteorological Observations made at Brunswick, Maine, between 1807 and 1859. By PARKER CLEAVELAND. Reduced and discussed by Charles A. Schott. May, 1867. 4to., pp. 60, 8 woodcus. (S. C. xvi.)
- 205. Circular Relating to Collections in Archæology and Ethnology. By JOSEPH HENRY. January, 1867. 8vo., pp. 2. (M. C. VIII.)
- 206. Smithsonian Contributions to Knowledge. Vol. XV. 1867. 40., pp. 620, 47 woodcuts, 13 plates, 4 maps.

NEWCOMB, S. Orbit of Neptune. No. 199.

WHITTLESEY, C. Fresh-water glacial drift of Northwestern States. No. 197.

Pumpelly, R. Geological researches in China, Mongolia, and Japar. No. 202.

HAYES, I. I. Physical observations in the Arctic seas. No. 196.

- 207. Suggestions Relative to Objects of Scientific Investigation in Russian America. By Joseph Henry. May, 1867. 8vo., pp. 10. 1. M. C. VIII.)
- 208. The Gliddon Mummy Case in the Museum of the Smithsonian Institution. By CHARLES PICKERING. June, 1867. 4to., pp. 6, 1 plate. S. C. xvi.)
- 209. Annual Report of the Board of Regents of the Smithsonian Indiatution, for the year 1865. 39th Congress, 1st Session, House of Representatives Mis. Doc. No. 102. 1866. 8vo., pp. 496, 139 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS. Proceedings of

BAIRD, S. F. Report of publications, exchanges, museum, and explorations.

List of addresses of foreign institutions since 1862.

List of meteorological stations and observers.

List of meteorological material contributed.

Correspondence:

HENRY, J. Letter to Secretary of Treasury on payment of interest in coin.

LEIDY, J.; TRYON, G. W. Report on shells presented to Academ. of Natural Sciences.

MEXICAN Soc. of GEOG. AND STATISTICS. Exchange system.

209. Report for 1865—Continued.

EVANS, J. Climate of Colorado.

GIVEN, W. F. Remarkable electric phenomenon.

HALE, C. Explorations of the Nile.

MINING DEPT., MELBOURNE. Exchange system.

CARLETON, J. H. Meteorites in Mexico.

CONRAD, T. A. Chalk found in United States.

REGENTS OF UNIVERSITY OF STATE OF NEW YORK. Acknowledgment for specimens.

McMasters, S. Y. Language of Navajos said to resemble the Welsh.

GIBBS, G. Indian languages.

LISBOA, M. M. Books on Brazil presented.

BRAZILIAN NAT. HIST. MUSEUM. Specimens from Brazil presented.

BERN MUSEUM. Request for Bison.

IMPERIAL LIBRARY OF VIENNA. Books presented.

BRUCE, F. W. A., British Minister. Introducing and recommending Dr. Berendt.

DE IRISARRI, A. T., Guatemalan Minister. Introducing and recommending Dr. Berendt.

Molina, L., Costa Rican Minister. Introducing and recommending Dr. Berendt.

Rosing, J. Exchange system.

GOVERNMENT OF BREMEN. Exchange system.

Hodgins, J. G. Metebrological system of Canada.

BRITISH MUSEUM. Acknowledgment of specimens.

TORONTO UNIVERSITY. Objects of the Museum.

PETITOT, E. Account of the Indians of British America.

Act of Congress to transfer Smithsonian Library to Library of Congress.

BARNARD, J. G. Eulogy on Gen. Jos. G. Totten.

FLOURENS, M. Memoir of Ducrotay de Blainville.

CHOSSAT, DR. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1863, to June, 1864.

PLANTAMOUR, M. E. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1864, to June, 1865.

LOOMIS, E. Aurora Borealis, or Polar Light; its phenomena and laws. The Senses. Sense of feeling; sense of smell.

MATTEUCCI, C. Lectures on electro-physiology.

DESOR, E. Palafittes or Lacustrian constructions of the Lake of Neuchatel.

PLATEAU, J. Experimental and theoretical researches on the figures of equilibrium of a liquid mass withdrawn from the action of gravity, etc. Part V.

LILLJEBORG, W. Outline of a systematic review of the classification of birds.

Prize Questions:

Royal Danish Society of Sciences.

Imperial Academy of Sciences of Vienna.

Pontifical Academy of the Nuovi Lincei.

Royal Scientific and Literary Institute of Lombardy.

Imperial Society of Science, Agriculture, and Arts of Lille.

Dunkirk Society for the encouragement of sciences, letters, and arts.

NEWTON, H. A. Metric system of weights and measures, with tables.

- 210. Arrangement of Families of Birds, adopted provisionally by the Smithsonian Institution. By S. F. BAIRD. June, 1866. 8vo., pp. 8. (M. C. viii.)
- 211. Smithsonian Contributions to Knowledge. Vol. XVI. 1870. 4to., pp. 498, 76 woodcuts, 18 plates.

DEAN, J. Gray substance of the medulla oblongata and trapezium. No. 178.

CLEAVELAND, P. Meteorological observations, Brunswick, Maine, 53 years. No. 204.

HILDRETH, S. P. Meteorological observations, Marietta, O. No. 120.

PICKERING, C. Gliddon mummy case. No. 208.

COFFIN, J. H. Orbit and phenomena of a meteoric fire ball. No. 221.

GOULD, B. A. Transatlantic longitude. No. 223. SWAN, J. G. Indians of Cape Flattery. No. 220.

212. Smithsonian Miscellaneous Collections. Vol. VIII. 1869. 8vo., pp. 921, 730 woodcuts, 4 plates.

CONTENTS.

OSTEN SACKEN, R. Monograph of the diptera of North America. Part IV. No. 219.

SCUDDER, S. H. Catalogue of the orthoptera of North America. No. 189.

BINNEY, W. G.; BLAND, T. Land and fresh-water shells of North America. Part I. No. 194.

BAIRD, S. F. Arrangement of families of birds. .No. 210.

HENRY, J. Circular to officers of the Hudson's Bay Company. No. 137. HENRY, J.; GIBBS, G.; BAIRD, S. F. Suggestions relative to scientific

investigations in Russian America. No. 207.

HENRY, J. Circular relative to archeology and ethnology. No. 205.

HENRY, J. Circular to entomologists. No. 178.

HENRY, J. Circular relative to collections of birds from Middle and South America. No. 168.

BAIRD, S. F. Smithsonian Museum miscellanea. No. 164.

213. Smithsonian Miscellaneous Collections. Vol. IX. 1869. 8vo., pp. 914.

BINNEY, W. G. Bibliography of North American conchology. Foreign authors. Part II. No. 174.

Catalogue of publications of societies and of periodicals in Smithsonian Library, 1866. No. 179.

214. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1866. 39th Congress, 2d Session, House of Representatives, Mis. Doc. No. 83. 1867. 8vo., pp. 470, 70 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations. BOARD OF REGENTS. Proceedings of

214. Report for 1866—Continued.

BAIRD, S. F. Report on publications, exchanges, museum, and explorations.

List of meteorological stations and observers.

Act of Congress to receive into the United States Treasury the residuary legacy of Smithson, etc.

Memorial of Board of Regents to Congress relative to the fund.

HENRY, J. Memoir of W. W. Seaton.

Report on system of accounts.

FLOURENS, M. Memoir of Magendie.

THE SENSES. Sense of taste; sense of hearing; sense of sight.

Huggins, W. Results of spectrum analysis applied to the heavenly bodies. External appearance of the sun's disk.

MOIGNO, ABBE. Accidental or subjective colors: Persistence of images, contrast, irradiation, daltonism, etc.

PLATEAU, J. Figures of equilibrium of a liquid mass withdrawn from the action of gravity. Part VI.

Gosse, Dr. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1865, to June, 1866.

GIBBS, G. Notes on the Tinneh or Chepewyan Indians of British and Russian America.
1. The Eastern Tinneh, by Bernard R. Ross, Esq.
2. The Loucheux Indians, by William L. Hardisty.
3. The Kutchin Tribes, by Strachan Jones, Esq.

Von Hellwald, F. The American migration, with notes by Prof. Henry.

RAU, C. Indian pottery.

BRINTON, D. G. Artificial shell deposits of the United States.

DILLE, I. Sketch of ancient earth-works of Ohio.

Pile-work antiquities of Olmutz.

ESTES, L. C. Antiquities on the banks of the Mississippi river and Lake Pepin.

GIBBS, G. Physical Atlas of North America.

DAVIS, E. H. On ethnological research.

SCHERZER; SCHWARZ. Table of anthropological measurements.

International Archeological Congress, organized by the Archeological Academy of Belgium, in concert with the French Society of Archeology. Antwerp, 1866.

HIGGINS, H. H. On vitality, with notes by Prof. Henry.

LEWIS, J. Instructions for collecting land and fresh-water shells.

WOOD, H. C. Instructions for collecting myriapods, phalangidse, etc.

WETHERILL, C. M. Plan of a research upon the atmosphere.

LEWIS; QUALE. Account of the cryolite of Greenland.

EXTEACTS FROM THE METEOROLOGICAL CORRESPONDENCE OF THE INSTI-TUTION.—

HENRY, J. Remarks by

DENNIS, W. C. Evaporation in Florida.

DENNIS, W. C. Fresh-water in the ocean.

DEWEY, C. The winds.

BALTZELL, J. Winds in Florida.

WARD, L. F. Barometer tube breaking suddenly.

Andrews, S. L. Meteors.

214. Report for 1866—Continued.

BALFOUR, J. The wind and fog.

HUNTINGTON, G. C. Climate of Kelley's Island, Ohio.

MALLINIKROOT, C. Changes of wind.

CAPEN, F. L. Meteorological discovery.

FENDLER, A. Meteorology of Colonia Towar.

HILLIER, S. L. Effect of lightning.

BANNISTER, H. M. Formation of clouds over Gulf stream.

Bannsster, H. M. Climate of Alaska.

VAILLANT, M. Horary variations of the barometer, with notes by Prof. Henry.

ENGLEHARDT, M. Formation of ice at the bottom of water.

SARTORIUS, C. Earthquake in Eastern Mexico, January, 1866. DREUTZER, O. E. Statistics relative to Norwegian mountains, lakes, and

215. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1867. 40th Congress, 2d Session, Senate Mis

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD of REGENTS. Proceedings of

BAIRD, S. F. Report of exchanges and museum.

Doc. No. 86. 1868. 8vo., pp. 506, 10 woodcuts.

BAIRD, S. F. List of expeditions and other sources from which the specmens in the Government Museum have been mainly derived.

List of meteorological stations and observers.

Meteorological material contributed in addition to the regular observation. Cluss, A. Report of the architect.

Correspondence:

the snow-line.

AMER. ACAD. OF ARTS AND SCIENCES. Exchange system.

KNIGHT, G. H. New system of weights and measures with 8 as the metrical number.

Bolles, E. C. Account of Portland Society of Natural History.

BATH AND WEST OF ENG. Soc. FOR ENCOUR. OF AGRIC., ARTS, ETC. Exchange system.

HAMBURG ZOOL. GARDENS. Exchange of specimens.

CHICAGO ACAD. OF SCIENCES. Acknowledgment for specimens.

TAYLOR, W. B. Report on improved system of numeration.

ST. PETERSBURG ACAD. OF SCIENCES. Exchange system.

ZISGENBALS, H. Schlagintweit ethnographic collection.

LLOYD, W. A. Sparrows sent to United States.

LLOYD, W. A. Exchange of specimens.

MUNEUM OF NAT. UNIV. OF GREECE. Exchange of specimens. LABOULAYE, M. Acknowledgment for books on education.

ACAD. OF SCIENCES, STOCKHOLM. Acknowledgment of birds. GOULD, J. Acknowledgment of birds.

MAYBERRY, S. P. Gradual approach of sea upon land. UNIVERSITY OF COSTA RICA. Exchange of specimens.

Guild, R. A. Biographical notice of Chas. C. Jewett.

GRAY, A, Biographical notice of W. H. Harvey.

AGASSIZ, L. Report on use of new hall in Smithsonian Building.

215. Report for 1867—Continued.

Memorial of Regents to Congress asking appropriations for Museum.

DEBEAUMONT, E. Memoir of Legendre.

PELTIER, F. A. Memoir of J. C. A. Peltier.

Scientific researches of Peltier.

MAILLY, E. History of Royal Institution of Great Britain.

DE LA RIVE, PROF. Michael Faraday, his life and works.

FLOUBERS, M. The Jussieus and the natural method.

MAREY, M. Natural history of organized bodies.

MATTEUCCI, C. On the electrical currents of the earth.

Considerations on electricity.

Electricity—Account of lightning discharges, with notes by Prof. Henry.

G. W. Dodge, O. M. Poe, J. C. Cresson, H. Haas, H. J. Kron, B. F. Mudge, New Haven Journal, S. D. Martin, G. Wright, C. G. Boerner, W. S. Gilman, C. W. Dean, J. A. Osborne.

DARWIN, C. Queries about expression for anthropological inquiry.

PETTIGREW, J. B. Modes of flight in relation to aeronautics.

Man as the contemporary of the mammoth and reindeer in Middle Europe. Jamin, M. Photo-chemistry.

ABBE, C. Description of the observatories at Dorpat and Poulkova.

TYLOR, E. B. On traces of the early mental condition of man.

ETHNOLOGY:

GUNN, D. Indian remains, Red River Settlement, Hudson's Bay Territory.

READ, M. C. Ancient mound near Chattanooga, Tenn.

PIDGEON, W. Ancient burial mound in Indiana.

BERTHOUD, E. L. Ancient remains in Colorado.

AGNEW, S. A. Mounds in Mississippi.

WHITNEY, J. D. Cave in Calaveras county, California.

Ethnological Department of the French Exposition, 1867.

HAYDEN, F. V. Notes on Indian history.

MEIGS, J. A. Description of a human skull from Rock Bluff, Ill.

KABIS, M. Introduction to the study of the Coptic language.

SMART, C. Notes on the Tonto Apaches.

BERENDT, C. H. Explorations in Central America.

Gunn, D. Notes of an egging expedition to Shoal Lake, Lake Winnipeg. ROTHROCK, J. T. Sketch of the flora of Alaska.

METEOROLOGY:

Hurricane in the island of St. Thomas.

LATIMER, G. A. Earthquakes in St. Thomas.

Maritime disasters of the Antilles.

Dickinson, A. B. Eruption of a volcano in Nicaragua.

Young, W. J. Cloud-bursts.

WOODWORTH, A. Meteorite in Mexico.

SIMSON, R. Meteorite in Mexico.

ERNST, G. A. Meteorology of Caracas, South America.

TUCKETT, F. F. On barometer tables.

LATIMER, G. A. Great hurricane at Tortola, St. Thomas, and Porto Rico.

PIKE, N. Cyclone in the Indian Ocean.

215. Report for 1867—Continued.

PRIZE QUESTIONS:

Royal Danish Society of Sciences.

Pontifical Academy of the Nuovi Lincei.

DE LA RUE, W. Abbreviations used in England, 1867.

- 216. List of Photographic Portraits of North American Indians in the Gallery of the Smithsonian Institution. 1867. 8vo., pp. 42. (M. C. xiv.)
- 217. Letter by M. Hoek in reference to the Meteoric Shower of November 13th, 1867. 8vo., pp. 4.
- 218. Systems of Consanguinity and Affinity of the Human Family. By Lewis H. Morgan. 1869. 4to., pp. 616, 6 woodcuts, 14 plates. (S. C. xvii.)
- 219. Monographs of the Diptera of North America. By R. OSTEN SACKEN. Part IV. 1869. 8vo., pp. 358, 7 woodcuts, 4 plates of 92 figures. (M. C. VIII.)
- 220. The Indians of Cape Flattery, at the entrance to the Strait of Fuca. Washington Territory. By James G. Swan. March, 1870. 4to., pp. 118, 44 woodcuts. (S. C. xvi.)
- 221. The Orbit and Phenomena of a Meteoric Fire Ball, seen July 20, 1860. By James H. Coffin. May, 1869. pp. 56, 3 woodcuts, 1 map. (S. C. xvi.)
- 222. Tables and Results of the Precipitation, in Rain and Snow, in the United States, and at some stations in adjacent parts of North America, and in Central and South America. By Charles A. Schott. March, 1872. 4to., pp. 178, 8 woodcuts, 5 plates, and 3 double charts. (S. C. XVIII.)
- 223. The Transatlantic Longitude, as determined by the Coast Survey Expedition of 1866. By Benjamin Apthorp Gould. October, 1869. 4to., pp. 110, 2 woodcuts. (S. C. xvi.)
- 224. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1868. 40th Congress, 3d Session. House of Representatives, Ex. Doc. No. 83. 1869. 8vo., pp. 474, 40 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

List of meteorological stations and observers.

List of meteorological material contributed to Smithsonian Institution.

DELAFIELD, R. Report of Committee of Regents on Washington canal.

FLOURENS, M. Memoir of Cuvier.

FLOURENS, M. History of the works of Cuvier.

DE BEAUMONT, E. Memoir of Oersted.

Notice of Christian Frederic Schoenbein, the discoverer of ozone.

234. Report for 1868—Continued.

HENRY, J. Appendix to notice of Schoenbein.

HAGEN, G. Memoir of Encke.

RAWSON, R. Memoir of Eaton Hodgkinson.

CAZIN, A. Recent progress in relation to the theory of heat.

MULLER, J. Principles of the mechanical theory of heat.

MAGRINI, L. Continuous vibratory movement of all matter, ponderable and imponderable.

TYNDALL, J. Radiation.

DAUBRÉE, M. Synthetic experiments relative to meteorites.

BRUSH, G. J. Catalogue of meteorites in the mineralogical collection of Yale College.

DE SAUSSURE, H. Observations on the electric resonance of mountains.

STEWART, B. Experiments on aneroid barometers made at the Kew Observatory.

ELLERY, R. L. J. Address of the president of the Royal Society of Victoria.

WARTMANN, E. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1867, to June, 1868.

BROCA, P. History of the transactions of the Anthropological Society of Paris, from 1865 to 1867.

RAU, C. Drilling in stone without metal.

RAU, C. Agricultural flint implements in southern Illinois.

Notice of the Blackmore Museum, Salisbury, England.

PRIZE QUESTIONS:

Holland Society of Sciences of Harlem.

Imperial Academy of Sciences, Belles Lettres, and Arts, of Bordeaux.

Pollock, J. Assay of gold and silver coins at the Mint of the United States.

Table of foreign gold and silver coins.

List of publications of the Smithsonian Institution up to July, 1869, with systematic and alphabetical index.

- 225. List of Foreign Correspondents of the Smithsonian Institution; corrected to January, 1870. April, 1870. 8vo., pp. 56.
- 226. A List of the Smithsonian Publications, from 1846 to 1869. November, 1869. 8vo., pp. 34.
- 227. Arrangement of the Families of Mollusks. By THEODORE GILL. February, 1871. 8vo., pp. 65. (M. C. x.)
- 228. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1869. 41st Congress, 3d Session. House of Representatives, Ex. Doc. No. 153. 1871. 8vo., pp. 430, 38 woodcuts, 1 map.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

List of meteorological stations and observers.

List of meteorological material contributed to Smithsonian Institution.

228. Report for 1869-Continued.

BERTHRAND, M. Kepler: his life and works.

ARAGO, M. Eulogy on Thomas Young.

DE BEAUMONT, E. Memoir of Auguste Bravais.

RAU, C. Memoir of C. F. P. Von Martius.

MATTEUCCI, C. Life and scientific labors of Stefano Marianini.

HUNT, T. S. Chemistry of the earth.

MATTEUCCI, C. Electrical currents of the earth.

MAREY, M. Phenomena of flight in the animal kingdom.

BABINET, M. The northern seas.

LOMBARD, H. C. Report on the transactions of the Society of Physics and of Natural History of Geneva, July, 1868, to June, 1869.

SIMPSON, J. H. Coronado's march in search of the "Seven Cities of Cibola."

LUBBOCK, J. Social and religious condition of the lower races of man.

Huxley, T. H. Principles and methods of palseontology.

SCHOTT, A. Remarks on the "Cara Gigantesca" of Yzamal, in Yucatantan.

BECQUEREL, M. Forests and their climatic influence.

Brenndecke, F. Meteorites.

ABICH, S. Remarkable forms of hail-stones in Georgia.

Sartorius, C. Eruption of the volcano of Colima.

229. Smithsonian Contributions to Knowledge. Vol. XVII. 1871. 4to., pp. 616, 6 woodcuts, 14 plates.

CONTENTS.

MORGAN. Systems of consanguinity and affinity of the human family. No. 218.

- 230. Arrangement of the Families of Mammals. By THEODORE GILL. November, 1872. 8vo., pp. 104. (M. C. xi.)
- 231. Memoranda of Inquiry relative to the Food Fishes of the United States. By S. F. BAIRD. 1871. 8vo., pp. 8. (M. C. x.)
- 232. The Secular Variations of the Elements of the Orbits of the Eight Principal Planets, Mercury, Venus, the Earth, Mars, Jupiter, Saturn. Uranus, and Neptune, with tables of the same; together with the obliquity of the ecliptic and the precession of the equinoxes in both longitude and right ascension. By John N. Stockwell. 1872. 4to., pp. 220. (S. C. XVIII.)
- 233. Converging Series expressing the Ratio between the Diameter and the Circumference of a Circle. By WM. FERREL. April, 1871. 4to., pp. 6. (S. C. xvIII.)
- 234. Questions Relative to the Food Fishes of the United States. By S. F. BAIRD. 1871. 8vo., pp. 14. (M. C. x.)
- 235. Instructions for Observations of Thunder Storms. By JOSEPH HENRY. 1871. 8vo., p. 1. (M. C. x.)

- 236. Circular Relative to Heights. By JOSEPH HENRY. 1871. 8vo., pp. 2. (M. C. x.)
- 237. Directions for constructing Lightning Rods. By JOSEPH HENRY. 1871. 8vo., pp. 4. (M. C. x.)
- 238. List of the Institutions, Libraries, Colleges, and other Establishments in the United States in correspondence with the Smithsonian Institution. By Wm. J. Rhees. July, 1872. 8vo., pp. 256. (M. C. x.)
- 239. Observations on Terrestrial Magnetism, and on the Deviations of the Compasses of the United States Iron Clad Monadnock during her cruise from Philadelphia to San Francisco, in 1865 and 1866. By WM. HARKNESS. 1872. 4to., pp. 226, 2 woodcuts. (S. C. XVIII.)
- 240. Problems of Rotary Motion presented by the Gyroscope, The Precession of the Equinoxes, and The Pendulum. By J. G. BARNARD. 1872. 4to., pp. 56, 6 woodcuts. (S. C. xix.)
- 241. A Contribution to the History of the Fresh Water Algo of North America. By Horatio C. Wood, Jr. 1872. 4to., pp. 272, 21 plates of 340 figures. (S. C. xix.)
- 242. Lucernariæ and their Allies. A Memoir on the Anatomy and Physiology of Haliclystus Auricula and other Lucernarians, with a Discussion of their Relations to other Acalephæ, to Beroids and Polypi. By H. J. CLARK. 1878. 4to., pp. 138, 11 plates of 149 figures. (S. C. xx11.)
- 243. List of Foreign Correspondents of the Smithsonian Institution; corrected to January, 1872. 1872. 8vo., pp. 66. (M. C. x.)
- 244. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1870. 42d Congress, 1st Session. House of Representatives, Ex. Doc. No. 20. 1871. 8vo., pp. 494, 28 woodcuts.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

List of meteorological stations and observers.

List of meteorological material contributed to Smithsonian Institution.

HENRY, J. Eulogy on Alexander Dallas Bache.

BACHE, A. D. Lecture on Switzerland.

HENRY, J. On a physical observatory.

ARAGO, M. The History of my Youth, an autobiography.

ARAGO, M. Eulogy on Herschel.

Life and labors of Henry Gustavus Magnus.

Anderson, M. B. Life of Prof. Chester Dewey.

TAYLOR, W. B. Thoughts on the nature and origin of force.

244. Report for 1870-Continued.

VON LIEBIG, J. Induction and deduction.

HAUGHTON, S. Relation of food to work, and its bearing on medicapractice.

REYNOLDS, J. E. Hydrogen as gas and as a metal.

WISEMAN, N. Identification of the artisan and artist.

BABINET, M. The diamond and other precious stones.

ETHNOLOGY:

GIBBS, G. On the language of the aboriginal Indians of America.

WILLIAMS, H. C. On antiquities in some of the Southern States.

GARDNER, W. H. Ethnology of the Indians of the valley of the Red River of the North.

FINCK, H. Account of antiquities in the State of Vers Cruz, Mexic

DUNNING, E. O. Account of antiquities in Tennessee.

STEPHENSON, M. F. Account of ancient mounds in Georgia.

DAYTON, E. A. Explorations in Tennessee.

HARWOOD, A. A. Account of the sarcophagus in the National Museum, now in charge of the Smithsonian Institution.

GRANT, E. M. Account of the discovery of a stone image in Tenresee, now in possession of the Smithsonian Institution.

BLYDEN, E. D. On mixed races in Liberia.

FOWLER, J. On shell-heaps.

PEALE, T. R. On the uses of the brain and marrow of anima.s among the Indians of North America.

Lyon, S. S. Report of an exploration of ancient mounds in Uniccounty, Kentucky.

BARRANDT, A. Sketch of ancient earthworks on the Upper Missour.

STELLE, J. P. Account of aboriginal ruins at Savannah, Tennesed Stelle, J. P. Account of aboriginal ruins in Hardin Co., Tennesed

TERRESTRIAL PHYSICS:

CAMPBELL, J. V. The earthquake in Peru, August 13, 1868.

PALMIERI, Prof. The electro-magnetic seismograph.

JOHNSON, W. W. On the distribution of forest trees in Montana. Idaho, and Washington.

SARGENT, W. D. Influence of the aurora on the telegraph.

METEOROLOGY:

Poëy, A. New classification of clouds.

TACCHINI, P. On the evaporation observed at Palermo, in 1865 and 1866.

ZANTEDESCHI, F. On the electricity of induction in the zerial strate of the atmosphere, which, in the shape of a ring, surround a cloud that is resolving into rain, snow, or hail.

PALMIERI, Prof. On the presence of electricity during the fall of rain.

ELLIOTT, R. S. Climate of Kansas.

PORTER, Commodore. Account of a hail storm on the Bosphorus. BACHE, G. M. Account of a hail storm in Texas.

245. Check List of Publications of the Smithsonian Institution to July. 1872. 1872. 8vo., pp. 21. (M. C. x.)

246. Smithsonian Contributions to Knowledge. Vol. XVIII. 1872. 4to., pp. 646, 10 woodcuts, 5 plates, 3 charts.

CONTENTS.

- SCHOTT, C. A. Tables of rain and snow. No. 222.
- STOCKWELL, J. N. Secular variations of the orbits of planets. No. 282.
- HARKNESS, W. Observations on terrestrial magnetism. No. 239.
- FERREL, W. Converging series, expressing the ratio between the diameter and the circumference of a circle. No. 238.
- 247. Arrangement of the Families of Fishes, or Classes Pisces, Marsipobranchii, and Leptocardii. By Theodore Gill. November, 1872. 8vo., pp. 96. (M. C. XI.)
- 248. On the Geology of Lower Louisiana and the Salt Deposit on Petite Anse Island. By EUGENE W. HILGARD. June, 1872. 4to., pp. 38, 4 woodcuts, 2 plates. (S. C. XXIII.)
- 249. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1871. 42d Congress, 1st Session. Senate Mis. Doc. No. 149. 1873. 8vo., pp. 473, 3 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

List of meteorological stations and observers.

Meteorological material contributed to Smithsonian Institution.

Meteorological articles received by the Institution, and deposited in the Library of Congress. Auroras, earthquakes, etc.

Dodge, N. S. Memoir of Sir John Frederick William Herschel.

ARAGO, M. Eulogy on Joseph Fourier.

ODLING, W. Professor Thomas Graham's scientific work.

HELMHOLTZ, **H**. On the relation of the physical sciences to science in general.

KORNHUBER, G. A. Alternate generation and parthenogenesis in the animal kingdom.

REICHARDT, W. Present state of our knowledge of cryptogamous plants. STOCKWELL, J. N. Recent researches on the secular variations of the planetary orbits.

DE FOREST, E. L. Methods of interpolation applicable to the graduation of irregular series, such as tables of mortality. Part I.

DE SAUSSURE, H. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1870, to June, 1871.

Expedition toward the North Pole:

HENRY, J.; HILGARD, J. E.; NEWCOMB, S.; BAIRD, S. F.; MEEK, F. B.; AGASSIZ, L. Scientific instructions to Captain Hall.

ETHNOLOGY:

COMFORT, A. J. Indian mounds near Fort Wadsworth, Dakota.

Berthoud, E. L. Antiquities on the Cache la Poudre River, Weld county, Colorado Territory.

Lyon, W. B. Antiquities in New Mexico.

249. Report for 1871-Continued.

SPAINHOUR, J. M. Antiquities in Lenoir county, North Carolina. McCONNELL, E. M. Account of the old Indian village, Kushkush-

kee, near Newcastle, Pennsylvania.

GROSSMANN, F. E. Pima Indians of Arizona. CROOK, G. Indian mode of making arrow-heads and obtaining fire.

PETER, R. Ancient mound near Lexington, Kentucky.

Brown, D. Shell-heap in Georgia.

SCHOTT, A. Remarks on ancient relic of Maya sculpture.

Ancient history of North America. MUCH, M. RCEHRIG, F. L. O. On the language of the Dakota or Sioux Indians.

METEOROLOGY, with notes by Professor Henry:

LATIMER, G. Meteorology of Porto Rico.

Collins, Colonel. Meteorology of the Green river country.

LAMARK. Distinction between tornadoes and tempests.

MEEK, J. B. Account of a tornado which occurred in Spruce Creek Valley, Centre county, Pennsylvania.

HENRY, J. Effect of the moon on the weather.

KNIGHT, R. T.; HENRY, J. Connection of gales of wind and appearance of the aurora.

HARRISON, W. Account of a storm in Butler county, Kansas. June 28, 1871.

250. Smithsonian Miscellaneous Collections. Vol. X. 1873. 913, 5 woodcuts.

CONTENTS.

CARPENTER, P. P. Mollusks of western North America. No. 252.

GILL, T. Arrangement of the families of mollusks. No. 227.

HENRY, J. Instructions for observations of thunder storms. No. 235.

HENRY, J. Circular relative to heights. No. 286.

HENRY, J. Directions for constructing lightning rods. No. 287.

HENRY, J. Queries relative to tornadoes. No. 190.

Questions relative to the food fishes of the United States. BAIRD, S. F. No. 284.

BAIRD, S. F. Memoranda of inquiry relative to food fishes. No. 234.

RHEES, W. J. List of institutions, etc., in the United States in correspondence with the Smithsonian Institution. No. 238.

List of foreign correspondents of the Smithsonian Institution, for 1872 No. 248.

Check list of publications of the Smithsonian Institution, 1872. No. 245.

251. Memoir of C. F. P. Von Martius. By C. RAU. 1871. 8vo., pp. 12.

252. The Mollusks of Western North America. By P. P. CARPENTER. December, 1872. 8vo., pp. 446, 4 woodcuts. (M. C. x.)

CONTENTS.

Supplementary report on the present state of our knowledge with regard to the mollusca of the west coast of North America.

Review of Professor C. B. Adams' "Catalogue of the Shells of Panama" from the type specimens.

- 252. Mollusks of Western America—Continued.
 - Diagnoses of new forms of mollusks collected at Cape St. Lucas, Lower California.
 - Contributions towards a monograph of the pandoridæ.
 - Diagnoses of new forms of mollusca from the Vancouver district.
 - Diagnoses of new species and a new genus of mollusks, from the Reigen Mazatlan collection; with an account of additional specimens presented to the British Museum.
 - Descriptions of new species and varieties of chitonidæ and acmæidæ, from the Panama collection of the late Prof. C. B. Adams.
 - Diagnoses of new species of mollusks from the west tropical region of North America.
 - Diagnoses of new forms of mollusca from the west coast of North America, first collected by Col. E. Jewett.
 - Diagnoses of new forms of mollusca, collected by Col. E. Jewett, on the west tropical shores of North America.
 - Diagnoses des mollusques nouveaux provenant de Californie et faisant partie du Musée de l'Institution Smithsonienne.
 - On the pleistocene fossils collected by Col. E. Jewett, at Santa Barbara, California; with descriptions of new species.
- 253. Land and Fresh Water Shells of North America. Part IV. Strepomatidæ (American Melanians). By GEO. W. TRYON, Jr. December, 1873. 8vo., pp. 490, 871 woodcuts. (M. C. xvi.)
- 254. Synopsis of American Wasps. Solitary Wasps. By Henry De Saussure. December, 1875. 8vo., pp. 430, 4 plates of 31 figures. (M.C. xiv.)
- 255. The Constants of Nature. Part I. Specific Gravities: Boiling and Melting Points; and Chemical Formulæ. By Frank Wigglesworth Clarke. December, 1873. 8vo., pp. 272. (M. C. XII.)
- 256. Monographs of the Diptera of North America. Part III. By H. Loew. December, 1873. 8vo., pp. 381, 4 plates of 116 figures. (M. C. xi.)
- 257. Systematic Index to List of Foreign Correspondents of the Smithsonian Institution. 1872. 8vo., pp. 30. (M. C. x.)
- 258. Bibliographical Index to North American Botany; or, Citations of Authorities for all the recorded Indigenous and Naturalized Species of the Flora of North America; with a Chronological Arrangement of the Synonymy. Part I. Polypetale. By Sereno Watson. March, 1878. 8vo., pp. 484. (M. C. xv.)
- 259. Explorations of the Aboriginal Remains of Tennessee. By Joseph Jones. October, 1876. 4to., pp. 181, 110 woodcuts. (S. C. xxII.)
- 260. Regulations of the Smithsonian Institution. January, 1872. 8vo., pp. 41, 1 woodcut.

- 261. Directions for Collecting and Preserving Insects. By A. S. PACKARD, Jr. 1873. 8vo., pp. 60, 55 woodcuts. (M. C. XL.)
- 262. An Investigation of the Orbit of Uranus, with General Tables of its Motion. By Simon Newcomb. August, 1873. 4to., pp. 296. (S. C. XIX.)
- 263. Circular of Instructions to Observatories relative to Telegraphic Announcements of Astronomical Discoveries. By Joseph Henry. May, 1873. 8vo., pp. 4. (M. C. XII.)
- 264. New Species of North American Coleoptera. Part II. By John L. Le Conte. 1873. 8vo., pp. 74. (pp. 169-240.) (M. C. xi.)
- 265. Classification of the Coleoptera of North America. Part II. By JOHN L. LE CONTE. June, 1873. 8vo., pp. 72. (pp. 279-348. (M. C. XI.)
- 266. On the Structure of Cancerous Tumors and the mode in which Adjacent Parts are Invaded. Toner Lecture No. I. Delivered March 28, 1873. By J. J. WOODWARD. November, 1873. 8vo., pp. 44. 5 woodcuts. (M. C. xv.)
- 267. The Haidah Indians of Queen Charlotte's Islands, British Columbia, with a brief description of their Carvings, Tattoo Designs, etc. By James G. Swan. July, 1874. 4to., pp. 22, 7 plates of 24 figures. (S. C. xxi.)
- 268. The Winds of the Globe; or, the Laws of Atmospheric Circulation over the Surface of the Earth. By James Henry Coffin. The tables completed and maps drawn by S. J. Coffin, with a discussion and analysis of the tables and charts by Alexander Weekof. December, 1875. 4to., pp. 781, 4 woodcuts, 26 plates of 221 figures. (S. C. xx.)
- 269. The Sculptures of Santa Lucia Cosumalwhuapa, in Guatemala, with an account of travels in Central America and on the Western Coast of South America. By S. Habel. 1878. 4to., pp. 94, 8 plates of 25 figures. (S. C. XXII.)
- 270, Catalogue of the described Diptera of North America. By C. R. OSTEN SACKEN. 1878. 8vo., pp. 324. (M. C. xvi.)
- 271. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1872. 42d Congress, 3d Session. House of Representatives, Mis. Doc. No. 107. 1873. 8vo., pp. 456, 109 woodcuts.

HENRY, J. Secretary's report of operations. BOARD OF REGENTS, Proceedings of

71. Report for 1872-Continued.

List of meteorological stations and observers.

HENRY, J. Notes relative to George Catlin.

Report of Committee of Regents on Corcoran Art Gallery.

AGASSIZ, L. Narrative of the Hassler expedition.

BACHE, A. D. Bequest to National Academy of Sciences.

CORCORAN, W. W. Deed of gift of Art Gallery.

TONER, J. M. Deed of foundation of Toner lectures.

TYNDALL, J. Trust fund for the promotion of science in the U.S.

Hamilton, J. Bequest of \$1,000 to Smithsonian Institution.

HENRY, J. Circular sent with specimens presented.

ARAGO, M. Eulogy on Ampère.

FISCHER, F. Scientific labors of Edward Lartet.

PEABODY, A. P. Scientific education of mechanics and artisans.

BAUER, A. Organic bases.

KLETZINSKY, Prof. Nitrogen bodies of modern chemistry.

EGLESTON, T. Scheme for the qualitative determination of substances by the blowpipe.

Blowpipe apparatus of Hawkins and Wale.

Suess, E. Boundary line between geology and history.

Brezina, A. Explanation of the principles of crystallography and crystallophysics.

WCEIKOFF, A. Meteorology in Russia.

DONATI, G. B. Phenomena in telegraphic lines during the aurora borealis.

ETHNOLOGY:

BROCA, P. The Troglodytes, or cave dwellers, of the valley of the Vézere.

RAU, C. Ancient aboriginal trade in North America.

RAU, C. North American stone implements.

BRUFF, J. G. Indian engravings on the face of rocks along Green River valley in the Sierra Nevadas.

LEE, J. C. Y. Ancient ruin in Arizona.

BARRANDT, A. Haystack mound, Lincoln county, Dakota.

BREED, E. E. Earthworks in Wisconsin.

DEAN, C. K. Mound in Wisconsin.

WARNER, J. Big elephant mound in Grant county, Wisconsin.

CUTTS, J. B. Ancient relics in northwestern Iowa.

PERRINE, T. M. Mounds near Anna, Union county, Illinois.

PETER, R. Ancient mounds in Kentucky.

STEPHENSON, M. F. Mounds in Bartow county, Georgia.

McKinley, W. Mounds in McIntosh and Early counties, Georgia.

HOTCHKISS, T. P. Indian remains in Caddo parish, Louisiana.

LOCKETT, S. H. Mounds in Louisiana.

PEALE, T. R. Prehistoric remains in vicinity of City of Washington, D. C.

DEVEREUX, J. H. Catalogue of cabinet of Indian relics presented to Smithsonian Institution.

DEVEREUX, J. H. Ancient pottery from Phillips County, Arkansas.

KIPP, J. On the accuracy of Catlin's account of the Mandan ceremonies.

272. Smithsonian Contributions to Knowledge. Vol. XIX. 1874. 4to., pp. 640, 6 woodcuts, 21 plates.

CONTENTS.

Barnard, J. G. Problems of rotary motion. No. 240. Wood, H. C. Fresh-water algo of North America. No. 241. Newcomb, S. Orbit of Uranus. No. 262.

273. Smithsonian Miscellaneous Collections. Vol. XI. 1874. 8vo., pp. 790, 55 woodcuts, 4 plates.

CONTENTS.

GILL, T. Arrangement of the families of mammals. No. 230.

GILL, T. Arrangement of the families of fishes. No. 247.

LOEW, H. Monograph of the diptera of North America. Part III. No. 256.

Packard, A. S. Directions for collecting and preserving insects. No. 261.

LE CONTE, J. L. New species of North America coleoptera. Part II. No. 264.

LE CONTE, J. L. Classification of North America coleoptera. Part II. No. 265.

274. Smithsonian Miscellaneous Collections. Vol. XII. 1874. 8vo., pp. 767, 86 woodcuts.

CONTENTS.

BAIRD, S. F. Review of American birds. Part I. No. 181.

CLARKE, F. W. The constants of nature. Part I.—Specific gravities. No. 255.

HENRY, J. Circular on Telegraphic announcements of astronomical discoveries. No. 268.

275. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1873. 43d Congress, 1st Session. Senate Mis. Doc. No. 130. 1874. 8vo., pp. 452, 33 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on Museum and exchanges.

ENDLICH, F. M. List of minerals in the National Museum.

Classified record of monthly meteorological reports preserved in the Smithsonian Institution.

Classified list of meteorological publications, and meteorological articles in periodicals deposited in Library of Congress in 1878.

GARFIELD, J. A. Biographical notice of S. P. Chase and L. Agaseiz.

HAMLIN, H. Biographical notice of S. P. Chase.

PARKER, P. Biographical notice of L. Agassiz.

HAMILTON, J. Bequest deposited in United States Treasury.

Dodge, N. S. Memoir of Charles Babbage.

275. Report for 1873—Continued.

HENRY, J. On the "Moon-Hoax."

BABBAGE, C. Extracts from writings of

STEBBINS, R. P. Memoir of Louis Agassiz.

GRAY, A. Memoir of John Torrey.

STEVENS, J. A. Memoir of George Gibbs.

DALTON, J. C. Origin and propagation of disease.

HELMHOLZ H.; MAXWELL, J. C. Later views of the connection of electricity and magnetism.

GOULD, B. A. Account of the astronomical observatory at Cordoba, Argentine Republic.

MAILLY, E. Estimate of the population of the world.

MORIN, A. Warming and ventilating buildings. Part I.

DEFOREST, E. Additions to a memoir on methods of interpolation.

Part II.

ETHNOLOGY:

SCHUMACHER, P. Remarks on the Kjökken-Möddings on the Northwest Coast of America.

BERENDT, C. H. On a grammar and dictionary of the Carib or Karif

language, with some account of the people by whom it is spoken. GILLMAN, H. The mound-builders and platycnemism in Michigan.

MASON, O. T. The Leipsic "Museum of Ethnology."

PERRINE, T. M. Antiquities of Union county, Illinois.

Patton, A. Antiquities of Knox county, Indiana, and Lawrence County, Illinois.

Miscellaneous Correspondence:

DALL, W. H. Explorations on the Western coast of North America.

PIERSON, W. M. Discovery of a large meteorite in Mexico.

BRUNOT, F. R. On the habits of the beaver.

JEVONS, W. S. On a national library.

PRIZE QUESTIONS OF SCIENTIFIC SOCIETIES:

Society for the Encouragement of Science, Literature, and Art, Dankirk, France.

Society of Science, Art, and Literature of Hainaut, Mons. Belgium.

Royal Institute for the Encouragement of the Natural, Economical, and Technological Sciences, Naples, Italy.

Royal Academy of Science, Literature, and the Fine Arts, Brussels, Belgium.

Society of Sciences of Haarlem, Holland.

- 276. The Constants of Nature. Part II. A Table of Specific Heats for Solids and Liquids. By Frank W. Clarke. April, 1876. 8vo., pp. 58. (M. C. xiv.)
- 177. Tables, Distribution, and Variations of the Atmospheric Temperature in the United States and some adjacent parts of America. By Charles A. Schott. April, 1876. 4to., pp. 360, 10 woodcuts, 2 plates, 3 maps. (S. C. XXI.)
- 278. Check-list of Publications of the Smithsonian Institution. July, 1874. 8vo., pp. 24.

- 279. On Strain and Over-action of the Heart. Toner Lecture No. III. Delivered May 14th, 1874. By J. M. Da Costa. August. 1874. 8vo., pp. 32, 2 woodcuts. (M. C. xv.)
- 280. Statement and Exposition of certain Harmonies of the Solar System. By Stephen Alexander. March, 1875. 4to., pp. 104, 20 woodcuts. (S. C. xxi.)
- 281. On the General Integrals of Planetary Motion. By SIMON NEWCOMB. December, 1874. 4to., pp. 40. (S. C. xxi.)
- 282. A Study of the Nature and Mechanism of Fever. Toner Lecture No. IV. Delivered January 20, 1872. By Horatio C. Wood. February, 1875. 8vo., pp. 50. (M. C. xv.)
- 283. Catalogue of the Fishes of the East Coast of North America. By Theodore Gill. 1875. 8vo., pp. 56. (M. C. xiv.)
- 284. Smithsonian Contributions to Knowledge. Vol. XX. 1876. 4to., pp. 794, 4 woodcuts, 26 plates.

COFFIN, J. H. The winds of the globe. No. 268.

285. Smithsonian Contributions to Knowledge. Vol. XXI. 1876. 4to., pp. 543, 30 woodcuts, 9 plates, 3 maps.

CONTENTS.

ALEXANDER, S. Harmonies of the solar system. No. 280.

NEWCOMB, S. Integrals of planetary motion. No. 281.

Swan, J. G. Haidah Indians of Queen Charlotte's Islands, British Columbia. No. 267.

Schott, C. A. Tables, atmospheric temperature in the United States.
No. 277.

286. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1874. 43d Congress, 2d Session. House Doc. No. 56. 1875. 8vo., pp. 416, 46 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD of REGENTS. Proceedings of

List of articles deposited by the Smithsonian Institution in the Corcoran Art Gallery.

List of Indian vocabularies received from the Wheeler Expedition.

MITCHELL, S. W. Inquiries relative to the disease known as "Chorea." Classified record of meteorological material preserved in the Smithsonian Institution. Auroras, instruments, meteors, rain, wind, general meteorology, local meteorology.

GRAY, A.; COPPÉE, H. Report of Committee of Regents on the Smithsonian Museum.

286. Report for 1874—Continued.

ARAGO, M. Eulogy on La Place.

MAILLY, M. Eulogy on Quetelet.

DUMAS, M. Eulogy on A. A. de la Rive.

HILGARD, J. E. Tides and tidal action in harbors.

LEMSTRÖM, S.; DE LA RIVE, A. A. Electricity of the atmosphere and the Aurora Borealis.

DE CANDOLLE, A.; GRAY, J. E. On a dominant language for science.

SCHOTT, C. A.; EVERETT, J. D. Underground temperature.

DUPRE, W.; HENRY, J. Earthquakes in North Carolina, 1874.

DE LA RIVE, A. A. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1872, to June, 1878.

MORIN, A. Warming and ventilating buildings. Part II.

ETHNOLOGY:

SCHUMACHER, P. Ancient graves and shell-heaps of California.

King, W. M. Account of the burial of an Indian squaw, San Bernardino county, California.

McWhorter, T. Ancient mounds of Mercer county, Illinois.

PRATT, W. H. Antiquities of Whitesides county, Illinois.

FARQUHARSON, R. J. A study of skulls and long bones, from mounds near Albany, Illinois.

TIFFANY, A. S. The shell-bed skull.

HILL, G. W. Antiquities of northern Ohio.

ROBERTSON, R. S. The age of stone, and the troglodytes of Breck-inridge county, Kentucky.

DAVIS, A. C. Antiquities of Isle Royale, Lake Superior.

SMITH, J. W. C. Antiquities of Yazoo county, Mississippi.

WRIGHT, D. F. Antiquities of Tennessee.

LAW, A. E. Antiquities of Blount county, Tennessee.

CUSHING, F. H. Antiquities of Orleans county, New York.

ROBERTSON, R. S. Antiquities of La Porte county, Indiana.

ROBERTSON, R. S. Antiquities of Allen and De Kalb counties, Ind.

HAILE, J.; McHENRY, J. W. Antiquities of Jackson county, Tennessee.

ANDERSON, W. Antiquities of Perry county, Ohio.

BRYAN, O. N. Antiquities of Charles county, Maryland.

Kron, F. J. Antiquities of Stanley and Montgomery counties, North Carolina.

MITCHELL, A. Antiquities of Florida.

BARTRAM, J. Antiquities of Florida.

287. The Archæological Collection of the United States National Museum, in charge of the Smithsonian Institution, Washington, D. C. By Charles Rau. 1876. 4to., pp. 118, 340 woodcuts. (S. C. xxII.)

288. The Constants of Nature. First supplement to Part I. Specific Gravities, Boiling Points, and Melting Points. By Frank W. Clarke. April, 1876. 8vo., pp. 62. (M. C. xiv.)

289. The Constants of Nature. Part III. Tables of Expansion by Heat for Solids and Liquids. By Frank W. Clarke. April, 1876. 8vo., pp. 58. (M. C. XIV.)

- 290. Circular for Distribution at the Centennial Exhibition, 1876, containing List of Smithsonian Publications and Rules of Distribution; List of Foreign Agents; Number of Foreign and Domestic Institutionin Correspondence; List of Regents, Officers, and Assistants of the Institution. 1876. 12mo., pp. 12.
- 291. Dual Character of the Brain. Toner Lecture No. II. Deliveral April 22, 1874. By C. E. Brown-Séquard. January, 1877. 8vo., pp. 26. (M. C. xv.)
- 292. Check-list of North American Batrachia and Reptilia; with a Sytematic List of the Higher Groups, and an Essay on Geographical Distribution based on the specimens contained in the United States National Museum. By EDWARD D. COPE. 1875. 8vo., pp. 108. (M. C. XIII.) Bulletin of the National Museum No. 1.
- 293. Contributions to the Natural History of Kerguelen Island, made in Connection with the American Transit-of-Venus Expedition, 1874—75.

 By J. H. Kidder. I.—Ornithology, by Elliott Coues. 1875. 8ve., pp. 61. (M. C. XIII.) Bulletin of the National Museum, No. 2.
- 294. Contributions to the Natural History of Kerguelen Island, made in connection with the United States Transit-of-Venus Expedition, 1874-75. By J. H. Kidder. II.—Oology, Botany, &c. 1876. 8vo., pp. 124. (M. C. XIII.) Bulletin of the National Museum, No. 3.

KIDDER, J. H.; Cours, E. Oology.

BOTANY

GRAY, ASA. A.—Phænogamia, filices, et lycopodiaces.

JAMES, T. P. B .- Musci.

TUCKERMAN, E. C .- Lichones.

FARLOW, W. G. Algæ.

. CROZET FLORA.

ENDLICH, F. M. Geology.

KIDDER, J. H. Mammals.

GILL, T. N. Fish.

DALL, W. H. Mollusks.

INSECTS:

OSTEN-SACKEN, C. C. Diptera.

HAGEN, H. A. Pseudo-neuroptera.

SMITH, S. I. Crustaceans.

VERRILL, A. E. Annelids, echinoderms, and anthozoa.

KERSHNER, E. Collection.

KIDDER, J. H.; COUES, E. A study of chionis minor.

295. Birds of Southwestern Mexico, collected by Francis E. Sumichrast for the United States National Museum. By George N. LAWRENCE.

- 1875. 8vo., pp. 56. (M. C. XIII.) Bulletin of the National Museum, No. 4.
- 296. Catalogue of the Fishes of the Bermudas. Based chiefly upon the collections of the United States National Museum. By G. Brown Goode. 1876. 8vo., pp. 84. (M. C. XIII.) Bulletin of the National Museum, No. 5.
- 297. Classification of the Collection to illustrate the Animal Resources of the United States. A List of Substances derived from the Animal Kingdom, with Synopsis of the Useful and Injurious Animals and a Classification of Methods of Capture and Utilization. By G. Brown Goode. 1876. 8vo., pp. 140. (M. C. XIII.) Bulletin of the National Museum, No. 6.
- 298. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1875. 44th Congress, 1st Session. Senate Mis. Doc. No. 115. 1876. 8vo., pp. 422, 354 woodcuts.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on Museum and explorations.

BAIRD, S. F. Report on the proposed plan of exhibit by the Smithsonian Institution at the International Centennial Exhibition at Philadelphia.

PARKER, P. Eulogy on Henry Wilson.

ARAGO, M. Eulogy on Alexander Volta.

DE CANDOLLE, A. Probable future of the human race.

DE CANDOLLE, A. Report on the transactions of the Society of Physics and Natural History of Geneva, from July, 1873, to June, 1874.

PRESTWICH, J. The past and future of geology.

WEX, H. G. Diminution of the water of rivers and streams.

TAYLOR, W. B. Refraction of sound.

HENRY, J. On the organization of local scientific societies.

ETHNOLOGY:

MORTILLET, G. DE; CHANTEE, E. International code of symbols for charts of pre-historic archeology.

GILLMAN, H. Characteristics pertaining to ancient man in Michigan. Abbott, C. C. The stone age in New Jersey.

299. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1876. 44th Congress, 2d Session. Senate Mis. Doc. No. 46. 1877. 8vo., pp. 488, 73 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on Museum and explorations.

BAIRD, S. F. Report on Centennial Exhibition of 1876.

Centennial awards to Smithsonian Institution.

299. Report for 1876—Continued.

GRAY, A.; SARGENT, A. A.; CLYMER, H. Report of Committee on the Museum.

BANCROFT, G. Memorial to Congress, in behalf of the Regents, for new museum.

List of collections presented by foreign commissioners to the United States. Arago, M. Eulogy on Gay Lussac.

FIALHO, A. Biographical sketch of Dom Pedro II.

TAYLOR, W. B. Kinetic theories of gravitation.

PILAR, G. The revolutions of the crust of the earth.

KIRKWOOD, D. The asteroids between Mars and Jupiter.

ETHNOLOGY:

MASON, O. T. The Latimer collection of antiquities from Porto Rico in the National Museum at Washington, D. C.

ROMER, F. F. Pre-historic antiquities of Hungary.

BLONDEL, S. Jade. A historical, archeological, and literary study of the mineral called Yu by the Chinese.

WILLIAMSON, G. Antiquities in Guatemala.

BERENDT, C. H. Collection of historical documents in Guatemala.

Strone, Moses. Observations on the pre-historic mounds of Grant county, Wisconsin.

SNYDER, J. F. Deposits of flint implements, Illinois.

SMITH, C. D. Ancient mica mines in North Carolina.

PEET, S. D. Double-walled earthwork in Ashtabula county, Ohio. ELLSWORTH, E. W. Ancient implement of wood, from Connecticut. Powers, S. Centennial mission to the Indians of Western Nevada and California.

DOYLE, W. E. Indian forts and dwellings, Indian Territory.

BRACKETT, A. G. The Sioux or Dakota Indians.

- 300. On the Surgical Complications and Sequels of the Continued Fevers. With a Bibliography of works on Diseases of the Joints, Bones, Larynx, the Eye, Gangrene, Haematoma, Phlegmasia, etc. Toner Lecture No. V. Delivered February 17, 1876. By WILLIAM W. Keen. March, 1876. 8vo., pp. 72, 5 woodcuts. (M. C. xv.)
- 301. List of Publications of the Smithsonian Institution, July, 1877. 1877 8vo., pp. 72. (M. C. xiv.)
- 302. Subcutaneous Surgery, its Principles, and its recent Extension in Practice. Toner Lecture No. VI. Delivered September 13, 1876. By WILLIAM ADAMS. April, 1877. 8vo., pp. 20. (M. C. xv.)
- 303. Contributions to the Natural History of the Hawaiian and Fanning Islands and Lower California, made in connection with the United States North Pacific Surveying Expedition, 1873-75. By Thos. H. STREETS. 1877. 8vo., pp. 172. (M. C. XIII.) Bulletin of the National Museum, No. 7.

303. Natural History of Hawaiian Islands—Continued.

ORNITHOLOGY.

HERPETOLOGY.

ICHTHYOLOGY:

- I. Fishes of Upper and Lower California.
- II. Fishes of the Hawaiian Islands.
- III. Fishes of the Fanning Islands.
- IV. Fishes from the Samoan Islands.

CRUSTACEA.

BOTANY.

- 304. Index to the Names which have been appplied to the Subdivisions of the Class Brachiopoda, excluding the *Rudistes*, previous to the year 1877. By W. H. Dall. 1877. 8vo., pp. 88. (M. C. XIII.) Bulletin of the National Museum, No. 8.
- 305. Contributions to North American Ichthyology. Based primarily on the collections of the United States National Museum. I.—Review of Rafinesque's Memoirs on North American Fishes. By David S. Jordan. 1877. 8vo., pp. 56. (M. C. XIII.) Bulletin of the National Museum, No. 9.
- 306. Contributions to North American Ichthyology. Based primarily on the Collections of the United States National Museum. II. A.—
 Notes on Cottidæ, Etheostomatidæ, Percidæ, Centrarchidæ, Aphododeridæ, Dorysomatidæ, and Cyprinidæ, with revisions of the genera and descriptions of new or little known species. B.—Synopsis of the Siluridæ of the Fresh Waters of North America. By David S. Jordan. 1877. 8vo., pp. 124, 45 plates of 100 figures. (M. C. XIII.) Bulletin of the National Museum, No. 10.
- 307. Report on the Centennial Exhibition, Philadelphia. By S. F. BAIRD. 1877. 8vo., pp. 22.
- 308. Contributions to North American Ichthyology, based primarily on the collections of the United States National Museum. III. A.—On the Distribution of the Fishes of the Alleghany Region of South Carolina, Georgia, and Tennessee, with descriptions of new or little known species. By David S. Jordan and Alembert W. Brayton. B.—A Synopsis of the Family Catostomidæ. By David S. Jordan. 1878. 8vo., pp. 237. (M. C. XXIII.) Bulletin of the National Museum, No. 12.
- 309. List of Foreign Correspondents of the Smithsonian Institution. Corrected to January, 1878. March, 1878. 8vo., pp. 120. (M. C. xv.)
- 310. On the Internal Structure of the Earth, considered as affecting the Phenomena of Precession and Nutation; supplementary to article under the above head, Smithsonian Contributions to Knowledge, Vol. XIX, No. 240, being the third of the Problems of Rotary Mo-

- tion. By J. G. BARNARD. August, 1877. 4to., pp. 19, 4 woodcuts. (S. C. XXIII.)
- 311. Index Catalogue of Books and Memoirs relating to Nebulse and Clusters, etc. By Edward S. Holden. November, 1877. 8vo., pp. 126. (M. C. XIV.)
- 312. Smithsonian Miscellaneous Collections. Vol. XIII. 1878. 8vo., pp. 982, 45 plates. Bulletins of the National Museum, Nos. 1-10.

COPE, E. D. North American batrachia and reptilia. No. 292.

KIDDER, J. H.; COUES, E. Birds of Kerguelen Island. No. 293.

KIDDER, J. H., and others. Oology, botany, &c., of Kerguelen Island.

LAWRENCE, G. N. Birds of Mexico. No. 295.

GOODE, G. B. Fishes of Bermuds. No. 296.

No. 294.

GOODE, G. B. Classification of animal resources. No. 297.

STREETS, T. H. Natural history of Hawaiian Islands, Fanning Islands, and Lower California. No. 308.

DALL, W. H. Index of brachiopoda. No. 304.

JORDAN, D. S. North American Ichthyology. Nos. 305 and 306.

- 313. The Flora of St. Croix and the Virgin Islands. By Baron H. F. A. EGGERS. 1879. 8vo., pp. 139. (M. C. XXIII.) Bulletin of the National Museum, No. 13.
- 314. Smithsonian Miscellaneous Collections. Vol. XIV. 1878. 8va, pp. 911, 4 plates.

CONTENTS.

DE SAUSSURE, H. Synopsis of American wasps. No. 254.

GILL, T. Catalogue of fishes. No. 288.

CLARKE, F. W. Specific gravity tables. Supp. I. No. 288.

CLARKE, F. W. Specific heat tables. No. 276.

CLARKE, F. W. Tables of expansion by heat. No. 289.

Catalogue of photograph portraits of North American Indians. No. 216.

Check list of Smithsonian publications to July, 1877. No. 301.

Holden, E. S. Catalogue of books relative to nebulse. No. 311.

315. Smithsonian Miscellaneous Collections. Vol. XV. 1878. 8vo., pp. 880, 53 woodcuts.

CONTENTS.

WATSON, S. Botanical index. No. 258..

WOODWARD, J. J. Toner Lecture I. Cancerous tumors. No. 266.

BROWN-SÉQUARD, C. E. Toner Lecture III. The brain. No. 291.

DA COSTA, J. M. Toner Lecture III. The heart. No. 279.

WOOD, H. C. Toner Lecture IV. Study of fever. No. 282.

KEEN, W. W. Toner Lecture V. Continued fevers. No. 300.

ADAMS, W. Toner Lecture VI. Subcutaneous surgery. No. 302.

List of foreign correspondents of the Smithsonian Institution to January, 1878. No. 309.

- 315. Miscellaneous Collections. Vol. XV—Continued.
 - Circular in reference to American archeology. No. 816.
 - Circular. Inquiries relative to crawfish and crustaces. No. 819.
 - Circular relating to collections of living reptiles. No. 820.
- 316. Circular in reference to American Archæology. 1878. 8vo., pp. 15, 38 woodcuts. (M. C. xv.)
- 317. A Classification and Synopsis of the Trochilidæ. By DANIEL GIRAUD ELLIOT. 1879. 4to., pp. 289, 127 woodcuts. (S. C. XXIII.)
- 318. On the Remains of Later Pre-historic Man, obtained from Caves in the Catherina Archipelago, Alaska Territory, and especially from the Caves of the Aleutian Islands. By W. H. Dall. 1878. 4to., pp. 44, 10 plates of 39 figures. (S. C. XXII.)
- 319. Circular of Inquiries relative to the Natural History of the American Crawfish and other Fresh Water Crustacea. 1878. 8vo., pp. 8, 2 woodcuts. (M. C. xv.)
- 820. Circular relating to Collections of Living Reptiles. 1878. 8vo., pp. 2. (M. C. xv.)
- 321. The Nature of Reparatory Inflammation in Arteries after Ligatures, Acupressure, and Torsion. Toner Lecture No. VII. Delivered June 27, 1878. By Edw. O. Shakespeare. March, 1879. 8vo., pp. 74, 7 plates of 11 figures. (M. C. xvi.)
- 322. Smithsonian Miscellaneous Collections. Vol. XVI. 1880. 8vo., pp. 950, 871 woodcuts, 7 plates.

TEVON, G. W. Land and fresh water shells. Strepomatides. No. 253.

OSTEN SACKEN, C. R. Catalogue of diptera. No. 270.

SHAKESPEARE, E. O. Nature of reparatory inflammation. No. 321.

Circular relative to scientific and literary exchanges. No. 824.

Business arrangements of the Smithsonian Institution. No. 825.

ELLIOT, D. G. List of species of humming birds. No. 884.

List of the principal scientific and literary institutions in the United States. No. 335.

List of publications of the Smithsonian Institution. No. 844.

323. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1877. 45th Congress, 2d Session. Senate Mis. Doc. No. 35. 1878. 8vo., pp. 500, 49 woodcuts.

CONTENTS.

HENRY, J. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

BAIRD, S. F. Report on Museum and explorations.

BAIRD, S. F. List of explorations furnishing collections to the National Museum, 1888 to 1877.

323. Report for 1877—Continued.

HOLMGREN, F. Color blindness in its relation to accidents by rail and sea.

HENRY, J. Color blindness.

PLANTAMOUR, E. Report on the transactions of the Geneva Society of Physics and Natural History, from June, 1874, to June, 1875.

MÜLLER, J. Report on the transactions of the Geneva Society of Physics and Natural History, from June, 1875, to June, 1876.

FAVRE, A. Report on the transactions of the Geneva Society of Physics and Natural History, from June, 1876, to June, 1877.

ETHNOLOGY:

Cannon, G. L. Antiquities of Jefferson and Clear Creek counties, Colorado.

STRONG, Moses. Antiquities in Wisconsin.

HART, J. N. DE. The mounds and osteology of the mound builders of Wisconsin.

BREED, E. E. Pits at Embarrass, Wisconsin.

MOULTON, M. W. Mounds in Delaware county, Iowa.

KNAPP, Mrs. G. Earthworks on the Arkansas river, sixteen miles below Little Rock.

LYKINS, W. H. R. Antiquities of Kansas City, Missouri.

SHAW, J. The mound builders in the Rock river valley, Illinois.

COCHRANE, J. Antiquities of Mason county, Illinois.

HILL, G. W. Ancient earthworks of Ashland county, Onio.

CASE, H. B. Flint implements in Holmes county, Ohio.

MILLER, F. Mound in Trumbull county, Ohio.

FRIEL, J. Antiquities of Hancock county, Kentucky.

ROBERTSON, R. S. Antiquities of Nashville, Tennessee.

CLARK, W. M. Antiquities of Tennessee.

Jones, Jr., C. C. Aboriginal structures in Georgia.

BAILEY, W. B. F. Antiquities of Spalding county, Georgia.

GAINES, A. S.; CUNNINGHAM, K. M. Shell-heaps on Mobile river.

RAU, C. The stock in trade of an aboriginal lapidary.

RAU, C. Observations on a gold ornament from a mound in Florida. HALDEMAN, S. S. On a polychrome bead from Florida.

HARRISON, A. M. Colored bead dug from a mound on the eastern coast of Florida.

MAYBERRY, S. P. Shell-heaps at the mouth of Saint John's river, Florida.

TAYLOR, W. M. Ancient mound in western Pennsylvania.

SHEPARD, E. M. Deposit of arrow heads near Fishkill, New York.

GIBBS, G. J. Stone celts in the West Indies and Africa.

GALT, F. L. The Indians of Peru.

Bowers, S. History and antiquities of Santa Rosa Island, California

McParlin, T. A. Notes on the history and climate of New Mexico.

WEISMANN, A. On the change of the Mexican axolotl to an amblystoms.

ABBE, C. Short memoirs on meteorological subjects, translated by, viz:

HANN, J. On the diminution of aqueous vapor with increasing altitude in the atmosphere.

HANN, J. On the influence of rain upon the barometer and upon the formation of precipitation in general.

333. Report for 1877—Continued.

HANN, J. Atmospheric pressure and rain-fall.

HANN, J. The laws of the variation of temperature in ascending currents of air.

SOHNCKE, L. The law of the variation of temperature in ascending moist currents of air.

REYE, T. Rain-fall and barometric minima.

HANN, J. On the relation between the difference of pressure and the velocity of the wind according to the theories of Ferrel and Colding.

FERREL, W. Reply to the criticisms of J. Hann.

COLDING, A. Some remarks concerning the nature of currents of air.
COLDING, A. On the whirlwind at St. Thomas on the 21st of August,
1871.

PESLIN, M. On the relation between barometric variations and the general atmospheric currents.

- 324. Circular relative to Scientific and Literary Exchanges. 1879. 8vo., pp. 2. (M. C. xvi.) Free.
- 325. Business Arrangements of the Smithsonian Institution. January, 1879. 8vo., pp. 7. (M. C. xvi.) Free.
- 326. Catalogue of the Collection to illustrate the Animal Resources and the Fisheries of the United States, exhibited at Philadelphia, in 1876, by the Smithsonian Institution and the United States Fish Commission, and forming a part of the United States National Museum. By G. Browne Goode. 1879. 8vo., pp. 367. (M. C. XXIII.) Bulletin of the National Museum, No. 14.
- 327. The Scientific Writings of James Smithson. Edited by W. J. Rhees. 1879. 8vo., pp. 166, 32 woodcuts, 1 plate. (Portrait.)
 (M. C. xxi.)

CONTENTS.

SMITHSON, J. Scientific writings.

Johnson, W. R. Memoir on the scientific character and researches of James Smithson.

IRBY, J. R. McD. Works and character of James Smithson.

328. The Smithsonian Institution. Documents relative to its Origin and History. By WILLIAM J. RHEES. 1879. 8vo., pp. 1027. (M. C. XVII.)

CONTENTS.

SMITHSON, J. Will of

Correspondence between attorneys in England, Department of State, Richard Rush, and others relative to the bequest of Smithson.

RUSH, R. The case stated.

Opinion of English counsel.

Decree of Chancery Court awarding bequest to the United States.

Account of case of the United States.

Bill of costs of the case of the United States.

RUSH, R. Accounts of

SMITHSON, J. Schedule of personal effects.

338. Smithsonian Institution. Documents-Continued.

SMITHSON, J. Residuary legacy of

Congressional proceedings in relation to the bequest. 1885 to 1877. Twenty-fourth to Forty-fourth Congress.

HENRY, J. Digest of act of Congress to establish the Smithsonian Institution.

ADAMS, J. Q. Extracts from Memoirs of, relative to Smithson bequest.

TREASURY OF U. S. Accounts of, with Smithson fund.

Proposed applications of Smithson bequest.

Report of Organization Committee of the Board of Regents of the Smithsonian Institution.

HENRY, J. Programme of organization of Smithsonian Institution. Letters relative to the programme of organization.

329. The Smithsonian Institution. Journals of the Board of Regents, Reports of Committees, Statistics, etc. By WILLIAM J. RHEES. 1879. 8vo., pp. 851. (M. C. xvIII.)

CONTENTS.

Journal of Proceedings of the Board of Regents from 1846 to 1876.

Report of the Special Committee of the Board of Regents on the distribution of the income.

Report of the Special Committee of the Board of Regents-Prof. Feltonon the present of the Greek album from Miss E. Contaxaki.

Report of the Special Committee of the Board of Regents-Prof. Feltonon the purchase of the Stanley Indian gallery.

HENRY, J. Communication relative to a publication by Prof. S. F. B Morse.

Report of the Special Committee of the Board of Regents on the invention of the electro-magnetic telegraph.

Report of the Special Committee of the Board of Regents relative to the Smithsonian fire of January 24, 1865.

Act of Congress to transfer custody of library to Library of Congress.

Act of Congress to receive into Treasury of the United States residuary legacy of Smithson, and to authorize increase of fund to a million dollars.

Report of the Special Committee of the Board of Regents on best use of new hall of Institution.

Report of Executive Committee of the Board of Regents on the Washington city canal.

Report of the Executive Committee of the Board of Regents relative to the Corcoran Art Gallery.

Report of the Special Committee of the Board of Regents on the museum. EULOGIES AND BIOGRAPHICAL SKETCHES:

Agassiz, L.; Bache, A. D.; Chase, S. P.; Cleaveland, P.; Douglas, S. A.; Espy, J. P.; Felton, C. C.; Harvey, W. H.; Irving. W.; Jewett, C. C.; Pearce, J. A.; Priestley, J.; Rush, R.;

Seaton, W. W.; Totten, J. G.; Turner, W. W.; Wilson, H.; Wurdeman, G.

AGASSIZ, L. Narrative of expedition through Straits of Magellan to California.

Will of, establishing Bache Scientific Fund. BACHE, A. D. CORCORAN, W. W. Deed of gift and trust of Corcoran Art Gallery. 329. Smithsonian Institution. Journals, &c .- Continued.

Toner, J. M. Deed of, establishing Toner Lectures for advance of medical science.

TYNDALL, J. Deed of, establishing trust for promotion of science in the United States.

Hamilton, J. Bequest of, to Smithsonian Institution.

Circular sent with specimens of natural history, etc., presented to institutions.

Journal of the Executive Committee from Sept. 12, 1846, to Dec. 21, 1849. Report of the Executive Committee, from 1847 to 1875.

Journal of the Building Committee, from Feb. 17, 1847, to Dec. 1, 1847.

Report of the Building Committee, 1847-1867.

General financial and statistical statements, receipts and expenditures, classified, 1846 to 1877.

Statistics of literary and scientific exchanges, additions to the library, etc., 1846 to 1877.

 Appropriations from National Treasury by Congress for Smithsonian Institution and National Museum, 1846-1876.

List of Regents of the Institution from 1846 to 1879, according to mode of appointment, residence, etc.

Act of Congress to establish the Smithsonian Institution, August 10, 1846.

The same, according to the Revised Statutes.

By-Laws of the "Establishment."

The "Establishment" of the Smithsonian Institution, organization and journal of proceedings, 1846 to 1877.

Examination of Professor Henry by the English Government Scientific Commission, June 28, 1870.

330. Smithson and his Bequest. By WILLIAM J. RHEES. 1880. 8vo., pp. 76, 9 plates. (M. C. XXI.)

CONTENTS.

Life of Smithson.

Legislation of Congress in relation to the bequest.

List of first Board of Regents.

Obituary notice of Smithson, March, 1880.

DUTENS, L. Account of the First Duke of Northumberland.

Notice of the First Duke in The Gentleman's Magazine, 1786.

Coffin-plate inscription of Hugh Smithson.

Account of Earl Percy.

Notice of Smithson's paper on "Tabasheer."

DAVY, SIR H. Notice of Smithson's paper on "Calamines."

Illustrations of presentations of books to Smithson.

BERZELIUS. Notice of Smithson's researches.

Extracts from Smithson's scientific writings.

Catalogue of the library of Smithson.

Notice of the city of Washington, in Harriott's travels.

Notice of the city of Washington, in Weld's travels.

331. The Palenque Tablet, in the United States National Museum, Washington, D. C. By Charles Rau. 1879. 4to., pp. 90, 2 plates. (S. C. XXII.)

332. Proceedings of the U. S. National Museum for 1878. Vol. I. 1879. 8vo., pp. 524, 8 cuts, 8 plates. (M. C. xix.)

CONTENTS.

BEAN, T. H. Description of a new sparoid fish, Sargus holbrookii, from Savannah bank.

On the occurrence of Stichæus punctatus (Fabr.) Kröyer, at St. Michael's, Alaska.

On the identity of Euchalarodus Putnami Gill, with Pleuronectes glaier (Storer) Gill, with notes on the habits of the species.

Description of a species of Lycodes (L. Turneri) from Alaska. See also under Goode and Bean.

BELDING, L. A partial list of the birds of Central California.

Cook, C. The manufacture of porpoise-oil.

Dall, W. H. Description of new forms of mollusks from Alaska contained in the collections of the National Museum.

Postpliocene fossils in the Coast Range of California.

Fossil mollusks from later tertiaries of California.

Note on shells from Costa Rica kitchenmidden, collected by Drs. Flint and Bransford.

Distribution of Californian tertiary fossils.

Descriptions of new species of shells from California in the collections of the National Museum.

Report on the limpets and chitons of the Alaskan and Arctic regions.

descriptions of genera and species believed to be new.

EDWARDS, V. N. On the occurrence of the Oceanic Bonito, Orcynis pelamys (Linné) Poey, in Vineyard Sound, Massachusetts.

Gill, T. Synopsis of the pediculate fishes of the Eastern Coast of Extratropical North America.

Note on the Antennariidæ.

On the proper specific name of the common pelagic antennariid Pterophryne.

Note on the Ceratiidæ.

Note on the Maltheidæ.

GOODE, G. B. The Clupea tyrannus of Latrobe.

The occurrence of Belone latimanus in Buzzard's bay, Massachusetts. The voices of Crustaceans.

A revision of the American species of the genus Brevoortia, with a description of a new species from the Gulf of Mexico.

The occurrence of Hippocampus antiquorum, or an allied form, on St. George's Banks.

The occurrence of the Canada porcupine in West Virginia.

On two fishes from the Bermudas mistakenly described as new by Dr. Günther.

GOODE, G. B.; BEAN, T. H. The Craig flounder of Europe, Glyptoer-halus cynoglossus, on the coast of North America.

The Oceanic Bonito on the coast of the United States.

Description of Caulolatilus microps, a new species of fish from the Gulf coast of Florida.

- 332. Proceedings of National Museum. Vol. I-Continued.
 - GOODE, G. B.; BEAN, T. H. On a new serranoid fish, Epinephelus Drummond-Hayi, from the Bermudas and Florida.
 - Descriptions of two new species of fishes, Lutjanus Blackfordii and Lutjanus stearnsii, from the Coast of Florida.
 - A note upon the Black Grouper (Epinephelus nigritus (Holbrook) Gill) of the Southern coast.
 - Descriptions of two gadoid fishes, Phycis Chesteri and Haloporphyrus viola, from the deep-sea fauna of the Northwestern Atlantic.
 - Description of Argentina syrtensium, a new deep-sea fish from Sable Island Bank.
 - The identity of Rhinonemus caudacuta, (Storer) Gill with Gadus cimbrius Linn.
 - Note on Platessa ferruginea D. H. Storer, and Platessa rostrata H. R. Storer.
 - On the identity of Brosmius brosme americanus Gill, with Brosmius brosme (Müller) White.
 - JACKSON, J. B. S. Arsenic acid for protecting anatomical preparations from insects.
 - JEFFERSON, J. P. On the mortality of fishes in the Gulf of Mexico in 1878.
 - JEFFERSON, J. P.; PORTER, J. Y.; MOORE, T. On the destruction of fish in the vicinity of the Tortugas during the months of September and October, 1878.
 - JOEDAN, D. S. Notes on a collection of fishes from Clackamas river, Oregon.
 - JORDAN, D. S.; GILBERT, C. H. Notes on the fishes of Beaufort harbor, North Carolina.
 - LAWRENCE, G. N. Catalogue of the birds of Dominica, from collections made for the Smithsonian Institution, by Frederick A. Ober, together with his notes and observations.
 - Catalogue of the birds of St. Vincent, from collections made by Fred.

 A. Ober, under the directions of the Smithsonian Institution.
 with his notes thereon.
 - Catalogue of the birds of Antigua and Barbuda, from collections made for the Smithsonian Institution, by Fred. A. Ober, with his observations.
 - Catalogue of the birds of Grenada, from a collection made by Fred.

 A. Ober for the Smithsonian Institution, including others seen by him, but not obtained.
 - Catalogue of the birds collected in Martinique, by Fred. A. Ober for the Smithsonian Institution.
 - Catalogue of a collection of birds obtained in Guadeloupe for the Smithsonian Institution, by Fred. A. Ober.
 - A general catalogue of the birds noted from the islands of the Lesser Antilles visited by Fred. A. Ober; with a table showing their distribution, and those found in the United States.
 - LUPTON, N.T. On the breeding habits of the sea-catfish (Ariopsis milbertif)

 MERRILL, J. C. Notes on the ornithology of Southern Texas, being a
 list of birds observed in the vicinity of Fort Brown, Texas, from
 February, 1876, to June, 1878.

382. Proceedings of National Museum. Vol. I-Continued.

POET, F. Notes on the American species of the genus Cybium.

PRATT, R. H. Catalogue of casts taken by Clark Mills of the heads of sixty-four Indian prisoners of various western tribes, and held at Fort Marion, St. Augustine, Florida, in charge of R. H. Pratt.

RIDGWAY, R. On a new humming bird (Atthis Ellioti) from Guatemala.

A review of the American species of the genus Scops, Savigny.

Descriptions of several new species and geographical races of birds contained in the collection of the United States National Museum.

Description of two new species of birds from Costa Rica, and notes on other rare species from that country.

Descriptions of new species and races of American birds, including a synopsis of the Genus Tyrannus, Cuvier.

STEARNS, S. A note on the Gulf menhaden, Brevoortia patronus, Goode. STEINDACHNER, F. Note on Perca flavescens.

WILMOT, S. Notes on the western gizzard shad, *Dorosoma cepedianum* heterurum (Raf.) Jordan.

833. Proceedings of the U. S. National Museum for 1879. Vol. II. 1881.
8vo., pp. 503, 2 wood cuts, 7 plates. (M. C. xix.)
.

CONTENTS.

BEAN, T. H. A list of European fishes in the collection of the United States National Museum.

On the species of Astroscopus of the Eastern United States.

On the occurrence of *Hippoglossus vulgaris*, Flem., at Unalashka and St. Michael's, Alaska.

Description of an apparently new species of Gasterosteus (G. atkinsii) from the Schoodic Lakes, Maine.

Description of a new fish from Alaska (Anarrhichas lepturus), with notes upon other species of the genus Anarrhichas.

Notes on collection of fishes from eastern Georgia.

Description of a new species of Amiurus (A. ponderosus) from the Mississippi river.

Descriptions of two species of fishes, collected by Prof. A. Duges in Central Mexico.

Descriptions of some genera and species of Alaskan fishes. See also under Goode and Bean.

BREWER, T. M. Notes on the nests and eggs of the eight North American species of empidonaces.

COURS, J. G. On the migrations and nesting habits of west coast birds. COURS, E. Fourth instalment of ornithological bibliography, being a list

of faunal publications relating to British birds. GERKE, H. On the birds of Heligolaud.

GOODE, G. B. A study of the trunk fishes (Ostraciontides), with notes upon the American species of the family.

A preliminary catalogue of the fishes of the St. John's river and the east coast of Florida, with descriptions of a new genus and three new species.

Description of a new species of amber fish (Seriola stearnsii) obtained near Pensacola, Fla., by Mr. Silas Stearns.

333. Proceedings of National Museum. Vol. II-Continued.

- GOODE, G. B.; BEAN, T. H. Description of Alepocephalus bairdii, a new species of fish from the deep-sea fauna of the western Atlantic.
 - Description of a species of Lycodes (L. paxillus), obtained by the United States Fish Commission.
 - Description of a new species of Liparis (L. ranula), obtained by the United States Fish Commission off Halifax, Nova Scotia.
 - Catalogue of a collection of fishes sent from Pensacola, Fla., and vicinity, by Mr. Silas Stearns, with descriptions of six new species.
 - Description of a new genus and species of fish, Lopholatilus chamœ-leonticeps, from the south of New England.
 - On the occurrence of Lycodes vahlii, Reinhardt, on La Have and Grand Banks.
 - Catalogue of a collection of fishes obtained in the Gulf of Mexico by Dr. J. W. Velie, with description of seven new species.
- HARGER, O. Notes on New England Isopoda.
- JORDAN, D. S. Notes on certain typical specimens of American fishes in the British Museum and in the Museum d'Histoire Naturelle, at Paris.
 - Description of new species of North American fishes.
 - Notes on a collection of fishes obtained in the streams of Guanajuato and in Chapala lake, Mexico, by Prof. A. Dugès.
- KIDDER, J. H. Report of experiments upon the animal heat of fishes, made at Provincetown, Mass., during the summer of 1879, in connection with operations of the United States Fish Commission.
- LOCKINGTON, W. N. Review of the Pleuronectidae of San Francisco.

 Descriptions of new genera and species of fishes from the coast.
- Descriptions of new genera and species of fishes from the coast of California.
- MERRILL, J. C. On the habits of the Rocky Mountain goat.
- PRATT, R. H.. List of names, ages, tribe, &c., of Indian boys and girls at Hampton Normal and Agricultural Institute, Virginia, plaster casts of whose heads were taken by Clark Mills, Esq., March, 1879.
- SMITH, S. I. Occurrence of *Chelura terebrans*, a crustacean destructive to the timber of submarine structures, on the coast of the United States.
 - Notice of a new species of the Willemæsia group of crustacea (recent Eryontidæ.)
- VERRILL, A. E. Notice of recent additions to the marine invertebrata of the northeastern coast of America, with descriptions of new genera and species and critical remarks on others.
- VERBILL, A. E.; RATHBUN, R. List of marine invertebrata from the New England coast, distributed by the United States Commission of Fish and Fisheries.
- WHITE, C. A. Descriptions of new species of carboniferous invertebrate fossils.
 - Descriptions of new cretaceous invertebrate fossils from Kansas and Texas.
 - Note on Endothyra ornata.
 - Note on Criocardium and Ethmocardium.

- 334. List of described species of Humming Birds. By DANIEL GIRAUD ELLIOT. 1879. 8vo., pp. 22. (M. C. XVI.)
- 335. List of the Principal Scientific and Literary Institutions in the United States, May, 1879. 1879. 8vo., pp. 6. (M. C. xvi.)
- 336. Smithsonian Miscellaneous Collections. Vol. XVII. 1880. 8vo., pp. 1034.

Documents relative to the origin and history of the Smithsonian Institution. No. 328.

337. Smithsonian Miscellaneous Collections. Vol. XVIII. 1880. 8vo., pp. 851.

CONTENTS.

Journals of the Board of Regents, Reports of Committees, Statistics, etc. No. 329.

- 338. Notes on the Life and Character of Joseph Henry. Read before the Philosophical Society of Washington, October 26, 1878. By James C. Welling. 1880. 8vo., pp. 30, 1 plate. (Portrait.) (M. C. XXI.)
- 339. A Memoir of Joseph Henry: a Sketch of his Scientific Work. Read before the Philosophical Society of Washington, October 26, 1878. By WILLIAM B. TAYLOR. 1880. 8vo., pp. 225, 1 plate. (Portrait. (M. C. XXI.)
- 340. Smithsonian Contributions to Knowledge. Vol. XXII. 1880. 4to, pp. 537, 474 woodcuts, 20 plates.

CONTENTS.

JONES, J. Explorations of aboriginal remains of Tennessee. No. 259. HABEL, S. Sculptures of Santa Lucia Cosumaluhuapa, in Guatemaia. No. 269.

RAU, C. Archæological collection of U.S. National Museum. No. 237 RAU, C. Palenque tablet in the U.S. National Museum. No. 381. DALL, W. H. Remains of later pre-historic man in Alaska. No. 318.

341. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1878. 45th Congress, 3d Session. Senate, Mis. Doc. No. 59. 1879. 8vo., pp. 575, 12 woodcuts.

CONTENTS.

BAIRD, S. F. Secretary's report of operations.

Report of Government explorations and surveys.

Acts and resolutions of Congress relative to the Smithsonian Institution and the National Museum. Forty-fifth Congress, 2d and 3d Sess. 1878, 1879.

BOARD OF REGENTS, Proceedings of

GRAY, A. Biographical memoir of Joseph Henry.

ABAGO, F. Biography of Condorcet.

FAVRE, E. Biographical notice of Louis Agassiz.

341. Report for 1878-Continued.

TAYLOR, W. B. Henry and the telegraph.

LAUTENBACH, B. F. Irritation of a polarized nerve.

Wood, H. C. Researches upon fever.

LECONTE, J. Constants of nature.

List of apparatus available for scientific researches.

OBER, F. A. Ornithology of the Carribee Islands.

Kumlien, L. Report of explorations in Greenland.

HENRY, J. Researches in sound.

342. Contributions to the Natural History of Arctic America, made in connection with the Howgate Polar Expedition, 1877-'78. By LUDWIG KUMLIEN. 1879. 8vo., pp. 179. (M. C. XXIII.) Bulletin of the National Museum, No. 15.

CONTENTS.

KUMLIEN, L. Ethnology, mammals, and birds.

BEAN, T. H. Fishes.

VERBILL, A. E. Annelides, molluscoids, and radiates.

DALL, W. H. Mollusks.

INSECTS:

EDWARDS, W. H. Diurnal lepidoptera.

Scudder, S. H. and others. Hymenoptera, nocturnal lepidoptera, diptera, coleoptera, neuroptera, and arachnida.

GRAY, A. Plants.

TUCKERMAN, E. Lichens.

FARLOW, W. G. Algæ.

- 343. Annual Reports of the Secretary of the Smithsonian Institution, JOSEPH HENRY, 1865 to 1877. 1880. 8vo., pp. 548.
- 344. Check-list of Publications of the Smithsonian Institution, July, 1879. 1879. 8vo., pp. 16. (M. C. xvi.)
- 345. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1879. 1880. 46th Congress, 2d Session. Senate Mis. Doc. No. 54. 8vo., pp. 631, 216 woodcuts.

CONTENTS.

BAIRD, S. F. Secretary's report of operations.

BOARD OF REGENTS, Proceedings of

Acts and resolutions of Congress relative to the Smithsonian Institution and the National Museum. 45th Congress, 3d Session; 46th Congress, 2d Session. 1878-1880.

Report of the National Museum Building Commission.

Report of the Architects.

RHEES, W. J. James Smithson and his bequest.

KNIGHT, E. H. A study of the savage weapons at the Centennial Exhibition, Philadelphia, 1876.

ANTHROPOLOGY:

WORSAAE, J. J. A. The preservation of antiquities and national monuments in Denmark.

HAVARD, V. The French half-breeds of the Northwest.

845. Report for 1879-Continued.

NORRIS, P. W. Prehistoric remains in Montana, between Fort Ellis and the Yellowstone river.

BRACKETT, A. G. The Shoshone or Snake Indians: their religion. superstitions, and manners.

BURR, R. T. Ruins in White river canon, Pima county, Arizona.

Armstrong, T. Mounds in Winnebago county, Wisconsin.

Anderson, W. G. Mounds near Quincy, Ill., and in Wisconsin.

EVANS, S. B. Notes on some of the principal mounds in Des Moises vallev.

DAHLBERG, R. N. and C. Composition of ancient pottery found near the mouth of Chequest creek, at Pittsburg, on the Des Moines river.

BROADHEAD, G. C. Prehistoric evidences in Missouri.

THOMPSON, T. Mounds in Muscatine county, Iowa, and Rock Island county, Illinois.

TŒLLNER, A. Antiquities of Rock Island county, Illinois.

OEHLER, A. Stone cists near Highland, Madison county, Illinois.

MITCHELL, B. Mounds in Pike county, Illinois.

ADAMS, W. H. Mounds in the Spoon River valley, Illinois.

QUICK, E. R. Mounds in Franklin county, Indiana.

JACKMAN, F. Mounds and earthworks of Rush county, Indians.

JONES, JR., C. C. Primitive manufacture of spear and arrow points along the line of the Savannah river.

GESNER, W. Mica beds in Alabama.

Hough, J. Mounds in Washington county, Mississippi.

BRODNAX, B. H. Mounds in Moorehouse Parish, Louisiana.

BEAUCHAMP, W. M. Wampum belts of the Six Nations.

ANDREWS, F. D. Indian relics from Schoharie, New York.

WALKER, S. T. Preliminary explorations among the Indian mounds in Southern Florida.

WALKER, S. T. Report on the shell heaps of Tampa bay, Florida. NUTTER, F. H. Mounds on Gideon's farm, near Excelsior, Hennepin county, Minnesota.

MASON, O. T. Summary of correspondence of the Smithsonian Institution previous to January 1, 1880, in answer to circular No. 316.

MASON, O. T. Anthropological investigations during the year 1879.

BOEHMER, G. H. Index to papers on anthropology published by the Smithsonian Institution, 1847 to 1878.

PISKO, F. J. On the present fundamental principles of physics.

VON BAUMHAUER, E. H. A universal meteorograph, designed for detached observatories.

HOLDEN, E. S. Reports of American observatories.

Smithsonian Contributions to Knowledge. Vol. XXIII. 1881. **346**. 4to., pp. 766, 160 woodcuts, 18 plates of 165 figures.

CONTENTS.

CLARK, H. J. Lucernarise and their allies. No. 242.

HILGARD, E. W. Geology of Lower Louisiana and Petite Anse Island. No. 248.

BARNARD, J. G. Internal structure of the earth. No. 310.

ELLIOT, D. G. Classification and synopsis of the trochilidse. No. 317.

WOOD, H. C. Fever; a study in physiology. No. 357.

- 347. Nomenclature of Clouds. Two lithographic plates. 1851. 8vo.
- 348. Report on the Fishes of the New Jersey Coast, as observed in the Summer of 1854. By Spencer F. Baird. 1855. 8vo., pp. 40.
- 349. Suggestions for the Sanitary Drainage of Washington City. Toner Lecture No. VIII. Delivered May 26, 1880. By George E. Waring, Jr. June, 1880. 8vo., pp. 26.
- 350. A Map of the Stars near the North Pole; for observations on the Aurora. Copied from the Map used in the Toronto Observations. 1856. 15 inches by 12 inches.
- 351. On the Distribution of the Forests and Trees of North America; with notes on its Physical Geography. By J. G. COOPER. 1859. 8vo., pp. 36, 1 woodcut.
- 352. Brief Abstract of a series of six Lectures on the Principles of Linguistic Science. Delivered at the Smithsonian Institution in March, 1864. By WILLIAM D. WHITNEY. 1864. 8vo., pp. 22.
- 353. Tables and Results of the Precipitation, in Rain and Snow, in the United States, and at some stations in adjacent parts of North America, and in Central and South America. Collected by the Smithsonian Institution, and discussed under the direction of Joseph Henry and Spencer F. Baird, Secretaries. By Charles A. Schott. May, 1881. 4to., pp. 269, 8 woodcuts, 5 plates, 5 charts.
- 354. Essay on the Velocity of Light. By M. Delaunay. Translated for the Smithsonian Institution by Alfred M. Mayer. 1864. 8vo., pp. 31.
- 355. Ozone and Antozone. By CHARLES M. WETHERILL. 1864. 8vo., pp. 12.
- 356. A Memorial of Joseph Henry. (Published by order of Congress.) 1880. 8vo., pp. 532, 1 plate.

INTRODUCTION. Proceedings in Congress relative to public commemoration.

PART I .- OBSEQUIES OF JOSEPH HENRY.

WAITE, M. R. Mortuary announcement.

Proceedings of the Board of Regents.

The funeral, May 16, 1878.

Hodge, C. Prayer at the funeral.

MITCHELL, S. S. Funeral sermon.

PART II .- MEMORIAL EXERCISES AT THE CAPITOL.

Announcement by Executive Committee of the Regents.

McCosн, J. Introductory prayer.

HAMLIN, H. Address.

356.—Memorial of Joseph Henry—Continued.

WITHERS, R. E. Address.

GRAY, A. Address.

CLYMER, H. Reading of telegrams.

ROGERS, W. B. Address.

GARFIELD, J. A. Address.

Cox, S. S. Address.

SHERMAN, W. T. Address.

SUNDERLAND, B. Concluding prayer.

PART III.—MEMORIAL PROCEEDINGS OF SOCIETIES.

Proceedings of the Philosophical Society of Washington.

Proceedings of the Albany Institute.

MEADS, O. Memorial minute.

Proceedings of the United States Light-House Board.

Don, S. B. Memorial discourse.

CAMERON, H. C. Reminiscences.

WELLING, J. C. Memorial address.

TAYLOR, W. B. Memorial address.

LOVERING, J. Obituary memoir.

NEWCOMB, S. Biographical memoir.

MAYER, A. M. Memorial address.

APPENDIX. Proceedings in Congress regarding the erection of a monument to Joseph Henry.

- 357. Fever: A Study in Morbid and Normal Physiology. By Horatio C. Wood. January, 1878. 8vo., pp. 263, 16 woodcuts, 5 plates of 16 figures. (S. C. XXIII.)
- 358. The Constants of Nature. Part IV. Atomic Weight Determinations:

 A Digest of the Investigations published since 1814. By George F.
 Becker. August, 1880. 8vo., pp. 152.
- 359. A Planisphere of the Visible Heavens, extending to 40° of south declination, for observations on Meteoric displays. Prepared by a Committee of the Connecticut Academy of Arts and Sciences. 1864. 30 inches by 24½ inches.
- 360. Palafittes, or Lacustrian Constructions of the Lake of Neuchard By E. Desor. Translated for the Smithsonian Institution. 1865. 8vo., pp. 53, 118 woodcuts.
- 361. An account of the Aboriginal Inhabitants of the California Peninsula as given by Jacob Baegert, a German Jesuit Missionary, who lived there seventeen years during the second half of the last century. Translated and arranged for the Smithsonian Institution, by Charles Rau. 1865. 8vo., pp. 41.
- 362. Artificial Shell-deposits in New Jersey. By Charles Rau. 1865. 8vo., pp. 6, 1 woodcut.
- 363. Instructions for collecting Land and Fresh Water Shells. By James Lewis. 1866. 8vo., pp. 8.

- 364. Outline of a Systematic Review of the Class of Birds. By W. LILLJE-BORG. Translated for the Smithsonian Institution. 1866. 8vo., pp. 16.
- 365. Notes on the Tinneh or Chepewyan Indians of British and Russian America. 1.—The Eastern Tinneh; by Bernard R. Ross. 2.—The Loucheux Indians; by William L. Hardisty. 3.—The Kutchin Tribes; by Strachan Jones. Compiled by George Gibbs. 1866. 8vo., pp. 25, 11 woodcuts.
- 366. Directions for Collecting, Preserving, and Transporting Specimens of Diatomacea, and other Microscopic Organisms. By ARTHUR M. EDWARDS. 1867. 8vo., pp. 7.
- 367. Sketch of the Flora of Alaska. By J. T. ROTHROCK. 1867. 8vo., pp. 33.
- 368. Indian Pottery. By Charles Rau. 1867. 8vo., pp. 11, 9 woodcuts.
- 369. Dorpat and Poulkova. By CLEVELAND ABBE. 1867. 8vo., pp. 23.
- 370. A Deposit of Agricultural Flint Implements found in Southern Illinois. By Charles Rau. 1868. 8vo., pp. 9, 5 woodcuts.
- 371. The Metric System of Weights and Measures, with Tables. Prepared for the Smithsonian Institution. By H. A. NEWTON. 1868. 8vo., pp. 23.
- 372. Drilling in Stone without the use of Metal. By CHARLES RAU. 1869. 8vo., pp, 11, 12 woodcuts.
- 373. Meteorological Stations, and Observers of the Smithsonian Institution, in North America and adjacent Islands, from the year 1849, to the end of the year 1868. 1869. 8vo., pp. 42.
- 374. Three Rain-Charts of the United States, showing the distribution by Isohyetal lines of the mean precipitation in rain and melted snow.

 1, for the Summer months; 2, for the Winter months; 3, for the Year. 1870. 20 inches by 14 inches.
- 375. Thoughts on the Nature and Origin of Force. By WILLIAM B. TAYLOR. 1870. 8vo., pp. 19.
- 376. On the Chemistry of the Earth. By T. STERRY HUNT. 1871. 8vo., pp. 26.
- 377. The Diamond and other Precious Stones. By M. BABINET. Translated for the Smithsonian Institution, by John Stearns. 1872. 8vo., pp. 33.
- 378. The Language of the Dakota or Sioux Indians. By F. L. O. RŒHRIG. 1872. 8vo., pp. 19.

- 379. Eulogy on Prof. Alexander Dallas Bache, late Superintendent of the U. S. Coast Survey, President of the National Academy of Sciences, etc. By Joseph Henry. 1872. 8vo., pp. 28.
- 380. The Scientific Education of Mechanics and Artizans. By Andrew P. Peabody. 1873. 8vo., pp. 13.
- 381. Temperature-Chart of the United States, showing the distribution, by Isothermal lines, of the mean temperature for the Year. 1873. 16! inches by 10! inches.
- 382. North American Stone Implements. By CHARLES RAU. 1873 8vo., pp. 16, 7 woodcuts.
- 383. Archæological Researches in Nicaragua. By J. F. Bransford. 1881. 4to., pp. 100, 202 woodcuts, 2 plates of 40 figures.
- 384. Circular in reference to Shipping Fresh Fish and other Animals. By S. F. BAIRD. 1881. 8vo., pp. 4.
- 385. Ancient Aboriginal Trade in North America. By CHARLES RAU. 1873. 8vo., pp. 49.
- 386. Explanation of the Principles of Crystallography and Crystallophysics. By Aristides Brezina. Translated for the Smithsonian Institution, by T. Egleston. 1874. 8vo., pp. 36.
- 387. Three Temperature-Charts of the United States, showing the distribution, by Isothermal curves, of the mean temperature of the lower atmosphere. 1, for the Summer months; 2, for the Winter months: 3, for the Year. 1874. 28½ inches by 19½ inches.
- 388. Temperature Chart of the United States, showing the distribution, by Isothermal lines, of the mean temperature for the year. 1874. 20 inches by 13½ inches.
- 389. An Account of Investigations relative to Illuminating Materials. (From the Light-House Board Report for 1875.) By JOSEPH HENRY. 1881. 8vo., pp. 25.
- 390. On Tides and Tidal Action in Harbors. By Julius E. Hilgard, 1875. 8vo., pp. 22, 15 woodcuts.
- 391. An act to establish the Smithsonian Institution, etc. Revised Statutes: Title LXXIII. 1875. 8vo., pp. 10.
- 392. The Prehistoric Antiquities of Hungary. An Address delivered by F. F. Romer, at the Opening of the International Anthropological Congress, held at Budapest, September, 1876. Translated for the Smithsonian Institution, by Charles Rau. 1877. 8vo., pp. 9.

- 393. The Mound Builders, and Platycnemism in Michigan. By Henry Gillman. 1874. 8vo., pp. 28, 12 woodcuts. Also, Certain Characteristics pertaining to Ancient Man in Michigan. By Henry Gillman. 1877. 8vo., pp. 13, 13 woodcuts.
- 394. The Stone Age in New Jersey. By C. C. Abbott. 1877. 8vo. pp. 136, 58 plates of woodcuts, containing 223 figures.
- 395. Kinetic Theories of Gravitation. By William B. Taylor. 1877. 8vo., pp. 80.
- 396. Notes on the History and Climate of New Mexico. By Thomas A. McParlin. 1877. 8vo., pp. 30.
- 397. The Latimer Collection of Antiquities from Porto Rico, in the National Museum at Washington, D. C. By Otis T. Mason. 1877. 8vo., pp. 23. 14 plates of woodcuts, containing 60 figures.
- 398. Short Memoirs on Meteorological Subjects: By Julius Hann; L. Sohncke; Theodore Reye; William Ferrel; A. Colding; and M. Peslin. Compiled and Translated for the Smithsonian Institution by CLEVELAND ABBE. 1878. 8vo., pp. 104.
- 399. Color Blindness in its Relation to Accidents by Rail and Sea. By FRITHIOF HOLMGREN. Translated for the Smithsonian Institution, by M. L. Duncan. 1878. 8vo., pp. 72, 5 woodcuts.
- 400. Aboriginal Structures in Georgia. By Charles C. Jones, Jr. 1878. 8vo., pp. 13, 5 woodcuts.
- 401. On the change of the Mexican Axolotl to an Amblystoma. By August Weismann. Translated for the Smithsonian Institution, by Henry M. Douglass. 1878. 8vo., pp. 29.
- 402. The Stock-in-Trade of an Aboriginal Lapidary. By CHARLES RAU. 1878. 8vo., pp. 9, 16 woodcuts.
- 403. Observations on a Gold Ornament found in a Mound in Florida.

 By CHARLES RAU. 1878. 8vo., pp. 6, 1 woodcut.
- 404. On a Polychrome Bead from Florida. By S. S. HALDEMAN. 1878. 8vo., pp. 6, 2 woodcuts.
- 405. An Historical Sketch of Henry's Contribution to the Electro-Magnetic Telegraph; with an account of the origin and development of Prof. Morse's invention. By WILLIAM B. TAYLOR. 1879. 8vo., pp 103.
- 406. Henry on Sound. A Summary of Researches in Sound conducted in the service of the United States Light-House Board, by JOSEPH HENRY, during the years 1865 to 1877. 1879. 8vo., pp. 106, 12 woodcuts.

- 407. Biographical Memoir of Joseph Henry. Prepared in behalf of the Board of Regents. By Asa Gray. 1879. 8vo., pp. 35.
- 408. Report of the Secretary of the Smithsonian Institution, for the year 1878. By S. F. Baird. 1879. 8vo., pp. 60.
- 409. Report of the National Museum Building Commission, and of the Architects, for 1879. 1880. 8vo., pp. 18, 2 folding plates.
- 410. Reports of Astronomical Observatories for 1879. By E. S. HOLDEN. 1880. 8vo., pp. 60.
- 411. The effect of Irritation of a Polarized Nerve. By B. F. LAUTEN-BACH. 1880. 8vo., pp. 59.
- 412. On the Zoological Position of Texas. By Edward D. Cops. 1880. 8vo., pp. 55. Bulletin of the National Museum No. 17.
- 413. Exhibit of the Fisheries and Fish Culture of the United States of America, at the Internationale Fischerei Austsellung, at Berlin, April 20, 1880, and forming a part of the collection of the National Museum, made by the United States Fish Commission. Prepared under the direction of G. Brown Goode. 1880. 8vo., pp. 279. Bulletin of the National Museum, No. 18.
- 414. Base Chart of the United States. Prepared by Charles A. Schott.
 1880. (28½ x 19½ inches.)
- 415. A Study of the Savage Weapons at the Centennial Exhibition, Philadelphia, 1876. By EDWARD H. KNIGHT. 1880. 8vo., pp. 90, 147 woodcuts.
- 416. Smithsonian Miscellaneous Collections. Vol. XIX. 1880. 8vo., pp. 1034.

Proceedings of the U. S. National Museum. Vol. I. 1878. No. 332. Proceedings of the U. S. National Museum. Vol. II. 1879. No. 333.

- 417. Henry as a Discoverer. A Memorial Address, delivered before the American Association for the Advancement of Science, August 26, 1880. By Alfred M. Mayer. 1880. 8vo., pp. 36.
- 418. Report of the Secretary of the Smithsonian Institution, for the year 1879. By Spencer F. Baird. 1880. 8vo., pp. 76.
- 419. Report of the Secretary of the Smithsonian Institution, for the year 1880. By Spencer F. Baird. 1881. 8vo., pp. 88.
- 420. Anthropological Investigations in 1879. By Oris T. Mason. 1881. 8vo., pp. 30.
- 421. Index to Anthropological Articles in Publications of the Smithsonian Institution. By George H. Bœhmer. 1881. 8vo., pp. 10.

- 422. Nomenclature of North American Birds chiefly contained in the United States National Museum. By Robert Ridgway. 1881. 8vo., pp. 94. Bulletin of the National Museum, No. 21.
- 423. Smithsonian Miscellaneous Collections. Vol. XX. 1881. 8vo. pp. 846.

Bulletin of the Philosophical Society of Washington:

Vol. I, March, 1871, to June, 1874, pp. 218.

Vol. II, October 10, 1874, to November 2, 1878, pp. 452.

· Vol. III, November 9, 1878, to June 19, 1880, pp. 169.

424. Smithsonian Miscellaneous Collections. Vol. XXI. 1881. 8vo., pp. 773, 42 woodcuts, 13 plates.

CONTENTS.

RHEES, W. J. James Smithson and his bequest. No. 380.

SMITHSON, J. Scientific writings. No. 827.

JOHNSON, W. R. Scientific character and researches of Smithson. No. 827.

IRBY, J. R. McD. The works and character of Smithson. No. 327. A memorial of Joseph Henry. No. 856.

425. Proceedings of the United States National Museum for 1880. Vol. III. 1881. 8vo., pp. 595, 5 cuts, 2 plates.

CONTENTS.

- BEAN, T. H. Description of a new hake (*Phycis earlii*), from South Carolina, and a note on the occurrence of *Phycis regius* in North Carolina.
 - Check-list of duplicates of North American fishes distributed by the Smithsonian Institution in behalf of the United States National Museum, 1877-'80.
- CATTIE, S. T. On the genitalia of male eels and their sexual characters. ENDLICH, F. M. List of species and varieties of minerals in the National Museum of the United States in 1879.
- GARMAN, S. Synopsis and descriptions of the American Rhinobatidæ.
- GILL, T. On the identity of the genus Leurynnis Lockington, with Lycodopsis Collett.
- GOODE, G. B. Descriptions of seven new species of fishes from deep soundings on the Southern New England coast, with diagnoses of two undescribed genera of flounders and a genus related to Merlucius.
 - Fishes from the deep waters on the South coast of New England obtained by the United States Fish Commission in the summer of 1880.
 - The frigate mackerel (Auxis Rochei) on the New England coast.
 - Notacanthus phasganorus, a new species of Notacanthidæ from the Grand Banks of Newfoundland.
- HAY, O. P. On a collection of fishes from Eastern Mississippi.

425. Proceedings of National Museum. Vol. III-Continued.

HEILPRIN, A. On some new species of eocene molluscs from the Southern United States.

JORDAN, D. S. Notes on a collection of fishes from East Florida, obtained by J. A. Henshall.

Notes on a collection of fishes from St. John's river, Florida, obtained by A. H. Curtiss.

Note on a forgotten paper of Dr. Ayres, and its bearing on the nonenclature of the cyprinoid fishes of the San Francisco markets.

Note on "Sema" and "Dacentrus."

Description of a new species of Caranx (Caranx Beani), from Beaufor'.

North Carolina.

JORDAN, D. S.; GILBERT, C. H. Notes on a collection of fishes from San Diego, California.

Description of new flounder (Xystreurys liolepis), from Santa Catalina Island, California.

Description of a new ray (Platyrhina triseriata), from the coast of California.

Description of a new species of "rock cod" (Sebastichthys services.

from the coast of California.

On the occurrence of Cephaloscyllium laticeps (Duméril) Gill on the

coast of California.

On the oil shark of Southern California (Galeorhinus galeus.)

Description of a new flounder (Pleuronichthys verticalis), from the

coast of California, with notes on other species.

Notes on sharks from the coast of California.

On the generic relations of Platyrhina exasperata.

Description of a new species of Schastichthys (Schastichthys miniatus from Monterey bay, California.

Description of a new species of "rock fish" (Sebastichthys carnatus from the coast of California.

Description of a new species of ray (Raia stellulata), from Monterey. California.

Description of a new species of Xiphister and Apodichthys, from Monterey, California.

Description of two new species of Sebastichthys (Sebastichthys entomelas and Sebastichthys rhodochloris), from Monterey bay, Cal.

Description of a new agonoid fish (Brachiopsis xyosternus), from Monterey bay, California.

Description of a new flounder (Hippoglossoides exilis), from the coast of California.

Description of a new species of ray (Raia rhina) from the coast of California.

Description of two new species of fishes (Ascelicthys rhodorus and Scytalina cerdale), from Neah Bay, Washington Territory.

Scytalina cerdate), from Neah Bay, Washington Territory.

Description of two new species of scopeloid fishes (Sudis ringers and

Myctophum crenulare), from Santa Barbara channel, California.

Description of two new species of flounders (Parophrys ischyurus and Hippoglossoides elassodon), from Puget Sound.

Description of seven new species of sebastoid fishes, from the coast of California.

435. Proceedings of National Museum. Vol. LII—Continued.

JORDAN, D. S.; GILBERT, C. H. Description of a new embiotocoid (Abeona aurora), from Monterey, California, with notes on a related species.

Description of a new flounder (*Platysomatichthys stomias*), from the coast of California.

Description of a new embiotocoid fish (Cymatogaster rosaceus), from the coast of California.

Description of a new species of deep-water fish (*Icichthys Lockingtoni*), from the coast of California.

Description of a new embiotocoid fish (Ditrema atripes), from the coast of California.

Description of a new scorpsenoid fish (Sebastichthys maliger), from the coast of California.

Description of a new scorpsonoid fish (Sebastichthys proriger), from Monterey Bay, California.

Description of a new agonoid (Agonus vulsus), from the coast of California.

Description of a new species of Hemirhamphus (Hemirhamphus Rosæ), from the coast of California.

Description of a new species of notidanoid shark (*Hexanchus corinus*), from the Pacific coast of the United States.

Description of a new species of Nomichthys (Nemichthys avocetta), from Puget Sound.

Description of a new species of Paralepis (Paralepis coruscans), from the Straits of Juan de Fuca.

List of the fishes of the Pacific coast of the United States, with a table showing the distribution of the species.

On the generic relations of Belone exilis (Girard).

Notes on a collection of fishes from Utah Lake.

Description of a new species of rockfish (Sebastichthys chrysomelas), from the coast of California.

LAWRENCE, G. N. Description of a new species of bird of the family Turdidæ, from the island of Dominica, W. I.

Description of a new species of parrot of the genus Chrysotis, from the island of Dominica.

Description of a new species of Icterus, from the West Indies.

LOCKINGTON, W. N. Remarks on the species of the genus Chirus found in San Francisco market, including one hitherto undescribed.

Description of a new fish from Alaska (Uranidea microstoma).

Description of a new species of Agonidas (Brachyopsis verrucosus), from the coast of California.

Description of a new genus and some new species of California fishes (Icosteus ænigmaticus and Osmerus attenuatus).

Description of a new chiroid fish (Myriolepis zonifer), from Monterey Bay, California.

Description of a new sparoid fish (Sparus brachysomus), from Lower California.

Note on a new flat fish (Lepidopsetta isolepis), found in the markets of San Francisco.

Description of a new species of Prionotus (Prionotus stephanophrys), from the coast of California.

RATHBUN, R. The littoral marine fauna of Provincetown, Cape Cod, Massachusetts.

425. Proceedings of National Museum. Vol. III—Continued.

RIDGWAY, R. Revisions of nomenclature of certain North American birds.

A catalogue of the birds of North America.

Catalogue of Trochilidæ in the collection of the United States National Museum.

RYDER, J. A. On Camaraphysema, a new type of sponge.

List of the North American species of myriapods belonging to the family of the *Lysiopetalida*, with a description of a blind form from Luray Cave, Virginia.

SMITH, R. On the occurrence of a species of Cremnobates at San Diego.
California.

SMITH, S. I. Preliminary notice of the crustacea dredged in 64 to 325 fathoms off the south coast of New England by the United States Fish Commission, in 1880.

SWAN, J. G. The surf smelt of the northwest coast, and the method of taking them by the Quillehute Indians, West Coast of Washington Territory.

The eulachon or candle fish of the northwest coast.

VERRILL, A. E. Notice of recent additions to the marine invertebrats of the northeastern coast of America, with descriptions of new

genera and species, and critical remarks on others.

Part II.—Mollusca, with notes on Annelida, Echinodermata, etc...

collected by the United States Fish Commission.

Part III.—Catalogue of mollusca recently added to the fauna of

Southern New England.

WHITE, U. A. Note on the occurrence of Productus giganteus in Cal.

Note on Acrothele,

Description of a new cretaceous Pinna from New Mexico. .

Note on the occurrence of Stricklandia salteri and S. davidsoni in Georgia.

Description of a very large fossil gasteropod, from the State of Puebla. Mexico.

Mexico.

Descriptions of new invertebrate fossils from the mesozoic and cenozoic rocks of Arkansas, Wyoming, Colorado, and Utah.

- 426. A Synopsis of the Scientific Writings of Sir William Herschel. By Edward S. Holden and Charles S. Hastings. 1881. 8vo., pp. 118.
- 427. Record of Recent Progress in Science, 1879 and 1880. Astronomy. By Edward S. Holden. 1881. 8vo., pp. 39.
- 428. Record of Recent Progress in Science, 1879 and 1880. Geology and Mineralogy. By George W. Hawes. 1881. 8vo., pp. 30.
- 429. Record of Recent Progress in Science, 1879 and 1880. Physics and Chemistry. By George F. Barker. 1881. 8vo., pp. 65.
- 430. Record of Recent Progress in Science, 1879 and 1880. Botany. By WM. G. FARLOW. 1881. 8vo., pp. 19.

- 431. Record of Recent Progress in Science, 1879 and 1880. Zoology. By THEODORE GILL. 1881. 8vo., pp. 62.
- 432. Record of Recent Progress in Science, 1880. Anthropology. By Otis T. Mason. 1881. 8vo., pp. 51.
- 433. Report of a visit to the Luray Cavern, in Page County, Virginia, under the auspices of the Smithsonian Institution, July 13 and 14, 1880. 1881. 8vo., pp. 12, 6 woodcuts.
- 434. Report of the National Museum Building Commission and of the Architects for 1880. 1881. 8vo., pp. 12.
- 435. Discussion of the Barometric Observations of E. S. Snell. By F. H. Loud. 1881. 8vo., pp. 23, 7 woodcuts.
- 436. List of Periodicals received by the Smithsonian Institution. 1881. 8vo., pp. 9.
- 437. Check-list of Publications of the Smithsonian Institution to December, 1881. 1881. 8vo., pp. 22.
- 438. Reports of Astronomical Observatories for 1880. By E. S. Holden and George H. Bæhmer. 1881. 8vo., pp. 128.
- 439. Warming and Ventilating Occupied Buildings. By ARTHUR MORIN. Translated for the Smithsonian Institution by Clarence B. Young. 1882. 8vo., pp. 92, 32 woodcuts.
- 440. Articles on Anthropological Subjects contributed to the Annual Reports of the Smithsonian Institution from 1863 to 1877. By CHARLES RAU. 1882. 8vo., pp. 180, 53 woodcuts.

RAU, C., Preface by. (February, 1882.)

BAEGERT, J. Account of the aboriginal inhabitants of the Californian peninsula. Translated and arranged for the Smithsonian by Chas. Rau. From Smithsonian Report for 1868-64. No. 361.

Agricultural implements of the North American stone period. From Report for 1863.

Artificial shell deposits in New Jersey. From Report for 1864. No. 862. Indian pottery. From Report for 1866. No. 368.

Drilling in stone without metal. From Report for 1868. No. 372.

Deposit of agricultural flint implements in Southern Illinois. From Report for 1868. No. 370.

Memoir of C. F. P. Von Martius. From Report for 1869. No. 251.

Ancient aboriginal trade in North America. From Report for 1872. No. 385.

North American stone implements. From Report for 1872. No. 882.

ROMER, F. F. Prehistoric antiquities of Hungary. Translated by Chas. Rau. From Report for 1876. No. 392.

Stock in trade of an aboriginal lapidary. From Report for 1877. No. 402. Observations on a gold ornament from a mound in Florida. From Report for 1877. No. 403.

- 441. The Constants of Nature. Part V. A Recalculation of the Atomic Weights. By Frank Wigglesworth Clarke. pp. 293.
- 442. Annual Report of the Board of Regents of the Smithsonian Institution, for the year 1880. Forty-sixth Congress, 3d Session, Senate Mis. Doc. No. 31. 1881. 8vo., pp. 782, 13 woodcuts.

BAIRD, S. F. Secretary's report of operations.

List of periodicals received by the Institution.

Rules for the examination of specimens submitted to the Institution.

TAYLOR, F. W. Report of the chemist.

Additions to the collections of the National Museum.

Additions to Museum by the Berlin International Fisheries Exhibition. STEVENSON, J. Report of explorations in New Mexico and Arizona. Receipts and distribution of specimens.

GOODE, G. B. The first decade of the U.S. Fish Commission.

Report of the Executive Committee on the Henry statue.

Report of the Executive Committee for 1880.

BOARD OF REGENTS, Proceedings of

Report of the National Museum Building Commission for 1880.

Report of the Superintending Architects of the Museum Building.

RECORD OF RECENT SCIENTIFIC PROGRESS:

BAIRD, S. F. Introduction.

Holden, E. S. Astronomy.

HAWES, G. W. Geology.

BARKER, G. F. Physics.

BARKER, G. F. Chemistry.

HAWES, G. W. Mineralogy.

FARLOW, W. G. Botany. GILL, T. Zoology.

MASON, O. T. Anthropology.

MASON, O. T. Bibliography of anthropology.

MASON, O. T. Abstracts of anthropological correspondence.

MASON, O. T. Report on Luray Cavern.

LOUD, F. H. Discussion of Prof. Snell's barometric observations.

HENRY, J. Investigation of illuminating materials.

HOLDEN, E. S.; HASTINGS, C. S. Synopsis of the scientific writings of William Herschel.

HOLDEN, E. S.; BEHMER, G. H. Reports of astronomical observatories.

- 443. Results of Meteorological Observations made at Providence, Rhode Island, extending over a period of forty-five years, from December. 1831, to December, 1876. By ALEXIS CASWELL. 1882. 4to, pp. 40.
- 444. Guide to the Flora of Washington and Vicinity. By LESTER F. WARD. 1881. 8vo., pp. 265. One map. Bulletin of the National Museum, No. 22.

- 445. Plan of Organization and Regulations of the United States National Museum. By G. Brown Goode. 1882. 8vo., pp. 58. Two woodcuts. (Circular of U. S. National Museum, No. 1.)
- 446. Circular addressed to Friends of the Museum. By Spencer F. Baird. 1882. 8vo., pp. 2. (Circular of U. S. National Museum, No. 2.)
- 447. Circular in reference to Petroleum Collections. By Spencer F. Baird. 1882. 8vo., pp. 4. (Circular of U.S. National Museum, No. 3.)
- 448. Circular concerning the Department of Insects. By Spencer F. Baird. 1882. 8vo., pp. 2. (Circular of U.S. National Museum, No. 4.)
- 449. Establishment and Officers of the Smithsonian Institution and National Museum, January 1, 1882. 1882. 8vo., pp. 2. (Circular of U. S. National Museum, No. 5.)
- 450. Classification and Arrangement of the Materia Medica Collection. By James M. Flint. 1882. 8vo., pp. 2. (Circular of U.S. National Museum, No. 6.)
- 451. A Classification of the Forms in which Drugs and Medicines appear, and are administered. By James M. Flint. 1882. 8vo., pp. 8. (Circular of U. S. National Museum, No. 7.)
- 452. Memoranda for Collectors of Drugs for the Materia Medica Section of the National Museum. By James M. Flint. 1882. 8vo., pp. 2. (Circular of U. S. National Museum, No. 8.)
- 453. Circular in reference to the Building-Stone Collection. By Spencer F. Baird. 1882. 8vo., pp. 6. (Circular of U. S. National Museum, No. 9.)
- 454. Two letters on the work of the National Museum. By BARNET PHILLIPS. 1882. 8vo., pp. 10. (Circular of U.S. National Museum, No. 10.)
- 455. A Provisional Classification of the Food Collections. By G. Brown Goode. 1882. 8vo., pp. 18. (Circular of U. S. National Museum, No. 11.)
- 456. Classification of the Collections of the U. S. National Museum to illustrate the Art of Taxidermy. By W. T. HORNADAY. 1882. 8vo., pp. 2. (Circular of U. S. National Museum, No. 12.)
- 457. Outline of a Scheme of Classification for the Collections in the United States National Museum. By G. Brown Goode. 1882. 8vo., pp. 4. (Circular of U. S. National Museum, No. 13.)

- 458. Circular requesting material for the Library of the United States National Museum. By Spencer F. Baird. 1882. 8yo., pp. 4. (Circular of U. S. National Museum, No. 14.)
- 459. The Organization and Objects of the United States National Museum. By G. Brown Goode. 1882. 8vo., pp. 4. (Circular of U. S. National Museum, No. 15.)
- **460.** Directions for Collecting and Preserving Plants. By LESTER F. WARD. 1882. 8vo., pp. 32.
- 461. Check-List. Flora of Washington and Vicinity. By LESTER F. WARD. 1882. 8vo., pp. 62.
- 462. Catalogue of Old World Birds in the U. S. National Museum. By ROBERT RIDGWAY. 1882. 8vo., pp. 20.
- 463. Bibliography of the Fishes of the Pacific Coast of the United States. to the end of 1879. By Theodore Gill. 1882. 8vo., pp. 77. (M. C. XXIII.) Bulletin of National Museum, No. 11.
- 464. Directions for Collecting and Preserving Fish. By TARLETON H. BEAN. 1881. 8vo., pp. 6.
- 465. List of Marine Invertebrates, mainly from the New England Coast, distributed by the U.S. National Museum. Series II. By RICHARD RATHBUN. 1881. 8vo., pp. 6.
- 466. Directory of Officers, Collaborators, Employés, etc., of the Smithsonian Institution, National Museum, Geological Survey, Bureau of Ethnology, and Fish Commission. 1882. 8vo., pp. 8.
- 467. Proceedings of the United States National Museum for 1881. Vol. IV. 1882. 8vo., pp. 600, 13 cuts, 2 plates. (M. C. xxxx.)

- BAIRD, S. F. Notes on certain aboriginal shell mounds on the coast of New Brunswick and of New England.
- BEAN, T. H. Descriptions of new species of fishes (Uranidea marginata.

 Potamocottus Bendirei) and of Myctophum crenulare J. and G.

Notes on some fishes from Hudson's Bay.

Descriptions of new fishes from Alaska and Siberia.

Directions for collecting and preserving fish.

A preliminary catalogue of the fishes of Alaskan and adjacent waters. A partial bibliography of the fishes of the Pacific coast of the United States and of Alaska, for the year 1880.

Notes on a collection of fishes made by Capt. Henry E. Nichols. U. S. N., in British Columbia and Southern Alaska, with descriptions of new species and a new genus (*Delolopis*).

See also under Goode and Bean.

BENDIRE, C. Notes on Salmonidæ of the upper Columbia.

467. Proceedings of National Museum. Vol. IV-Continued.

BOYD, C. H. Remains of the walrus (?) in Maine.

CARLIN, W. E. Observations of Siredon lichenoides.

DALL, W. H. On the genera of chitons.

On certain limpets and chitons from the deep waters off the eastern coast of the United States.

ENDLICH, F. M. An analysis of water destructive to fish in the Gulf of Mexico.

FARLOW, W. G. Report on the contents of two bottles of water from the Gulf of Mexico, forwarded by the Smithsonian Institution.

GILBERT, C. H. See under Jordan and Gilbert.

GILL, T. Note on the Latiloid genera.

GLAZIER, W. C. W. On the destruction of fish by polluted waters in the Gulf of Mexico.

GOODE, G. B. The Taxonomic relations and geographical distribution of the members of the swordfish family, (Xiphiidæ.)

GOODE, G. B.; BEAN, T. H. Description of a new species of fish, Apogon pandionis, from the deep water off the mouth of Chesapeake Bay.

Benthodesmus, a new genus of deep-sea fishes, allied to Lepidopus.

HAWES, G. W. On the mineralogical composition of the normal mesozoic diabase upon the Atlantic border.

On the determination of feldspar in thin sections of rocks.

INGERSOLL, E. On the fish mortality in the Gulf of Mexico.

JAPANESE LEGATION, Washington. Catalogue of a collection of Japanese cotton fibre presented to the U.S. National Museum by the Government of Japan, together with the amount of the annual crop of Japan and the price of cotton.

JOHNSON, S. H. Notes on the mortality among fishes of the Gulf of Mexico.

JORDAN, D. S.; JOUY, P. L. Check-list of duplicates of fishes from the Pacific coast of North America, distributed by the Smithsonian Institution in behalf of the United States National Museum, 1881.

JORDAN, D. S.; GILBERT, C. H. Notes on the fishes of the Pacific coast of the United States.

Description of Sebastichthys mystinus.

Description of a new species of Ptychochilus (Ptychochilus Harfordi) from Sacramento river.

Note on Raia inornata.

Notes on a collection of fishes, made by Lieut. Henry E. Nichols, U. S. N., on the west coast of Mexico, with descriptions of new species.

List of fishes collected by Lieut. Henry E. Nichols, U. S. N., in the Gulf of California and on the west coast of Lower California, with descriptions of four new species.

Descriptions of thirty-three new species of fishes from Mazatlan, Mexico.

Description of a new species of *Pomadasys* from Mazatlan, with a key to the species known to inhabit the Pacific coasts of tropical America.

467. Proceedings of National Museum. Vol. IV—Continued.

Description of a new species of Xenichthys (Xenichthys ocyurus) from the west coast of Central America.

Descriptions of five new species of fishes from Mazatlan, Mexico.

JOUY, P. L. Description of a new species of Squalius (Squalius alicia) from Utah Lake.

LAWRENCE, G. N. Description of a new sub-species of Lozigilla from the island of St. Christopher, West Indies.

LOCKINGTON, W. N. Description of a new genus and species of Cottida. LUGGER, O. The occurrence of the Canada porcupine in Maryland.

Mckar, C. L. A review of the genera and species of the family Centrarchidæ, with a description of one new species.

MOORE, M. A. Fish mortality in the Gulf of Mexico.

PIRZ, A. Methods of making and preserving plaster casts.

PLATEAU, F. The rapid preparation of large myological specimens.

PORTER, J. Y. On the destruction of fish by poisonous water in the Gulf of Mexico.

RATHBUN, R. List of marine invertebrates, mainly from the New England coast, distributed by the United States National Museum.
(Series II.)

List of Marine invertebrates from the New England coast, distributed by the United States National Museum. (Series III.)

RAU, C. List of anthropological publications by

RIDGWAY, R. On a duck new to the North American fauna.

On Amazilia Yucatanensis, (Cabot.) and A. cerviniventris, Gould.

A review of the genus Centurus, Swainson.

List of species of Middle and South American birds not contained in the United States National Museum.

List of special desiderata among North American birds.

Catalogue of Old World birds in the United States National Museum.

Notes on some Costa Rican birds.

Description of a new fly-catcher, and a supposed new petrel, from the Sandwich Islands.

Description of a new owl from Porto Rico.

Descriptions of two new thrushes from the United States.

On two recent additions to the North American bird fauna, by L. Belding.

RYDER, J. A. On Semper's method of making dry preparations.

SHUFELDT, R. W. Remarks upon the osteology of Opheosaurus ventralis.

SMITH, R. Description of a new Gobioid fish, (Othomops cos.) from Sar. Diego, California.

Description of a new species of Gobiesox, (Gobiesox rhessodon,) from San Diego, California.

SMITH, S. B. On the Chinook names of the salmon in the Columbia River.

STEJNEGER, L. Description of two new races of Myadestes obscurus, Laft. Translation. Metallic castings of delicate natural objects.

TRUE, F. W. On the North American land tortoises of the genus Xero-bates.

On the rare rodent Cricetodipus parvus (Baird) Coues.

467. Proceedings of National Museum. Vol. IV-Continued.

WARD, L. F. Catalogue of a collection of Japanese woods presented to the U. S. National Museum by the University of Tokio, Japan.

WHITE, C. A. On certain cretaceous fossils from Arkansas and Colorado.

ANONYMOUS. The comparative action of dry heat and sulphurous acid upon putrefactive bacteria.

APPENDICES-Circulars of the U. S. National Museum:

- No. 1. Plan of organisation and regulations of the Museum.
- No. 2. Circular addressed to the friends of the Museum.
- No. 3. Circular in reference to petroleum collections.
- No. 4. Circular concerning the department of insects.
- No. 5. Establishment and officers of the Smithsonian Institution and Museum.
- No. 6. Classification and arrangement of the materia medica collection.
- No. 7. A classification of the forms in which drugs and medicines appear and are administered.
- No. 8. Memoranda of collectors of drugs for the materia medica section of the National Museum.
- No. 9. Circular in reference to the building-stone collection.
- No. 10. Two letters on the work of the National Museum.
- No. 11. A provisional classification of the food collections.
- No. 12. Classification of the collections to illustrate the art of taxidermy.
- No. 13. Outline of a scheme of Museum classification
- No. 14. Circular requesting material for the library of the Museum.
- No. 15. The organization and objects of the National Museum.
- No. 16. Plans for the installation of collections.
- No. 17. Contributions and their acknowledgment.
- No. 18. List of publications of the United States National Museum.
- 468. Smithsonian Miscellaneous Collections. Vol. XXII. 1882. 8vo., pp. 1200, cuts 18, plates 4.

CONTENTS.

Proceedings of the U. S. National Museum. Vol. III. 1880. No. 425. Proceedings of the U. S. National Museum. Vol. IV. 1881. No. 467.

- 469. List of Foreign Correspondents of the Smithsonian Institution. Corrected to January, 1882. 1882. 8vo., pp. 174.
- 470. Nomenclator Zoologicus. An Alphabetical List of all Generic Names that have been employed by Naturalists for Recent and Fossil Animals from the earliest times to the close of the year 1879. In two parts. Part I.—List of Generic Names employed in Zoology and Paleontology to the close of the year 1879, chiefly supplemental to those catalogued by Agassiz and Marschall, or indexed in the Zoological Record. By Samuel H. Scudder. 1882. 8vo., pp. 398. Bulletin of the National Museum, No. 19.
- 471. List of Marine Invertebrates from the New England Coast, distributed by the U. S. National Museum. Series III.—Educational Series. By RICHARD RATHBUN. 1881. 8vo., pp. 4.
- 472. Plans for the Installation of Collections in the U. S. National Museum. By G. Brown Goode. 1882. 8vo., pp. 2. (Circular of U. S. National Museum, No. 16.)

- 473. Contributions to the United States National Museum, and their Acknowledgement. 1882. 8vo., pp. 2. (Circular of U. S. National Museum, No. 17.)
- 474. List of Publications of the United States National Museum. 1882. 8vo., pp. 12. (Circular of U. S. National Museum, No. 18.)
- 475. Smithsonian Miscellaneous Collections. Vol. XXIII. 1882. 8vo., pp. 1003. Bulletins of the National Museum, Nos. 11 to 15.

- GILL, T. Bibliography of fishes of Pacific coast. Bulletin No. 11. No. 468.
- JORDAN, D. S.; BRAYTON, A. W. Fishes of Alleghany region of South Carolina, Georgia, and Tennessee, and synopsis of Catostomidæ. Bulletin No. 12. No. 808.
- EGGERS, H. F. A. Flors of St. Croix and the Virgin Islands. Bulletin No. 18. No. 318.
- GOODE, G. B. Catalogue of collection illustrating animal resources and fisheries of the United States. Bulletia No. 14. No. 326.
- Kumlien, L. Natural history of Arctic America. Bulletia No. 15. No. 342.
- 476. First Annual Report of the Bureau of Ethnology to the Secretary of the Smithsonian Institution, 1879-80. By J. W. Powell, Director. 1881. Royal 8vo., pp. 638. 343 cuts, 54 plates, 1 folded map.

CONTENTS.

POWELL, J. W. Report of the Director of the Bureau.

Powell, J. W. On the evolution of language.

POWELL, J. W. Mythology of the North American Indians.

Powell, J. W. Wyandot government.

Powerl, J. W. Limitations to the use of some Anthropological dats.

YARROW, H. C. Study of the mortuary customs of the North American Indians.

HOLDEN, E. S. Studies in Central American picture-writing.

ROYCE, C. C. Cessions of land by Indian tribes to the United States.

MALLERY, GARRICK. Sign language among North American Indians.

compared with that among other peoples and deaf-mutes.

Pilling, J. C. Catalogue of linguistic manuscripts in the library of the

Bureau of Ethnology.

Dorsey, J. O. Illustration of the method of recording Indian languages.

How the rabbit caught the sun in a trap. (An Omaha myth.)

GATSCHET, A. S. Details of an Indian conjurer's practice. (In the Klamath Lake dialect.)

GATSCHET, A. S. The relapse. (In the Klamath Lake dialect.)

RIGGS, S. R. . A dog's revenge. (A Dakota Fable.)

- 477. Report on International Exchange, with list of Official Publications of the U.S. Government, between 1868 and 1881. By George H. BEHMER. 1882. 8vo., pp. 113.
- 478. Catalogue of Publications of the Smithsonian Institution up to July
 1, 1882, with Index to all Articles in the "Smithsonian Contributions
 to Knowledge," "Miscellaneous Collections," "Annual Reports,"
 "Bulletins and Proceedings of the U. S. National Museum," etc.
 By WILLIAM J. RHEES. 1882. 8vo., 1882, pp. .

•

CLASSIFIED LIST OF SEPARATE PUBLICATIONS

OF THE

SMITHSONIAN INSTITUTION.

SUBJECTS.

- I. ANATOMY, PHYSIOLOGY, MEDICINE, and SURGERY.
- II. ANTHROPOLOGY. (See also Philology.)
- III. ARCHÆOLOGY. (See Anthropology.)
- IV. ARCHITECTURE.
 - V. ASTRONOMY.
- VI. BIBLIOGRAPHY.
- VII. BIOGRAPHY.
- VIII. BIOLOGY. (See also Anatomy, Botany, Microscopy, Palsontology, and Zoology.)
 - IX. BOTANY.
 - X. CHEMISTRY and TECHNOLOGY.
 - XI. ELECTRICITY and MAGNETISM.
 - XII. ETHNOLOGY. (See Anthropology, also Philology.)
- XIII. GEOLOGY. (See also Palsontology.)
- XIV. MAGNETISM. (See Electricity, etc.)
 - XV. MATHEMATICS.
- XVI. MEDICINE and SURGERY. (See Anatomy, etc.)
- XVII. METEOROLOGY.
- XVIII. MICROSCOPY.
 - XIX. MINERALOGY.
 - XX. MISCELLANEOUS. XXI. NATURAL HISTORY. (See Biology.)
- XXII. PALEONTOLOGY.
- XXIII. PHILOLOGY.
- XXIV. PHYSICAL GEOGRAPHY.
- XXV. PHYSICS. (See also Terrestrial Physics.)
- XXVI. PHYSIOLOGY. (See Anatomy, etc.)
 XXVII. TECHNOLOGY. (See Chemistry, etc.)
- XXVIII. TERRESTRIAL PHYSICS.
 - XXIX. ZOOLOGY.—Birds.

Fishes.

Insects.

Mammals.

Mollusks.

Radiates.

Reptiles, and Batrachians.

Shells. (See Mollusks.)

I.-ANATOMY, PHYSIOLOGY, MEDICINE, AND SURGERY.

No Cata
Adams. Subcutaneous surgery (Toner Lecture, No. 6)
Brown-Séquard. Dual character of the brain (Toner Lecture, No. 2)
DACOSTA. The heart (Toner Lecture, No. 8)
DEAN. Medulla oblongata (s. c.)
FLINT. Classification of drugs and medicines
FLINT. Classification of materia medica collection
FLINT. Memoranda for collection of drugs
Holmgren. Color-blindness.
JONES. Chemical and physiological investigations (S. C.)
KEEN. Continued fevers (Toner Lecture, No. 5)
LAUTENBACH. Effect of irritation of a polarized nerve
LEIDY. Flora and fauna within living animals (s. c.)
MITCHELL and MOREHOUSE. Chelonia (s. c.)
MITCHELL. Venom of rattlesnake (s. c.)
SHAKESPEARE. Reparatory inflammation (Toner Lecture, No. 7)
WARING. Sanitary drainage of Washington (Toner Lecture, No. 8)
WEISMANN. Change of Mexican axolotl
Wood. Morbid and normal physiology of fever (s. c.)
Wood. Study of fever (Toner Lecture, No. 4)
WOODWARD. Cancerous tumors (Toner Lecture, No. 1)
WYMAN. Rana pipiens (s. c.)
price (si vijeene in the contract of the contr
·
II.—ANTHROPOLOGY. (See also Philology.)
Abbott. Stone age in New Jersey
Archæology, circular on205.
BEGERT. Aboriginal inhabitants of California
BŒHMER. Index to archeological articles in Smithsonian publications
Bransford. Archæological researches in Nicaragua (s. c.)
Dall. Prehistoric man, Aleutian Islands (s. c.)
DESOR. Palafittes of the Lake of Neuchatel
GIBBS. Ethnological and philological instructions
GIBBS. Tinneh, or Chepewyan Indians
GILLMAN. Characteristics of ancient man in Michigan.
GILLMAN. Mound builders and platycnemism
HABEL. Sculptures in Guatemala (s. c.)
HALDENAN Polychrome beed from Floride

ANTHROPOLOGY—Continued.	No. i Satalog
HARDISTY. Loucheux Indians	
Haven. American archæology (s. c.)	
Indians, photographic portraits of, list of	
Jones. Aboriginal structures in Georgia	
JONES. Antiquities of Tennessee (s. c.)	
Jones. Kutchin Indians	3
Knight. Savage weapons	
LAPHAM. Antiquities of Wisconsin (s. c.)	
Mason. Anthropological investigations in 1879	
Mason, " 1880	4
Mason. Latimer antiquities from Porto Rico	3
MASON. Recent progress in anthropology	4
MAYER. Mexican archeology (s. c.)	
Morgan. Circular relative to relationship	
Morgan. Systems of consanguinity (s. c.)	2
Pickering. Gliddon mummy case (s. c.)	_ 2
Powell. First annual report of Bureau of Ethnology	_ 4
RAU. Ancient aboriginal trade in North America	3
RAU. Archæological collection in Smithsonian Institution (s. c.)	
RAU. Articles on anthropological subjects, (from Reports S. I.)	_ 4
RAU. Artificial shell deposits in New Jersey	_ 3
RAU. Deposit of agricultural flint implements	3
RAU. Drilling in stone without metal	3
RAU. Gold ornament in mound in Florida	. 4
RAU. Indian Pottery	3
RAU. North American stone implements	_ 3
RAU. Palenque tablet (s. c.)	_ 3
RAU. Stock-in-trade of aboriginal lapidary	4
ROMER. Prehistoric antiquities of Hungary (translated by RAU)	3
Ross. Eastern Tinneh Indians	_ 3
Squire and Davis. Ancient monument of Mississippi valley (s. c.)	-
SQUIER and DAVIS. Aboriginal monuments of New York (s. c.)	- :
STANLEY. Catalogue of Indian Portraits	_
SWAN. Indians of Cape Flattery (s. c.)	_ 2
Swam. Haidah Indians (s. c.)	2
WHITTLESEY. Ancient mining at Lake Superior (s. c.)	
WHITTI TERY Ancient works in Ohio (s. c.)	

III.—ARCHÆOLOGY. (See Anthropology.)

IV.—ARCHITECTURE.	
C m	io. in Mogue.
Building-stones, circular relative to	453
OWEN. Hints on public architecture	P
Report of National Museum building commission and architects for 1879	409
Report of National Museum building commission and architects for 1880	434
V.—ASTRONOMY.	
A Ol (4 D 4 D 1)	
ABBE. Observatories of Dorpat and Poulkova	
ALEXANDER. Harmonies of the solar system (s. c.)	
Astronomical telegram circular	
Downes. Occultations 1848 (s. c.)	8
Downes. " 1849 (s. c.)	9
Downes. " 1850 (s. c.)	10
Downes. " 1851 (s. c.)	11
Downes. " 1852 (s. c.)	29
Downes. " 1858 (s. c)	54
DEAPER. Silvered-glass telescope (s. c.)	180
Eclipse, annular, 1854	66
GILLISS. Solar eclipse, Peru (s. c.)	100
Gould. History of Neptune	18
Gould. Transatlantic longitude (8. c.)	223
HILL. Map solar eclipse	101
HOLDEN and BEHMER. Reports of observatories for 1880	438
Holden and Hastings. Synopsis of Sir William Herschel's writings	426
HOLDEN. Index catalogue of Nebulæ	311
Holden. Recent progress in astronomy	427
Holden. Reports of observatories for 1879	410
KANE. Astronomical observations arctic seas (s. c.)129	, 198
Newcomb. Integrals of planetary motion (s. c.)	281
NEWCOMB. Orbit of Neptune (s. c.)	199
Newcomb. Orbit of Uranus (s. c.)	262
RUNKLE. Planetary tables (s. c.)	79
Runkle. Planetary tables, supplement	94
Star-map, circumpolar	350
Star-map, of the visible heavens	359
STOCKWELL. Orbits of planets (s. c.)	232

WALKER. Researches Neptune (s. c.)

	A.S	STRONOMY—Continued.	No. in atalogue.
WALKER.	Ephemeris Neptu	ine, 1848 (s. c.)	
WALKER.	"	1795-1849 (s. c.)	_ 5
WALKER.	"	1850 (s. c.)	. 6
WALKER.	"	1851_ (s. c.)	
WALKER.	"	1852 (s. c.)	_ 24
	•	VI.—BIBLIOGRAPHY.	
Binney.	• . •	onchology, part I	142
Binney.	44	" " II	_ 174
Exchanges,	list of Smithsonia	an, part I	_ 73
Exchanges,		, " II	. 85
Exchanges,		to 1858	_ 117
Gill. Bib	oliography of fishes	s of Pacific coast	. 463
GIRARD.	Bibliography of na	atural history, 1851	. 48
Holden an	d Hastings. Syr	nopsis of Sir William Herschel's writings	. 42 6
Holden.	Index catalogue of	f books and memoirs on Nebulse	. 311
		and scientific in United States	
Jewett. (Datalogue system		. 47
Publication:	s, Smithsonian In	stitution, list of (successive issues,)74, 203	
		245, 278, 290, 301, 344, 43	
		nstitution, list of	
		ies in library Smithsonian Inst	
		of Smithsonian publications	
		110	
Watson.	Bibliography of N	Torth American botany	. 258
		VII.—BIOGRAPHY.	
		enry	
Henry. E	ulogy on Alex. D	allas Bache	. 37 9
Henry, Mer	norial of (By nur	merous contributors.)	. 356
		entific writings of Sir William Herschel	
HAYER. H	lenry as a discover	er	. 417
RAU. Men	oir of Von Marti	us	251
Reres. Li	fe and bequest of	James Smithson	330
laylor. B	femoir of and scie	entific work of Joseph Henry	339
Welling.	Life and characte	or of Joseph Henry	338

VIII.—BIOLOGY. (See also Anatomy, Microscopy, Palkontology, Botas	Ŧ,
and Zoology.)	כ
Catalog	
r, and a second	84
	.68
	19
	.63
Circular for collecting shells176, 3	
	107
	37
	L78
	112
Directions for collecting and preserving specimens	34
Directions for collecting and preserving specimens of nests and eggs1	L39
EDWARDS. Directions for collecting diatomacea	366
GILL. Recent progress in zoology	131
GIRARD. Bibliography of natural history	48
	326
GOODE. Classification of animal resources	297
	294
	342
· · · · · · · · · · · · · · · · · · ·	44
	164
	261
Proceedings of the U. S. National MuseumI, 332; II, 333; III, 425; IV. 4	
Periodical phenomena register	
•	470 50
,	303
WEISMANN. Change of Mexican axolotl	40 1
IX.—BOTANY.	
COOPER. Forests and trees of North America	351
EGGERS. Flora of St. Croix	313
FARLOW. Recent progress in botany	130
	22
	42
	32
	43
	95

BOTANY—Continued. No. Catalo	in
HARVEY. North American marine algae, complete (s. c.)	96
LEIDY. Flora within animals (s. c.)	44
Periodical phenomena, register of65,	148
	367
Torrey. Batis maritima (s. c.)	60
Torrey. Darlingtonia Californica (s. c.)	61
HARVEY. Plantæ Frémontianæ (s. c.)	46
	461
WARD. Directions for collecting and preserving plants	160
	444
WATSON. Index to North American botany	258
Wood. Fresh water alge (s. c.)	241
X.—CHEMISTRY AND TECHNOLOGY.	
BARKER. Recent progress in chemistry	429
BECKER. Atomic weight determinations	358
BOOTH and MORFIT. Chemical arts	27
CLARKE. Expansion tables	289
CLARKE. Recalculation of atomic weights	441
CLARKE. Specific gravity tables	255
CLARKE. " " 1st supplement	288
CLARKE. Specific heat tables	276
GIBBS and GENTH. Ammonia-cobalt bases (s. c.)	88
HARE. Explosiveness of nitre (s. c.)	17
gg	3 89
• • • • • • • • • • • • • • • • • • • •	376
JONES. Chemical and physiological investigations (S. C.)	82
	439
	371
6	327
	405
WETHERILL. Ozone and antozone	355
XI.—ELECTRICITY AND MAGNETISM.	
BACHE. Magnetic discussion, part I (s. c.)	113
	121
	132

ELECTRICITY AND MAGNETISM—Continued. No.	in KCC.
	162
	175
•	186
	195
· · · · · · · · · · · · · · · · · · ·	166
	223
	239
HAYES. Arctic observations (s. c.)	196
	115
KANE. Arctic observations (s. c.)97,	198
Locke. Observations on terrestrial magnetism (s. c.)	35
SECCHI. Researches on electrical rheometry (s. c.)	36
SONNTAG. Terrestrial magnetism, Mexico (8. C.)	114
· · · · · · · · · · · · · · · · · · ·	405
XII.—ETHNOLOGY. (See Anthropology and Philology.)	
XIII.—GEOLOGY. (See also PALEONTOLOGY.)	
Cooper. Physical geography of North America	
COOPER. Physical geography of North America	428
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (8. C.)	428
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (s. c.) HITCHCOCK. Surface geology (s. c.)	428 248 90
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (8. C.)	428 248 90
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (s. c.) HITCHCOCK. Surface geology (s. c.) MASON. Luray Cavern in Virginia PUMPELLY. Geology of China, Mongolia, and Japan (s. c.)	428 248 90 433 202
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (s. c.) HITCHCOCK. Surface geology (s. c.) MASON. Luray Cavern in Virginia PUMPELLY. Geology of China, Mongolia, and Japan (s. c.) WHITTLESEY. Fluctuations of lakes (s. c.)	428 248 90 433 202 119
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (s. c.) HITCHCOCK. Surface geology (s. c.) MASON. Luray Cavern in Virginia PUMPELLY. Geology of China, Mongolia, and Japan (s. c.)	428 248 90 433 202 119
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (s. c.) HITCHCOCK. Surface geology (s. c.) MASON. Luray Cavern in Virginia PUMPELLY. Geology of China, Mongolia, and Japan (s. c.) WHITTLESEY. Fluctuations of lakes (s. c.)	428 248 90 433 202 119
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (s. c.) HITCHCOCK. Surface geology (s. c.) MASON. Luray Cavern in Virginia PUMPELLY. Geology of China, Mongolia, and Japan (s. c.) WHITTLESEY. Fluctuations of lakes (s. c.)	428 248 90 433 202 119
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (s. c.) HITCHCOCK. Surface geology (s. c.) MASON. Luray Cavern in Virginia PUMPELLY. Geology of China, Mongolia, and Japan (s. c.) WHITTLESEY. Fluctuations of lakes (s. c.)	428 248 90 433 202 119
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (s. c.) HITCHCOCK. Surface geology (s. c.) MASON. Luray Cavern in Virginia PUMPELLY. Geology of China, Mongolia, and Japan (s. c.) WHITTLESEY. Fluctuations of lakes (s. c.) WHITTLESEY. Glacial drift (s. c.)	428 248 90 433 202 119
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (s. c.) HITCHCOCK. Surface geology (s. c.) MASON. Luray Cavern in Virginia PUMPELLY. Geology of China, Mongolia, and Japan (s. c.) WHITTLESEY. Fluctuations of lakes (s. c.) WHITTLESEY. Glacial drift (s. c.) XIV.—MAGNETISM. (See Electricity.)	428 248 90 433 202 119
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (s. c.) HITCHCOCK. Surface geology (s. c.) MASON. Luray Cavern in Virginia PUMPELLY. Geology of China, Mongolia, and Japan (s. c.) WHITTLESEY. Fluctuations of lakes (s. c.) WHITTLESEY. Glacial drift (s. c.) XIV.—MAGNETISM. (See Electricity.) XV.—MATHEMATICS. ALVORD. Tangencies of circles (s. c.)	428 90 433 202 119 197
COOPER. Physical geography of North America HAWES. Recent progress in geology HILGARD. Geology of lower Louisiana (s. c.) HITCHCOCK. Surface geology (s. c.) MASON. Luray Cavern in Virginia PUMPELLY. Geology of China, Mongolia, and Japan (s. c.) WHITTLESEY. Fluctuations of lakes (s. c.) WHITTLESEY. Glacial drift (s. c.) XIV.—MAGNETISM. (See Electricity.) XV.—MATHEMATICS. ALVORD. Tangencies of circles (s. c.)	428 90 433 202 119 197 80 233

XVII.-METEOROLOGY.

	No. in Catalogue
ABBE. Meteorological memoirs	398
Bache. Girard College observations (s. c.) 113, 121, 132, 162, 175, 1	.86, 195
Caswell. Observations, Providence, Rhode Island, 1881-1860 (s. c.)	103
CASWELL. Observations, Providence, Rhode Island, 1831-1876 (s. c.)	443
CHAPPELSMITH. Tornado (s. c.)	59
Clouds, nomenclature of	347
COFFIN. Meteoric fire-ball (s. c.)	221
Coffin. Psychrcmetrical tables	
COFFIN. Winds, northern hemisphere (s. c.)	52
COFFIN. Winds of the globe (s. c.)	268
Colding. The nature of air currents	398
COLDING. The whirlwind of St. Thomas in 1871	398
COOPER, J. G. Forests and trees of North America	351
Directions for meteorological observations	19, 148
Ferrel. Reply to strictures of J. Hann	
Force. Record of auroras (s. c.)	84
GUYOT. Meteorological directions	19, 148
Guyor. Meteorological and physical tables	31, 153
HANN. Diminution of vapor with altitude	398
HANN. Influence of rain on the barometer	398
HANN. Atmospheric pressure and rain-fall	398
HANN. Laws of temperature in ascending currents	398
HANN. Relation between pressure and wind	398
HAYES. Arctic observations (s. c.)	196
HENRY. Circular, altitudes	236
HENRY. " directions for constructing lightning rods	237
Henry. "thunder-storms	235
Henry. " tornadoes	190
HILDRETH AND WOOD. Observations, Marietta, Ohio (s. c.)	120
Hork. Letter on meteoric shower	217
Loud. Discussion of Snell's barometric observations	435
LOOMIS. Storms in 1836 (s. c.)	127
Map of the stars near the north pole for aurora	_ 350
McClintock. Arctic observations (s. c.)	. 146
McParlin. Climate of New Mexico	. 396
MEECH. Intensity sun's heat (s. c.).	83
CLEAVELAND. Meteorological observations, Brunswick, Maine (s. c.)	
KANE. Meteorological arctic observations (s. c.)1	04, 198
Meteorological observations, 1855	93.

METEOROLOGY—Continued.	No. in Catalogue.
Meteorological results, 1854-59, vol. 1	_
Meteorological results, 1854-59, vol. 2	
Meteorological stations and observers 1849-1868	373
Meteors, November	
OLMSTEAD. Aurors (s. c.)	
Periodical phenomena, registry of	.65, 1 4 8
PESLIN. Relation between barometric pressure and air currents	398
Planisphere of the visible heavens, for observation of meteors	359
Rain charts for summer, winter, and year	374
REYE. Rainfall and barometric minima	398
SCHOTT. Base chart of the United States	414
SCHOTT. Tables of rain and snow in the United States (s. c.)	222, 353
Schott. Tables of temperature (s. c.)	 27 7
SMITH. Observations, Washington, Arkansas (s. c.)	131
SOHNCER. Variation of temperature in ascending currents of moist air	398
Temperature chart of U. S. for year.	381, 388
Temperature charts of the U.S. for summer, winter, and year	387
XVIII.—MICROSCOPY.	
BAILEY. Microscopic observations, Southern States (s. c.)	
BAILEY. New microscopic organisms (s. c.)	
BAILEY. Soundings, coast survey (s. c.)	
DEAN. Medulla oblongata (8. C.)	
EDWARDS. Directions for collecting microscopic organisms LEIDY. Flora and fauna within living animals (8. c.)	
LEIDY. Flora and fauna within living animals (s. c.) WOODWARD. Cancerous tumors	
WOODWARD, Cancerous tumors	
XIX.—MINERALOGY.	
BABINET. Diamond and precious stones	
Brezina. Crystallography and crystallophysics	
EGLESTON. Catalogue of minerals	
HAWES. Recent progress in mineralogy	428
XX.—MISCELLANEOUS.	
Relative to the Smithsonian Institution.	
Act of Congress to establish the Smithsonian Institution	
BOEHMER. Report on international exchanges	477

		MISCELL	ANEO	US—Cont		lo. m alogue.
Business	arrangemen	its of the Smith	sonian I	Institutio	n	325
Commit	tee on Organ	nization, report	of			B, L
						K
•	-					324
		-				290
DALLAS.	Address,	laying corner-st	one of S	B miths oni	an Institution	D
Digest of	f act of Con	gress relative to	Smiths	onian Ins	titution	C
Director	y of officers,	collaborators as	nd empl	oyés		466
			_	-	and Museum	449
HENRY.						B
History	of Smithson	ian Institution .	- 			328
					69	, 238
Instituti	ons, list of f	oreign			64, 154, 225, 243	, 469
Journals	of the Boar	d of Regents, s	tatistics,	etc		329
Officers a	and Regents	and act of orga	nization			N
	_	_				380
Proceedi	ngs, Regent	s, 1846				A
						J
	_					260
Bair d .	Circular re	Relative to petrole				447
Baird.			•			446
BAIRD.					al, 1876	307
	•	ed States Nation			•	445
16	11	16	66		2	446
**	44	44	44	44	8	447
: 6	"	44	14	46	4	448
. 6	"	"		"	5	449
	u ·	44	**	"	6	450
44	44	"	46	"	7	451
"	ii	44	"	"	8	452
44	"	"	44	4.6	9	453
14	46	44	44	46	10	454
		"	44	"	11	455
"	"	44	"	"	12	456
14	11	"	"	"	18	457
14	"	16	"	"	14	458
44	"	~ 66	"	44	15	459
16	"	"	"	. "	16	472

MISCELLANEOUS—Continued. No.	. in logue.
Circular of the United States National Museum, No. 17	473
" " " 18	474
Circular requesting material for Museum library	458
Classification of animal resources	297
Contributions to Museum, and acknowledgement	473
GOODE. Animal resources and fisheries	326
GOODE. Classification of Museum collections	457
GOODE. Classification of the food collections	455
GOODE. Installation of collections in Museum	472
GOODE. Organization and objects of the Museum	459
GOODE. Organization and regulations of the Museum	445
HORNADAY. Classification of the collections in taxidermy	456
Hudson's Bay circular	137
PHILLIPS. Letters on the work of the National Museum	454
Publications of National Museum, list of	474
XXI.—NATURAL HISTORY. (See BIOLOGY.) XXII.—PALÆONTOLOGY. Conrad. Check list of eocene fossils	200 14 58
LEIDY. Cretaceous reptiles (s. c.)	192
LEIDY. Extinct sloths (s. c.)	72
LEIDY. Fossil ox (s. c.)	41
MEEK. Check list of cretaceous and jurassic fossils	177
MEEK. Check list of miocene fossils	183
MEEK and HAYDEN. Palsontology of Nebraska (s. c.)	172
XXIII.—PHILOLOGY. Bowen. Yoruba grammar and dictionary (s. c.)	96
GIBBS. Chinook jargon	161
GIBBS. Comparative Vocabulary	170
GIBBS. Ethnological and philological instructions.	160
LIEBER. Vocal sounds of Laura Bridgman (s. c.) MITCHELL and TURNER. Vocabulary of jargon	12 68

	No. 1-
Ca	No. in talogue.
Morgan. Systems of consanguinity (s. c.)	
Riogs. Dakota grammar and dictionary (s. c.)	
RCHRIG. Language of the Dakota or Sioux Indians	
WHITEEN. Lectures on Linguistics	352
XXIV.—PHYSICAL GEOGRAPHY.	
AAIV.—IHIBIOAD GEOGRAIHI.	
Coores. Physical geography of North America	351
ELLET. Physical geography of the Mississippi valley (s. c.)	13
HAYES. Observations in the Arctic seas (s. c.)	196
HENRY. Circular relative to heights	236
Mason. Luray Cavern in Virginia	483
XXV.—PHYSICS. (See also Terrestrial Physics.)	
BARKER. Recent progress in physics	429
BARNARD. Internal structure of the earth (s. c.)	310
BARNARD. Problems of rotary motion (s. c.)	240
Brezina. Crystallography and crystallophysics	. 386
COFFIE. Winds of the globe (s. c.)5	2, 268
DAVIS. Law of deposit of flood-tide (s. c.)	. 33
DELAUNAY. Essay on velocity of light	354
DRAPER. Telescope in photography (s. c.)	. 180
GOULD. Transatlantic longitude (S. C.)	. 223
GUYOT. Physical and meteorological tables	1, 153
HAYES. Physical observations, Arctic seas (s. c.)	
Henry. Electro-magnetic telegraph	. 115
Henry. Investigation of illuminating materials	. 389
Henry. Lightning rods	. 237
Henry. Researches on sound	406
HILGARD. Tides and tidal action	. 390
Holmsen. Color-blindness	
Hunt., Chemistry of the earth	376
KANE. Physical observations, Arctic seas (S. C.)	. 198
Merch. Heat and light of the sun (s. c.)	
Newton. Metric weights and measures	371
TAYLOR. Henry and the telegraph	405
TAYLOR. Kinetic theories of gravitation	395
TAYLOR. Nature and origin of force	375

PHYSICS—Continued. No Came	o. in
SCHOTT. Base chart of the United States	41
SECCHI. Electrical rheometry (s. c.)	3
WHITTLESEY. Fluctuations, level of lakes (s. c.)	_
WHITEHOUTE FINDING (6. 0.)	
	
XXVI.—PHYSIOLOGY. (See ANATOMY, etc.)	
XXVII.—TECHNOLOGY. (See CHEMISTRY, etc.)	
XXVIII.—TERRESTRIAL PHYSICS.	
BACHE. Discussion of magnetic observations (s. c.)I, 113; II, 121; III, IV_VI, 162; VII-IX, 175; X-XII, 186; I-XII,	
BACHE. Magnetic survey of Pennsylvania (8. C.)	
BARNARD. Problems of rotary motion, and precession of equinoxes (s. c.)	
COFFIN. Winds of the globe (8. C.)	
COFFIN. Winds of the northern hemisphere (s. c.)	5
· · · · · · · · · · · · · · · · · · ·	
DAVIS. Dynamical action, etc., of the flood tide (s. c.)	3
Force. Record of auroral phenomena (s. c.)	84
HAYES. Physical observations in Arctic seas (s. c.)	190
HENRY. Circular relative to altitudes	230
HILGARD. Tides and tidal action in harbors	390
	370
KANE. Magnetical observations in the Arctic seas (8. c.)	91
KANE. Physical observations in the Arctic seas.(s. c.)	196
KANE. Tidal observations in the Arctic seas (s. c.)	130
LOCKE. Observations on terrestrial magnetism in United States (s. c.)	35
MÜLLER. Observations on terrestrial magnetism in Mexico (s. c.)	114
OLMSTRAD. Secular period of the aurora (s. c.)	81
SONNTAG. Observations on terrestrial magnetism in Mexico (s. c.)	114
WHITTLESEY. Fluctuations of level in the North American lakes (s. c.)	119
XXIX.—ZOOLOGY.	
Birds.	
BAIRD. Arrangement of birds	A1^
	210
_	106
BAIRD. " 8vo	108

ZOOLOGY—Continued. No Case
BAIRD. Instructions for collecting bird's nests and eggs
BAIRD. Review of birds, part I
Brewer. North American cology, part I (s. c.)
Circular, birds of South America
Desiderata of birds of Mexico, Central America, etc
ELLIOT. List of Trochilids, (humming birds)
ELLIOT. Monograph of Trochilidse (s. c.)
KIDDER and COURS. Birds of Kerguelen Island
LAWRENCE. Birds of Mexico
LILLJEBORG. Outline of classification of birds
RIDGWAY. Catalogue of old world birds in Museum
RIDGWAY. Nomenclature of North American birds
Fishes.
BAIRD. Circular for shipping fresh fish
BAIRD. Circular relative to food fishes
BAIRD. Fishes of New Jersey coast
BAIRD. Inquiry relative to food-fishes
BEAN. Directions for collecting and preserving fish
GILL. Arrangement of fishes
GILL. Bibliography of fishes of Pacific coast
GILL. Catalogue of fishes
GILL. Fishes of Western North America
GIRARD. Monograph, cottoids (s. c.)
GOODE. Berlin exhibit of fisheries and fish culture
GOODE. Fishes of Bermuda
JORDAN and BRAYTON. Fishes of South Carolina, Georgia, and Tennessee
JORDAN. Notes on Rafinesque's memoirs, North American fishes
Insects.
AGASSIZ. Classification of insects (S. C.)
Circular to entomologists
Circular concerning department of insects
DE SAUSSURE. Monograph of wasps, hymenoptera
HAGEN. Synopsis of neuroptera
LE CONTE. Classification of coleoptera, part I
LE CONTE. " part II
LE CONTE. Kancas and New Mexico coleoptera (s. c.)

ZOOLOGY—Continued.	alogue.
LE CONTE. List of coleoptera	140
LE CONTE. New species of coleoptera, part I	167
LE CONTE. " part II	264
Lorw. Monograph of diptera, part I	141
LOEW. " part II	171
LOEW. " part III	256
MELSHEIMER. Catalogue of coleoptera	· 62
MORRIS. Catalogue of lepidoptera	118
MORRIS. Synopsis of lepidoptera	133
OSTEN SACKEN. Monograph of diptera, part IV	219
OSTEN SACKEN. Catalogue of diptera	
PACKARD. Directions for collecting insects	261
SOUDDER. Catalogue of orthoptera	189
·	
Mammals.	
aaa	
ALLEN. Monograph, North American bats	165
BAIRD. Catalogue, mammals	105
GILL. List of families of mammals	230
Gibl. Diet of families of mammais	200
Mollusks.	
BINNEY. Bibliography North American conchology, part I	142
3	
DIRREI.	
BINNEY. Land and fresh-water shells, part I. Pulmonata geophila	194
DIARE. I Unidonate Dimitophils	143
Dinnai.	144 252
CARPENTER. Mollusks of Western North America	251
CARPENTER. Lectures on mollusca.	152
Check list of shells	128
Circular for collecting shells	176
DALL. Index of names applied to brashiopoda	304
Exploring expedition, duplicate shells	193
GILL. Families of mollusks	227
LEWIS. Instructions for collecting land and fresh-water shells	363
	145
PRIME. Corbiculads	201
Throw I and and fresh mater shalls next IV Strongeride	201

-		
7	41	17
	u	

ZOOLOGY—Continued. Radiates.	No. in Catalogue.
BAIRD. Circular for collecting cray-fish	319
CLARK. Lucernariæ (s. c.)	242
RATHBUN. List of marine invertebrates distributed, series II	
RATHBUN. " " " " III	471
STIMPSON. Marine invertebrata of Grand Manan (s. c.)	50
Reptiles.	•
BAIRD and GIRARD. Catalogue of serpents	49
BAIRD. Circular on collecting reptiles	320
COPE. North American batrachia and reptilia	292
MITCHELL and MOREHOUSE. Chelonia (s. c.)	159
•	

Shells. (See Mollusks.)

SMITHSUNIAN CONTRIBUTIONS TO KNOWLEDGE. (Quarto.)

Vol.	DATE.	Pages.	Wood- curs.	PLATES.	Марв.	No. 11 Series.	CONTAINS NOS. OF CATALOGUE.
I	1848	860	207	48		2	1.
, II	1851	572	.89	24		26	8, 12, 20, 18, 14, 16, 17, 23, 15
III	1852	562		85		88	4, 5, 6, 7, 11. 85, 86, 80, 82, 22, 88, 37, 24, 29.
ΙV	1852	426				89	40.
v	1858	588	4	45		55	44, 41, 45, 43, 42.
VI	1854	484	9	53		56	46, 60, 61, 50, 52, 58, 54.
VII	1855	260	74	72	2	76	59, 63, 70, 72, 78.
VIII	1856	564	52	9		78	71, 81, 80, 82, 84, 85.
IX	1857	480	45	22		92	88, 90, 86, 88, 79, 94.
x	1858	462	1	16	1	99	95, 97, 98.
ΧI	1859	502	20	28		111	89, 100, 118, 104, 126, 114, 127.
XII	1860	58 8	15	8		112	129, 119, 103, 131, 185.
XIII	1868	558	80	4	8	151	180, 146, 155, 121, 182, 162, 163.
XIV	1865	490	158	25		184	159. 175, 186, 180, 172, 192.
X∇	1867	620	47	18	4	206	199, 197, 202, 196.
XVI	1870	498	76	18		211	173, 204, 120, 208, 221, 223, 220.
XVII	1871	616	6	14		229	218.
XVIII	1872	646	10	5	8,	246	222, 282, 289, 233.
XIX	1874	640	6	21		272	240, 241, 262.
ХX	1876	7 94	4	26		284	268.
XXI	1876	5 4 8	80	9	8	285	280, 281, 267, 277.
IIXX	1880	587	474	20		840	259, 269, 287, 381, 318.
IIIXX	1881	766	160	18		846	242, 248, 310, 817, 857.

SMITHSONIAN MISCELLANEOUS COLLECTIONS. (Octavo.)

Vol	DATE.	Pages.	Curs.	PLATES.	No. in Series.	CONTAINS NOS. OF CATALOGUE.
I	1862	788	28		122	148, 87, 158.
п	1862	714	88		128	27, 115, 58, 108, 49, 128, 84, 187, 189,
щ	1862	776	49		124	168, 176, 188. 102, 118, 186, 117.
ΙV	1862	762	80		125	184, 188.
7	1864	774			158	142, 74, 15 4 .
VI	1867	888	15	7	169	141, 171, 140, 167.
VII	1867	878	818		191	165, 148, 144, 201, 145, 200, 188, 177,
VIII	1869	921	780	4	212	156, 161, 160, 208. 219, 189, 194, 210, 187, 207, 205, 178,
IX	1869	914			218	168, 164. 174, 179.
x	1878	918	5		250	252, 227, 235, 236, 287, 190, 284, 288,
XI	1874	790	55	4	278	248, 245. 280, 247, 256, 261, 264, 265.
XII	1874	767	86		274	181, 255, 268.
XIII	1878	982	45		812	292, 298, 294, 295, 296, 297, 808, 804,
XIV	1878	911	 	4	814	305, 306. 254, 283, 288, 276, 289, 216, 301, 311.
XV	1878	880	58		815	258, 266, 291, 279, 282, 800, 802, 809,
XVI	1880	950	871	7	822	816, 819, 820. 258, 270, 321, 824, 825, 884, 885, 844.
XVII	1880	1084			886	828.
XVIII	1880	851			887	829.
XIX	1880	1084			416	882, 888.
XX	1881	846			428	
XXI	1881	778	42	18	424	880, 827, 866.
XXII	1882	1200	18	4	468	425, 467.
IXIII	1882	1008			475	463, 808, 818, 826, 842.

ANNUAL REPORTS OF THE SMITHSONIAN INSTITUTION. (Octore)

				
	Pages.	Woodcuts.	No. in Smithsonian Catalogue.	When Published
1st, for 1846	88		G	1847
2d, for 1847	208		H	1848
8d, for 1848	64		I	1849
4th, for 1849	272		21	1850
5th, for 1850	826		28	1851
6th, for 1851	10 4		51	1852
7th, for 1852	96		57	1853
8th, for 1858	810		67	1854
9th, for 1854	464	4	75	1855
10th, for 1855	440	79	77	1856
11th, for 1856	4 68	69	91	1857
12th, for 1857	438	100	107	1858
18th, for 1858	448	48	109	1859
14th, for 1859	450	57	110	1860
15th, for 1860	448	78	147	1861
16th, for 1861	464	25	149	1862
17th, for 1862	446	94	150	1868
18th, for 1868	420	56	187	1864
19th, for 1864	450	50	188	1865
20th, for 1865	496	189	209	1866
21st, for 1866	470	70	214	1867
22d, for 1867	506	10	215	1868
28d, for 1868	474	40	224	1869
24th, for 1869	480	88	. 228	1871
25th, for 1870	494	28	244	1871
26th, for 1871	478	8	249	1873
27th, for 1872	456	119	271	1873
28th, for 1878	452	88	275	1874
29th, for 1874	416	46	286	1875
80th, for 1875	422	85 4	298	1876
81st, for 1876	488	78	299	1877
82d, for 1877	500	49	828	1878
38d, for 1878	575	12	841	1879
84th, for 1879	- 681	216	845	1880
85th, for 1880	782	18	442	1881
	I	1	1	1

PUBLICATIONS OF THE U. S. NATIONAL MUSEUM. (Octavo.)

No. of Museum Series.		TITL	Е.		In Vol. of Miscellaneous Collections.	No. in Smithsonian Catalogue.
1	Bulletin of th	ne U. S. Nati	onal Muse	ım, No. 1	XIII	292
2	"	u	16	No. 2	XIII	298
8	46	".	"	No. 8	XIII	294
4	46	**	44	No. 4	XIII	295
5	**	44	44	No. 5	XIII	296
6	٠,,	"	"	No. 6	XIII	297
7	"	"	11	No. 7	XIII	808
8	"	"	"	No. 8	XIII	80 4
9	"	"	"	No. 9	XIII	805
10	4.6	"	**	No. 10	XIII	806
11	44	•"	**	N o. 11	XXIII	4 63
12	46	"	"	No. 12	IIIXX	808
18	11	. 44	44	No. 18	ххін	818
14	44	44	"	No. 14	XXIII	826
15	"	"	"	No. 15	XXIII	842
16	46	"	**	No. 16	<i>;,</i>	
17	Proceedings	Nat. Mus.,	Vol. I		XIX	882
18	Bulletins Na	t. Mus., Vol	. I, Nos. 1	to 10	XIII	812
19	Proceedings	Nat. Mus.,	V ol. II		XIX	883
20	Bulletin of th	ae U. S. Nati	ional Muse	m, No. 17		412
21	46	"	4.6	No. 18		4 18
22	"	"	46	No. 19		470
28	"	u	"	No. 20		
24	"	u	44	No. 21		422
25	Proceedings	Nat. Mus.,	Vol. III		XXII	425
26	Bulletin of tl	ne U.S. Nati	ional Muse	ım, No. 22		444
27	Proceedings	Nat. Mus.,	Vol. IV		XXII	4 67
28	Bulletins Na	t. Mus., Vol	. II, Nos.	11 to 15	XXIII	4 75

PUBLICATIONS OF THE U. S. NATIONAL MUSEUM—Continued. SEPARATE LIST OF BULLETINS.

No.	When Published.	Pages.	Cuts.	Plates.	Maps.	In Vol. of Miscellaneous Collections.	No. in Museum Series.	No. in Smithsonisa Catalogue.
1	1875	108				XIII	1	292
2	1875	61				XIII	2	293
8	1876	124				XIII	8	294
4	1875	56				XIII	4	295
5	1876	84				XIII	5	296
6	1876	140				XIII	6	297
7	1877	172				XIII	7	908
8	1877	88				XIII	8	804
9	1877	56				XIII	9	305
10	1877	124		45		XIII	10	306
11	1882	77				ıfıxx	11	463
12	1878	287				XXIII	12	308
18	1879	189				XXIII	18	813
14	, 1879	867				XXIII	14	826
15	1879	179				XXIII	15	842
16							16	
17	1880	55					20	412
18	·1880	279	1				21	418
19	1882	898					22	470
20							28	
21	1881	94					24	423
22	1881	265			1		26	444

THE BULLETINS COLLECTED IN VOLUMES.

Vol.	Conten	ts.	Published.	Pages.	Outs.	Plates.	In Vol. of Miscellaneous Collections.	No. in Museum Series.	No. in Smithsonian Catalogue.
I	Nos. 1	l –10	1878	982		45	XIII	18	812
II	Nos. 11	l -1 5	1882	1008			XXIII	28	475

PUBLICATIONS OF THE U.S. NATIONAL MUSEUM-Continued.

SEPARATE LIST OF PROCEEDINGS.

Vol.	For Year.	Pages.	Cuts.	Plates.	When Published.	In Vol. of Miscellaneous Collections.	No. in Museum Series.	No. in Smithsonian Catalogue.
I	1878	524	8	8	1879	XIX	17	882
11	1879	503	2	7	1880	XIX	19	388
III	1880	594	5	2	1881	XXII	25	425
1 V	1881	600	18	2	1862	XXII	27	467

PUBLICATIONS OF THE BUREAU OF ETHNOLOGY OF THE SMITHSONIAN INSTITUTION. (Imperial Octavo.)

Title.	Published.	Pages.	Cuts.	Plates.	Maps.	No. in Smithsonian Catalogue.
First Annual Report, for 1879-80	1881	688	348	54	1	476

PUBLICATIONS HAVING SEPARATE NUMBERS IN THE SMITH-SONIAN SERIES, BUT INCLUDED IN THE ANNUAL REPORTS OF THE INSTITUTION.

		Vol. 1 Report
A	Journal of Regents	1847
В	Report of Organization Committee	1847
D	Address at laying corner-stone. DALLAS	1847
E	Exposition of Bequest. HENRY	1847
F	First Report of Secretary. HENRY	1848
J	Programme of Organization. HENRY	1848
K	Correspondence, Squier and Davis	1849
L	First Report of Organization Committee	1547
M	Reports of Institution to 1849	1849
N	Officers and Regents	1547
25	Public Libraries. JEWETT	1850
152	Lectures on Mollusca. CARPENTER.	1850
251	Memoir of Von Martius. RAU	1869
307	Report on Centennial. BAIRD	1876
343	Annual Reports. HENRY	865-77
3 4 8	Fishes of New Jersey. BAIRD	1854
351	Forests of North America. Cooper	1836
352	Lectures on Linguistics. WHITNEY	186
354	Essay on Velocity of Light. DELAUNAY	1874
355	Ozone and antozone. WETHERILL	1564
360	Palafittes. DESOR.	1865
361	Aborigines of California. BAEGERT	1863-4
362	Artificial Shell Deposits in New Jersey. RAU	1864
364	Classification of Birds. LILLJEBORG	195
365	Tinneh Indians. Ross and others	1866
367	Flora of Alaska. Rothrock	1867
368	Indian Pottery. RAU	1866
369	Dorpat and Poulkova. ABBE	1867
37 0	Flint Implements in Illinois. RAU	1868
371	Metric Tables. Newton	1865
372	Drilling in Stone without Metal. RAU	
373	Meteorological Stations and Observers	1868
375	Origin and Nature of Force. TAYLOR	1870
376	Chemistry of the Earth. Hunt	1869
377	Diamond and Precious Stones. BABINET	1870
378	Dakota Language. RŒHRIG	1871
379	Eulogy on A. D. Bache. HENRY	1870

	PUBLICATIONS HAVING SEPARATE NUMBERS—Continued.	
	V R	ol. of eport.
380	Scientific Education of Mechanics. PRABODY	1872
382	North American Stone Implements. RAU	1872
385	Ancient Aboriginal Trade. RAU	1872
386	Crystallography. Brezina	1872
389	Investigation of Illuminants. HENRY	1880
390	Tides. HILGARD	1874
392	Prehistoric Antiquities of Hungary. ROMER	1876
393	Mound Builders and Ancient Man in Michigan. GILLMAN1873 and	1875
394	Stone Age in New Jersey. Abbott	1875
3 95	Kinetic Theories of Gravitation. TAYLOR	1876
396	History and Climate of New Mexico. McParlin	1877
397	Latimer Antiquities. MASON	1876
398	Meteorological Memoirs. ABBE	1877
399	Color-blindness. Holmgren	1877
400	Aboriginal Structures in Georgia. Jones	1877
401	Mexican Axolotl. Weismany	1877
4 02	Stock-in-trade of Aboriginal Lapidary. RAU	1877
403	Gold Ornament from Florida. RAU	1877
404	Polychrome Bead from Florida. 'HALDEMAN	1877
4 05	Henry and the Telegraph. TAYLOR	1878
406	Researches in Sound. HENRY	1878
407	Memoir of Joseph Henry. GRAY	1878
408	Report of Secretary for 1878. BAIRD.	1878
109	Report of Museum Building Commission and Architects	1879
4 10	Reports of Observatories. Holden	1879
\$ 11	Irritation of Polarized Nerve. LAUTENBACH	1878
1 15	Savage Weapons. KNIGHT	1879
118	Report of Secretary for 1879. BAIRD	1879
119	Report of Secretary for 1880. BAIRD	1880
120	Anthropological Investigations. Mason	1879
121	Index to Anthropological Articles. BEHMER	1879
1 26	Synopsis of Herschel's Writings. HOLDEN and HASTINGS	1880
127	Recent Progress in Astronomy. Holden	1880
128	" Geology and Mineralogy. HAWES	1880
129	" Physics and Chemistry. BARKER	1880
130	. " Botany. FARLOW	1880
l31	" Zoology. GILL	1880
132	" " Anthropology. MASON	1880
133	Report of Visit to Luray Cave. MASON	1880
134	Report of Museum Building Commission and Architects	1880

	PUBLICATIONS HAVING SEPARATE NUMBERS—Continued.
	Rep
435	Snell's Barometric Observations. Loud 18
436	List of Periodicals received by Smithsonian Institution 18
438	Reports of Observatories. Holden and BEHMER 18
439	Warming and Ventilating Buildings. MORIN1873
440	
770	Anthropological Articles. RAU1863-
	
PUB	LICATIONS NOT IN REGULAR SERIES OF "CONTRIBUTIONS
	"COLLECTIONS," OR "REPORTS.
•	•
C	Digest of Act of Congress.
P	Hints on Public Architecture. OWEN.
Q	Check-list of Periodicals.
18	Report on the Discovery of the Planet Neptune. Gould.
47	On Construction of Catalogues of Libraries. JEWETT.
48	Bibliography of American Natural History for 1851. GIRARD.
62	Catalogue of Described Coleoptera of the United States. MELSHEIMER.
64	List of Foreign Institutions.
65	Registry of Periodical Phenomena.
66 68	The Annular Eclipse of May 26, 1854. Vocabulary of the Jargon. MITCHELL and TURNER.
69	List of Domestic Institutions.
93	Meteorological Observations for 1855.
101	Map of the Solar Eclipse.
105	Catalogue of North American Mammals. BAIRD.
106	Catalogue of North American Birds. BAIRD.
116	List of Public Libraries. RHEES.
157	Results of Meteorological Observations, 1854-1859. Vol. I.
170	Comparative Vocabulary.
182	Results of Meteorological Observations, 1854-1859. Vol. II.
185	List of Birds of Mexico, &c. BAIRD.
193	List of Duplicate Shells collected by Wilkes.
217	Letter of M. Hoek on Meteoric Shower.
225	List of Foreign Correspondents.
226	List of Smithsonian Publications. Regulations of the Smithsonian Institution.
260 278	Check-list of Publications of Smithsonian Institution.
290	Circular for distribution at Centennial.
290 34 7	Nomenclature of Clouds.
350	Map of Stars near the North Pole.
3 59	Planisphere of the Visible Heavens.
366	Directions for collecting Diatomacea, &c. EDWARDS.
414	Base Chart.
437	Check-list of Smithsonian Publications.
4 66	Directory of Officers and Employés.

LIST OF PAPERS PUBLISHED IN THE "SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE."

	_ 		i
Hanial. Number.	AUTHOR.	TITLE OF WORK.	Commissions of Reference.
1	SQUIER, E. G., AND DAVIS, E. H.	Ancient Monuments of Mississippi Valley.	E. Robinson, D. D., J. R. Bart- lett, Prof. W. W. Turner, S. G. Morton, M. D., G. P. Marsh.
3	WALKER, S. C	Researches, Planet Neptune.	Morwit, M. D., G. F. Marsii.
4	WALKER, S. C	Ephemeris of Neptune for 1848.	
5	Walker, 8. C	Ephemeris of Neptune for 1849.	•
6	WALKER, S. C	Ephemeris of Neptune for 1850.	
7	WALKER, S. C	Ephemeris of Neptune for 1851.	
8	Downes, John	Occultations in 1848.	
9	Downes, John	Occultations in 1849.	
10	Downes, John	Occultations in 1830.	
\mathbf{n}	Downes, John	Occultations in 1851.	·
12	LIEBER, FRANCIS	Vocal Sounds of L. Bridgman	Col. W. W. S. Bliss, Miss D. L. Dix.
13	Ellet, Charles	Physical Geography of U. S	Maj. A. Mordecai, Capt. F. A. Smith, Lieut. M. L. Smith.
14	GIBBES, R. W	Memoir on Mosasaurus	Prof. Louis Agassiz, Prof. Henry D. Rogers.
15	SQUIER, E. G	Aboriginal Monuments of New York	Brantz Mayer, Wm. W. Turner.
16	Agassiz, Louis,	Classification of Insects	Prof. C. D. Meigs, Thaddeus W. Harris, M. D.
17	HARE, ROBERT	Explosiveness of Nitre	John Torrey, M. D., Col. J. J. Abert.
20	BAILEY, J. W	Microscopic Examination of Soundings	Prof. Lewis R. Gibbes, Prof. Wm. B. Rogers.
22	Gray, Apa	Plantæ Wrightianæ. Part I	Prof. John Torrey, John Carey.
z;	Bailey, J. W	Microscopic Observations in South Carolina, Georgia, and Florida.	Prof. Wm. B. Rogers, Prof. Lewis R. Gibbes.
2 4 ;	WALKER, S. C	Ephemeris of Neptune, 1852. Appendix I.	
29	Downes, John	Occultations in 1852.	•
:.0	GIRARD, CHARLES	Fresh-Water Fishes of North America	Dr. Jared P. Kirtland, Dr. J. E. Holbrook.
32	HARVEY, WM. H	Marine Algæ of North America. Part I	Prof. J. W. Bailey, Dr. Asa Gray.
33	DAVIS, CHAS. H	Law of Deposit of Flood Tide	Prof. L. Agassiz, Prof. A. Guyot.
35	LOCKE, JOHN	Observations on Terrestrial Magnetism	Prof. Elias Loomis, Prof. Julius E. Hilgard.
26	Secchi, A	Researches on Electrical Rheometry	J. H. Lane, Prof. Jas. Curley.
37	WHITTLESEY, CH	Ancient Works in Ohio	Brantz Mayer, E. G. Squier.
40	Riggs, S. R	Dakota Grammar and Dictionary	Minnesota Historical Society, Prof. W. W. Turner, Prof. C. C. Felton.
41 '	LEIDY, JOSEPH	Extinct American Ox	Paul B. Goddard, M. D., Joseph Pancoast, M. D.
42	GRAY, ASA	Plantæ Wrightianæ. Part II	Prof. John Torrey, John Carey.
43	HABVEY, WM. H	Marine Algae of North America. Part II	Prof. J. W. Bailey, Dr. Asa Gray.
41	LEIDY, JOSEPH	Flora and Fauna within Living Animals	Prof. J. W. Bailey, Charles F. Beck, M. D. J. B. S. Jackson, M. D., Joseph
45	WYMAN, JEFFRIES	Anatomy of Rana Pipiens	Leidy, M. D.
46	TORREY, JOHN	Plantæ Fremontianæ	Dr. W. Darlington, Dr. Asa Gray-

PAPERS PUBLISHED IN SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE-Continued.

- #			
SERIAL NUMBER.	Author.	TITLE TO WORK.	COMMISSIONS OF REPERENCE.
5 0	Stimpson, Wm	Marine Invertebrata of Grand Manan	A. A. Gould, M. D., Prof. J. b
52	Coppin, Jas. H	Winds of the Northern Hemisphere	Dana. Prof. W. B. Rogers, Prof E
54	Downes, John	Occultations in 1853.	Loomis.
5 8	LEIDY, JOSEPH	Ancient Fauna of Nebraska	Prof. J. Hall, J. L. Leconte, M.
59	Chappelsmith, J	Tornado in Indiana	Prof. J. H. Coffin, Prof. A. (3-
60	Torrey, John	Batis Maritima	well. Prof. A. Gray, John Carey.
61	Torrey, John	Darlingtonia Californica	Prof. A. Gray, John Carey.
63	Bailey, J. W	New Species of Microscopic Organisms	Prof. C. R. Gilman, M. D., W I. Burnett, M. D.
70	Lарнам, I. A	Antiquities of Wisconsin	American Antiquarian Soc.
71	HAVEN, S. F	Archæology of the United States	Peter Force, Brants Mayer.
72	LEIDY, JOSEPH	Extinct Sloth Tribe of North America	Isaac Hays, M. D., Prof. W. L. Horner.
73		Publications of Societies in Smithsonian Library.	norner.
79	Runkle, John D	Tables for Planetary Motion	Prof. Benj. Peirce, Com. Char E Davis.
80	ALVORD, BENJ	Tangencies of Circles and Spheres	Prof. A. E. Church, Prof. L. F. Gibbes.
81	Olmsted, D	Secular Period of Aurora Borealis	Prof. J. B. Cherriman, Prof. J.F.
82	Jones, Joseph	Investigation on Amer. Vertebrata	S. Jackson, M. D., J. Leidy, M.P. J. Wyman, M. D. Prof. B. Peirce, Dr. B. A. Gould
.8 3	Мевсн, L. W	Relative Intensity of Heat and Light of the Sun.	Prof. B. Peirce, Dr. B. A. Goult Jr.
·84	FORCE, PETER	Auroral Phenomena in North Latitudes.	
85	·····	Publications of Societies in Smithsonian Library. Part II.	
86	MAYER, BRANTZ	Mexican History and Archæology	S. F. Haven, E. H. Davis, M. P.
88	GIBBS, W., AND GENTH, F. A.	Ammonia Cobalt Bases	Prof. John F. Frazer, Prof. Jel Torrey.
89	BREWER, TH. M	North American Oology. Part I	John Gould, John Cassin.
90	Нітенсоск, Е	Illustrations of Surface Geology.	
94	Runkle, John D	Asteroid Supplement to New Tables for $b = \frac{(i)}{s}$.	Prof. Benj. Peirce, Com. Cl.as. H. Davis.
95	HARVEY, WM. H	Marine Algæ of North America. Part III.	Dr. John Torrey, Dr. Asa Gray
96	HARVEY, WM. H	Marine Algæ of North America. 3 parts complete.	Prof. J. W. Bailey, Dr. Asa Gray
97	KANE, E. K	Magnetic Observations in the Arctic Seas	
98	Bowen, T. J	Yoruba Grammar and Dictionary	Prof. J. W. Gibbs, Prof. W. P. Whitney R. Anderson.
100	Gilliss, J. M	Eclipse of the Sun, Sept. 7, 1858	Prof. J. W. Gibbs, Flox. Whitney, R. Anderson. Prof. J. S. Hubbard, J. H. Lane.
103	Caswell, A	Meteorological Observations, Providence, R. I.	
104	KANE, E. K	Meteorological Observations in Arctic Seas.	Prof. A. D. Bache.
113	Васне, А. Д	Magnetic and Meteorological Observa- tions at Girard College. Part I.	Prof. Benjamin Peirce, Prof. W. Chauvenet.

PAPERS PUBLISHED IN SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE-Continued.

Stutat.	AUTHOR.	Title of Work.	Commissions of Reference.
114	SONNTAG, A	Terrestrial Magnetism in Mexico	Prof. Stephen Alexander, Prof. Arnold Guyot.
119	WHITTLESEY, CH	Fluctuations of Level in N. A. Lakes	Capt. A. A. Humphreys, Capt. A. W. Whipple.
1.50	HILDRETH, S. P., AND WOOD, J.	Meteorological Observations, Marietta, Ohio.	w. wnippie.
121	Васпе, А. D	Magnetic and Meteorological Observa- tions at Girard College. Part II.	
126	Le Conte, John L.	Coleoptera of Kansas and New Mexico	C. Zimmerman, F.E. Melsheimer.
127	LOOMIS, E	Storms in Europe and America, Dec. 1836.	Dr. S. P. Hildreth, Prof. A. Caswell.
129	KANE, E. K	Astronomical Observations in the Arctic Seas.	Prof. A. D. Bache.
13)	KANE, E. K	Tidal Observations in the Arctic Seas	Prof. A. D. Bache.
131	SRITE, N. D	Meteorological Observations in Arkansas from 1840 to 1850.	•
132	Bacne, A. D	Magnetic and Meteorological Observa- tions at Girard College. Part III.	
135	MITCHELL, S. W	Venom of the Rattlesnake	Franklin Bache, M. D., Robley Dunglison, M. D.
146	M'CLINTOCK, SIR F. L.	Meteorological Observations in the Arctic Seas.	Dunguison, M. D.
155	WHITTLESEY, CH	Ancient Mining on Lake Superior	D. Wilson, LL. D., E. H. Davis,
159	Mitchell, S. W., & Morehouse, G. R.	Anatomy and Physiology of Respira- tion in Chelonia.	M. D. Prof. J. Wyman, Prof. J. Leidy.
162	Васив, А. Д	Magnetic and Meteorological Observa- tions at Girard Coll. Pt. IV, V, & VI.	
106	BACHE, A. D	Magnetic Survey of Pennsylvania.	
172	Meek, F. B., and Hayden, F. V.	Palmontology of the Upper Missouri. Part I.	Isaac Lea, Prof. Jas. D. Dana.
173	DEAN, JOHN	Gray Substance of the Medulla Oblongata.	Dr. W. A. Hammond, Prof. Jeff- ries Wyman.
175	Васив, А. Д	Mag. and Met. Observ. at Girard College. Parts VII, VIII, and IX.	
180	Draper, H	Construction of a Silvered Glass Telescope.	Prof. Wolcott Gibbs, Com. J. M. Gilliss.
186	Васпи, А. D	Mag. and Met. Observ. at Girard College. Parts X, XI, and XII.	
192	LEIDY, JOSEPH	Cretaceous Reptiles of the U.S	Prof. L. Agassiz, Prof. E. D. Cope.
195	Васив, А. D	Girard College Observations. Complete. Parts I to XII.	Prof. Benj. Peirce, Prof. W. Chauvenet.
146	HATES, I. I	Physical Observations in the Arctic Seas.	
197	WHITTLESEY, CH	Glacial Drift of Northwestern States	Prof. L. Agassiz, Prof. J. P. Les-
198	KANE, E. K	Physical Observations in the Arctic Seas. Complete.	ley. Prof. A. I). Bache.
190	Newcomb, S	Orbit of Neptune	Adm'l C. H. Davis, Prof. Stephen
20:2	PUMPELLY, R	Geological Researches in China, Mongolia, &c.	Alexander. National Academy of Sciences.
3H '	CLEAVELAND, P	Meteorological Observations, Brunswick, Me., 1807-1859.	

PAPERS PUBLISHED IN SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE-Continued

		•		
SEELAL NUMBER.	Author.	Title of Work.	COMMISSIONS OF REPERENCE.	
208	PICKERING, CHAS	Gliddon Mummy Case in Smithsonian Institution.		
218	Morgan, L. H	Systems of Consanguinity and Affinity	Prof. J. A. Hadley, J. H. Tran- bull, Prof. W. D. Whitney.	
220	Swan, Jas. G	Indians of Cape Flattery	George Gibbs, Jeffries Wymar	
221	COFFIN, JAMES H	Orbit, &c., of Meteoric Fire Ball, July 20, 1860.	Prof. H. A. Newton, C. A. Schot	
222	SCHOTT, CHAS. A	Tables of Rain and Snow in United States.		
223	GOULD, B. A	Transatlantic Longitude.		
232	STOCKWELL, J. N	Secular Variations of Orbits of Planets	Prof. J. H. C. Coffin, Prof. S. Newcomb.	
233	FERREL, WM	Converging Series, Ratio of Diameter, and Circumference of Circles.	Nowcomb.	
239	Hareness, Wm	Magnetic Observations on the Monad- nock.	Pres. Acad. Sciences. Prof. J. H. C. Coffin, Prof. F. Rogers	
240	BARNARD, J. G	Problems of Rotary Motion.		
241	W00D, H. C	Fresh-Water Algos of North America	Dr. J. Torrey, Dr. F. A. P. Far-	
242	Clark, H. J	Lucernariæ.	27002 04-	
248	Hilgard, E. W	Geology of Lower Louisiana.		
259	Jones, Jos	Antiquities of Tennessee	Dr. G. A. Otis, Prof. O. T. Mac :	
262	NEWCOMB, S	Orbit of Uranus	Prof. J. H. C. Coffin, Prof. A-a- Hall.	
267	Swan, J. G	Haidah Indians	J. C. Welling, LL. D., Dr. 6-4 Otis.	
268	Coffin, J. H	Winds of the Globe.		
269	Habel, Simeon	Sculptures of Santa Lucia Cosumal- whuapa in Guatemala.	Prof. W. D. Whitney, J. H. Trumbull.	
277	Вснотт, С. А	Temperature Tables.		
280	ALEXANDER, S	Harmonies of Solar System.		
281	Newcomb, S	Planetary Motion	Prof. H. A. Newton, G. W. Hill.	
287	Rau, Charles	Archæological Collection, Nat. Museum.		
310	Barnard, J. G	Internal Structure of the Earth.		
317	ELLIOT, D. G	Classification and Synopsis of Trochilidæ.		
318	DALL, WM. H	Remains of Man from Caves in Aleutian Islands.		
331	RAU, CHARLES	The Palenque Tablet	S. F. Haven, H. H. Bancroft.	
353	Зснотт, С. А	Tables of Rain Fall, (2d edition.)		
357	W00D, H. C	Researches on Fever	S. Weir Mitchell, M. D. 1 Woodward, M. D.	
383	Bransford, J. F	Archæological Researches in Nicaragua	Garrick Mallery, H. C. Yarr 4.	
443	Caswell, A	Meteorological Observations, Providence, R. I., 1831 to 1876.		

ALPHABETICAL INDEX OF ARTICLES

IN THE

SMITHSONIAN CONTRIBUTIONS TO KNOWLEDGE, Vols. 1-23. 1848-1881.

SMITHSONIAN MISCELLANEOUS COLLECTIONS, Vols. 1-23. 1862-1882.

> SMITHSONIAN ANNUAL REPORTS, Vols. 1-35. 1846-1880.

BULLETINS OF THE U. S. NATIONAL MUSEUM,

Nos. 1-22, (except 16 and 20,) 1875-1881,

PROCEEDINGS OF THE U. S. NATIONAL MUSEUM, Vols. 1-4. 1878-1881.

AND IN THE

FIRST ANNUAL REPORT OF THE BUREAU OF ETHNOLOGY OF THE . SMITHSONIAN INSTITUTION,

Vol. 1. 1879-1880.

• • . • .

ALPHABETICAL INDEX.

- NOTE.—The heavy-faced figures indicate the specific number in the Catalogue of a separate publication; the ordinary, or light-faced figures indicating the number of a publication of which the paper is only a constituent part.
- "R." represents the annual Smithsonian Report, immediately followed by the year of the report, and the number given to it in the Catalogue.
- "P." represents the Proceedings of the National Museum, followed by the year of publication and the number given to the volume in the Catalogue.
- "E." represents Report of the Bureau of Ethnology.

A

ABBE, C. Description of Observatories at Dorpat and Poulkova. (R, 1867) 215, 36
ABBE, C. Meteorological memoirs, translated by. (R. 1877) 828, 39
Abbott, C. C. Stone age in New Jersey. (R. 1875) 298, 39
Abbreviations of names of States, etc., for labelling collections 16
Abbreviations used in England in 1867—W. DE LA RUE. (R. 1867) 21
Abeons aurora—Jordan and Gilbert. (P. 1880)
ABICH, S. Remarkable forms of hail-stones. (R. 1869) 22
Aboriginal indians of America, language of—G. GIBBS. (R. 1870) 24
Aboriginal inhabitants of Cal.—J. BAEGERT. (R. 1868; R. 1864)_ 187, 188, 440, 36
Aboriginal lapidary, Stock-in-trade of-C. RAU. (R. 1877) 823, 440, 40:
Aboriginal monuments of State of New York-E. G. Squier. 1
Aboriginal remains of Tennessee—J. Jones 25
Aboriginal ruins at Savannah, Tenn.—J. P. STELLE. (R. 1870) 24
Aboriginal ruins in Hardin Co., TennJ. P. STELLE. (R. 1870) 24
Aboriginal shell-mounds, New Brunswick & New Eng.—S. F. BAIRD. (P. 1881). 46
Aboriginal structures in Georgia-C. C. Jones, Jr. (R. 1877) 828, 400
Abstracts of anthropological correspondence—O. T. Mason. (R. 1879; R. 1880)
Academy, Pontifical, Rome—See Prize questions.
Academy of Sciences—See National.
Academy of Sciences, Bordeaux—See Prize questions.
Academy of Sciences, Chicago, acknowledging specimens. (R. 1867) 216
Academy of Sciences, St. Petersburg, exchange system. (R. 1867) 210
Academy of Sciences, Vienna—See Prize questions.

Acalephæ, anatomy and physiology of—H. J. CLARK	242
Accidental or subjective colors—Abbe Moigno. (R. 1866)	214
Account of—	
aboriginal ruins at Savannah, Tenn.—J. P. Stelle. (R. 1870)	244
aboriginal ruins in Hardin Co., Tenn.—J. P. STELLE. (R. 1870)	
aborigines of California-J. BARGERT. (R. 1868; R. 1864). 187, 188, 440), 361
ancient mounds, Georgia-M. F. STEPHENSON. (R. 1870)	. 244
antiquities, Tennessee—E. O. DUNNING. (R. 1870)	244
antiquities, Vera Cruz, Mexico—H. Finck. (R. 1870)	. 244
astronomical observatory at Cordoba—B. A. Gould. (R. 1878)	275
balloon ascensions—J. GLAISHER. (R. 1868)	. 187
barometer, new-J. Green. (R. 1855)	- 77
burial of Indian squaw, California-W. M. King. (R. 1874)	286
Carib or Karif language and people. (R. 1878)	. 275
discovery of stone image in Tennessee-E. M. GRANT. (R. 1870)	. 244
Draper's telescope—T. W. WEBB. (R. 1864)	- I**
eclipse of sun in Peru, 1858-J. M. GILLISS	_ 100
hail storm in Texas—G. M. BACHE. (R. 1870)	
hail storm on Bosphorus—Com. PORTER. (R. 1870)	244
Indians of British America—E. Petitor. (R. 1865)	_ 209
lightning discharges. (R. 1867)	_ 215
Lindheimer's botanical explorations—A. GRAY. (R. 1849)	_ 21
marble used in extension of U. S. Capitol—J. HENRY. (R. 1856)	_ 91
"occultator" for indicating moon's position—T. HILL	_ 101
old Indian village, Kushkushkee, Pa.—E. M. McConnell. (R. 1871)	_ 249
Portland Society of Natural History—E. C. Bolles. (R. 1867)	
Priestley's lens—J. HENRY. (R. 1859)	
remarkable accumulation of bats—M. Figanierre. (R. 1863)	
sarcophagus in National Museum—A. A. HARWOOD. (R. 1870)	
storm in Butler Co., Kans., June 28, 1871—W. HARRISON. (R. 1871)	
tornado in Spruce Creek Valley, Pa.—J. B. MEEK. (R. 1871)	
tornado near New Harmony, Indiana—J. CHAPPELSMITH	
travels in Central and South America—S. HABEL	
Acid, sulphurous, action of, upon putrefactive bacteria. (P. 1881)	
Acknowledgment of birds—Academy of Sciences, Stockholm. (R. 1867)	
Acknowledgment of birds—J. Gould. (R. 1867)	
Acknowledgment of contributions to National Museum. (P. 1881) 46	
Acknowledgment of ferns—G. METTENIUS. (R. 1862)	
Acknowledgment of perennibranchiates—J. G. FISHER. (R. 1863)	_
Acknowledgment of shells, &c.—W. H. PEASE. (R. 1861) Acknowledgment of specimens—British Museum. (R. 1865)	
Acknowledgment of specimens—British Museum. (R. 1865)	
Acknowledgment of specimens—Unlorgo Academy of Sciences. (K. 1867)	_ 215

Acknowledgment of specimens—Regents of Univ. of State of N. Y. (R. 1865). 209
Acmæidæ, from Panama, new species of—P. P. CARPENTER 252
Acoustics applied to public buildings—J. HENRY. (R. 1856)
Acoustics, different kinds of refraction—W. B. TAYLOR. (R. 1875) 298
Acoustics, lectures on architecture in relation to—D. B. REID. (R. 1856) 91
Acrothele, note on—C. A. WHITE. (P. 1880) 425
Acts of Congress—See Congress.
ADAMS, C. B. Catalogue of shells of Panama, review of—P. P. CARPENTER. 252
ADAMS, C. B. Shells collected by, description of 252
ADAMS, W. Subcutaneous surgery. Toner lecture No. VI
ADAMS, W. H. Mounds in Spoon River Valley, Illinois. (R. 1879) 345
Additions to Museum by Berlin Fisheries Exhibition. (R. 1880) 442
Address at laying of corner-stone of Smithsonian building-G. M. Dallas. D, 329
Address on the Smithsonian Institution-J. HENRY. (R. 1858) B 67
Address, Pres. Royal Society of Victoria-R. L. J. Ellery. (R. 1868) 224
Addresses at memorial services of J. Henry-See Henry.
Addresses, list of, foreign institutions, since 1862. (R. 1865) 209
Administration of drugs and medicines-J. M. FLINT. (P. 1881) 467, 451
Aerial navigation—J. HENRY. (R. 1860) 147
Aerial navigation, currents of the atmosphere and-J. HENRY. (R. 1860) 147
Aeronautic voyage across the Atlantic, Memorial relative to Lowe's. (R. 1860). 147
Aeronautic voyage, reply to memorial on Lowe's-J. HENRY. (R. 1860) 147
Aeronautic voyages—F. Arago. (R. 1863) 187
Aeronautic voyages-J. Glaisher. (R. 1863)
Aeronautics, modes of flight in relation to-J. B. Pettigrew. (R. 1867) 215
Affinity, systems of—L. H. Morgan 218
Africa, stone celts in—G. J. GIBBS. (R. 1877)
Africa, Yoruba, account of, and grammar and dictionary 98
AGASSIZ, L., biographical notice of-J. A. GARFIELD. (R. 1878) 275, 829
AGASSIZ, L., biographical notice of-P. PARKER. (R. 1878)
Agassiz, L., biography of—E. FAVRE. (R. 1878)
Agassiz, L., memoir of-R. P. Stebbins. (R. 1873)
Agassiz, L., supplemental list of generic names employed by-S. H. Scudder. 470
Agassiz, L., papers by—
Classification of insects from embryological data
Narrative of the Hassler expedition. (R. 1872) 271, 829
On the formation of a museum. (R. 1849)
Report on use of new hall in Smithsonian building. (R. 1867) 215, 329
Scientific instructions to Captain Hall. (R. 1871) 249
Age of stone—R. S. Robertson. (R. 1874)
Acnew, S. A. Mounds in Mississippi. (R. 1867)
Agonidæ, new species of, from California—W. N. Lockington. (P. 1880) 425

Agonus vulsus—Jordan and Gilbert. (P. 1880)
Agricultural Association of Milan, on exchanges. (R. 1863)
Agricultural chemistry, lectures on-S. W. Johnson. (R. 1859)
Agricultural flint implements in Illinois—C. RAU. (R. 1868) 224, 440, 370
Agricultural flint implements, Southern Illinois—C. RAU. (R. 1868) 224, 370
Agricultural implements of N. American stone period:—C. RAU. (R. 1863) 187. 44
AINSA, SANTIAGO. Tucson meteorite. (R. 1863)
Air, temperature in ascending currents of—J. HANN. (R. 1877) 323.9
Air, temperature in ascending moist currents of—L. Sohneke. (R. 1877) 323.
Air-currents, nature of—A. COLDING. (R. 1877) 323,3
AIRY, G. B. Correcting measures of the sun's distance. (R. 1859)
Alabama, mica beds in—W. GESNER. (R. 1879)
Alabama, shell-heaps—A. S. GAINES and K. M. CUNNINGHAM. (R. 1877)
Alaska, bibliography of fishes of, for 1880—T. H. BEAN. (P. 1881)
Alaska, climate of—H. M. BANNISTER. (R. 1866)
Alaska, description of Lycodes Turneri from-T. H. BEAN. (P. 1878)
Alaska, descriptions of new fishes from—T. H. BEAN. (P. 1881)
Alaska, flora of—J. T. ROTHROCK. (R. 1867)
Alaska, limpets and chitons of, report on-W. H. Dall. (P. 1878)
Alaska, new fish from-T. H. BEAN. (P. 1879)
Alaska, new fish from—W. N. Lockington. (P. 1880)
Alaska, new forms of mollusks from-W. H. Dall. (P. 1878)
Alaska, notes on Nichols' collection of fishes of—T. H. BEAN. (P. 1881)
Alaska, occurrence of Hippoglossus vulgaris at—T. H. BEAN. (P. 1879) "
Alaska, occurrence of Stichæus punctatus at—T. H. BEAN. (P. 1878)
Alaska, preliminary catalogue of fishes of—T. H. BEAN. (P. 1881)
Alaska, remains of prehistoric man from caves in—W. H. Dall
Alaska, some genera and species of fishes from—T. H. BEAN. (P. 1879) **
Albany, Illinois, mound skull and bones—R. J. FARQUHARSON. (R. 1874)
Albany Institute, memorial proceedings relative to Joseph Henry
Album from Greece, presented by Miss E. B. Contaxaki. (R. 1857) 107.
Alepocephalus Bairdii, from deep sea Atlantic—G. B. Goode; T. H. Beas.
A leutian islands, remains of predistoric man from—w. H. Dall.
ALEXANDER, B. S. Report of architect. (R. 1854)
ALEXANDER, C. A. Origin and history of Royal Society of London. (R. 1863)
ALEXANDER () A (translations by)
Accidental or subjective colors—ABBE Moiono. (R. 1866)
American migration—R. VON HELLWALD. (R. 1866)
Biography of Peltier by his Son. (R. 1867)
Carlsruhe Scientific Congress—M. J. NICKLES. (R. 1860)
Catalytic force—T. L. Phipson. (R. 1862)

-	\sim	
1	•,	•
	- 24	- 1

OF SMITHSONIAN PUBLICATIONS.

Electr	cal currents of the earth—C. MATTEUCCI. (R. 1867)
Electr	eity. (B. 1867)
Electr	-physiology—C. MATTEUCCI. (R. 1865)
Extern	al appearance of sun's disk. (R. 1866)
Figure 18	s of equilibrium of a liquid mass—J. PLATEAU. (R. 1865; R
Forma	tion of ice at bottom of water—M. Englehabdt. (R. 1866)
Histor	of Academy of Sciences of Paris-M. FLOURENS. (R. 1862)
Histor	of Royal Institution of Great Britain—E. MAILLY. (R. 1867)
Horar	variations of barometer-M. VAILLANT. (R. 1866)
Intern	ational Archæological Congress. (R. 1866)
Jussie	s, the, and the natural method—M. FLOURENS. (R. 1867)
	Faraday—De LA Rive. (R. 1867)
	Kepler-M. Berthrand. (R. 1869)
Man tl	e contemporary of the mammoth. (R. 1867)
	r of A. Bravais-E. DE BEAUMONT. (R. 1869)
_	r of D. de Blainville-M. FLOURENS. (R. 1865)
Memo	r of von Buch-M. FLOURENS: (R. 1862)
	r of A. de Candolle-M. FLOURENS. (R. 1859)
	of Delambre—J. Fourier. (R. 1864)
	r of R. J. Häuy-M. CUVIER. (R. 1860)
	r of Legendre—E. DE BEAUMONT. (R. 1867)
	of L. J. Thenard—M. FLOURENS. (R. 1862)
	of Magendie-M. Flourens. (R. 1866)
	of Priestley-M. Cuvier. (R. 1858)
	r of Geoffroy St. Hilaire-M. Flourens. (R. 1861)
	eope. (R. 1860)
	ent of the stars—M. MAEDLER. (R. 1859)
	l History of organized beings-M. MARKY. (R. 1867)
	es or lacustrian constructions—E. DESOR. (R. 1865)
	chemistry—M. Jamin. (R. 1867)
	ork antiquities of Olmutz. (R. 1866)
_	ration of wood. (R. 1864)
	c Congress of Carlsruhe—M. J. Nickles. (R. 1860)
	stion of the stars—C. D. Kämtz. (R. 1861)
	f feeling: sense of smell. (R. 1865)
	f hearing. (R. 1866)
	f sight. (R. 1866)
	f taste. (R. 1866)
	planets between Mars and Jupiter—LESPIAULT. (R. 1861)

ALEXANDER, C. A., (translations by)—Continued.
State of ethnology in relation to form of human skull—A. Pelzius. (R. 1859)
Transactions of Geneva Society of Physics, &c. (R. 1859; R. 1864; R. 1865; R. 1866)
ALEXANDER, S. Harmonies of the solar system 280
ALEXANDER, S. Lecture on relations of time and space. (R. 1861)
ALEXANDER, S. Lecture on vastness of visible creation. (R. 1857)
Alge-J. W. BAILEY
Algæ, Arctic, list of-W. H. HARVEY
Algæ, Arctic America—W. G. FABLOW
Alga, bibliography of-H. C. Wood.
Algæ, Kerguelen Island-W. G. FARLOW
Algæ, lecture on marine-W. H. HARVEY. (R. 1855)
Algæ, marine. Part 1, Melanospermeæ-W. H. HARVEY 32
Algæ, marine. Part 2, Rhodospermeæ—W. H. HARVEY
Algse, marine. Part 8, Chlorospermes-W. H. HARVEY 95
Algæ, marine. Parts 1, 2, 8, complete—W. H. HARVEY 96
Algæ, North American fresh water, history of-H. C. Wood 241
Algæ, Rhode Island, list of—S. T. OLNEY
ALLEN, H. Monograph of bats of North America
ALLEN, Z. Explosibility of coal oils. (R. 1861) 146
Alphabetical index of genera of birds—S. F. BAIRD
Alphabetical index to Smithsonian publications
Alternate generation and parthenogenesis-G. A. KORNHUBER. (R. 1871) 24
Altitude, diminution of vapor with increasing-J. HANN. (R. 1877) 323. 300
Altitudes, circular relative to—J. HENRY
ALVORD, B. Tangencies of circles and of spheres 80
Amazilia cerviniventris-R. RIDGWAY. (P. 1881)
Amazilia yucatanensis—R. Ridgway. (P. 1881)
Amblystoms, change of axolotl to—A. WEISMANN. (R. 1877) 323, 401
Amblystoma, luridum-P. R. Hoy. (R. 1854)
America-See Arctic, British, Central, Middle, North, Russian, South.
America, ancient history of—M. MUCH. (R. 1871)
America, collection of charts and maps of-J. G. Kohl. (R. 1856)
America, grasshoppers and locusts of—A. S. TAYLOR. (R. 1858)
America, language of aboriginal Indians of-G. GIBBS. (R. 1870) 24
America, storms in, December, 1886—E. Loomis127
American Academy of Arts and Sciences-
On Smithsonian exchange system. (R. 1867) 215
Report on organization of Smithsonian. (R. 1858)
Thanks for exchanges. (R. 1855)

OF SMITHSONIAN PUBLICATIONS.

American Antiquarian Society. Antiquities of Wisconsin	70
American archæology circular	
American Association for Advancement of Science—	
Combined meteorological observations proposed. (R. 1851)	51
Memorial address on Joseph Henry. A. M. MAYER	417
American authors, North American conchology-W. G. BINNEY	
American Board of Commissioners for Foreign Missions. Grammar and	
dictionary of Dakota language	40
American corbiculads, monograph of—T. PRIME	145
American correspondents, list of 69,	
American Ethnological Society, report on Squier and Davis' work. (R. 1847) I	ł, K
American migration—F. Von Hellwald; Jos. Henry. (R. 1866)	214
American natural history, bibliography for 1851. C. GIRARD	48
Amiurus ponderosus, from Mississippi river—T. H. BEAN. (P. 1879)	888
Ammonia-cobalt bases—W. GIBBS; F. A. GENTH.	88
Ampère, eulogy on-F. Arago. (R. 1872)	271
Amphorosteus, memoir on—R. W. GIBBES	14
Ampullariids W. G. BINNEY	144
Analyses of Chinese and Japanese coals—J. A. MACDONALD	202
Analysis, chemical, of the sun—A. LAUGEL. (R. 1861)	149
Analysis of blood—J. Jones	82
Analysis of water destructive to fish in Gulf of Mexico—F. M. ENDLICH. (P. 1881)	467
Analysis, spectrum—W. Huggins. (R. 1866)	214
Anarrhichas lepturus, from Alaska, and other species—T. H. Bean. (P. 1878)	888
Anatomical preparations, arsenic acid for protecting, from insects—J. B. S. JACKSON. (P. 1878)	882
Anatomy and physiology of chelonia—S. W. MITCHELL; G. R. MOREHOUSE.	159
Anatomy and physiology of lucernarians—H. J. CLARK	242
Anatomy and physiology of the rattlesnake—S. W. MITCHELL	135
Anatomy of the nervous system of Rana pipiens-J. WYMAN	45
Ancient aboriginal trade in North America—C. RAU. (R. 1872) 271, 440,	385
Ancient and modern purple dyeing. (R. 1863)	187
Ancient burial mound in Indiana—W. Pidgeon. (R. 1867)	215
Ancient earthworks of Ashland county, Ohio-G. W. Hill. (R. 1877)	828
Ancient earthworks of Ohio, sketch of—I. DILLE. (R. 1866)	214
Ancient earthworks on Upper Missouri-A. BARRANDT. (R. 1870)	244
Ancient fauna of Nebraska-J. Leidy	58
Ancient fort and burial ground, Tomp. Co., N. Y.—D. TROWBRIDGE. (R. 1863)	187
Ancient graves and shell-heaps of California—P. Schumacher. (R. 1874)	286
Ancient history of North America—M. Much. (R. 1871)	249
Ancient implement of wood found in Conn.—E. W. Ellsworth. (R. 1876)	299

Ancient man in Michigan, characteristics of—H. GILLMAN. (R. 1875) 298, 3	193
Ancient mica mines North Carolina—C. D. SMITH. (R. 1876)	
Ancient mining on Lake Superior-C. WHITTLESEY	L55
Ancient mining on Lake Superior, circular on-J. HENRY. (R. 1861)	149
Ancient monuments in the Miss. val., correspondence relative to. (R. 1847) H,	, K
Ancient monuments in the Mississippi valley—E. G. SQUIER; E. H. DAVIS	1
Ancient mound in western Pennsylvania-W. M. TAYLOR. (R. 1877)	823
Ancient mound near Chattanooga, Tennessee-M. C. READ. (R. 1867)	215
Ancient mound, near Lexington, Ky.—R. PETER. (R. 1871)	249
Ancient mound, St. Louis, MoT. R. PEALE. (R. 1861)	149
Ancient mounds in Georgia-M. F. Stephenson. (R. 1870)	344
Ancient mounds in Kentucky—R. Peter. (R. 1872)	271
Ancient mounds of Mercer county, Illinois—T. McWhorter. (R. 1874)	286
Ancient mounds of Union county, Kentucky-S. S. Lyon. (R. 1870)	244
Ancient pottery from Phillips county, Arkansas—J. H. DEVEREUX. (R. 1872)	271
Ancient pottery on Des Moines river-R. N. and C. DAHLBERG. (R. 1879)	34 5
Ancient relic of Maya sculpture—A. Schott. (R. 1871)	249
Ancient relics in Missouri-J. W. FOSTER. (R. 1868)	187
Ancient relics in northwestern Iowa—J. B. Cutts. (R. 1872)	271
Ancient remains in Canada West, near Prescott—W. E. Guest. (B. 1856)	91
Ancient remains in Colorado—E. L. BERTHOUD. (R. 1867)	215
Ancient ruin in Arizona—J. C. Y. LEE. (R. 1872)	271
Ancient town in Minnesota-O. H. KELLEY. (R. 1868)	187
Ancient works in Ohio, description of—C. WHITTLESEY	37
Anderson, M. B. Life of Professor Chester Dewey. (R. 1870)	244
ANDERSON, W. Antiquities of Perry county, Ohio. (R. 1874)	286
ANDERSON, W. G. Mounds in Illinois and Wisconsin. (R. 1879)	345
Andrews, F. D. Indian relics from Schoharie, New York. (R. 1879)	345
Andrews, S. L. Meteors. (R. 1866)	
Anemometer, description of Smithsonian—J. Henry. (R. 1860)	
Aneroid barometers, experiments—B. STEWART. (R. 1868)	
Animal kingdom, list of substances derived from—G. B. Goode	20
Animal kingdom, parthenogenesis in—G. A. KORNHUBER. (R. 1871)	249
Animal resources of United States, classification of collection—G. B. Goode.	297
Animal resources of United States, collection at Centennial—G. B. Goods	326
Animals, circular for shipping—S. F. BAIRD	384
Animals, flora and fauna within living—J. LEIDY	
Animals, list of all generic names of—S. H. SCUDDER	
Animals, synopsis of useful and injurious—G. B. GOODE	
Anna, Illinois, mounds near—T. M. PERRIN. (R. 1872)	
Annals of Philosophy, list of Smithson's papers in. (B. 1858)67	, 330
Annelids of Arctic America—A. E. VERRILL	342

Annelids of Kerguelen Island—A. E. Verrill	294
Annelids of northeastern coast of America—A. E. VERRILL. (P. 1879; P.	
1880) 883	, 425
Annular eclipse of May 26, 1864	66
Antennariidæ, note on—T. GILL. (P. 1878)	882
Anthozoa of Kerguelen Island-A. E. VERBILL.	294
Anthozoa of northeast coast of America—A. E. VERRILL. (P. 1879)	888
Anthropological articles, Smithsonian, index to-G. H. BOEHMER. (R. 1879) 845,	421
Anthropological Congress, address before—F. F. ROMER. (R. 1876) 299,	392
Anthropological correspondence, abstracts of—O. T. Mason. (R. 1880)	442
Anthropological correspondence, summary of—O. T. MASON. (R. 1879)	345
Anthropological data, limitations to use of-J. W. Powell. (E. 1880)	476
Anthropological inquiry, queries for—C. DARWIN. (R. 1867)	215
Anthropological investigations in 1879-O. T. MASON. (R. 1879) 845,	420
Authropological measurements, table of-Scherzer; Schwarz. (R. 1866)	214
Anthropological publications, list of C. RAU's. (P. 1881)	467
Anthropological Society of Paris, transactions of-P. Broca. (R. 1868)	224
Anthropological Society of Vienna, address on ancient history of North	
America—M. Much. (R. 1871)	249
Anthropology, articles on. (R. 1879)	845
Authropology, bibliography of-O. T. MASON. (R. 1880) 442,	432
Anthropology, index to Smithsonian papers on-G. H. BORHMER. (R. 1879).	845
Anthropology, progress of, in 1880—O. T. Mason. (R. 1880) 442,	432
Authropology, recent progress in-O. T. Mason. (R. 1880) 442,	
Antigua, catalogue of Ober's collection of birds of-G. N. LAWRENCE. (P. 1878)	882
Antilles, maritime disasters of the. (R. 1867)	215
Antilles—See Lesser Antilles.	
Antiquarian Society, American. Antiquities of Wisconsin	70
Antiquities, California, Santa Rosa Island—S. Bowers. (R. 1877)	82 8
Antiquities, Col., Jefferson and Clear Creek Cos.—G. L. CANNON. (R. 1877)	828
Antiquities, Colorado, Weld county—E. L. BERTHOUD. (R. 1871)	249
Antiquities, Denmark, preservation of-J. J. A. WORSAAE. (R. 1879)	8 4 5
Antiquities, Florida—J. BARTRAM. (R. 1874)	286
Antiquities, Florida—A. MITCHELL. (R. 1874)	286
Antiquities, Georgia, Spalding county-W. B. F. BAILEY. (R. 1877)	828
Antiquities, Guatemala—G. WILLIAMSON. (R. 1876)	299
Antiquities, Hungary, prehistoric-F. F. ROMER. (R. 1878) 299, 440,	392
Antiquities, Illinois, Lawrence county—A. Patton. (R. 1878)	275
Antiquities, Illinois, Mason county—J. Cochrane. (R. 1877)	828
Antiquities, Illinois, Rock Island county—A. Toellner. (R. 1879)	845
Antiquities, Illinois, Union county—T. M. PERRINE. (R. 1878)	275
Antiquities, Illinois, Whitesides county—W. H. PRATT. (R. 1874)	286
• • • • • • • • • • • • • • • • • • • •	

Antiquities, Indiana, Allen and DeKalb counties—R. S. Robertson. (R. 1874) 296
Antiquities, Indiana, Knox county—A. Patton. (R. 1878)
Antiquities, Indiana, Laporte county—R. S. Robertson. (R. 1874)
Antiquities, Kentucky-S. S. Lyon. (R. 1858)
Antiquities, Kentucky, Hancock county—J. FRIEL. (B. 1877)
Antiquities, Maryland, Charles county-O. N. BRYAN. (R. 1874) 286
Antiquities, Mexico, Vera Cruz-H. Finck. (R. 1870) 24
Antiquities, Michigan, Isle Royale—A. C. Davis. (R. 1874) 288
Antiquities, Mississippi river and Lake Pepin-L. C. Estes. (R. 1866) 214
Antiquities, Mississippi, Yazoo county—J. W. C. Smith. (R. 1874) 296
Antiquities, Missouri and Tennessee-I. DILLE. (R. 1862)
Antiquities, Missouri, Kansas City-W. H. R. LYKINS. (R. 1877) 324
Antiquities, New Mexico-W. B. Lyon. (R. 1871) 249
Antiquities, New York, Orleans county—F. H. Cushing. (R. 1874)
Antiquities, Nicaragua-E. G. SQUIER. (R. 1850)
Antiquities, North Carolina, Lenoir county-J. M. SPAINHOUR. (R. 1871) 249
Antiquities, N. C., Stanley and Montgomery counties-F. J. KRON. (R. 1874)
Antiquities, Ohio-G. W. HILL. (R. 1874) >
Antiquities, Ohio, Perry county—W. Anderson. (B. 1874)
Antiquities, Olmutz pile-work. (R. 1866)
Antiquities, Porto Rico, Latimer collection—O. T. MASON. (R. 1876) 299, 397
Antiquities, Southern States—H. C. WILLIAMS. (R. 1870) 244
Antiquities, Tennessee-W. M. CLARK. (R. 1877)
Antiquities, Tennessee—E. O. Dunning. (B. 1870)
Antiquities, Tennessee—J. Jones 259
Antiquities, Tennessee-R. S. Robertson. (R. 1877)
Antiquities, Tennessee-D. F. WRIGHT. (R. 1874)
Antiquities, Tennessee, Blount county-Miss A. E. Law. (R. 1874)
Antiquities, Tennessee, Jackson county-J. Halle; J. W. McHenry. (B. 1874) 26
Antiquities, United States, proposed map of-A. J. HILL. (R. 1861)
Antiquities, Wisconsin—I. A. LAPHAM
Antiquities, Wisconsin-M. Strong. (R. 1877)
Antiquity in Europe, the study of high—A. Morlot. (R. 1862; R. 1864) 150.100
Antiquity in United States, vestiges of S. F. HAVEN
Antozone and ozone—C. M. WETHERILL. (R. 1864)
Apaches, Tonto, notes on the-C. SMART. (R. 1867)
Aphododeridæ, notes on—D. S. JORDAN
Apodichthys-Jordan and Gilbert. (P. 1880)
Apogon pandionis, description of-G. B. GOODE; T. H. BEAN. (P. 1881)
Apparatus available for research, list of. (R. 1878)
Apparatus, blow-pipe—Hawkins and Wale. (R. 1872)
Apparatus for testing perspiration and respiration—Perrengoges (R 1861-19

Apparatus, letter relative to gift of-R. HARE. (R. 1848)	I
Appropriations for Museum, memorial of regents asking for. (R. 1867) 215, 32	29
Aqueous vapor, diminution of, with altitude—J. HANN. (B. 1877) 828, 89	98
	42
Arago, F. Aeronautic voyages. (R. 1868)18	87
	44
	41
	71
	4 9
•	99
Eulogy on Herschel. (R. 1870) 24	44
Eulogy on La Place. (R. 1874) 26	86
Eulogy on Alex. Volta. (R. 1875) 20	98
Eulogy on Thomas Young. (R. 1869) 22	28
Archæological Academy of Belgium, Archæological Congress organized by.	
	14
Archæological collection of the U. S. National Museum—C. RAU 28	37
Archæological Congress, International, Antwerp. (R. 1866)	14
Archeological investigations, instructions for-G. GIBBS. (R. 1861) 149, 16	j0
Archæological researches at Concise—F. TROYON. (R. 1861)	49
Archæological researches in Nicaragua—J. F. Bransford	13
Archæological study of jade—S. BLONDEL. (R. 1876) 26	99
Archæology, circular relative to American	16
Archæology, French Society of. (R. 1866)	14
Archæology, general views on—A. MorLor. (R. 1860; R. 1861) 147, 14	49
Archæology, Mexican—B. MAYER	36
Archæology, North American—J. Lubbock. (R. 1862)	50
Archæology of the United States-S. F. HAVEN 7	1
Archæology, prehistoric, international code of symbols for charts of. (R. 1875) 29	98
Archæology and ethnology, circular for collecting specimens of 20)5
Architect's report—B. S. ALEXANDER. (R. 1854) 75, 82	29
Architect's report—A. CLUSS. (R. 1867) 215, 82	29
Architects' report for 1879—Cluss & Schulze. (R. 1879) 345, 40	
Architects' report for 1880—Cluss & Schulze. (R. 1880) 442, 46	
A A	P
Architecture in relation to ventilation, warming, lighting, fire-proofing, acoustics,	
	91
A street and the street are set	95
Arctic America, natural history of—L. Kumlien	
A	96
and the second s	34
Arctic expedition, Howgate-L. Kumlikn	12

ALPHABETICAL INDEX

Arctic explorations, lecture on—I. I. HAYES. (R. 1861)	149
Arctic latitudes and longitudes—E. K. KANE	129
Arctic observations—See Force, Hall, Hayes, Kane, Kumlien, McClintock, Sch	ott.
Arctic regions, limpets and chitons of, report on-W. H. Dall. (P. 1878)	862
Arctic regions, record of voyage of yacht "Fox"	146
Arctic seas, astronomical observations—E. K. KANE	129
Arctic seas, magnetical observations—E. K. KANE	97
Arctic seas, meteorological observations—E. K. KANE	104
Arctic seas, meteorological observations—F. L. McClintock	146
Arctic seas, physical observations in—I. I. HAYES.	196
Arctic seas, physical observations in—E. K. KANE	198
Arctic seas, tidal observations—E. K. KANE	130
Arctic temperature—I. I. HAYES	196
Arctic winds—I. I. HAYES.	196
Argentina syrtensium, new deep-sea fish-Goods and Bean. (P. 1878)	332
Argentine Republic, astronomical observatory—B. A. Gould. (R. 1873)	275
Ariopsis Milberti, breeding habits of-N. T. Lupton. (P. 1878)	33.)
Arizona, ancient ruin in-J. C. Y. Lez. (R. 1872)	271
Arizona, explorations in—J. STEVENSON. (R. 1880)	442
Arizona, Pima county, ruins in-R. T. Buer. (R. 1879)	845
Arizona, Pima Indians of-F. E. Grossman. (R. 1871)	249
Arizona, Tonto Apaches of-C. SMART. (R. 1867)	215
Arkansas, cretaceous fossils from—C. A. WHITE. (P. 1881)	417
Arkansas, earthworks in-Mrs. G. KNAPP. (R. 1877)	325
Arkansas, invertebrate fossils from—C. A. WHITE. (P. 1880)	427
Arkansas, meteorological observations near Washington-N. D. Smith	
Arkansas, Phillips county, ancient pottery-J. H. DEVEREUX. (R. 1872)	271
Armstrong, T. Mounds in Winnebago county, Wisconsin. (R. 1879)	34.,
Arrangement of families of birds—S. F. BAIRD	210
Arrangement of families of fishes—T. GILL	247
Arrangement of families of mammals—T. GILL	230
Arrangement of families of mollusks—T. Gill	227
_	149
Arrangement of the materia medica collection-J. M. FLINT. (P. 1881) 467,	
Arrangements, business, of the Smithsonian Institution	325
Arrow-heads, deposit of, near Fishkill, New York-E. M. SHEPARD. (R. 1877)	325
Arrow-heads, Indian mode of making and obtaining fire—G. Crook. (R. 1871)	249
Arrows, lectures on wounds of poisoned—D. Brainard. (R. 1854)	7.5
Arsenic acid for protecting anatomical preparations—J. B. S. JACKSON. (P. 1878)	3
Art, on tables of constants of nature and—C. BABBAGE. (B. 1856)	91
Art Gallery, Corcoran, deed of gift of. (R. 1872) 271,	\$29
Art Gallery, Corcoran, deposits in hy Smithsonian Institution (R. 1974)	286

Art Gallery, Corcoran, report of Committee of Regents on. (R. 1872) 271, 329
Arteries, reparatory inflammation in. Toner lecture No. VII-E. O. SHAKE-
8PEARE 321
Artificial shell deposits in New Jersey-C. RAU. (R. 1864) 188, 440, 362
Artificial shell deposits of the United States-D. G. BRINTON. (B. 1868) 214
Artisan and artist, identification of the-N. WISEMAN. (R. 1870) 244
Artisans, scientific education of mechanics and—A. P. Prabody. (R. 1872) 271, 380
Arts, chemical, recent improvements in—J. C. BOOTH; C. MORFIT
Ascelicthys rhodorus—Jordan and Gilbert. (P. 1880) 425
Ascensions—See Balloons.
Assay of coins at mint of the United States-J. Pollock. (R. 1868) 224
Asteroid supplement to new tables—J. D. RUNKLE 94
Asteroids between Mars and Jupiter—D. KIRKWOOD. (R. 1876) 299
Astronomical discoveries, telegraphic announcements of-J. Henry 263
Astronomical observations—W. HARKNESS
Astronomical observations, Arctic—I. I. HAYES
Astronomical observations, discussion of Piazzi's-B. A. Gould. (R. 1868) 187
Astronomical observations in the Arctic Seas—E. K. KANE 129
Astronomical observatories, reports of—E. S. Holden; G. H. Boehmer. (R. 1880)442, 438
Astronomical observatories, reports of, 1879—E. S. Holden. (R. 1879) 845, 410
Astronomical observatories, reports of, 1880—E. S. Holden; G. H. Boehmer.
(R. 1880)
Astronomical observatory at Cordoba, Argentine Republic, account of—B. A.
GOULD. (R. 1878) 275
Astronomical photography, progress of—Dr. LEE. (R. 1861) 149
Astronomical tables—J. D. RUNKLE
Astronomical telegram circular—J. Henry 263
Astronomy, lecture on—A. CASWELL. (R. 1858)
Astronomy, progress of, in 1879 and 1880—E. S. Holden. (R. 1880) 442, 427
Astronomy, recent progress in—E. S. Holden. (R. 1880) 442, 427
Astroscopus, species of, in Eastern United States.—T. H. BEAN. (P. 1879) 388
Atlantic coast, microscopic examination of soundings-J. W. BAILEY 20
Atlantic, Lowe's aeronautic voyage across. (R. 1860) 147
Atlas, physical, of North America, proposed—G. Gibbs. (R. 1866) 214
Atmosphere, currents of the, and aerial navigation—J. HENRY. (R. 1860) 147
Atmosphere, diminution of aqueous vapor with altitude—J. HANN. (R. 1877) 323, 398
Atmosphere, plan of research upon—C. M. WETHERILL. (R. 1866) 214
Atmosphere, vegetation and the—J. Jamin. (R. 1864) 188
Atmospheric circulation over the earth, laws of—J. H. Coffin 268
Atmospheric currents, relation between barometric variations and-M. Peslin.
(R 1877) 828, 398

Atmospheric electricity—M. F. DUPBEZ. (R. 1858)	109
Atmospheric pressure and rain fall—J. HANN. (R. 1877)	8, 398
Atmospheric pressure, Arctic—I. I. HAYES	196
Atmospheric pressure, Arctic-E. K. KANE	104
Atmospheric pressure, Arctic-F. L. McCLINTOCK.	146
Atomic volume, bibliography of—F. W. CLABKE	2.5
Atomic weight determinations—G. F. BECKER.	358
Atomic weights, recalculation of-F. W. CLARKE	441
Atoms—J. Herschel. (R. 1862)	150
Atthis Ellioti, a humming bird from Guatemala—R. RIDGWAY. (P. 1878)	302
Aurora and electricity—S. Lemström; A. A. De La Rive. (R. 1874)	256
Aurora, connection of gales of wind and—R. T. KNIGHT. (R. 1871)	249
Aurora, directions for observing-J. HENRY. (R. 1855)	77
Aurora, effect of, on magnetic declination—A. D. BACHE	175
Aurora, influence of, on the telegraph—W. D. SARGENT. (R. 1870)	244
Aurora, instructions for observations of—Toronto Observatory	148
Aurora, magnetic effect of—A. D. BACHE	175
Aurora, map of stars for observations on	350
Aurora, observations on—S. Masterman. (R. 1857)	107
Aurora, or Polar light, its phenomena and laws—E. Loomis. (R. 1865)	209
Aurora, phenomena in telegraphic lines during-G. B. Donati. (R. 1872)	271
Aurora, recent secular period of—D. Olmsted	81
Auroras, articles on, received by the Institution. (R. 1871; R. 1874) 249	, 286
Auroras, records of, in high northern latitudes—P. Force	84
Autobiography, the history of my youth—F. ARAGO. (R. 1870)	244
Auxis Rochei, frigate mackerel, on New England coast—G. B. Goode. (P. 1880)	125
Awards to Smithsonian Institution at the Centennial Exhibition. (R. 1876; R. 1878)	, 341
Axolotl changed to amblystoma—A. Weismann. (R. 1877) 323,	401
AVERS Dr. Notes on fishes (P. 1880)	425

B.

BABBAGE, C., extracts from writings of. (R. 1873)	16
BABBAGE, C., memoir of, by N. S. Dodge. (R. 1878)	/5
BABBAGE, C. On constants of nature and art. (R. 1856))1
BABINET, J. The diamond and other precious stones. (R. 1870) 244, 37	7
BABINET, J. The northern seas. (R. 1869)	:8
Васне, А. D.—	
Bequest to National Academy of Sciences. (R. 1872) 271, 32	9
Discussion of magnetic and meteorological observations. Part I 11	3
Discussion of magnetic and meteorological observations. Part II 12	1
Discussion of magnetic and meteorological observations. Part III 13	2
Discussion of magnetic and meteorological observations. Pts. IV, V, VI_ 16	2
Discussion of magnetic and meteorological observations. Pts. vii, viii, ix. 17	5
Discussion of magnetic and meteorological observations. Pts. x, xI, xII 18	6
Discussion of magnetic and meteorological observations. Parts I-XII 19	5
Eulogy on James A. Pearce. (R. 1862) 15	0
Lecture on Switzerland. (R. 1870) 24	4
Magnetic survey of Pennsylvania	6
Notice of James P. Espy. (R. 1859) 11	0
Notice of G. Würdemann. (R. 1859)	0
BACHE, A. D., eulogy on, by J. HENRY. (R. 1870) 244, 37	9
BACHE, A. D., list of scientific papers of-B. A. Gould. (R. 1870) _ 244, 329, 37	6
BACHE, G. M. Hail storm in Texas. (R. 1870)	4
Bacteria, putrefactive, action of dry heat and sulphurous acid upon. (P. 1881) 46	7
BAEGEBT, J. Aboriginal inhabitants of California peninsula. (R. 1868; R.	
1864) 187, 188, 440, 36	1
Baffin's Bay, meteorological observations in-F. L. McCLINTOCK	6
Bailey, J. W.—	
Algæ2	8
Diatomacea	8
	8
Microscopical examination of soundings, Atlantic coast	0
Microscopical observations in South Carolina, Georgia, Florida 2	3
New species and localities of microscopical organisms	3
BAILEY, W. B. F. Antiquities of Spalding county, Georgia. (R. 1877) 32	3
Baird, S. F.—	_
•Alphabetical index of genera of birds	
Arrangement of families of birds 21	
Catalogue of North American birdsoctavo, 108; quarto, 10	
Catalogue of North American mammals 10	5

BAIRD, S. F.—Continued.	
Circular concerning department of insects. (P. 1881) 467, 44	8
Circular in reference to building-stone collection. (P. 1881) 467, 45	
Circular in reference to petroleum collections. (P. 1881) 467, 44	7
Circular in reference to shipping fresh fish and other animals 38	4
Circular relative to scientific and literary exchanges 32	4
Circular requesting material for the library of the National Museum. (P. 1881) 467, 45	8
Circular to friends of the Museum. (P. 1881) 467, 44	6
Directions for collecting, preserving, and transporting specimens of natu-	
ral history. (R. 1856) 91, 3	4
Fishes on the coasts of New Jersey and Long Island. (R. 1854) 75, 34	
Instructions for collecting insects. (R. 1858)	Ŷ
Instructions for collecting nests and eggs of North American birds. (R. 1858)	9
List of explorations furnishing collections to National Museum, 1838 to	
1877. (R. 1877)	3
List of sources from which specimens have been derived, 1888 to 1867. (R. 1867)	5
Memoranda of inquiry relative to food fishes	1
Museum miscellanea	ŧ
Notes on aboriginal shell-mounds on coast of New Brunswick and of	
37 77 1 1 1 47 4001)	_
New England. (P. 1881) 46	7
New England. (P. 1881) 40 Notes on Henry's researches on sound. (R. 1878) 341, 400	
	6
Notes on Henry's researches on sound. (R. 1878) 341, 400	6 ≟
Notes on Henry's researches on sound. (R. 1878) 341, 400 On reports of recent scientific progress. (R. 1880) 44	6 <u>-</u>
Notes on Henry's researches on sound. (R. 1878) 341, 40 On reports of recent scientific progress. (R. 1880) 44 Questions relative to food fishes 23	6 4 8
Notes on Henry's researches on sound. (R. 1878) 341, 40 On reports of recent scientific progress. (R. 1880) 44 Questions relative to food fishes 23 Register of periodical phenomena 65, 144 Report of Secretary for 1878. (R. 1878) 341, 404	6
Notes on Henry's researches on sound. (R. 1878)	6
Notes on Henry's researches on sound. (R. 1878) 341, 40 On reports of recent scientific progress. (R. 1880) 44 Questions relative to food fishes 23 Register of periodical phenomena 65, 14 Report of Secretary for 1878. (R. 1878) 341, 404 Report of Secretary for 1879. (R. 1879) 845, 414	6 2 4 8 8
Notes on Henry's researches on sound. (R. 1878) 341, 40 On reports of recent scientific progress. (R. 1880) 44 Questions relative to food fishes 23' Register of periodical phenomena 65, 14 Report of Secretary for 1878. (R. 1878) 341, 40 Report of Secretary for 1879. (R. 1879) 345, 41 Report of Secretary for 1880. (R. 1880) 442, 41	6 2 8 8 8
Notes on Henry's researches on sound. (R. 1878)	6 2 4 8 8 8 7
Notes on Henry's researches on sound. (R. 1878) 341, 40 On reports of recent scientific progress. (R. 1880) 44 Questions relative to food fishes 23 Register of periodical phenomena 65, 14 Report of Secretary for 1878. (R. 1878) 341, 40 Report of Secretary for 1879. (R. 1879) 845, 41 Report of Secretary for 1880. (R. 1880) 442, 41 Report on Centennial Exhibition. (R. 1876) 299, 30 Report on and statistics of British Museum. (R. 1850) 29	6 2 4 8 8 9 7 5
Notes on Henry's researches on sound. (R. 1878)	6 2 4 8 8 9 7 2 5
Notes on Henry's researches on sound. (R. 1878)	6 2 4 8 8 8 9 7 2
Notes on Henry's researches on sound. (R. 1878)	6 2 4 8 8 9 7 2 3
Notes on Henry's researches on sound. (R. 1878)	6 2 4 8 8 8 9 7 2 5

Balloon ascensions—See Aeronautics.
BALTZELL, J. Winds in Florida. (R. 1866)
BANCROFT, G. Memorial to Congress for new Museum building. (R. 1876) 299
BANCROFT, G. On publication of Spanish works on New Mexico. (R. 1855). 77
BANNISTER, H. M. Climate of Alaska. (B. 1866) 214
Bannister, H. M. Formation of clouds over Gulf stream. (R. 1866) 214
Baptist Missionary Society. Yoruba grammar and dictionary 98
Barbuda, catalogue of Ober's collection of birds of—G. N. LAWBENCE. (P. 1878)
BARKER, G. F. Recent progress in chemistry. (R. 1880) 442, 429
BARKER, G. F. Recent progress in physics. (R. 1880) 442, 429
BARNARD, F. A. P. Lectures on the undulatory theory of light. (R. 1862) 150
BARNARD, J. G. Eulogy on Gen. Joseph G. Totten. (R. 1865) 209
BARNARD, J. G. Internal structure of the earth
BARNARD, J. G. Problems of rotary motion 240
BARNARD, V. Birds of Chester county, Pennsylvania. (R. 1860) 147
Barometer, account of Green's. (R. 1855)
Barometer, Green's standard 148
Barometer, horary variations of the-F. Vaillant; J. Henry. (R. 1866) 214
Barometer, influence of rain upon the-J. HANN. (R. 1877) 323, 398
Barometer, rain and snow gauges—R. H. GARDINER. (R. 1858) 109
Barometer tables, on construction of—F. F. TUCKETT. (R. 1867) 215
Barometer tube breaking suddenly—L. F. WARD. (R. 1866)
Barometer tubes, on filling-J. GREEN; W. WÜRDEMANN. (R. 1859) 110
Barometers at the Kew Observatory, verification of-J. WELSH. (R. 1859) 110
Barometers, experiments on aneroid—B. STEWART. (R. 1868) 224
Barometric minima and rainfall—J. T. REYE. (R. 1877)
Barometric observations, discussion of Snell's-F. H. Loud. (R. 1880) 442, 435
Barometric observations, results of 157
Barometric variations and atmospheric currents, relation between-M. PESLIN.
(R. 1877) 328, 898
Barometrical tables—A. Guyor
BARRANDT, A. Ancient earthworks on the upper Missouri. (R. 1870) 244
BARRANDT, A. Haystack mound, Dakota. (R. 1872)
BARTLETT, J. R. On publication of Squier and Davis' Ancient Monuments.
(R. 1847) H, K
BARTRAM, J. Antiquities of Florida. (R. 1874) 286
Base chart of the United States—C. A. Schott
Bases, ammonia-cobalt, researches on—W. Gibbs; F. A. Genth
Bases, organic—A. BAUER. (R. 1872)
Batavian Society of Experimental Philosophy, Rotterdam. Prize questions.
(R 1861) 149

Bath and West of England Society for encouragement of Agriculture, Arts, etc. Exchanges. (R. 1867) 2	215
Batis maritima—J. Torrey	60
Batrachia, North American—E. D. Cope2	92
Bats, account of remarkable accumulation of—M. FIGANIEBRE. (R. 1863) 1	187
Bats of North America, monograph of—H. ALLEN 1	65
BAUER, A. Organic bases. (R. 1872)	271
Bay of Fundy, marine invertebrata of—W. STIMPSON	50
Bead, colored, from a mound at Black Hammock, Florida—A. M. HARRISON. (R. 1877)	323
Bead, polychrome, from Florida-S. S. HALDEMAN. (R. 1877) 323, 4	104
Bean, T. H.—	
	425
Description of a new hake (Phycis Earlii) and note on Phycis regius. (P. 1880)	425
Description of new fish from Alaska, Anarrhichas lepturus, and notes on genus. (P. 1879)	833
Description of new sparoid fish, Sargus Holbrookii, from Savannah bank. (P. 1878)	g.,
Description of new species of Amiurus from the Mississippi. (P. 1879).	333
Description of new species of Gasterosteus from Schoodic Lakes, Maine. (P. 1879)	\$3 3
Description of species of Lycodes (L. Turneri) from Alaska. (P. 1878).	332
Descriptions of genera and species of Alaskan fishes. (P. 1879)	373
Descriptions of new fishes from Alaska and Siberia. (P. 1881)	467
Descriptions of new species of fishes (Uranidea marginata and Potamo-cottus Bendirei,) and of Myctophum crenulare. (P. 1881)	4 5
Descriptions of species of fishes collected by Dugès in Mexico. (P. 1879)	333
Directions for collecting and preserving fish. (P. 1881) 467,	464
Fishes of Arctic America. (Bull. 15, U.S. National Museum)	342
Identity of Euchalarodus Putnami with Pleuronectes glaber and notes	
on habits. (P. 1878)	1,2
List of European fishes in National Museum. (P. 1879)	333
Notes on collection of fishes from eastern Georgia. (P. 1879)	3.3
Notes on fishes collected by H. E. Nichols in British Columbia and Alaska, with descriptions of new species and new genus (<i>Delolepis.</i>) (P. 1881)	407
Notes on fishes from Hudson's Bay. (P. 1881)	467
Occurrence of Hippoglossus vulgaris in Alaska. (P. 1879)	GB.
Occurrence of Stichaus punctatus at St. Michael's, Alaska. (P. 1878).	320
Partial bibliography of fishes of Pacific coast of United States and Alaska for 1880. (P. 1881)	
Preliminary catalogue of fishes of Alaskan waters. (P. 1881)	
Species of Advancement of France II-ited States (B. 1970)	

BEAN, T. H.; GOODE, G. B.—	
Catalogue of Stearns's fishes of Florida and descriptions of new species.	
(P. 1879)	388
Catalogue of Velie's fishes in Gulf of Mexico and description of new species. (P. 1879)	888
Craig flounder of Europe (Glyptocephalus cynoglossus) on coast of North	
• America. (P. 1878)	3 82
Description of Alepocephalus Bairdii, new deep-sea fish, western Atlantic. (P. 1879)	888
Description of Argentina systemsium, new deep-sea fish, Sable Island bank. (P. 1878)	882
Description of Caulolatilus microps from Gulf coast of Florida. (P. 1878)	882
Description of Lycodes paxillus. (P. 1879)	388
Description of new genus and species of fish, Lopholatilus chamæleonticeps, from New England. (P. 1879)	888
Description of new species of fish, Apogon pandionis, from mouth of Chesapeake Bay. (P. 1881)	
Description of new species of fish (Lutjanus Blackfordii and Lutjanus Stearnsii) from coast of Florida. (P. 1878)	882
Descriptions of gadoid fishes, Physis chesteri and Haloporphyrus viola, from northwestern Atlantic. (P. 1878)	882
Identity of Brosmius brosme Americanus, Gill, with Brosmius brosme (Müller) White. (P. 1878)	882
Identity of Rhinonemus caudacuta with Gadus cimbrius. (P. 1878)	832
New genus of fishes, Benthodesmus. (P. 1881)	4 67
New serranoid fish, Epinephelus Drummond-Hayi, from Bermudas and Florida. (P. 1878)	882
New species of Liparis (L. ranula) from Halifax. (P. 1879)	888
Note on Platessa ferruginea and Platessa rostrata. (P. 1878)	882
(P. 1878)	382
Occurrence of Lycodes Vahlii on La Have and Grand Banks. (P. 1879.)	338
Oceanic bonito on coast of United States. (P. 1878)	882
BEAUCHAMP, W. M. Wampum belts of the Six Nations. (R. 1879)	345
Beaufort, (N. C.,) library, deposit of, by E. M. Stanton. (R. 1862)	150
Beaufort, N. C., notes on fishes of D. S. Jordan. (P. 1880)	425
Beaufort harbor, notes on fishes of—D. S. JORDAN; C. H. GILBERT. (P. 1878)	832
BEAUMONT—See Do Beaumont. REALIZEMENT REALIZEMENT C. F. mornoir of hy Five Dr Realizement (P. 1962)	187
Beautemps-Beaupré, C. F., memoir of, by Elie de Beaumont. (R. 1863) Beaver, on the habits of the—Felix R. Brunot. (R. 1878)	
Beaver Island, Mich., natural history of—J. J. STRANG. (R. 1854)	215 75
Becker, G. F. Atomic weight determinations	

BECQUEREL, A. E. Forests and their climatic influence. (R. 1869) 228
BECQUEREL, A. E. Preservation of copper and iron in salt water. (R. 1864.) 188
Beirut, Syria, account of sarcophagus from-A. A. HARWOOD. (R. 1870) 244
BELDING, L. Additions to North American bird fauna by—R. RIDGWAY. (P. 1881)
BELDING, L. Partial list of birds of central California. (P. 1878)
Belgium, Archæological Academy of. Archæological congress organized by. (R. 1866)
Belgium—See Prize questions.
Belgium, Society of Science, etc.—See Prize questions.
Belone exilis, generic relations of-D. S. JORDAN; C. H. GILBERT. (P. 1880) 425
Belone latimanus, occurrence of, in Buzzard's bay-G. B. Goode. (P. 1878) 32
BENDIRE, C. Notes on Salmonida of Columbia river. (P. 1881)
Benthodesmus, a new genus of deep-sea fishes-G. B. GOODE; T. H. BRAN.
(P. 1881) 467
Bequest of A. D. Bache to National Academy of Sciences. (R. 1872) 271, 329
Bequest of S. Habel. (R. 1879)
Bequest of J. Hamilton to Smithsonian Institution. (R. 1872; R. 1873) 271, 273
Bequest of Smithson, act of Congress, July 1, 1886, accepting. (R. 1853) 67. M
Bequest of Smithson, exposition of—JHENRY
Bequest of Smithson, history of 328. 33
Bequest, Smithson and his-W. J. RHEES. (R. 1879)
BEBENDT, C. H.—
Collection of historical documents in Guatemala. (R. 1876) 299
Explorations in Central America. (R. 1867) 215
On the Carib language and people. (R. 1873) 275
recommendation of, by British Minister. (R. 1865)
recommendation of, by Guatemalan Minister. (R. 1865) 209
recommendation of, by Guatemalan Minister. (R. 1865) 209 recommendation of, by L. Molina, Costa Rican Minister. (R. 1865) 209
recommendation of, by Guatemalan Minister. (R. 1865) 209 recommendation of, by L. Molina, Costa Rican Minister. (R. 1865) 209 Berlandier, L. Catalogue of collection of historical and geographical manu-
recommendation of, by Guatemalan Minister. (R. 1865)
recommendation of, by Guatemalan Minister. (R. 1865)
recommendation of, by Guatemalan Minister. (R. 1865)
recommendation of, by Guatemalan Minister. (R. 1865) 209 recommendation of, by L. Molina, Costa Rican Minister. (R. 1865) 299 BERLANDIER, L. Catalogue of collection of historical and geographical manuscripts, maps, &c., of Mexico. (R. 1854)
recommendation of, by Guatemalan Minister. (R. 1865)
recommendation of, by Guatemalan Minister. (R. 1865)
recommendation of, by Guatemalan Minister. (R. 1865)

BERRIEN, J. McPherson. Construction of act establishing Smithsonian. (R. 1853)	(
BERTHOUD, E. L. Ancient remains in Colorado. (R. 1867)	2
BERTHOUD, E. L. Antiquities of Colorado. (R. 1871)	2
BERTHRAND, M. Life and works of Kepler. (R. 1869)	2
Best hours for temperature observations—C. DEWEY. (R. 1860)	14
BETHUNE, C. J. Acknowledgment of books. (R. 1861)	14
Bibliographia Americana Historico-Naturalis for 1851—C. GIRARD	4
Bibliographia Americana, prospectus of a-H. Stevens. (R. 1848)	•
Bibliographical index of North American Botany—S. WATSON	25
Bibliographical sketches of vestiges of antiquity in United States—S. F. HAVEN	-
Bibliography of alge—H. C. Wood	24
Bibliography of American natural history for 1851—C. GIRARD	4
Bibliography of anthropology—O. T. Mason. (R. 1880) 442,	_
Bibliography of atomic volume and specific gravity, expansion, and boiling and	70
melting—F. W. CLARKE	2
Bibliography of Chinook jargon—G. GIBBS	16
Bibliography of Dakota language—S. R. RIGGS	4
Bibliography of diptera—C. R. OSTEN SACKEN	2
Bibliography of diseases of joints, bones, larŷnx, eye, gangrene, etc.—W. W. Keen	80
Bibliography of expansion—F. W. CLARKE	28
Bibliography of fishes—T. GILL	24
Bibliography of fishes of Pacific coast of United States and Alaska for 1880—	
T. H. BEAN. (P. 1881)	40
Bibliography of fishes of Pacific coast of United States—T. GILL	46
Bibliography of mammals—T. GILL.	28
Bibliography of mollusks—T. GILL	22
Bibliography of nebulæ, clusters, milky way, nebular hypothesis, etc.—E. S. Holden	81
Bibliography of North American conchology previous to 1860. Part I. American authors. W. G. BINNEY	14
Bibliography of North American conchology previous to 1860. Part II. Foreign authors—W. G. BINNEY	
Bibliography of ornithology: faunal publications relative to British birds—E. COUES. (P. 1879)	8
Bibliography of orthoptera—S. H. SCUDDER	18
Bibliography of rattlesnakes, serpent venom, etc.—S. W. MITCHELL	18
	2
Bibliography of specific heat—F. W. CLARKE	
Bibliography of Yoruba language—T. J. Bowen	1
Bibliography, plan of—J. FRIEDLÄNDER. (R. 1858)	10
Biela's comet, investigations of—J. S. HUBBARD. (R. 1862)	15
Big elephant mound in Grant county, Wisconsin-J. WARNER. (R. 1872)	2

Benney, W. G.—
Bibliography of North American conchology previous to 1860. Part I. American authors
Bibliography of North American conchology previous to 1860. Part II. Foreign authors 1
Check-list of shells of North America
Land and fresh-water shells of North America. Part I
Land and fresh-water shells of North America. Part II 1
Land and fresh-water shells of North America. Part III 1
Biographical memoir of Joseph Henry—A. GRAY. (R. 1878) 341, 356, 4
Biographical notice of L. Agassiz—E. FAVRE. (R. 1878)
Biographical notice of L. Agassiz—J. A. GARFIELD. (R. 1873)
Biographical notice of L. Agassiz-P. PARKER. (R. 1873)
Biographical notice of S. P. Chase—J. A. GARFIELD. (R. 1878)
Biographical notice of S. P. Chase—H. Hamlin. (R. 1873)
Biographical notice of W. H. Harvey-A. GRAY. (R. 1867)
Biographical notice of Charles C. Jewett-R. A. Guild. (R. 1857)
Biographical sketch of Dom Pedro II-A. FIALHO. (R. 1876)
Biography of Condorcet—F. Arago. (R. 1878)
Biography—See Memoirs.
Bird, a new humming, from Guatemala—R. RIDGWAY. (P. 1878) 3
Birds, acknowledgment of—Academy of Sciences, Stockholm. (R. 1867) 2
Birds, acknowledgment of—J. Gould. (R. 1867)
Birds, acknowledgment of—Hungarian National Museum. (R. 1863)
Birds, alphabetical index of genera of—S. F. BAIRD
Birds, American, instructions for collecting nests and eggs of—S. F. BAIRD. (R. 1858)
Birds and mammals of Missouri river, list of—E. HARRIS. (R. 1850)
Birds, arrangement of families of -S. F. BAIRD 2
Birds, bibliography of British-E. Coues. (P. 1879)
Birds, catalogue of North American—S. F. BAIRD. Octavo1
Birds, catalogue of North American—S. F. BAIRD. Quarto 10
Birds, catalogue of Old World, in Nat. Museum -R. RIDGWAY. (P. 1881) 467, 4
Birds, circular relative to collections of
Birds, dates of first.appearance of—F. B. Hough
Birds, description of new species of, from Costa Rica—R. RIDGWAY. (P. 1878) 3
Birds, descriptions of new-R. RIDGWAY. (P. 1881)
Birds, descriptions of several new species and races of, in National Museum— R. Ridgway. (P. 1878)
Birds, desiderata among North American—R. RIDGWAY. (P. 1881) 4
Birds' eggs, suggestions for forming collections of—A. NEWTON
Birds, humming, list of-D. G. ELLIOT 3

OF SMITHSONIAN PUBLICATIONS.

Birds, list of described, of Mexico, Central America, and the West Indies, not in Smithsonian collections	185
Birds' nests and eggs, instructions for collecting	189
Birds, new species and races of American, and synopsis of genus Tyrannus—R. BIDGWAY. (P. 1878)	882
Birds, new species of Turdidæ, from Dominica—G. N. LAWRENCE. (P. 1880)	425
Birds; nomenclature and catalogue of—R. RIDGWAY	422
Birds. North American oölogy—T. M. Brewer.	89
Birds, notes on some Costa Rican—R. RIDGWAY, (P. 1881)	467
Birds of Antigua, catalogue of Ober's collection of-G. N. LAWRENCE. (P. 1878)	882
Birds of Arctic America—L. Kumlien	842
Birds of Barbuda, catalogue of Ober's collection—G. N. LAWRENCE. (P. 1878)	882
Birds of Bermuda-T. BLAND; J. R. WILLIS. (R. 1858)	109
Birds of Caribbee Islands—F. A. OBER. (R. 1878)	841
Birds of Central California—L. BELDING. (P. 1878)	882
Birds of Chester county, Pennsylvania-V. BARNARD. (R. 1860)	147
Birds of District of Columbia, list of-E. Cours; S. S. Prentiss. (R. 1861)	149
Birds of Dominica, catalogue of Ober's collection of—G. N. LAWRENCE. (P. 1878)	882
Birds of Grenada, catalogue of Ober's collection—G. N. LAWRENCE. (P. 1878)	882
Birds of Guadeloupe, catalogue of Ober's collection—G. N. LAWBENCE. (P. 1878)	882
Birds of Heligoland—H. GÄRKE. (P. 1879)	888
Birds of Kerguelen Island—J. H. KIDDER; E. COUES	293
Birds of Lesser Antilles, catalogue of Ober's collection of—G. N. LAWRENCE, (P. 1878)	882
Birds of Martinique, catalogue of Ober's collection of—G. N. LAWRENCE. (P. 1878)	882
Birds of Middle and South America not in National Museum—R. RIDGWAY. (P. 1881)	467
Birds of Missouri river—E. HARRIS. (R. 1850)	28
Birds of North America, nomenclature and catalogue of—R. RIDGWAY. (P. 1880)	425
Birds of Nova Scotia-Blackiston; T. Bland; J. R. Willis. (R. 1858)	109
Birds of Saint Vincent, catalogue of Ober's collections of—G. N. LAWRENCE. (P. 1878)	832
Birds of South America, circular relative to collections of	168
Birds of Southern Texas—J. C. MERRILL. (P. 1878)	882
Birds of Southwestern Mexico, collected by F. E. Sumichrast—G. N. Law-	295
Birds of West coast, migrations and nesting habits of-J. G. Cooper. (P. 1879)	338
Birds of West Indies, catalogue of Ober's collections from—G. N. LAWRENCE. (P. 1878)	882
Birds, review of American. Part I-S. F. BAIRD	181
Birds, revisions of nomenclature of—R. RIDGWAY. (P. 1880)	425
10	

Birds, suggestions for saving skeletons of—A. NEWTON. (R. 1860) 147
Birds, systematic review of classification of—W. LILLJEBORG. (R. 1865.) 209, 364
Birds, two new North American, added by L. Belding—R. RIDGWAY. (P. 1881) 467
BISHOP, W. D. Commissioner of Patents, Meteorological observations 157
Bison, request of Bern Museum for. (B. 1865)209
Bites of serpents, nature and cure of-D. Brainard. (R. 1854) 75
Black bass of the Ohio, habits of the J. EOFF. (R. 1854) 75
Black Hammock, colored bead from—A. M. HARRISON. (R. 1877) 323
BLACKISTON, -; BLAND, T.; WILLIS, J. R. Birds of Nova Scotia. (R. 1858) 109
Blackmore Museum, Salisbury, England, notice of the. (R. 1868) 224
Black Mountain, North Carolina, topography of-T. L. CLINGMAN. (R. 1855) 77
BLAINVILLE, D. DE, memoir of-M. FLOURENS. (R. 1865) 209
BLAND, T.; BINNEY, W. G. Land and fresh-water shells of North America. Part I.—Pulmonata geophila
BLAND; WILLIS, J. R. Birds of Bermuda. (R. 1858)
BLAND, T.; BLACKISTON, -; WILLIS, J. R. Birds of Nova Scotia. (R. 1858) 103
Blank check-lists, labels, figures, &c
Blind deaf mute, vocal sounds of Laura Bridgman, the—F. LIEBER 12
BLODGET, L. List of meteorological observers. (R. 1853)
BLONDEL, S. Historical, archeological, and literary study of jade. (R. 1876) 29
Blood, analysis of—J. Jones &
Blossoming of plants, dates of—F. B. Hough 182
Blowpipe apparatus of Hawkins and Wale. (R. 1872) 271
Blowpipe, qualitative determination by the-T. Egleston. (R. 1872) 271
BLYDEN, E. D. Mixed races in Liberia. (R. 1870) 24
Board of Commissioners for Foreign Missions—See American Board, etc.
Board of Regents, Journals of. (See each annual report) 829. A
Board of Regents-See Regents.
BOEHMER, G. H. Index to anthropological articles in Smithsonian publications. (R. 1879)
BOEHMER, G. H. Report on International Exchange477
BOEHMER, G. H. Reports of astronomical observatories, 1880. (R. 1880) 442, 438
BOEHMER, G. H.; HOLDEN, E. S. Reports of observatories, 1880. (R. 1880) 442, 436
Boerner, C. G. Lightning discharges. (R. 1867) 213
Boiling point, bibliography of-F. W. CLARKE
Boiling points, tables of—F. W. CLARKE 255, 28
Bolles, E. C. Account of Portland Society of Natural History. (R. 1867) 215
Bologna, Academy of Sciences of the Institute of. Prize questions. (R. 1862)
Bones, bibliography of diseases of the-W. W. KEEN 300
Bonito, oceanic, description of—V. N. EDWARDS. (P. 1878) 32
Renits occanic on coast of United States

OF SMITHSONIAN PUBLICATIONS.	147
Books, acknowledgment of-C. J. BETHUNE. (R. 1861)	149
Books, acknowledgment of—J. A. Codd. (R. 1861)	149
Books, acknowledgment of-M. LABOULAYE. (R. 1867)	215
Books, copyright, from 1846 to 1849—C. C. JEWETT. (R. 1850)	28
Books on Brazil, presented—M. M. LISBOA. (R. 1865)	209
Books on Egypt, presentation of—R. LEPSIUS. (R. 1860)	147
Books, on the classification of—J. P. LESLEY. (R. 1862)	150
Books, presentation of—Duke of Northumberland. (R. 1859)	110
Books, presented—Imperial Library of Vienna. (R. 1865)	209
BOOTH, J. C.; MORFIT, C. Recent improvements in the chemical arts	27
Bordeaux, Chamber of Commerce of. Exchange of publications. (R. 1863)	187
Bordeaux, Imperial Academy of Sciences. Prize questions. (R. 1868)	224
Bosphorus, account of a hailstorm on the—Com. PORTER. (R. 1870)	244
Boston, Mass., American Academy—See American Academy.	
Botanical explorations in New Mexico and California—A. GRAY. (R. 1849)	21
Botany, index to North American—S. WATSON	258
Botany of Hawaiian and Fanning Islands and California—T. H. STREETS	808
Botany of Kerguelen Island-A. GRAY; T. P. JAMES; E. TUCKERMAN; W.	
G. FABLOW	294
Botany, progress in, in 1879 and 1880-W. G. FARLOW. (R. 1880) 442,	430
Botany—See Plants.	
Boundary line between geology and history—E. Surss. (R. 1872)	271
Bowen, T. J. Yoruba grammar and dictionary	98
Bowers, S. Explorations in Santa Rosa Island, California. (R. 1877)	828
BOYD, C. H. Remains of walrus (?) in Maine. (P. 1881)	467
Brachiopoda, index of-W. H. DALL	304
Brachyopsis verrucosus, from California—W. N. Lockington. (P. 1880)	425
Brachyopsis xyosternus—D. S. JOBDAN; C. H. GILBERT. (P. 1880)	425
BRACKETT, A. G. Shoshone or Snake Indians, religion, superstitions, manners,	
etc. (R. 1879)	34 5
Brackett, A. G. Sioux or Dakota Indians. (R. 1876)	299
Brain and marrow of animals, uses of, among Indians—T. R. PEALE. (R. 1870)	244
Brain, dual character of the-C. E. BROWN-SEQUARD. Toner lecture No. II.	291
Brainard, D. Lecture on bites of serpents and wounds of poisoned arrows. (R. 1854)	75
Bransford, J. F. Archæological researches in Nicaragua	
Bransford, J. F., note on shells from Costa Rica kitchen-midden, collected	300
by—W. H. Dall. (P. 1878)	832
BRAVAIS, AUGUSTE, memoir of, by E. DE BEAUMONT. (R. 1869)	228
BRANDON A W Distribution of the School of South Coroline Consults and	

BRAYTON, A. W. Distribution of the fishes of South Carolina, Georgia, and Tennessee.....

Brayton, A. W. North American ichthyology

808

808

Brazil, books on, presented—M. M. Lisboa. (R. 1865) 209
Brazil, specimens from, presented. Brazilian Nat. Hist. Museum. (B. 1865) 39
Brazilian Nat. Hist. Museum, specimens from Brazil presented. (R. 1865) 209
Breed, E. E. Earthworks in Wisconsin. (R. 1872) 271
BREED, E. E. Pits at Embarrass, Wisconsin. (R. 1877)
Breeding habits of sea-catfish—N. T. Lupron. (P. 1878)
Bremen, government of, exchange system. (R. 1865) 209
Brenndecke, F. Meteorites. (R. 1869)
Brevoortia patronus, note on—S. STEARNS. (P. 1878)
Brevoortia, revision of American species of—G. B. Goode. (P. 1878)
BREWER, T. M. Instructions for collecting and preserving nests and eggs 139
BREWER, T. M. Nests and eggs of North American species of empidonaces. (P. 1879)
BREWER, T. M. North American oölogy. Part 1. Raptores and Fissirostres. 89
BREZINA, A. Explanation of principles of crystallography and crystallophysics. (R. 1872)
Bridges, lectures on roads and—F. ROGERS. (R. 1860; R. 1861) 147, 149
BRIDGMAN, L., vocal sounds of—F. LIEBER12
BRINTON, D. G. Artificial shell deposits in the United States. (R. 1866) 214
British America, account of the Indians of—E. PETITOT. (R. 1865)
British America, Tinneh or Chepewyan Indians of-G. GIBBS. (R. 1866.) 214. 365
British Association, report to, on radiant heat—B. Powell. (R. 1859) 136
British birds, list of faunal publications relative to—E. Cours. (P. 1879) \$\frac{3}{2}\$
British Columbia, Haidah Indians of-J. G. SWAN 267
British Columbia, notes on fishes of-T. H. BEAN. (P. 1881)
British Minister, F. W. A. Bruce, recommending Dr. Berendt. (R. 1865) 39
British Museum, acknowledgment of specimens. (R. 1865)
British Museum, electrotypes of shells granted by. (R. 1868)
British Museum, notes on typical American fishes in—D. S. JORDAN. (P. 1879) 33
British Museum, statistics of—S. F. BAIRD. (R. 1850)
British Provinces of North America, list of libraries, institutions, and societies
in-W. J. Rhees
BROADHEAD, G. C. Prehistoric evidences in Missouri. (R. 1879)
Broca, P. History of Anthropological Society of Paris, 1865 to 1867. (R. 1868)
BROCA, P. Troglodytes, or cave dwellers, of the valley of the Vézère. (R. 1872)
BRODNAX, B. H. Mounds in Moorehouse parish, Louisiana. (R. 1879) 345
Brosmius brosme Americanus, Gill, identity of, with Brosmius brosme (Müller) White—G. B. Goode; T. H. Bean. (P. 1878)
Brown, D. Shell heap in Georgia. (R. 1871)
BROWN-SÉQUARD, C. E. Dual character of the brain. Toner lecture No. II . 291
REVERS I G. Indian anguarings along Green sires wellow (P. 1979)

BRUNOT, F. R. Habits of the beaver. (R. 1878)	275
Brunswick, Maine, meteorological observations—P. CLEAVELAND	204
BRUSH, G. J. Catalogue of meteorites in Yale College. (R. 1868)	224
Brussels, Belgium, Royal Academy of Sciences, Literature, and the Fine Arts. See Prize questions.	
BRYAN, O. N. Antiquities of Charles county, Maryland. (R. 1874)	286
Buch, L. von, memoir of, by M. Flourens. (R. 1862)	150
Budapest, address before Anthropological Congress at—F. F. ROMER. (R. 1876)	392
Building Commission, National Museum, report of, for 1879. (R. 1879) 845,	
Building Commission, National Museum, report of, for 1880. (R. 1880) 442,	434
Building Committee of Regents of S. I., journal and reports of, 1847-1867	829
Building Committee of Smithsonian, report of, on architecture	P
Building materials—R. D. OWEN	P
Building materials, mode of testing—J. HENRY. (R. 1856)	91
Building materials, report on. (R. 1847)	329
Building, new, for museum, memorial to Congress for. (R. 1876)	299
Building, report on use of new hall in—L. Agassiz. (R. 1867)	215
Building-stone collection, circular in reference to—S. F. BAIRD. (P. 1881) 467,	
Building stones, report on—D. D. OWEN. (R. 1847)	•
Buildings, public, acoustics applied to—J. Henry. (R. 1856)	91
Buildings, warming and ventilating—A. MORIN. Part I. (R. 1878) 275,	
Buildings, warming and ventilating—A. MORIN. Part II. (R. 1874) 286,	439
Bulletin National Museum— No. 1. Batrachia and reptilia—E. D. Cope	202
No. 2. Birds of Kerguelen Island—J. H. Kidder; E. Cours	
No. 3. Oölogy, plants, mammals, etc., Kerguelen Island—J. H. Kidder	230
and others	294
No. 4. Birds of Mexico-G. N. LAWRENCE	
No. 5. Catalogue of fishes of Bermudas-G. B. Goods	
No. 6. Classification of animal resources of United States-G. B. Goode.	297
No. 7. Natural history, Hawaiian Islands, etc.—T. H. STREETS	303
No. 8. Index of brachiopoda—W. H. DALL	304
No. 9. Review of Rafinesque's fishes—D. S. JORDAN	305
No. 10. Notes on cottidæ, etc.; synopsis of siluridæ—D. S. JORDAN	
No. 11. Bibliography of Pacific fishes—T. GILL	463
No. 12. Fishes of South Carolina, Georgia, Tennessee; synopsis of catostomide—D. S. Jordan; A. N. Brayton	308
No. 18. Flora of St. Croix and Virgin Islands-H. F. A. EGGERS	313
No. 14. Catalogue Centennial exhibit, animal resources and fisheries— G. B. Goode	326
No. 15. Natural history and ethnology, Arctic America-L. Kumlien.	342
No. 16. (Not published.)	

Bulletin National Museum—Conunued.	
No. 17. Zoölogical position of Texas—E. D. Cope	412
No. 18. Fishery exhibit of United States at Berlin-G. B. Goode	413
No. 19. List of genera in zoölogy and paleontology—S. H. SCUDDER	47 0
No. 20. (Not published.)	
No. 21. Nomenclature of North American birds-R. RIDGWAY	422
No. 22. Guide to flors of Washington, D. CL. F. WARD	444
Bulletin of Philosophical Society of Washington, Vols. I, II, III	423
Bulletin, Smithsonian, No. 1. Vocabulary of the jargon	68
Bulletins of the U.S. National Museum, catalogue and index of	475
Bureau of Ethnology, directory of officers and employés of	
Burial caves—J. Jones	
Burial ground, ancient, in Tompkins Co., N. YD. TROWBRIDGE. (R. 1863)	
Burial, modes of-J. Jones	
Burial of an Indian squaw, California-W. M. KING. (R. 1874)	
BURB, R. T. Ruins in Pims county, Arizona. (R. 1879)	345
Business arrangements of the Smithsonian Institution	
BUTLER, A. P. Report of Senate Judiciary Committee on management of Smithsonian. (R. 1855)	
Buzzard's bay, Mass., occurrence of Belone latimanus in—G. B. Goode. (P. 1878)	
By-laws of the Smithsonian Institution. (R. 1853) 67	
	•
C.	
Cache la Poudre river, Colorado, antiquities on—E. L. Berthoud. (B. 1871) California—	249
account of botanical explorations in-A. Grav. (R. 1849)	21
ancient graves and shell heaps of-P. SCHUMACHER. (R. 1874)	286
	286
cave in Calaveras county—J. D. WHITNEY. (R. 1867)	215
climate of—A. GIBBONS. (R. 1854)	75
climate of M. Logan. (R. 1855)	77
description of Gobiesox rhessodon from San Diego-R. SMITH. (P. 1881)	467
description of new gobioid fish from San Diego—R. SMITH. (P. 1881)	
	332
distribution of tertiary fossils of—W. H. Dall. (P. 1878)fishes of—See D. S. JORDAN; C. H. GILBERT.	832
,	30\$

California—Continued.	
fossil mollusks from later tertiaries of—W. H. Dall. (P. 1878)	882
history and antiquities of Santa Rosa Island—S. Bowers. (R. 1877)	828
Indians of—S. Powers. (R. 1876)	299
list of fishes of, collected by Lt. H. E. Nichols-D. S. JORDAN; C. H. GILBERT. (P. 1881)	467
meteorological observations at Sacramento—F. W. HATCH. (R. 1854)	75
meteorological observations at Sacramento—T. M. Logan. (R. 1854)	75
meteorology of Sacramento-T. M. Logan. (R. 1857)	107
natural history of—T. H. STREETS	303
new genera and species of fishes from coast of—W. N. Lockington. (P. 1879)	388
new pitcher plant from—J. Torrey	61
notes on a collection of fishes from San Diego—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
occurrence of Cremnobates at San Diego-R. SMITH. (P. 1880)	425
occurrence of Productus giganteus in—C. A. WHITE. (P. 1880)	425
plants collected by J. C. Frémont in-J. TORREY	46
pleistocene fossils from-P. P. CARPENTER	252
postpliocene fossils in coast range of—W. H. Dall. (P. 1878)	882
wingless grasshopper of—E. P. Vollum. (R. 1860)	147
California, central, partial list of birds of—L. Belding. (P. 1878)	882
California, Gulf of, lectures on shells of—P. P. CARPENTER. (R. 1859)	110
California peninsula, aboriginal inhabitants of—J. BAEGERT. (R. 1863; R. 1864) 187, 188, 440,	361
California, upper, explorations in, 1860—J. FEILNER. (R. 1864)	188
Camaraphysema, a new type of sponge—J. A. RYDER. (P. 1880)	425
Camel, lecture on the-G. P. Marsh. (R. 1854)	75
CAMERON, H. C. Reminiscences of Joseph Henry	856
CAMPBELL, J. V. Earthquake in Peru, August 18, 1868. (R. 1870)	241
Canada, meteorological system of—J. G. Hodgins. (R. 1865)	209
Canada East, description of observatory at St. Martin, Isle Jesus—C. SMALL- WOOD. (R. 1856)	91
Canada porcupine, occurrence of, in Maryland-O. LUGGER. (P. 1881)	467
Canada porcupine, occurrence of, in West Virginia—G. B. GOODE. (P. 1878)	882
Canada West, ancient Indian remains near Prescott—W. E. Guest. (R. 1856)	91
Canada West, meteorological stations of—J. G. Hodgins. (R. 1858)	109
Canal, Washington, report of committee of Regents on. (R. 1868)	224
CANBY, S. Deposit of Bishop John's library. (R. 1862)	150
Cancerous tumors, structure of—J. J. WOODWARD	266
Candle fish of northwest coast—J. G. SWAN. (P. 1880)	
CANDOLLE, P. DE, memoir of, by M. FLOUBENS. (R. 1859)	110
CANNON, G. L. Antiquities of Colorado. (R. 1877)	828

CANUDAS, A. Earthquakes in Guatemala. (R. 1858)	109
Caoutchoue and gutta percha. (R. 1864)	188
Cape Cod, littoral marine fauna of—R. RATHBUN. (P. 1880)	425
Cape Flattery, Indians of—J. G. SWAN	220
CAPEN, F. L. Meteorological discovery. (R. 1866)	214
Capitol, U. S., memorial exercises at	356
Capitol, U. S., account of marble used in extension of-J. HENRY. (R. 1856)	91
Cara gigantesca of Yzamal, in Yucatan—A. Schott. (R. 1869)	223
Caracas, South America, meteorology of—G. A. Ernst. (R. 1867)	215
Caranx Beani, from Beaufort, N. C., description of—D. S. JORDAN. (P. 1880)	425
Carib language and people—C. H. BERENDT. (R. 1873)	275
Caribbee Islands, ornithology of—F. A. OBER (R. 1878)	34 1
CARLETON, J. H. Excursion to Abó, Quarra, and Gran Quivira, New Mexico. (R. 1854)	75
CARLETON, J. H. Meteorites in Mexico. (R. 1865)	
CARLIN, W. E. Observations of Siredon lichenoides. (P. 1881)	467
Carlsruhe, Scientific Congress of, 1858—F. J. NICKLES. (R. 1860)	147
CARPENTER, P. P.—	
Check-list of the shells of North America	128
Lectures on mollusca, or shell-fish, and their allies. (R. 1860) 147,	152
Lectures on shells of Gulf of California. (R. 1859)	
Mollusks of western North America	
CASELLA, L. Description of meteorological instruments. (R. 1859)	110
CASS, L., Secretary of State. Circular to diplomatic agents relative to L. H. Morgan's research	135
Castings, metallic, of delicate natural objects. (P. 1881)	467
Casts, catalogue of, of Indian boys and girls at Hampton Normal Institute, Vir-	
ginia—R. H. Pratt. (P. 1879)	3 33
Casts, catalogue of, of Indian prisoners at St. Augustine, Florida—R. H. PRATT. (P. 1878)	332
Casts of statues, etc., proposed by W. J. Stone. (R. 1855)	7.
Casts, plaster, methods of making and preserving—A. Pirz. (P. 1881)	467
Caswell, A.—	
Lecture on astronomy. (R. 1858)	109
Meteorological observations at Providence, R. I., 1831-1860	103
Meteorological observations at Providence, R. I., 1831-1876	
Catalogue, general stereotype, of public libraries, report of commission on. (R.	28
1850)	•
Catalogue of— birds of Antigua and Barbuda, collected by F. A. Ober—G. N. Law-	
BENCE. (P. 1878)	332
birds of Dominica, collected by F. A. Ober-G. N. LAWRENCE. (P. 1878)	337

Catalogue of—Continued.	
birds of Grenada, collected by F. A. Ober-G. N. LAWBENCE. (P. 1878)	382
birds of Guadeloupe, collected by F. A. Ober—G. N. LAWRENCE. (P. 1878)	382
birds of Lesser Antilles, collected by F. A. Ober-G. N. LAWRENCE. (P. 1878)	332
birds of Martinique, collected by F. A. Ober—G. N. LAWRENCE. (P. 1878)	882
birds of North America—R. RIDGWAY. (P. 1880) 425,	
birds of St. Vincent, collected by F. A. Ober—G. N. LAWRENCE. (P. 1878)	832
books and memoirs relating to nebulæ and clusters—E. S. Holden	
casts taken by Clark Mills of Indian boys and girls at Hampton, Virginia—R. H. Pratt. (P. 1879)	838
casts taken by Clark Mills of Indian prisoners at St. Augustine, Florida— R. H. Pratt. (P. 1878)	382
collection at International Exhibition, 1876. (R. 1876)	
collection of Japanese cotton fibre presented by Government of Japan. (P. 1881)	
collection of Japanese woods presented by University of Tokio—L. F. WARD. (P. 1881)	467
collection to illustrate animal resources and fisheries of United States exhibited at Philadelphia, 1876—G. B. Goode	
described coleopters of the United States-F. E. Melsheimer	62
described dipters of North America—C. R. OSTEN SACKEN 102,	
described lepidopters of North America-J. G. Morris	
engravings presented to Smithsonian—C. B. KING. (R. 1861)	149
fishes from Pensacola, Florida, collected by S. Stearns-G. B. Goode;	388
fishes obtained in the Gulf of Mexico by J. W. Velie-G. B. GOODE;	
	338
fishes of the Bermudas—G. B. Goode	
fishes of the east coast of North America—T. GILL 2	283
,	888
geological specimens—D. D. Owen. (R. 1854)	75
historical and geographical manuscripts, maps, etc.—L. BERLANDIER. (R. 1854)	75
• ,	271
	880
,	476
meteorites in Yale College—G. J. BRUSH. (R. 1868)	224
minerale with their formulas T. For more	TEE

Catalogue of—Continued.
mollusca added to fauna of southern New England—A. E. VERRILL. (P. 1880)
North American birds—S. F. BAIRDoctavo, 108
North American birds—S. F. BAIRDquarto, 106
North American mammals—S. F. BAIRD 105
North American reptiles. Part I—Serpents—S. F. BAIRD; C. GIRARD. 49
Old World birds in United States National Museum—R. RIDGWAY. (P. 1881)
orthoptera of North America—S. H. SCUDDER
photographic portraits of Indians in gallery of Smithsonian Institution 216
portraits of North American Indians, painted by J. M. Stanley 53
publications of Smithsonian Institution 74, 203, 226, 245, 278, 478
publications of societies in Smithsonian library 73, 85, 117, 179
rocks, minerals, and ores in Michigan—C. T. Jackson. (R. 1854)
rocks, minerals, etc.—J. W. Foster. (R. 1854)
rocks, minerals, etc.—J. D. WHITNEY. (R. 1854)
rocks, minerals, ores, and fossils—J. Locke. (R. 1854)
serpents—S. F. BAIRD; C. GIRARD
trees of the United States-J. G. Cooper. (R. 1858) 109, 35
trochilids in collection of National Museum—R. RIDGWAY. (P. 1880) 42
Catalogue system for libraries—C. C. JEWETT. (R. 1849; R. 1850) 21, 28. 47
Catalogues of libraries, construction of—C. C. JEWETT
Catalytic force, or phenomena of contact—T. L. Phipson. (R. 1862) 15
Catfish, sea, breeding habits of—N. T. LUPTON. (P. 1878)
Catherina Archipelago, Alaska, remains of prehistoric man from—W. H. Dall.
CATLIN, G., notes relative to life and paintings of-J. HENRY. (R. 1872) 2
CATLIN, G., account of Mandan ceremonies by, accuracy of-J. KIPP. (R. 1872)
Catostomidæ, synopsis of—D. S. Jordan 35
CATTIE, S. T. Genitalia of male eels and their sexual characters. (P. 1880) 42
Caulolatilus microps, from Gulf coast of Florida—G. B. Goode; T. H. Bran. (P. 1878)
Causes which limit vegetable species—A. DE CANDOLLE. (R. 1858) 10
Cave dwellers of the valley of the Vézère—P. Broca. (R. 1872) 27
Cave in Calaveras county, California—J. D. WHITNEY. (R. 1867) 21
Cavern, Luray, in Virginia, report of visit to—O. T. Mason and others. (R. 1880)42, 43
Caves in Alaska, remains of prehistoric man from-W. H. Dall314
CAZIN, A. Recent progress in relation to the theory of heat. (R. 1868)
Celts, stone, in West Indies and Africa—G. J. GIBBS. (R. 1877)
Cenozoic rocks, fossils from—C. A. WHITE. (P. 1880)
Centennial awards to Smithsonian Institution. (R. 1876; R. 1878) 299.34

Centennial Exhibition, catalogue of collection of animal resources and fisheries at—G. B. Goods	326
Centennial Exhibition, circular for distribution at	
Centennial Exhibition, plan of proposed Smithsonian exhibit at—S. F. BAIRD. (R. 1875)	298
Centennial Exhibition, report on-S. F. BAIRD. (R. 1876) 299,	307
Centennial Exhibition, savage weapons at—E. H. KNIGHT. (R. 1879) 845,	
Centennial mission to Indians of Western Nevada and California—S. Powers. (R. 1876)	299
Centigrade degrees, tables for conversion of, to Fahrenheit. (R. 1863)	187
Central America—	
circular relative to collections of birds of	168
description of new fishes from—D. S. JORDAN; C. H. GILBERT. (P. 1881)	467
explorations in—C. H. BERENDT. (R. 1867)	215
list of birds of, not in Smithsonian collection	185
rain-fall in—C. A. Schott 222,	353
review of birds of—S. F. BAIRD	181
studies in picture writing of—E. S. HOLDEN. (E. 1879-80)	476
travels in—S. Habel	269
Centrarchidæ, notes on—D. S. JORDAN	806
Centrarchidse, review of family—C. L. McKAY. (P. 1881)	467
Centurus, review of genus—R. RIDGWAY. (P. 1881)	467
Cephaloscyllium laticeps—D. S. JOBDAN; C. H. GILBERT. (P. 1880)	425
Ceratiidæ, note on—T. GILL. (P. 1878)	882
Ceremonies at laying of corner-stone of Smithsonian Institution. (R. 1847). H	, 829
Ceremonies, Mandan, accuracy of Catlin's account of-J. KIPP. (R. 1872)	271
Cessions of land by Indian tribes to United States—C. C. ROYCE. (E. 1879-80)	476
CHACE, G. I. Lecture on oxygen and its combinations. (R. 1855)	77
Chalk found in United States—T. A. CONRAD. (R. 1865)	209
Chamber of Commerce of Bordeaux. Exchange of publications. (R. 1863)	187
Change of Mexican axolotl to amblystoma—A. Weismann. (R. 1877) 328,	401
CHANNING, W. F. Lecture on the American fire-alarm telegraph. (R. 1854)	75
CHANTEE, E. Code of symbols for prehistoric archeology. (R. 1875)	298
Chapala Lake, notes on Dugès' fishes from—D. S. JORDAN. (P. 1879)	888
CHAPPELSMITH, J. Tornado near New Harmony, Indiana, April 80, 1852	59
Characteristics of ancient man in Michigan-H. GILLMAN. (R. 1875) 298,	393
Chart, base, of the United States—C. A. SCHOTT	414
Chart, temperature, of the United States, year—C. A. SCHOTT 381,	388
Charts and maps of America, on a collection of—J. G. Kohl. (R. 1856)	91
Charts of prehistoric archeology, international code of symbols for—G. DE MORTILLET; E. CHANTEE. (R. 1875)	298
Charts, rain, of the United States, summer, winter, year—C. A. Schott	

Charts, temperature, of United States, summer, winter, year—C. A. Schott 387
Charts, wind—J. H. Coffin; S. J. Coffin
CHASE, S. P., biographical notice of, by J. A. GARFIELD. (R. 1878) 275
CHASE, S. P., biographical notice of, by H. HAMLIN. (R. 1873) 275
CHASE, S. P., statement of, relative to the telegraph. (R. 1857) 107, 115
Chattanooga, Tenn., ancient mound near—M. C. READ. (R. 1867) 215
Check-list of—
duplicates of fishes from Pacific coast, distributed by Smithsonian Insti-
tution in 1881—D. S. JORDAN; P. L. JOUY. (P. 1881) 467
fishes, distributed by Smithsonian Institution for National Museum-
T. H. Bran. (P. 1880)
fossils of North America, miocene—F. B. MEEK
invertebrate fossils of North America, cretaceous and jurassic-F. B.
Meek 177
invertebrate fossils of North America, eocene and oligocene—T. A. Con-
RAD 200
North American batrachia and reptilia—E. D. Cope 292
periodicals received by Smithsonian Institution, 1853 Q
plants of Washington and vicinity-L. F. WARD 460
publications of Smithsonian Institution to July, 1872 245
publications of Smithsonian Institution to July, 1874
publications of Smithsonian Institution to July, 1877 301
publications of Smithsonian Institution to July, 1879 344
publications of Smithsonian Institution to December, 1881 437
shells—Lea, Carpenter, and others
Smithsonian publications
Dheck-lists, blank
Chelonia, anatomy and physiology of—S. W. MITCHELL; G. R. MOREHOUSE 159
Chelonia, extinct, of Nebraska—J. LEIDY &
Chelura terebrans, occurrence of, on coast of U. S.—S. I. SMITH. (P. 1879) 333
Chemical analysis of the sun—A. LAUGEL. (R. 1861)
Chemical and physiological investigations of vertebrata—J. Jones82
Chemical arts, recent improvements in—J. C. BOOTH; C. MORFIT
Chemical equivalents of sixty-three elements, table of. (R. 1864) 188
Ohemical formulæ, tables of—F. W. CLABKE
Chemical types, modern theory of—C. M. WETHERILL. (R. 1863) 187
Chemist, report of—F. W. TAYLOB. (B. 1880) 42
Chemistry, lectures on agricultural—S. W. Johnson. (R. 1859)
Chemistry of the earth—T. S. Hunt. (R. 1869) 228, 376
Chemistry, photographic—M. JAMIN. (R. 1867)
Chemistry, recent progress in, 1879 and 1880—G. F. BARKER. (R. 1880). 442, 429
Chepewyan Indians, notes on the—G. GIBBS and others. (R. 1866) 214, 365

Cherbourg, Imperial Society of Natural Sciences of. Prize questions. (R. 1864) Chesapeake Bay, new fish from mouth of—G. B. Goode; T. H. Bean. (P.	188
1881)	467
Chester county, Pennsylvania, birds of—V. Barnard. (R. 1860)	147
Chester county, Pennsylvania, history of—W. Darlington. (R. 1862)	150
Chicago Academy of Sciences. Acknowledging specimens. (R. 1867)	215
China, geological researches in—R. Pumpelly	202
Chinese coal-bearing rocks, fossil plants from—J. S. NEWBERRY	202
Chinese coals, analyses of—J. A. MACDONALD	202
Chinese mineral called yu, study of—S. BLONDEL. (R. 1876)	299
Chinook jargon—H. HALE	161
Chinook jargon-M. LIONNET; B. R. MITCHELL	68
Chinook jargon, bibliography of-G. GIBBS	161
Chinook jargon, dictionary of-G. GIBBS	161
Chinook names of salmon in Columbia river—S. B. SMITH. (P. 1881)	467
Chionis minor, study of-J. H. KIDDER; E. COUES.	294
Chiroid fish (Myriolepis zonifer) from California, new-W. N. LOCKINGTON. (P. 1880)	425
Chirus, remarks on genus—W. N. Lockington. (P. 1880)	425
Chitonidse, new species of, from Panama-P. P. CARPENTER.	252
Chitons of Alaskan and Arctic regions, report on-W. H. Dall. (P. 1878)	882
Chitons from deep waters off eastern coast of U. SW. H. Dall. (P. 1881.)	467
Chitons, on genera of—W. H. Dall. (P. 1881)	467
Chlorospermeze, marine algze of North America-W. H. HARVEY	95
Chorea, inquiries relative to disease known as-S. W. MITCHELL. (R. 1874)	286
CHOSSAT, Dr. Transactions of Geneva Society of Physics and Natural History, 1863 to 1864. (R. 1865)	209
Christiania, Norway, University of, ethnological specimens presented. (R. 1868)	187
Chrysotis, new species of, from Dominica-G. N. LAWRENCE. (P. 1880)	425
Cibola, Coronado's march in search of cities of-J. H. SIMPSON. (R. 1869)	228
Circle, ratio between diameter and circumference of-W. FERREL	233
Circles and spheres, tangencies of—B. ALVORD	80
Circular—	
ancient mining in Lake Superior copper region. (R. 1861)	149
for distribution at the Centennial Exhibition	290
giving directions for constructing lightning-rods-J. Henry	237
in reference to collecting North American shells	176
in reference to history of North American grasshoppers-P. R. UHLER-	163
of inquiries relative to crawfish and other fresh water crustacea	319
of Institute of Rupert's Land. (R. 1861)	149
of instructions for observations of thunder storms—J. HENRY	235
of instructions to observatories, relative to telegraph announcements of	263

Circular—Continued.	
philological—G. Gibbs. (B. 1862)	
relating to collections in archæology and ethnology	
relating to collections of living reptiles	
relative to American archeology—O. T. Mason. (R. 1879) 345,	
relative to collections of birds from Middle and South America	168
relative to food fishes	234
relative to heights	236
relative to observations in Russian America	207
relative to scientific and literary exchanges	324
relative to shipping fresh fish and other animals	
respecting copyright. (R. 1854)	57
respecting new report on libraries. (R. 1854)	75
respecting system of relationship—L. H. MORGAN	138
sent with specimens presented. (R. 1872)	271
to entomologists	178
to officers of Hudson's Bay Company	137
Circulars of United States National Museum:	
No. 1. Plan of organization and regulations of the Museum-G.	
B. GOODE. (P. 1881) 467,	
No. 2. To friends of the Museum-S. F. BAIRD. (P. 1881) 467,	
No. 8. Petroleum collections—S. F. BAIRD. (P. 1881) 467,	
No. 4. Department of insects—S. F. BAIRD. (P. 1881) 467,	
No. 5. Establishment and officers of S. I. and Nat. Mus. (P. 1881) 467,	
No. 6. Materia-medica collection—J. M. FLINT. (P. 1881) 467,	
No. 7. Forms of drugs and medicines—J. M. FLINT. (P. 1881) 467,	, 4 51
No. 8. Collections of drugs for materia-medica section—J. M.	
FLINT. (P. 1881)	
No. 9. Building-stone collection—S. F. BAIRD. (P. 1881) 467,	
No. 10. Letters on the National Museum—B. PHILLIPS. (P. 1881) 467,	454
No. 11. Provisional classification of the food collections—G. B. GOODE. (P. 1881)467,	AEE
No. 12. Classification of the collections to illustrate taxidermy—	733
W. T. Hornaday. (P. 1881) 467,	456
No. 18. Scheme of classification for the collections in the Museum-	
G. B. GOODE. (P. 1881) 467,	457
No. 14. Asking for contributions to library of Museum—S. F. BAIRD. (P. 1881)467,	458
No. 15. Organization and objects of the Museum—G. B. Goode. (P. 1881)467,	459
No. 16. Plans for installation of collections—G. B. Goode. (P. 1881) 467,	
No. 17. Contributions to the Museum and their acknowledgment.	
(P. 1881) 467,	473
No. 18. List of publications of the Museum. (P. 1881) 467,	

Cists, stone, Madison county, Illinois—A. OEHLER. (R. 1879) 846
Citizens of Philadelphia, memorial of, relative to aeronautic voyage of Lowe.
(R. 1860)
Civil Engineers, London Institution of. Prize questions. (R. 1862) 150
Clackamas river, Oregon, notes on fishes from—D. S. Jordan. (P. 1878) 882
CLARK, H. J. Lucernarise and their allies, anatomy, physiology, and relations of 242
CLARK, H. J., memoir of, by A. S. PACKARD242
CLARK, W. M. Antiquities of Tennessee. (R. 1877) 828
CLARKE, FLADGATE, and FINCH. Residuary legacy of Smithson. (R. 1861) 149, 828
CLARKE, F. W.—
Bibliography of atomic volume, specific gravity, &c 255
Recalculation of atomic weights41
Specific gravities, boiling and melting points, chemical formulæ, tables 255
Specific gravities, &c. Supplementary tables 288
Specific heat tables for solids and liquids 276
Tables of expansion by heat, for solids and liquids 289
Classification and arrangement of materia medica collection—J. M. FLINT.
(P. 1881) 467, 450
Classification and synopsis of trochilidæ—D. G. ELLIOT
Classification of—
birds, review of the-W. LILLJEBORG. (R. 1865) 209, 364
books, on the—J. P. LESLEY. (R. 1862) 150
clouds-A. Pory. (R. 1870) 244
coleoptera. Part I-J. L. LE CONTE
coleoptera. Part II—J. L. LE CONTE 265
collection to illustrate animal resources of United States—G. B. Goode 297
collections of U.S. National Museum—G. B. Goode. (P. 1881) 467, 457
collections of United States National Museum to illustrate taxidermy—
W. T. Hobnaday. (P. 1881) 467, 456
food collections—G. B. Goode. (P. 1881) 467, 455
forms in which drugs and medicines appear and are administered—J. M.
FLINT. (P. 1881)
insects from embryological data—L. Agassız 16
· -
methods of capture and utilization of animals—G. B. Goods
Classified list of meteorological publications and articles in periodicals received
by Smithsonian Institution. (R. 1873) 275
Classified record of meteorological material preserved in Smithsonian. (R.
1874) 286
Classified record of monthly meteorological reports. (R. 1878) 275
CLEAVELAND, P. Meteorological observations at Brunswick, Maine 204
CLEAVELAND, P., notice of, by J. Henry. (R. 1859)
CLEMENS, B. Instructions for collecting hymenopters. (R. 1858) 109

CLEMENS, B. Instructions for collecting lepidoptera. (R. 1858)	109
CLEMENS, B. On lepidoptera	
Climate and history of New Mexico-T. A. McParlin. (B. 1877) 823,	396
Climate of Alaska—H. M. BANNISTER. (R. 1866)	
Climate of California—T. M. LOGAN. (R. 1855)	77
Climate of Colorado—J. Evans. (R. 1865)	200
Climate of Kansas—R. S. Elliott. (R. 1870)	244
Climate of Kelley's Island, Ohio-G. C. Huntington. (R. 1866)	
Climate of San Francisco—H. GIBBONS. (R. 1854)	•;;
Climate—See Meteorology.	
Climatic influence of forests—M. BECQUEREL. (R. 1869)	23
CLINGMAN, T. L. Topography of Black Mountain, N. C. (R. 1855)	77
Clinton, N. Y.—See Hamilton College.	
Closing of Kennebec river, Maine-R. H. GARDINER. (R. 1858)	103
Cloud-bursts-W. J. YOUNG. (R. 1867)	215
Cloud, dispersion of, by an electrical discharge—D. W. NAILL. (R. 1858)	109
Clouds, electricity of induction in—F. ZANTEDESCHI. (R. 1870)	
Clouds, formation of, over Gulf Stream-H. M. BANNISTER. (R. 1866)	
Clouds, new classification of—A. Porv. (R. 1870)	
Clouds, nomenclature of	
Clupea tyrannus of Latrobe-G. B. Goode. (P. 1878)	200
CLUSS, A. Report of architect. (R. 1867)	
Cluss & Schulze, report of, for 1879. (R. 1879) 345.	409
Cluss & Schulze, report of, for 1880. (R. 1880) 442,	434
CLYMER, H. Memorial of J. Henry	356
CLYMER, H., and others. Report on Museum. (R. 1876)	
Coal, Chinese and Japanese, analyses of—J. A. MACDONALD	
Coal, Chinese, fossil plants in—J. S. NEWBERRY	240
Coal, lecture on—J. LE CONTE. (R. 1857)	167
Coal oils, explosibility of—Z. Allen. (R. 1861)	145
Coast Survey expedition of 1866, to determine transatlantic longitude—B. A. GOULD	2:3
Coast Survey soundings on Atlantic coast, examination of-J. W. BAILEY	
Cobalt (ammonia-) bases—W. GIBBS; F. A. GENTH	88
COCHRANE, J. Antiquities of Mason county, Illinois. (R. 1877)	327
CODD, J. A. Acknowledgment for books. (R. 1861)	149
Code of symbols for charts of archeology—G. DE MORTILLET; E. CHANTEE.	
(R. 1875)	205
Coppin, J. H.—	
Orbit and phenomena of meteoric fire ball	22 1
Psychrometrical tables	87
Storms of 1859	152

OF SMITHSONIAN PUBLICATIONS.

,
COPPIN, J. H.—Continued.
Winds of the globe 268
Winds of the Northern Hemisphere52
COFFIN, S. J. Tables and maps of winds of the globe 268
COGSWELL, J. G. Report on Jewett's general stereotype catalogue of public libraries
Coins, assay of gold and silver, at United States mint-J. Pollock. (R. 1868) 224
Coins, table of foreign, gold and silver. (R. 1868)224
Colding, A. Meteorology. (R. 1877)
Colding, A. Nature of currents of air. (R. 1877) 828, 898
COLDING, A. Theory of relation between pressure and the velocity of the wind. (R. 1877)828, 898
COLDING, A. Whirlwind at St. Thomas in August, 1871. (R. 1877) 828, 398
Coleoptera—
catalogue of the described, of the United States—F. E. MELSHEIMER 62
classification of. Part I-J. L. Le Conte 136
classification of. Part II—J. L. LE CONTE 265
instructions for collecting—J. L. LE CONTE. (R. 1858) 109
new species of North American. Part I-J. L. LE CONTE 167
new species of North American. Part II.—J. L. LE CONTE 264
of Arctic America—S. H. Scudder 842
of Kansas and eastern New Mexico—J. L. LE CONTE 126
of North America, list of-J. L. LE CONTE 140
Colima, eruption of the volcano of—C. SARTORIUS. (R. 1869)
Collaborators of Smithsonian Institution, directory of
Collecting—
and preserving diptera, directions for—C. R. OSTEN SACKEN
and preserving fish, directions for—T. H. BEAN. (P. 1881) 467, 464
and preserving plants, directions for—L. F. WARD 460
insects, directions for—A. S. PACKARD 261
insects, instructions for—S. F. BAIRD. (R. 1858)
land and fresh-water shells, instructions for—J. Lewis. (R. 1866) 214, 363
microscopic organisms, directions for 68
myriapods, phalangidse, etc.—H. C. Wood. (R. 1866) 214
nests and eggs, instructions for—T. M. BREWER
nests and eggs of birds—S. F. BAIRD. (R. 1858) 109
North American shells, circular in reference to 176
specimens, directions for—S. F. BAIRD. (R. 1856)
specimens of diatomaces, directions for—A. M. EDWARDS 366
specimens of natural history—S. F. BAIRD 34
Collection—
and preservation of marine invertebrates—W. STIMPSON 84
archeological, of United States National Museum—C. RAU 287

Collection—	
of historical documents in Guatemala—C. H. BERENDT. (R. 1876)	299
of meteorological tables—A. Guyor31, 1	L53
to illustrate animal resources and fisheries of United States, exhibited at Philadelphia, in 1876, catalogue of—G. B. Goode	326
to illustrate animal resources of United States, classification of—G. B. GOODE	
Collections—	
additions to. (See each annual report.)	
furnished by explorations from 1838 to 1877—S. F. BAIRD. (B. 1877)	323
in archeology and ethnology, circular relative to-J. HENRY	
made by Lt. G. K. Warren, report on-F. B. MEEK; F. V. HAYDEN_	
of living reptiles, circular relating to	320
presented to United States by foreign commissioners at Centennial Exhibition, list of. (R. 1876)	299
to illustrate art of taxidermy-W. T. HORNADAY. (P. 1881) 467,	456
Collections—See Ethnographical.	
Collectors of customs to receive and transmit specimens to the Smithsonian	84
Collectors of drugs, memoranda for-J. M. FLINT. (P. 1881) 467,	452
College—See Girard, Hamilton, Liberia.	
Colleges in United States in correspondence with Smithsonian Inst., list of	238
COLLINS, Colonel. Meteorology of the Green river country. (R. 1871)	249
Colonia Tovar, Venezuela, meteorology and ethnology of—A. FENDLER. (R. 1857)	107
Colonia Toyar, Venezuela, meteorology of-A. FENDLER. (R. 1866)	214
Colorado—	
ancient remains in—E. L. BERTHOUD. (R. 1867)	215
climate of—J. Evans. (R. 1865)	209
cretaceous fossils from—C. A. WHITE. (P. 1881)	467
descriptions of new invertebrate fossils from-C. A. WHITE. (P. 1880)_	425
heights of mountains in-G. Engelmann. (R. 1862)	150
Jefferson and Clear Creek Cos., antiquities—G. L. CANNON. (R. 1877)	323
Weld county, antiquities of—E. L. BERTHOUD. (R. 1871)	249
Color-blindness-J. Henry. (R. 1877)	323
Color-blindness—A. Moigno. (R. 1866)	214
Color-blindness in its relations to accidents by rail and sea—F. Holmgren. (R. 1877)823,	3 99
Colored bead from mound in Florida—A. M. HARRISON. (R. 1877)	323
Colors, accidental or subjective—A. Moigno. (R. 1866)	214
	467
Columbia river, Salmonida of -C. Bendire. (P. 1881)	467
Combustion of gun-cotton and gunpowder, products of—Lieut. von Karolyi; R F Craig (R 1864)	188

OF SMITHSONIAN PUBLICATIONS.	163
met, investigations of Biela's—J. S. Hubbard. (R. 1862)	
mets, directions for telegrams of discovery of	. 263
MFORT, A. J. Indian mounds near Fort Wadsworth, Dakota. (R. 1871)	. 249
mmemoration of J. Henry	. 356
mmission, National Museum—See Museum.	
mmissioner of Patents. Meteorological observations, Vol. I	. 157
mmissioner of Patents. Meteorological observations, Vol. II	. 182
mmissioners for Foreign Missions—See American Board.	
mmissioners, report of, on plan of forming general stereotype catalogue for public libraries in United States. (R. 1850)	
mmittee of American Association for Advancement of Science. Report or meteorology. (R. 1851)	ı
mmittee of Regents—	
report on architecture	. P
report on fire at Smithsonian—R. WALLACH; J. HENRY. (R. 1864) 18	8, 829
report on income—J. A. PEARCE. (R. 1858)	7, 829
report on income—J. MEACHAM. (R. 1858)	
report on Museum—A. GRAY; A. A. SARGENT; H. CLYMER. (R. 1876) 26	9, 329
report on organization 1	B, 828
report on organization, (first)	L , 828
mmittee—See Report.	
mparative action of dry heat and sulphurous acid upon putrefactive bacteria (P. 1881)	
mparative vocabulary—A. GALLATIN	
mparative vocabulary, English, Spanish, French, Latin—G. GIBBS	
mpasses, deviations of, on iron-clad—W. HARKNESS	
nchology, North American, bibliography of. Part 1. American authors-	
W. G. BINNEY	. 142
w. G. Binner.	
enchology—See Mollusks, Shells.	•
encise, archeological researches at—F. Trovon. (R. 1861)	149
NDORCET, biography of, by F. Arago. (R. 1878)	. 841
one-in-cone—H. Poole. (R. 1868)	187
ongress, International Anthropological, address at—F. F. ROMEE. (R. 1876) 296	, 392
ongress, International Archæological, Antwerp, 1866. (R. 1866)	214
ongress of Carlsruhe, 1858, scientific—F. J. NICKLÉS. (R. 1860)	147
ongress, United States—	
act of, accepting bequest, July 1, 1886. (R. 1858) 67, 8	28, N
act of, to establish Smithsonian Institution. (R. 1853) 67, 829, 391, B,	N, O
act of, to establish Smithsonian Institution, construction of-J. McP.	
Berrien. (R. 1858)	67

.

.

Congress, United States—Continued.	
act of, to receive residuary legacy and increase fund. (R. 1866) 2	4, 325
act of, to transfer Smithsonian library to Library of Congress. (R. 1865) 2	09, 32
acts of, (45th Congress,) relative to Smithsonian. (R. 1878)	. 34
acts of, (45th and 46th Congresses,) relative to Smithsonian, 1878-1880 (R. 1879)	
memorial of J. Henry, published by order of	
memorial of Regents to, asking appropriations for Museum. (R. 1866 R. 1867)214, 2:	
memorial of Regents to, relative to new Museum building. (R. 1876).	. 29
memorial of Regents to, relative to Smithson fund. (R. 1850)	. 2
proceedings of, relative to appropriations for Museum	. 32
proceedings of, relative to monument of J. Henry	. 35
proceedings of, relative to new Museum building	
proceedings of, relative to Smithson bequest	. 32
proceedings of, relative to Smithsonian Institution, 1835-1877	
Connecticut Academy of Arts and Sciences, planisphere of the heavens by	. 359
Connecticut, ancient implement of wood from—E. W. Ellsworth. (B. 1876	299
Conosaurus, memoir on—R. W. GIBBES	. 14
CONRAD, C. M., Secretary of War. Authority to officers of Quartermaster's Department toreceive and transmit specimens to Smithsonian	
CONRAD, T. A. Chalk found in the United States. (R. 1865)	
CONBAD, T. A. Check-list of invertebrate fossils of North America. Eccent	
Consanguinity and affinity of the human family, circular-L. H. MOBGAN	
Consanguinity and affinity of the human family, systems of-L. H. MORGAN	218
Considerations on electricity. (R. 1867)	215
Constantinople, account of hail storm at-Com. PORTER. (R. 1870)	244
Constants of Nature—	
Part 1. Specific gravities, boiling and melting points, and chemical formulæ—F. W. CLARKE	
Part II. Table of specific heats for solids and liquids—F. W. CLARKE.	276
Part III. Tables of expansion by heat for solids and liquids—F. W. CLARKE	
Part IV. Atomic weight determinations-G. F. BECKEB.	358
Part v. Recalculation of atomic weights-F. W. CLARKE	
First supplement to Part 1. Specific gravities, boiling points and melting points—F. W. CLARKE	
Constants of nature—J. LE CONTE. (R. 1878)	
Constants of nature and art, tables of—C. BABBAGE. (R. 1856)	91
Construction of act establishing Institution—J. McP. Berrien. (B. 1858)	67
Construction of catalogues of libraries—C. C. JEWETT	47
Construction of silvaned class telescope II Dr. Burn	180

Contact, catalytic force or phenomena of—T. L. PHIPSON. (R. 1862)	150
CONTAXARI, Miss E. B., present of album from Greece. (R. 1857) 107	
Contrast of colors—A. Moigno. (R. 1866)	
Contribution to history of fresh-water algs of North America—H. C. Wood	
Contributions to Annals of Philosophy—J. Smithson. (R. 1858)	
Contributions to history of marine algo of North America—See W. H. Harvey	
Contributions to Knowledge, vols. I-XXIII 2, 26, 38, 39, 55, 56, 76, 78, 92 111, 112, 151, 184, 206, 211, 229, 246, 272, 284, 285, 340,	, 99, 346
Contributions to Knowledge, catalogue of, and index	
Contributions to National Museum, and their acknowledgment. (P. 1881.) 467,	
Contributions to natural history of—	770
Arctic America—L. Kumlien	342
Fanning Islands—T. H. STREETS	
fresh-water fishes of North America—C. GIRARD	
Hawaiian Islands—T. H. Streets	
Kerguelen Island. I. Ornithology—E. Coues	
•	
Kerguelen Island. 11. Oölogy, botany, etc.—J. H. KIDDER and others Lower California—T. H. STREETS	308
	5U 5
Contributions to North American ichthyology—	
Part I. Review Rafinesque's memoirs—D. S. JORDAN	
Part II. Notes on Cottidæ; sysopsis Siluridæ—D. S. JORDAN	
Part III. Fishes Alleghany region; synopsis Catostomida—D. S. Jor-	
DAN; A. W. BRAYTON	
Contributions to physical geography of United States—C. Eller, Jr	
Contributions toward monograph of the pandorids—P. P. CARPENTER	
Converging series expressing ratio between diameter and circumference of circle—W. Ferrel	
COOK, C. Manufacture of porpoise oil. (P. 1878)	882
Cooper, J. G.—	
Distribution of forests and trees of North America, and catalogue of native trees of United States. (R. 1858) 109,	
Forests and trees of Florida and Mexican boundary. (R. 1860)	
Migrations and nesting habits of west coast birds. (P. 1879)	388
COPE, E. D.—	
Genera and species of rattlesnakes	185
North America batrachia and reptilia	292
Zoölogical position of Texas	
Coppée, H. Report of Committee on Smithsonian Museum. (R. 1874)	286
Copper and iron in salt water, preservation of—A. E. BECQUEREL. (R. 1864.)	188
Copper region, circular, ancient mining in Lake Superior-J. HANN. (R.	
1861)	149
Coptic language, introduction to the study of the—M. KABIS. (R. 1867)	215
Copyright books from 1846-1849, list of—C. C. JEWETT. (R. 1850)	28

Copyright system and Smithsonian library—C. C. JEWETT. (R. 1851) 51
Copyrights, circular respecting—J. HENRY. (R. 1854)75
Corbiculadæ, American, monograph of—T. PRIME145
Corcoran Art Gallery, list of deposits in, by Smithsonian Institution. (R. 1874) 286
Corcoran Art Gallery, report of Committee of Regents on. (R. 1872) 271
CORCORAN, W. W. Deed of gift establishing Art Gallery. (R. 1872) 271, 329
Cordoba, Argentine Republic, account of astronomical observatory at—B. A. GOULD. (R. 1878)
Corner-stone of Smithsonian building, address at laying-G. M. Dallas D, 329
Corner-stone of Smithsonian building, ceremonies at laying of. (R. 1847) H, 329
Coronado's march in search of the "seven cities of Cibola"—J. H. SIMPSON.
(R. 1869) 228 Correspondence—
abstracts of anthropological—O. T. Mason. (R. 1880) 42
extracts from. (R. 1855, 1858-1868, 1865-1867, 1878) 77, 109, 110, 147, 149,
150, 187, 209, 214, 215, 275
meteorological—E. Foreman. (R. 1851) 31
relative to publication of memoir by Squier and Davis
relative to Smithson and his bequest 228
summary of anthropological. (R. 1879) 345
Correspondents—
3
domestic, list of. (R. 1858; R. 1872) 67, 271, 69, 238
foreign, list of 64, 154, 225, 243, 309
foreign, list of64, 154, 225, 243, 309
foreign, list of 64, 154, 225, 243, 309 foreign, systematic index of 257
foreign, list of 64, 154, 225, 243, 309 foreign, systematic index of 257 Costa Rica—
foreign, list of

OF SMITHSONIAN PUBLICATIONS.	167
CRAIG, B. F. Nitrification. (R. 1861)	149
CRAIG, B. F.; Lieut. VON KABOLYI. Products of combustion of gun-cotton and gunpowder. (R. 1864)	188
Craig flounder of Europe (Glyptocephalus cynoglossus) on coast of North America—G. B. Goode; T. H. Bean. (P. 1878)	882
Crania Helvetica—F. Troyon. (R. 1864)	188
Craw-fish, circular of inquiries relative to	319
Creation, visible, lecture on vastness of—S. ALEXANDEB. (R. 1857)	107
Cree Indians, system of relationship of—E. A. WATKINS. (R. 1862)	150
Cremnobates, occurrence of, at San Diego, California—R. SMITH. (P. 1880)	425
CRESSON, J. C. Lightning discharges. (B. 1867)	216
Cretaceous fossils, check-list of—F. B. MREK	
Cretaceous fossils from Arkansas and Colorado—C. A. WHITE. (P. 1881)	467
Cretaceous reptiles of United States—J. Leidy. (R. 1864)	
·	
Cricetodipus parvus, a rare rodent—F. W. TRUE. (P. 1881)	
Criocardium, note on—C. A. WHITE. (P. 1879)	
Criticisms of Dr. J. Hann, replies to the—W. FERREL. (R. 1877) 828	
CROOK, G. Indian mode of making arrow-heads and obtaining fire. (R. 1871.)	
Crozet flora—J. H. KIDDER	294
CRUMMELL, A. Facts respecting Liberia College. (R. 1861)	149
Crust of the earth, revolutions of the—G. PILAR. (B. 1876)	299
Crustacea—	
fresh water, circular of inquiries relative to	319
notice of, dredged off south coast of New England by Fish Commission— S. I. Smith. (P. 1880)	425
notice of new species of Willemssia group of-S. I. SMITH. (P. 1879.)	888
of Hawaiian and Fanning Islands and California—T. A. STREETS	808
Crustacean destructive to submarine timber on coast of United States—S. I. SMITH. (P. 1879)	888
Crustaceans of Kerguelen Island—S. I. SMITH	294
Crustaceans, voices of—G. B. Goode. (P. 1878)	382
Cryolite of Greenland—Lewis; Quale. (R. 1866)	214
Cryptogamous plants, present state of knowledge of—W. REICHARDT. (R. 1871)	249
Crystalline rocks, metamorphism and the formation of—G. A. DAUBRÉE. (R.	
1861)	149
Crystallography, explanation of principles of—A. Brezina. (R. 1872) 271,	
Crystallophysics, explanation of principles of—A. Brezina. (R. 1872) 271,	386
Culbertson, T. A.—	
Expedition to the Mauvaises Terres and Upper Missouri. (R. 1850)	28
Indian tribes in Upper Missouri. (R. 1850)	28
Sioux of the Upper Missouri. (R. 1850)	28
Culture, fish—See Fish.	
CUNARD, E. Free freight between United States and England. (R. 1859.) 110	, 829

Dakota-

CUNNINGHAM, K. M.; GAINES, A. S. Shell-heaps on Mobile river. (R. 1877.)	223
Currents, electrical, of the earth—C. MATTEUCCI. (R. 1867)	215
Currents of air—	
and aerial navigation—J. HENBY. (R. 1860)	147
law of variation of temperature in ascending-J. HANN. (R. 1877.) 323,	898
law of variation of temperature in ascending moist—L. SOHNCKE. (R. 1877)	, ၁५६
nature of—A. Colding. (R. 1877)	, 396
relation between barometric variations and—M PESLIN. (R. 1877). 323,	298
Curtiss, A. H., notes on collection of fishes made by-D. S. JORDAN. (P. 1880)	425
CUSHING, F. H. Antiquities of Orleans county, New York. (B. 1874)	286
Customs, collectors of, to receive and transmit specimens to Smithsonian	34
CUTTS, J. B. Ancient relics in northwestern Iowa. (R. 1872)	271
CUVIER, Baron G.—	
history of the works of, by M. FLOURENS. (R. 1868)	224
memoir of, by M. Flourens. (R. 1868)	224
Memoir of René Just Hauy. (R. 1860)	147
Memoir of Priestley. (R. 1858)	109
Cybium, notes on American species of—F. Porv. (P. 1878)	
Cyclone in the Indian Ocean—N. PIKE. (R. 1867)	
Cyclophoridæ—W. G. BINNEY	144
Cymatogaster rosaceus—D. S. Jordan; C. H. Gilbert. (P. 1880)	425
Cyprinidse, notes on—D. S. JORDAN	306
Cyprinoid fishes of San Francisco market—D. S. JORDAN. (P. 1880).	425
D.	
DAA, L. K. Ethnological museum of Norway. (R. 1862)	150
Dacentrus, note on—D. S. JORDAN. (P. 1880)	425
DA COSTA, J. M. On strain and over-action of the heart. Toner lecture No. 111.	279
DAHLBERG, R. N. and C. Ancient pottery at Pittsburg, on Des Moines river. (R. 1879)	84 5

Dakota, Lincoln county, haystack mound in—A. BARRANDT. (R. 1872) 271
Dakota or Sioux Indians—A. G. Brackett. (R. 1876) 299
Dakota or Sioux Indians, language of—F. L. O. Roehrig. (R. 1871)——— 249, 878
Dall, W. H.—
Distribution of California tertiary fossils. (P. 1878) 882
(
Index of Brachiopoda
Limpets and chitons of Alaskan and Arctic regions. (P. 1878)
Mollusks of Arctic America
Mollusks of Kerguelen Island 294
New mollusks from Alaska in National Museum. (P. 1878)
New species of shells from California in National Museum. (P. 1878) 382
Note on shells from Costa Rica kitchen-midden collected by Drs. Flint
and Bransford. (P. 1878) 382
On genera of chitons. (P. 1881)467
Postpliocene fossils in coast range of California. (P. 1878) 382
Remains of later prehistoric man from caves in Alaska 318
Dallas, G. M. Address at laying corner-stone of Smithsonian building D, 829
Dalton, J. C. Origin and propagation of disease. (R. 1878)
Daltonism, etc.—A. Moigno. (R. 1866) 214
Danilsen, A. F. Mound in East Tennessee. (R. 1868) 187
Danish Royal Society of Arts and Sciences. Prize questions. (R. 1862, 1865,
1867) 150, 209, 215
Darlington, W. History of Chester county, Pennsylvania. (R. 1862) 150
Darlingtonia Californica—J. Torrey
Darwin, C. Queries about expression for anthropological inquiry. (R. 1867) 215
DAUBRÉE, G. A. Metamorphism and formation of crystalline rocks. (R.
1861) 149
DAUBRÉE, G. A. Synthetic experiments relative to meteorites. (R. 1868) 224
Davis, A. C. Antiquities of Isle Royale, Lake Superior. (R. 1874) 286
DAVIS, C. H. Law of deposit of flood-tide 33
DAVIS, C. H. Occultations in United States, 1852 29
DAVIS, E. H. On ethnological research. (R. 1866) 214
DAVIS, E. H.; SQUIER, E. G. Ancient monuments of. Mississippi valley 1
DAYTON, E. A. Explorations in Tennessee. (R. 1870) 244
Deaf mutes, sign language of—G. MALLERY. (E. 1879-80) 2
DEAN, C. K. Mound in Wisconsin. (R. 1872) 271
DEAN, C. W. Lightning discharges. (R. 1867) 215
DEAN, J. Gray substance of medulla oblongata173
DE BEAUMONT, E.—
Memoir of C. F. Beautemps-Beaupré. (R. 1868) 187

DE BEAUMONT, E.—Continued.
Memoir of Auguste Bravais. (R, 1869) 22
Memoir of Legendre. (R. 1867) 21
Memoir of Oersted. (R. 1868) 22
DE BLAINVILLE, D., memoir of, by M. FLOURENS. (R. 1865) 20
DE CANDOLLE, A.—
Causes which limit vegetable species towards the north. (R. 1868) 100
Probable future of the human race. (R. 1875) 29
Report on transactions of Society of Physics and Natural History of Geneva for 1862. (R. 1864)
Report on transactions of Society of Physics and Natural History of Geneva, 1878-1874. (R. 1875)
DE CANDOLLE, A.; GRAY, J. E. On a dominant language for science. (R. 1874)
DE CANDOLLE, P., memoir of, by M. FLOUBENS. (R. 1859)
Deduction and induction—J. VON LIEBIG. (R. 1870) 24
Deed of foundation of Toner lectures—J. M. TONER. (R. 1872) 271, 32
Deed of foundation of Tyndall scientific fund-J. TYNDALL. (R. 1872) 271, \$2
Deed of gift of Art Gallery-W. W. CORCORAN. (R. 1872) 271, 32
DE FOREST, E. L. Methods of interpolation. Part I. (R. 1871) 24
DE FOREST, E. L. Additions to memoir on methods of interpolation. Part II. (R. 1878)
DE HART, J. N. Mounds and osteology of mound-builders of Wisconsin. (R. 1877)
DE IRISARRI, Guatemalan minister, recommendation of Dr. Berendt by. (R. 1865)
DELAFIELD, R. Report of Committee of Regents on Washington canal. (R. 1868)
DELAMBRE, J. B. J., memoir of, by J. FOURIER. (R. 1864)
DE LA RIVE, A. A.—
Eulogy on, by J. B. Dumas. (B. 1874)
Michael Faraday, his life and works. (R. 1867) 21
Phenomena accompanying propagation of electricity in rarefled elastic fluids. (R. 1868)
Report on transactions of Society of Physics and Natural History of Geneva, 1858-1859, 1872-1878. (R. 1859; R. 1874)
DE LA RIVE, A. A.; LEMSTRÖM, S. Electricity of the atmosphere and the aurora borealis. (R. 1874)
DE LA RUE, W. Abbreviations used in England in 1867. (R. 1867) 213
DELAUNAY, C. Essay on the velocity of light. (R. 1864) 188. 354
Delolepis, new genus of fishes, description of—T. H. BEAN. (P. 1881) 457
Denmark, preservation of antiquities in—J. J. A. Worsaae. (R. 1879) 345
DENNIS, W. C. Evaporation in Florida. (R. 1866)
DENNIS, W. C. Fresh water in the ocean. (R. 1866) 214

Deposit of—
arrow-heads near Fishkill, New York-E. M. SHEPARD. (R. 1877) 323
articles by Smithsonian in Corcoran Art Gallery. (R. 1874) 286
Beaufort Library—E. M. STANTON. (R. 1862) 150
Bishop Johns' Library—E. CANBY. (R. 1862)
flint implements in Illinois—J. F. SNYDER. (R. 1876) 299
flint implements in Southern Illinois-C. RAU. (R. 1868) 224, 440, 370
flood-tide, law of—C. H. DAVIS33
Deposit, shell, in New Jersey—C. RAU. (R. 1864) 188, 440, 362
Deposits—See Shell deposits.
DE PRADOS, Baron. Eclipse of the sun, April 25, 1865. (R. 1864) 188
DE SAUSSURE, H.—
Electric resonance of mountains. (R. 1868) 224
Hymenopters. (R. 1862) 150
Report on transactions of Society of Physics and Natural History of
Genevá, 1870–1871. (R. 1871)249
Synopsis of Vespidæ, (American wasps) 254
DE SCHLAGINTWEIT, H. Ethnographical collections. (R. 1862)
Description of—
a new fish, Apogon pandionis—G. B. GOODE; T. H. BEAN. (P. 1881) 467
a new fly-catcher and new petrel from Sandwich Islands—R. RIDGWAY. (P. 1881)
a new genus and species of Cottidæ—W. N. LOCKINGTON. (P. 1881) 467
а new gobioid fish (<i>Othonops cos</i>) from San Diego, California—R. Sмітн. (P. 1881) 467
a new owl from Porto Rico-R. Bidgway. (P. 1881) 467
birds—See Proceedings National Museum.
chitonidæ and acmæidæ from Panama—P. P. CARPENTER 252
country and people of Yoruba, Africa-T. J. Bowen 98
fishes-See Proceedings of National Museum.
fossil plants from China—J. S. NEWBERRY 202
Gobiesox rhessodon from San Diego, California—R. SMITH. (P. 1881)., 467
human skull from Rock Bluff, Illinois-J. A. MEIGS, (R. 1867) 215
magnetic observatory at Smithsonian. (R. 1859) 110
meteorological instruments—L. Casella. (R. 1859) 110
new fishes—T. H. BEAN. (P. 1881) 467
new fishes-See D. S. JORDAN; C. H. GILBERT.
new species of Centrarchidæ—C. L. McKAY. (P. 1881) 467
new subspecies of Loxigilla from island of St. Christopher, West Indies—G. N. LAWRENCE. (P. 1881)467
new thrushes—R. RIDGWAY. (P. 1881)467
observatories at Dorpat and Poulkova—C. ABBE. (R. 1867) 215, 369

Description of—Continued.
observatory at St. Martin, Isle Jesus, Canada East—C. SMALLWOOD. (R.
1856) 91
plants collected by J. C. Frémont in California—J. Torrey 46
remains of extinct mammalia and chelonia from Nebraska—J. LEIDY 58
Descriptions of—
ancient works in Ohio—C. WHITTLESEY 37
several new races of American birds—R. Ridgway. (P. 1881) 467
Smithsonian anemometer—J. HENRY. (R. 1860)
Squalius aliciæ from Utah Lake—P. L. Jouy. (P. 1881) 467
two new races of Myadestes obscurus—L. STEJNEGER. (P. 1881) 457
Desiderata, special, among North American birds, list of—R. RIDGWAY. (P. 1881)
Des Moines river, ancient pottery from-R. N. and C. Dahlberg. (R. 1879.) 347
Des Moines valley, mounds in—S. B. Evans. (R. 1879) 345
DESOR, E. Palafittes or lacustrian constructions, Lake Neuchâtel. (R. 1865.) 209, 360
Destruction of fish in Gulf of Mexico-J. Y. PORTER. (P. 1881)
Destruction of fish—See also Fish.
Destructive effect of iron rust. (R. 1861)
DEVEREUX, J. H. Ancient pottery from Arkansas. (R. 1872)
DEVEREUX, J. H. Catalogue of Indian relics presented to Smithsonian Insti-
tution. (R. 1872)
DEWBY, C.—
Best hours for temperature observations. (R. 1860)14:
Best hours to find mean temperature. (R. 1857)lot
life of, by M. B. Anderson. (R. 1870)244
The winds. (R. 1866)214
Diagnoses des mollusques nouveaux provenant de Californie et faisant partie du Musée de l'Institution Smithsonienne—P. P. CARPENTER
Diagnoses of new forms of mollusks collected at Cape St. Lucas, Lower Cali-
fornia—P. P. CARPENTER
Diagnoses of new mollusks-
collected on west tropical shores of North America—P. P. CARPENTER 23
from Reigen Mazatlan collection—P. P. CARPENTER
from Vancouver district—P. P. CARPENTER.
from west coast of North America—P. P. CARPENTER 2:
from west tropical region of North America—P. P. CARPENTER 22
Diamond and other precious stones—J. Babiner. (R. 1870) 244, 377
Diary of an excursion in New Mexice—J. H. Carleton. (R. 1854)
Diatomacea—J. W. Bailey 23, 6:
Diatomacea, instructions for collecting, preserving, and transporting—A. M.
EDWARDS
Digerment A R Reuntion of volceno in Niceregue (R 1867)

BERENDT. (R. 1878)	275
° Oregon—G. Gybbs 1	
	40
	98
N. W. TURNER	98
. ag Smithsonian Institution—J. HENBY_ C,	
	150
` ,	214
ger with increasing altitude—J. HANN. (R. 1877) 828,	
-	298
•	188
The draft to j Totality to Marigan & Totality	.00
i. of North America, catalogue of-R. OSTEN SACKEN 102, 2	70
ons for collecting and preserving—R. OSTEN SACKEN	102
structions for collecting-H. LOEW; R. OSTEN SACKEN. (R. 1858)	109
of Arctic America—S. H. SCUDDER	842
ra of North America, monographs of-	
Parts I-III—H. LORW	56
Part IV-R. OSTEN SACKEN 2	19
Directions for—	
auroral observations—J. HENRY. (R. 1855)	77
collecting and preserving dipters-R. OSTEN SACKEN	102
collecting and preserving fish-T. H. BEAN. (P. 1881) 467, 4	64
collecting and preserving insects—A. S. PACKARD 2	61
collecting and preserving plants-L. F. WARD	60
collecting microscopic organisms-J. W. BAILEY	63
collecting, preserving, and transporting specimens of Diatomacca—A. M.	
Edwards3	66
collecting, preserving, and transporting specimens of natural history— S. F. Baibb. (R. 1856)	24
constructing lightning-rods—J. HENRY 2	
earthquake observations—J. HENRY. (R. 1855)	77
meteorological observations—A. Guyor; J. Henry. (R. 1855). 77, 19, 1	
Directory of officers, collaborators, employés, etc., of the Smithsonian Institu-	.10
tion, National Museum, Geological Survey, Bureau of Ethnology, and Fish Commission	:66
	356
	214
Discovery of—	
•	275
(18
•	244

Discovery relative to magnetism, history of. (R. 1868)
Discussion of magnetic and meterological observations made at Girard College Observatory—A. D. BACHE:—
Part I. Eleven year period, solar diurnal variation ll
Part II. Solar diurnal variation and annual inequality 12
Part III. Influence of moon on declination13
Parts IV, V, VI. Horizontal component of magnetic force, etc 16
Parts VII, VIII, IX. Vertical force
Parts x, xI, xII. Dip and total force18
Parts I-XII. Complete19
Discussion of—
meteorological observations—See Schott.
Piazzi's astronomical observations—B. A. Gould. (R. 1868)
relations of lucernarians to other acalephse, beroids, and polypi—H. J. CLARK
Snell's barometric observations—F. H. Loud. (R. 1880) 442, 43
tables and charts of winds—A. Workor
Disease known as "chorea," inquiries relative to—S. W. MITCHELL. (R. 1874) 28
Disease, origin and propagation of—J. C. Dalton. (R. 1878)
Diseases of joints, bones, etc., bibliography of works on—W. W. KEEN 80
Dispersion of cloud by an electrical discharge—D. W. NAILL. (R. 1858)
Distinction between tornadoes and tempests—J. B. LAMABCK. (R. 1871) 245
Distribution of—
duplicate fishes of Pacific coast, check-list of-D. S. JORDAN; P. L. JOUX. (P. 1881)
fishes of Alleghany region of South Carolina, Georgia, and Tennessee- D. S. JORDAN; A. W. BRAYTON
forest trees in Montana, Idaho, and Washington-W. W. Johnson. (R. 1870)
forests and trees of North America-J. G. Cooper. (R. 1858) 109, 351
marine invertebrates from National Museum, list of—R. RATHBUN. (P. 1881)
Smithson income, majority and minority reports of Committee of Regents
on—J. A. Pearce; J. Meacham. (R. 1858)67, 329
specimens—See each annual report.
District of Columbia—
flora of-L. F. WARD. (P. 1881)
list of birds of-E. Cours; S. S. Prentiss. (R. 1861)149
meteorology of—J. WISSNER. (R. 1857)
prehistoric remains in -T. R. PEALE. (R. 1872) 271
Ditrema atripes, new embiotocoid fish from coast of California—D. S. JORDAN; C. H. GILBERT. (P. 1880)
Documents, collection of historical, in Guatemala—C. H. BERENDT. (R. 1876) 299

Documents relative to origin and history of Smithsonian Institution—W. J. Rhees.	328
Dod, S. B. Memorial discourse on J. Henry	856
Dodge, G. W. Lightning discharges. (R. 1867)	215
Dodge, N. S. Memoir of C. Babbage. (R. 1878)	275
Dodge, N. S. Memoir of Sir John F. W. Herschel. (R. 1871)	249
Dog's revenge, a Dakota fable—S. R. Riggs. (E. 1879–80)	476
Dolichopodidæ, monograph of—H. Loew	171
Domestic institutions in correspondence with Smithsonian Institution, list of 69,	
Domestic institutions in correspondence with Smithsonian Institution, number of	290
Dominant language for science—A. DE CANDOLLE; J. E. GRAY. (R. 1874)	286
Dominica, catalogue of Ober's collections of birds from—G. N. LAWRENCE. (P. 1878)	882
Dominica, new species of Turdidæ from—G. N. LAWRENCE. (P. 1880)	425
DONATI, G. B. Phenomena in telegraphic lines during aurora. (R. 1872)	271
Donations to library from foreign institutions, list of. (R. 1864)	188
Dorosoma cepedianum heterurum, (western gizzard shad,) notes on—S. Wil-	200
мот. (Р. 1878)	832
Dorpat and Poulkova, description of observatories at—C. Abbe. (R. 1867) 215, Dorsey, J. O. Illustrations of method of recording Indian languages. (E. 1879-80)	369 4 76
Dorysomatidæ, notes on-D. S. JORDAN	806
DOUGLAS, S. A., eulogy on, by S. S. Cox. (R. 1861)	149
Douglass, H. M. Translation of Weismann on change of Mexican axolotl. (R. 1877)823,	401
Downes, J. Occultations of planets and stars by the moon, 1858	54
Downes, J. Occultations visible in the United States, 1848-1852 8, 9, 10, 11	., 29
DOYLE, W. E. Indian forts and dwellings, Indian Territory. (R. 1876)	299
Drainage, sanitary, of Washington city, suggestions for-G. E. WARING	849
DRAPER, H., account of telescope of, by T. W. WEBB. (R. 1864)	188
DRAPER, H. Construction of silvered glass telescope and its use in celestial photography	180
DREUTZER, O. E. Statistics relative to Norwegian mountains, lakes, and snow line. (R. 1866)	214
Drift, fresh water glacial, of the Northwestern States—C. WHITTLESEY	197
Drilling in stone without metal—C. RAU. (R. 1868) 224, 440,	372
Drugs, classification of forms in which medicines and, appear and are adminis-	
tered—J. M. FLINT. (P. 1881) 467,	451
Drugs, memoranda for collectors of—J. M. FLINT. (P. 1881) 467,	
Dual character of the brain. Toner lecture No. 11—C. E. Brown-Séquard	
Duby, P. A. Report on transactions of Society of Physics and Natural History	
of Geneva, 1861. (B. 1864)	188

Duck, new to North American fauna-R. RIDGWAY. (P. 1881)	
DUDLEY, T. Earthquake at New Madrid, Missouri. (R. 1858)	109
DUPOUR, C.; Kämtz. Scintillation of the stars. (R. 1861)	149
Ducks, A., notes on fishes collected by, in Mexico-T. H. Bran. (P. 1879)	333
Ducks, A., notes on fishes collected by, in Mexico-D. S. Jordan. (P. 1879).	333
Duke of Northumberland, account of	330
Duke of Northumberland, books presented by. (R. 1859)	110
DUMAS, J. B. Eulogy on A. A. De La Rive. (R. 1874)	286
DUNCAN, M. L. Translation of Holmgren on color-blindness. (B. 1877.) 323, 3	199
Dunkirk Society for the Encouragement of Sciences. Prize questions. (R. 1865; R. 1873)	27.
DUNNING, E. O. Antiquities in Tennessee. (R. 1870)	244
Duplicate fishes distributed by Smithsonian-T. H. BEAN. (P. 1880)	125
Duplicate shells collected by the United States Exploring Expedition under C.	
Wilkes 1	
DUPRÉ, W.; HENRY, J. Earthquakes in North Carolina, 1874. (R. 1874)	26
DUPBEZ, F. Atmospheric electricity. (R. 1858)	
Dwellings and forts, Indian, in Indian Territory-W. E. DOYLE. (R. 1876)	
Dyeing purple, ancient and modern. (R. 1868)	187
H1.	
Earth— chemistry of the—T. S. Hunt. (R. 1869) 228, 3	76
electrical currents of the—C. MATTEUCCI. (R. 1867; R. 1869) 215, 2	
figure of the—St. M. Merino. (R. 1868)1	
internal structure of the—J. G. Barnard.	
laws of atmospheric circulation over the—J. H. Coffin2	
revolutions of crust of the—G. PILAR. (R. 1876)	
secular variation of elements of orbit of—J. N. STOCKWELL	
Karthquake—	
at New Madrid, Missouri—T. Dudley. (R. 1858)	09
directions—J. Henry. (R. 1855)	
in Eastern Mexico, January, 1866—C. SARTORIUS. (R. 1866) 2	
	:44
phenomena, on observations of—R. MALLET. (R. 1859)1	10
Earthquakes—	
articles on, received by Smithsonian Institution and deposited in Library	49

circular relative to—J. HENRY

Earthquakes—Continued.
in Guatemala—A. Canudas. (R. 1858)
in North Carolina, 1874—W. Du Pré; J. Henry. (R. 1874) 28
in St. Thomas—G. A. LATIMEB. (R. 1867) 21
Earthwork, double-walled, in Ashtabula county, Ohio-S. D. PEET. (R. 1876) 29
Earthworks—
ancient, of Ashland county, Ohio-G. W. Hill. (R. 1877) 82
ancient, of Ohio, sketch of—I. DILLE. (R. 1866) 21
ancient, on upper Missouri—A. BARRANDT. (R. 1870)24
in Tennessee—J. JONES
in Wisconsin—E. E. Breed. (R. 1872) 27
on Arkansas river—Mrs. G. KNAPP. (R. 1877)
Earthworks—See Mounds.
Echinoderms of Kerguelen Island—A. E. VERRILL 29
Echinoderms of northeastern coast of America—A. E. VERRILL. (P. 1879; P. 1880)
Eclipse-
annular, of May 26, 1854
of the sun, April 25, 1865—Baron DE PRADOS. (R. 1864)
of the sun, March 15, 1858, map of—T. HILL
of the sun, July 18, 1860—J. LAMONT. (R. 1864)
of the sun, September 7, 1858, in Peru-J. M. GILLISS 10
Reliptic, obliquity of the-J. N. STOCKWELL 28
Economic geology of Trinidad-G. P. Wall; J. G. Sawkins. (R. 1856) 9
Education—
acknowledgment of books on—E. LABOULAYE. (R. 1867) 21
in the United States, project of an outline history of—F. A. PACKARD. (B. 1863) 18
scientific, of mechanics and artisans—A. P. PEABODY. (R. 1872) 271, 380
Education, Illinois State Board of. Meteorological system for every State. (R. 1855)
Educational series of marine invertebrates distributed by United States National
Museum—R. Rathbun. (P. 1881) 467, 465, 47.
EDWARDS, A. M. Directions for collecting, preserving, and transporting diatomacea
EDWARDS, A. M. Results of examination under microscope of Japanese infusorial earths
Edwards, V. N. Occurrence of oceanic bonito (Orcynus pelamys) in Vineyard Sound, Massachusetts. (P. 1878) 38:
EDWARDS, W. H. Diurnal lepidoptera of Arctic America
EDWARDS, W. H. On lepidopters18
Rels, genitalia of male—S. T. CATTIE. (P. 1880) 42
Effect of irritation of polarized nerve—B. F. LAUTENBACH. (R. 1878) 841, 411
Effect of lightning—S. L. HILLIER. (R. 1866)

Effect of moon on the weather—J. HENRY. (R. 1871)	249
EGGERS, H. F. A. Flora of St. Croix and Virgin Islands	
Egging expedition to Shoal Lake, Lake Winnipeg-D. Gunn. (R. 1867)	
Eggs—	
instructions for collecting—S. F. BAIRD. (R. 1858)	139
instructions for collecting—A. NEWTON	139
instructions for collecting and preserving—T. M. Brewer	
North American—T. M. Brewer	89
of the eight North American species of <i>Empidonaces</i> —T. M. Brewer. (P. 1879)	333
suggestions for forming collections of-A. NEWTON	
Egleston, T.—	
Catalogue of minerals with their formulas	156
Scheme for qualitative determinations by blowpipe. (R. 1872)	
translation by, of Brezina on crystallography	
Egypt, presentation of books on—R. LEPSIUS. (R. 1860)	
Elastic force of aqueous vapor, table for determining—J. H. COFFIN	87
Electric phenomenon—W. F. GIVEN. (R. 1865)	
Electric resonance of mountains, observations on—H. DE SAUSSURE. (R. 1868.)	
Electrical-	
currents of the earth—C. MATTEUCCI. (R. 1867; R. 1869) 215,	
discharge, dispersion of a cloud, by D. W. NAILL, (R. 1858)	109
rheometry, researches on—A. SECCHI	36
Electricity—	
accounts of lightning discharges. (R. 1867)	215
and galvanism, report on recent progress in—J. MÜLLER. (R. 1857)	107
and magnetism, later views of connection of—H. Helmholtz; J. C. Maxwell. (R. 1878)	27.5
atmospheric—F. DUPREZ. (R. 1858)	
considerations on. (R. 1867)	
of atmosphere and the aurora borealis-S. Lemström; A. A. De La	286
of induction in clouds resolving into rain, snow, and hail—F. Zante- DESCHI. (R. 1870)	244
phenomena accompanying propagation of, in rarefied elastic fluids—A. A.	187
presence of, during fall of rain-Prof. PALMIERI. (R. 1870)	244
	91
Electro-magnetic—	
seismograph—Prof. PALMIERI. (R. 1870)	244
telegraph, Henry's contribution to-W. B. TAYLOR. (R. 1878) 341,	
telegraph—See Henry; Telegraph.	
Electro-physiology, lectures on—C. MATTEUCCI. (B. 1865)	209

OF SMITHSONIAN PUBLICATIONS.	179
Electrotypes of engravings of shells granted by British Museum. (R. 1808)	187
Elements, table of chemical equivalents of. (R. 1864)	188
Elephant mound in Grant county, Wisconsin—J. WARNER. (R. 1872)	271
ELLERY, R. L. J. Address of the President of the Royal Society of Victoria.	
(R. 1868)	224
ELLET, C. Physical geography of Mississippi valley	13
Ellior, D. G. Classification and synopsis of Trochilidæ	317
Ellior, D. G. List of described species of humming birds	
ELLIOTT, R. S. Climate of Kansas. (R. 1870)	244
ELISWORTH, E. W. Ancient implement of wood from Connecticut. (R. 1876)	299
Embarrass, Wisconsin, pits at—E. E. Breed. (R. 1877)	828
Embryology of insects—L. Agassiz	16
Empidonaces, notes on nests and eggs of the eight North American species of—	
T. M. Brewer. (P. 1879)	888
Employés of Smithsonian Institution, National Museum, Geological Survey,	
Bureau of Ethnology, and Fish Commission, directory of	466
ENCKE, J. F., memoir of, by G. HAGEN. (R. 1868)	224
Endlich, F. M.—	
Analysis of water destructive to fish in Gulf of Mexico. (P. 1881)	467
Geology of Kerguelen Island	294
List of minerals in National Museum, 1873, 1879. (R. 1873; P. 1880) 276	i, 42 5
Endothyra ornata, note on—C. A. WHITE. (P. 1879)	888
ENGELHARDT, M. Formation of ice at the bottom of water. (R. 1866)	214
ENGELMANN, G. Heights of mountains in Colorado. (R. 1862)	150
Engineers, Civil, London Institution of. Prize questions. (R. 1862)	150
England—	
sbbreviations used in—W. DE LA RUE. (R. 1867)	215
free freights between the United States and—E. Cunard. (R. 1859)	110
Salisbury, notice of Blackmore Museum at. (R. 1868)	224
English and French weights and measures—See Tables.	
English vocabulary with comparative words in Spanish, French, and Latin	170
Engravings—	
of shells, electrotypes of, granted by British Museum. (R. 1868)	187
on the face of rocks in the Sierra Nevada—J. G. BRUFF. (R. 1872)	271
presented to Smithsonian Institution, catalogue of—C. B. KING. (R. 1861.)	149
Entomologists, circular to	178
Entomologists, instructions to	261
Eccene and oligocene invertebrate fossils of North America, check-list of—T.	
A. CONBAD	
Edge, J. Habits of black bass of the Ohio. (R. 1854)	75
Ephemeris of the planet Neptune, 1848-1852—S. C. WALKER 4, 5, 6, 5	, 24
Epinephelus Drummond-Hayi, a new serranoid fish from the Bermudas—G. B. GOODE; T. H. BEAN. (P. 1878)	882
VVVII ; 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	

Epinephelus nigritus of the southern coast, note upon—G. B. Goode; T. H. BEAN. (P. 1878)
Equilibrium of a liquid mass withdrawn from the action of gravity—See J. Plateau.
Equinoxes, precession of the—J. G. BARNARD
Equinoxes, precession of the-J. N. STOCKWELL 22
Equivalents, chemical, of sixty-three elements, table of. (R. 1864)188
ERNST, G. A. Meteorology of Caracas, South America. (R. 1867) 215
Erosions of the earth's surface by rivers, etc.—E. HITCHCOCK
Eruption of volcano in Nicaragua—A. B. Dickinson. (R. 1867)
Eruption of volcano of Colima—C. Sartobius. (R. 1869)
Eryontidæ, notice of recent—S. I. SMITH. (P. 1879)
ESPY, J. P., notice of, by A. D. BACHE. (R. 1859)
Espy, J. P. On meteorology. (R. 1847)
Essay on geographical distribution of batrachia and reptilia—E. D. Cope 22
Essay on velocity of light—C. Delaunay. (R. 1864) 188, 354
Establishment and officers of the Smithsonian Institution and National Museum. 449
Establishment of the Smithsonian Institution, Journal and By-Laws of. (R. 1858)
ESTES, L. C. Antiquities of Minnesota, Mississippi river, and Lake Pepin. (R. 1866)
Estimate of population of the world—E. MAILLY. (R. 1878) 275
Etheostomatidæ, notes on—D. S. JORDAN
Ethmocardium, note on—C. A. WHITE. (P. 1879)
Bthnographical-
collections—T. LYMAN. (B. 1862)
collections—H. DE SCHLAGINTWEIT. (R. 1862)
collections, the Schlagintweit—H. ZISGENBALS. (R. 1867) 215
Ethnological—
collections of museum at Lausanne, report on—F. Troyon. (R. 1861). 149
department of the French Exposition, 1867. (R. 1867)215
map of North America, suggestions relative to—L. H. MORGAN. (R. 1861)
map of the United States—G. GIBBS. (R. 1862)
memoir of Squier and Davis, correspondence relative to acceptance of, for publication
museum of Norway—L. K. DAA. (R. 1862)
research—E. H. DAVIS. (R. 1866)
specimens presented by University of Christiania, Norway. (R. 1863) 187
suggestions for Russian America—G. GIBBS
Ethnology and archeology, circular relative to collections in205
Ethnology and philology, instructions for—G. GIBBS
Ethnology, articles on. (R. 1867–1870–1877, 1879, 1880) 215, 244, 249, 271, 275, 286, 298, 299, 328, 345, 442

Ethnology—
Leipsic Museum of—O. T. Mason. (R. 1873) 27
of Arctic America—L. Kumlien
of Colonia Tovar, Venezuela, South America—A. FENDLER. (R. 1857) 10
of Indians of Red river of the North-W. H. GARDNER. (R. 1870) 24
physical, lectures on-D. Wilson. (R. 1862) 18
present state of, in relation to form of human skull—A. Retzius. (R. 1859)
Ethnology, Bureau of—
catalogue of linguistic manuscripts in library of—J. C. Pilling. (B. 1879-80)
directory of officers and employés of 46
first annual report of—J. W. Powell
Euchalarodus Putnami, identity of, with Pleuronectes glaber—T. H. BEAN. (P. 1878)
Eulachon or candle fish of northwest coast—J. G. Swan. (P. 1880) 42
Eulogy on—
Ampère, by F. Arago. (R. 1872)27
A. D. Bache, by J. HENRY. (R. 1870) 244, 37
A. A. De La Rive, by J. B. Dumas. (R. 1874) 28
Stephen A. Douglass, by S. S. Cox. (R. 1861) 14
Cornelius C. Felton, by T. D. Woolsey. (R. 1861)
Joseph Fourier, by F. Arago. (R. 1871) 24
Herschel, by F. Arago. (R. 1870)24
Gay-Lussac, by F. Arago. (R. 1876)
La Place, by F. Arago. (R. 1874)
Quetelet, by A. MAILLY. (R. 1874)
James A. Pearce, by A. D. BACHE. (R. 1862) 15
Gen. J. G. Totten, by J. G. BARNARD. (R. 1865) 20
Alexander Volta, by F. Arago. (R. 1875) 29
Henry Wilson, by P. PARKER. (R. 1875)
Thomas Young, by F. Arago. (R. 1869) 22
Eulogy—See Memoir, Life, Biography.
Europe
causes which limit vegetable species towards the north, in—A. DE CAN- DOLLE. (R. 1858)
certain storms in America and, December, 1836—E. LOOMIS 12
man as the contemporary of the mammoth and reindeer in. (R. 1867) 21
study of high antiquity in—A. Morlor. (R. 1862; R. 1864) 150, 18
Europe, middle, fauna of, during stone age—L. RÜTIMEYER. (R. 1861) 14
EVANS, J. Climate of Colorado. (R. 1865) 20
EVANS, S. B. Notes on mounds in Des Moines valley. (R. 1879) 84
Evaporation in Florida—W. C. DENNIS. (R. 1866)

Evaporation observed at Palermo in 1865 and 1866—P. TACCHINI. (R. 1870.)	944
EVERETT, E. Report on Jewett's general stereotype catalogue of public libraries.	47
EVERETT, E. Report on organization of Smithsonian Institution. (R. 1853.)	
EVERETT, J. D. Under-ground temperature. (R. 1874)	
Evolution of language—J. W. Powell. (E. 1879-80)	
Examination—	110
microscopical, of soundings—J. W. BAILEY	20
of J. Henry by English Scientific Commission	
of specimens, rules for. (R. 1880)	
of Spencer's telescope for Hamilton College. (R. 1855)	
Excelsior, Minnesota, mounds near—F. H. NUTTER. (R. 1879)	
Exchange of publications—	
Agricultural Association of Milan. (R. 1868)	187
Chamber of Commerce of Bordeaux. (R. 1863)	187
Royal Horticultural Society, London. (R. 1861)	149
Exchange of specimens—	
Hamburg Zoölogical Gardens. (R. 1867)	
Hamilton College, Clinton, New York. (R. 1861)	
W. A. LLOYD. (R. 1867)	
Museum of National University of Greece. (R. 1867)	
University of Costa Rica. (R. 1867)	215
Exchanges—	
American Academy of Arts and Sciences, Boston, thanks for. (R. 1855; R. 1867)	
Bath and West of England Society for Encouragement of Agriculture, Arts, etc. (R. 1867)	
circular relative to scientific and literary	
Government of Bremen. (R. 1865)	
W. Hincks. (R. 1860)	
international, report on-G. H. BOEHMER	
list of. Parts I, II73	
list of, to 1858	
Mexican Society of Geography and Statistics. (R. 1861, 1865) 149	
Mining Department, Melbourne. (R. 1865)	
F. MÜLLER. (R. 1860)	
reports on. (R. 1853–1867, 1878) 67, 75, 77, 91, 107, 109, 110, 147,	
150, 187, 188, 209, 214, 215	, 275
J. Rosing. (R. 1865)	
Royal Academy of Science, Madrid. (R. 1861)	
St. Petersburg Academy of Sciences. (R. 1867)	315
statistics of, 1846-1877	\$29
Excursion in New Mexico, diary of—J. H. Carleton. (R. 1854)	75
Executive Committee of Regents—See also each annual report.	

Executive Committee of Regents, journal and reports of, 1846-1876	29
Exhibit by the Smithsonian Institution at Centennial Exhibition, report on	
proposed plan of—S. F. BAIRD. (R. 1875) 29	98
Exhibit of fisheries of United States at Berlin Fisheries Exhibition—G. B.	
Goode 41	.3
Exhibition, Centennial, report on—S. F. BAIRD. (R. 1876) 299, 30)7
Expansion, bibliography of—F. W. CLARKE 20	55
Expansion by heat, tables of—F. W. CLARKE 28	19
Expedition—	
Coast Survey, for transatlantic longitude—B. A. Gould 2	28
Hassler, narrative of the-L. Agassiz. (R. 1872) 2	71
North Pacific Surveying, contributions to natural history made in con-	
nection with the—T. H. STREETS 8	60
to Arctic Seas-See Hayes, Kane, McClintock.	
to Kerguelen Island—J. H. KIDDER and others 298, 2	94
to Lake Winnipeg-D. GUNN. (R. 1867)	15
	88
to the Mauvaises Terres and Upper Missouri-T. A. CULBERTSON. (R.	
1850)	28
toward the North pole, scientific instructions for—J. HENRY and others.	
(R. 1871) 2	49
Expeditions, account of—See each annual report.	
Expeditions contributing specimens, list of—S. F. BAIRD. (R. 1867) 2	15
Experimental and theoretical researches on figures of equilibrium—See J. Plateau.	
Experiments	
on aneroid barometers at Kew Observatory—B. STEWART. (R. 1868) 2	24
relative to meteorites—G. A. DAUBRÉE. (R. 1868) 2	24
upon animal heat of fishes-J. H. Kidder. (P. 1879)	88
Explanation of principles of crystallography and crystallophysics—A. Brezina.	
(R. 1872) 271, 38	36
Exploration of ancient mounds in Union county, Ky.—S. Lyon. (R. 1870) 2	44
Explorations—	
among Indian mounds in southern Florida—S. T. WALKER. (R. 1879.) 3	45
and surveys, Government, report of—S. F. BAIRD. (R. 1878) 8	41
Arctic, lecture on—I. I. HAYES. (R. 1861) 1	49
articles on—See each annual report.	
botanical, in New Mexico and California, account of—A. Grav. (R. 1849)	21
furnishing collections to National Museum, 1838-1877, list of-S. F.	28
•	15
·	41
·	42
,	

ALPHABETICAL INDEX

Explorations—Continued.	
in New York—E. G. Squier	15
in Tennessee-E. A. DAYTON. (R. 1870)	244
in upper California in 1860—J. FEILNEB. (R. 1864)	188
Kennicott's-Hudson's Bay Company. (R. 1863)	187
of aboriginal remains of Tennessee-J. JONES	259
of the Nile-C. HALE. (R. 1865)	209
of western Missouri in 1854—P. R. Hov. (R. 1864)	188
of John Xantus in Mexico-M. Romero. (R. 1862)	150
on western coast of North America-W. H. Dall. (R. 1878)	275
reports on—S. F. BAIRD. (R. 1851-1866, 1875-1877) 51, 57, 75, 77, 91,	107.
109, 110, 147, 149, 150, 187, 188, 209, 214, 298, 299	, 323
scientific, in America, in 1852—S. F. BAIRD. (R. 1852)	57
scientific, in Mexico. (R. 1864)	188
Exploring expedition under Capt. C. Wilkes, duplicate shells collected by the.	193
Explosibility of coal oils—Z. Allen. (R. 1861)	149
Explosiveness of nitre—R. HARE	17
Exposition of harmonies in the solar system—S. ALEXANDER	280
Exposition—See French.	
Expression for anthropological inquiry, queries about—C. DARWIN. (R. 1867)	215
External appearance of the sun's disk. (R. 1866)	214
Extinct—	
mammalia and chelonia of Nebraska—J. LEIDY	58
10/1110-0. 2222	192
GIVE BIDE OF AVIET AMORICA	72
species of minerical va—v. main a constant and a co	41
Eye, bibliography of diseases of the-W. W. KEEN	300

F.

Facts respecting Liberia College—A. CRUMMELL. (R. 1861) 149
Fahrenheit's scale, tables for conversion of centigrade degrees to. (R. 1863) 187
Families—
of fishes, arrangement of—T. GILL 247
of mammals, arrangement of—T. GILL230
of mollusks, arrangement of—T. GILL 227
Family, human, systems of consanguinity and affinity of—L. H. Morgan 188, 218
Fanning Islands, natural history of—T. H. Streets
Faraday, M., his life and works-A. A. DE LA RIVE. (R. 1867)
FARLOW, W. G.—
Alge of Arctic America
Algæ of Kerguelen Island 294
Recent progress in botany, 1879, 1880. (R. 1880) 442, 430
Report on water from Gulf of Mexico. (P. 1881) 467
Farming and gardening, natural history as applied to-J. G. Morris. (R. 1855.) 77
Farce Isles, vegetable colonization of-C. MARTINS. (R. 1858) 109
FARQUHARSON, R. J. Study of skull and long bones from mound in Illinois. (R. 1874) 286
Fauna-
and flora within living animals—J. LEIDY
littoral marine, of Provincetown, Mass.—R. RATHBUN. (P. 1880) 425
of middle Europe during the stone age—L. RÜTIMEYER. (R. 1861) 149
of Nebraska, ancient—J. Leidy 58
FAVRE, A. Report on transactions of Geneva Society of Physics and Natural History, July, 1876, to June, 1877. (R. 1877)
FAVRE, E. Biographical notice of Louis Agassiz. (R. 1878)
Feeling, sense of. (R. 1865) 209
FEILNER, J. Explorations in upper California in 1860. (R. 1864) 188
Feldspar, determination of, in thin sections of rocks-G. W. HAWES. (P. 1881) 467
Felton, C. C., eulogy on, by T. D. Woolsey. (R. 1861) 149
Felton, C. C.—
Notice of Washington Irving. (R. 1859) 110
Notice of W. W. Turner. (R. 1859) 110
Report on Prof. Henry and the telegraph. (R. 1857) 107
Fendler, A., botanical explorations by, in New Mexico and California—A. GRAY. (R. 1849)
Fendler, A.—
Meteorology and ethnology of Colonia Tovar, Venezuela. (R. 1857) 107 Meteorology of Colonia Tovar. (R. 1866) 214
Temperature of St. Louis, Missouri. (R. 1860)

Ferns, acknowledgment for—G. METTENIUS. (R. 1862)
FERREL, W. Converging series expressing ratio between diameter and cir-
cumference of a circle233
FERREL, W. Reply to the criticisms of Dr. J. Hann. (R. 1877) 823, 398
Ferrel, W., theories of, as to relation between difference of pressure and velocity of wind-J. Hann. (R. 1877) 823, 398
Fever—
a study in morbid and normal physiology—H. C. Wood
researches on—H. C. Wood. (R. 1878)
study of the nature and mechanism of. Toner lecture No. IV—H. C. WOOD
Fevers, surgical complications and sequels of. Toner lecture No. v—W. W. KEEN
Fialho, A. Biographical sketch of Dom Pedro II. (R. 1876)
FIGANIERRE, M. Account of remarkable accumulation of bats. (R. 1868) 187
Figure of the earth—St. M. MERINO. (R. 1868)
Figures of equilibrium of a liquid mass withdrawn from the action of gravity— See Plateau.
Figures, series of, for labels 164
Filices of Kerguelen Island—A. Grav
Finances of the Institution, 1846-1877
FINCH, FLADGATE, and CLARKE. Residuary legacy of Smithson. (B. 1861.) 149, 325
FINCK, H. Antiquities in the State of Vera Cruz, Mexico. (R. 1870) 24
Fire at Smithsonian, January, 1865, report of Committee of Regents relative to—R. WALLACH; J. HENRY. (R. 1864)
Fire, Indian mode of obtaining—G. CROOK. (R. 1871) 249
Fire-alarm telegraph, American, lecture on-W. F. CHANNING. (R. 1854) 5
Fire ball, meteoric, orbit and phenomena of-J. H. Coffin
Fire-proofing buildings, architecture in relation to-D. B. REID. (R. 1856) 9
First decade of United States Fish Commission—G. B. Goode. (R. 1880) 442
FISCHER, F. Scientific labors of Edward Lartet. (R. 1872) 27!
Fish Commission—
account of work of—S. F. BAIRD. (R. 1880)
catalogue of exhibit of, at Centennial, 1876—G. B. Goods 326
directory of officers and employés of466
exhibit by, at Berlin Fisheries Exhibition—G. B. Goods
experiments on animal heat in fishes, made in connection with—J. H. KIDDEB. (P. 1879)
first decade of—G. B. GOODE. (R. 1880)
A. E. Verrill; R. Rathbun. (P. 1879)
mollusca collected by—A. E. VERRILL. (P. 1880)
notice of crustacea dredged by, off south coast of New England—S. I. SMITH. (P. 1880)
•

Fish-	
analysis of water destructive to, in Gulf of Mexico-F. M. Endlich. (P. 1881)	
candle, of northwest coast—J. G. SWAN. (P. 1880)	425
dates of first appearance of-F. B. Hough	182
description of new species of, (Apogon pandionis)—G. B. GOODE; T. H. BEAN. (P. 1881)	467
destruction of, by poisonous water in Gulf of Mexico—J. Y. PORTER. (P. 1881)	467
destruction of, by polluted waters in Gulf of Mexico—W. C. W. GLA- ZIEB. (P. 1881)	467
directions for collecting and preserving—T. H. BEAN	464
fresh, circular on shipping—S. F. BAIRD	
from deep-sea fauns of western Atlantic, description of new species of, (Alepocephalus Bairdii)—G. B. GOODE; T. H. BEAN. (P. 1879)	888
mortality of, in Gulf of Mexico—E. INGERSOLL. (P. 1881)	467
mortality of, in Gulf of Mexico-M. A. Moore. (P. 1881)	467
new, from Alaska, with notes on genus Anarrhichas—T. H. BEAN. (P. 1879)	883
new, (Lopholatilus chamæleonticeps,) from south of New England—G. B. Goode; T. H. Bean. (P. 1879)	888
new gobioid, (Othonops cos.) from San Diego, Cal.—R. SMITH. (P. 1881)	467
new serranoid, (Epinephelus Drummond-Hayi,) from Bermudas and Florida—G. B. Goode; T. H. Bean. (P. 1878)	882
new sparoid, (Sargus Holbrookii,) from Savannah Bank, description of— T. H. BEAN. (P. 1878)	882
Fish-culture of United States, exhibit of, at Berlin Fisheries Exhibition— G. B. Goode.	418
FISHER, J. G. Acknowledgment of perennibranchiates. (R. 1868)	187
Fisheries—	
of United States, catalogue of collection to illustrate, at Philadelphia— G. B. GOODE	326
of United States, exhibit of, at Berlin Fisheries Exhibition—G. B. Goode.	413
Fisheries Exhibition, Berlin, additions to Museum from. (R. 1880)	442
Fishes—	
Alaskan, catalogue of—T. H. BEAN. (P. 1881)	467
American, in British Museum and Museum d'Histoire Naturelle, Paris,	
notes on typical—D. S. JORDAN. (P. 1879)	388
arrangement of families of—T. GILL	
bibliography of—T. GILL	247
check-list of duplicates of, distributed by Smithsonian Institution—T. H. BEAN. (P. 1880)	425
T. H. BEAN. (P. 1881)	467

Fishes—Continued.
cottoids, monograph of the-C. GIRARD
description of new species of North American—D. S. JORDAN. (P. 1879.)
descriptions of and notes on-See D. S. JORDAN; C. H. GILBERT.
descriptions of deep-sea species from New England, diagnoses of two undescribed genera of flounders and genus related to Merlucius—G. B. GOODE. (P. 1880)
descriptions of new species of, (Uranidea marginata, Potamocottus Bendirei,) and of Myctophum crenulare—T. H. Bean. (P. 1881)
directions for collecting and preserving-T. H. BEAN. (P. 1881) 4
European, in National Museum, list of-T. H. BEAN. (P. 1879) 3
experiments upon animal heat of—J. H. KIDDER. (P. 1879)
from deep waters on south coast of New England, obtained by Fish Commission in 1880—G. B. Goode. (P. 1880)
from eastern Georgia, notes on—T. H. BEAN. (P. 1879) 3
from eastern Mississippi, collection of—O. P. HAY. (P. 1880)
from Hudson's Bay, notes on—T. H. BEAN. (P. 1881)
from Utah Lake, notes on a collection of—D. S. JORDAN; C. H. GIL- BERT. (P. 1880)
mortality of, in Gulf of Mexico-J. P. JEFFERSON. (P. 1878) 3
mortality of, in vicinity of Tortugas—J. P. JEFFERSON; J. Y. POETER; T. MOORE. (P. 1878)
new genus of, (Benthodesmus)—G. B. GOODE; T. H. BEAN. (P. 1881.)
new genus of, (Delolepis)—T. H. BEAN. (P. 1881)
North American, check-list of duplicates distributed by Smithsonian— T. H. Bean. (P. 1880)
North American, review of Rafinesque's memoirs on-D. S. JORDAN
notes on—D. S. JORDAN 305, 306. 3
notes on collection of, from Clackamas river, Oregon—D. S. JORDAN. (P. 1878)
of Alaska, some genera and species of—T. H. BEAN. (P. 1879) 3
of Alaska and Siberia, descriptions of new-T. H. BEAN. (P. 1881) 4
of Alleghany region of South Carolina, Georgia, and Tennessee, distribution of—D. S. Jordan; A. W. Brayton
of Arctic America—T. H. BEAN34
of Beaufort Harbor, North Carolina, notes on-D. S. Jordan; C. H. Gilbert. (P. 1878)
of Bermuda, catalogue of—G. B. Goode29
of Bermuda mistakenly described as new by Günther—G. B. Goods. (P. 1878)
of California—T. H. Streets
of California, descriptions of new genera and species of—W. N. Lock- INGTON. (P. 1879)
Standard of North Association of M. Com.

٠	on

OF SMITHSONIAN PUBLICATIONS.

Fishes—Continued.
of Fanning Islands-T. H. STREETS
of Florida, description of new species of, (Seriola Stearnsii)—G. B. Goode. (P. 1879)833
of Florida, preliminary catalogue of, with new genus and three new species—G. B. Goode. (P. 1879)
of Gulf of Mexico, collected by J. W. Velie, catalogue of and description of seven new species—G. B. Goode; T. H. Bean. (P. 1879) 838
of Hawaiian Islands—T. H. STREETS
of Kerguelen Island—T. GILL
of Mexico, descriptions of two species of, collected by Dugès—T. H. BEAN. (P. 1879)
of Mexico, notes on Dugès collection of-D. S. JORDAN. (P. 1879) 383
of New York—T. Gill. (R. 1856) 91
of Pacific coast of United States, bibliography of-T. GILL468
of Pacific coast and Alaska, bibliography of-T. H. BEAN. (P. 1881) 467
of Pacific coast, check-list of duplicates of, distributed by Smithsonian Institution in 1881—D. S. JORDAN; P. L. JOUY. (P. 1881) 467
of Pacific coast, notes on-D. S. JORDAN; C. H. GILBERT. (P. 1881) 467
of Pacific coast of United States, list of—D. S. JORDAN; C. H. GILBERT. (P. 1880)425
of Pensacola, Florida, catalogue of Stearn's collection and description of new species—G. B. Goode; T. H. Bean. (P. 1879)
of St. John's river—G. B. Goode. (P. 1879)
of Samoan Islands—T. H. STREETS 303
on coast of New Jersey and Long Island—S. F. BAIRD. (R. 1854) 75, 348
pediculate, of eastern coast of extratropical North America, synopsis of— T. GILL. (P. 1878)832
Potamocottus Bendirei, description of-T. H. BEAN. (P. 1881) 467
trunk, Ostraciontidæ, a study of, with notes on American species—G. B. Goode. (P. 1879)
Uranidea marginata, description of—T. H. BEAN. (P. 1881)
Fishes, food—See G. B. Goode, T. H. Bean, D. S. Jordan, S. Stearns, F. Steindachner, S. Wilmot.
Fishes, food—
circular relative to—S. F. BATRD 284
memoranda of inquiry relative to—S. F. BAIRD 231
questions relative to—S. F. BAIRD 234
Fishkill, New York, deposit of arrow-heads near—E. M. SHEPARD. (R. 1877.) 323
Fissirostres, oölogy of—T. M. Brewer.
FLACHENECKER, G. Indian languages. (R. 1862) 150
FLADGATE, CLARKE, and FINCH. Smithson's residuary legacy. (R. 1861.) 149, 328
Flamingo from south Florida, specimens of—G. WURDEMAN. (R. 1860) 147
Flight in the animal kingdom, phenomena of-E. J. MAREY. (R. 1869) 228

Flight, modes of, in relation to aeronautics—J. B. Pettigrew. (R. 1867) 215
FLINT, E., shells from Costa Rica kitchen-midden collected by—W. H. Dall. (P. 1878)
FLINT, J. M.—
Classification and arrangement of the materia medica collection 450
Classification of forms of drugs and medicines451
Memoranda for collectors of drugs 452
Flint implements—
in Illinois, agricultural—C. RAU. (R. 1868) 224, 440, 370
in Illinois, deposits of—J. F. SNYDER. (R. 1876)
in Ohio, Holmes county—H. B. CASE. (R. 1877)
Flood-tide, law of deposit of—C. H. DAVIS
Flora—
and fauna within living animals—J. LEIDY 44
of Alaska, sketch of the—J. T. ROTHROCK. (R. 1867) 215, 367
of Crozet and Kerguelen Islands—J. H. KIDDER and others 29
of North America, index of authorities for species of—S. WATSON 258
of St. Croix and the Virgin Islands—H. F. A. EGGERS
of Washington and vicinity—L. F. WARD
Florida—
antiquities of—J. BARTRAM. (R. 1874)
antiquities of—A. MITCHELL. (R. 1874)
catalogue of casts of Indian prisoners in—R. H. PRATT. (P. 1878) 322
colored bead from mound in—A. M. HABRISON. (R. 1877)
Curtiss' collection of fishes from—D. S. JORDAN. (P. 1880)425
evaporation in—W. C. DENNIS. (R. 1866)
fishes of—G. B. GOODE; T. H. BEAN. (P. 1878) \$52
fishes of, preliminary catalogue and new genus and species of—G. B. GOODE. (P. 1879)
forests and trees of—J. G. Cooper. (R. 1860)
gold ornament from mound in—C. RAU. (R. 1877) 323, 440, 403
Henshall's collection of fishes from—D. S. Jordan. (P. 1880) 425
Indian mounds in—S. T. WALKER. (R. 1879)
microscopical observations in—J. W. BAILEY
new fish, Caulolatilus microps, from-G. B. Goode. (P. 1878)
polychrome bead from—S. S. HALDEMAN. (R. 1877) \$23, 404
shell-heaps at mouth of St. John's river—S. P. MAYBERRY. (R. 1877).
shell-heaps of Tampa Bay—S. T. Walker. (R. 1879) 845
specimens of flamingo from—G. WURDEMAN. (R. 1860)
winds in—J. Baltzell. (R. 1866)
Flounders, diagnoses of two undescribed genera of—G. B. Goode. (P. 1880). 425
FLOURENS, M. J.—
Historical sketch of the Academy of Sciences, Paris. (R. 1862)

OF SMITHSONIAN PUBLICATIONS.	191
FLOURENS, M. J.—Continued.	
History of the works of Cuvier. (R. 1868)	224
Memoir of Ducrotay De Blainville. (R. 1865)	209
Memoir of Leopold von Buch. (R. 1862)	150
Memoir of Pyramus de Candolle. (R. 1859)	110
Memoir of Cuvier. (R. 1868)	22
Memoir of Magendie. (R. 1866)	214
Memoir of Geoffroy Saint-Hilaire. (R. 1861)	149
Memoir of Louis Jacques Thénard. (R. 1862)	150
The Jussieus and the natural method. (R. 1867)	21
Fluctuations of level in North American lakes—C. WHITTLESEY	119
Fluids, elastic, electricity in—A. A. DE LA RIVE. (R. 1868)	187
Fly-catcher from Sandwich Islands, description of a new—R. RIDGWAY. (P. 1881)	467
Fog, the wind and—J. BALFOUR. (R. 1866)	214
Folsom, C. Report on Jewett's general stereotype catalogue of public libraries.	47
Food, relation of, to work, and its bearings on medical practice—S. HAUGHTON. (R. 1870)	244
Food collections, provisional classification of—G. B. GOODE	455
Food-fishes, inquiry relative to—S. F. BAIRD	231
Food-fishes, questions relative to—S. F. BAIRD.	234
Force, thoughts on the nature and origin of-W. B. TAYLOR. (R. 1870) 244,	375
FORCE, P. Record of auroral phenomena in high northern latitudes	84
Foreign—	
authors. North American conchology-W. G. BINNEY	174
Commissioners, collections presented to United States by. (R. 1876)	299
correspondents of Smithsonian Institution, list of 154, 225, 309,	469
correspondents of Smithsonian Institution, number of	290
correspondents of Smithsonian Institution, systematic index of	257
gold and silver coins, table of. (R. 1868)	224
institutions in correspondence with Smithsonian Inst., list of 64, 225	243
institutions, list of addresses of, since 1862. (R. 1865)	209
institutions making donations to Smithsonian library, list of. (R. 1864)	188
works in library of Smithsonian—See Library.	
FOREMAN, E. Meteorological system and correspondence of Smithsonian In-	
stitution. (R. 1851)	51
FOREMAN, E. Report on meteorological system. (R. 1852)	57
Forest trees, distribution of, in Montana, Idaho, and Washington—W. W.	244

Forests and their climatic influence—A. E. Becquerel. (R. 1869) ______ 228
Forests and trees of Florida and the Mexican boundary—J. G. Cooper. (R.

Forests and trees of North America, distribution of, and catalogue of native trees of the United States—J. G. Cooper. (R. 1858) 109, 351
Formation—
of a museum—L. Agassiz. (R. 1849)
of clouds over Gulf Stream-H. M. BANNISTER. (B. 1866) 214
of ice at bottom of water—M. ENGELHARDT. (R. 1866) 214
of precipitation—J. HANN. (R. 1877) 823, 398
Forms in which drugs and medicines appear—J. M. FLINT. (P. 1881) 467, 451
Formulas, chemical, tables of—F. W. CLARKE
Fort, ancient, and burial ground in Tompkins Co., N. Y.—D. TROWBEIDGE. (R. 1868)
Fort Brown, Texas, list of birds from-J. C. MERRILL. (P. 1878) 32
Fort Ellis, prehistoric remains near—P. W. Norris. (R. 1879) 345
Fort Marion, Florida, list of casts of heads of Indian prisoners in—R. H. PRATT. (P. 1878)
Fort Ripley, Minn., natural history of country about—J. E. HEAD. (R. 1854.) 75
Fort Wadsworth, Dakota, Indian mounds near—A. J. Comfort. (R. 1871) 249
Forts and dwellings, Indian, in Indian Territory-W. E. DOYLE. (R. 1876) 299
Fossil—
corbiculadæ—T. PRIME
gasteropod, from Mexico, description of-C. A. WHITE. (P. 1880) 425
mollusks from later tertiaries of California-W. H. Dall. (P. 1878) \$32
ox—J. Leidy
plants from China, description of—J. S. NEWBERRY 202
Fossils—
carboniferous invertebrate, description of new species of—C. A. WHITE. (P. 1879)
catalogue of rocks, minerals, ores, and—J. Locke. (R. 1854)
cretaceous, from Arkansas and Colorado—C. A. WHITE. (P. 1881) 45:
cretaceous invertebrate, from Kansas and Texas, description of new— C. A. White. (P. 1879)
from Nebraska, report on—J. LEIDY. (R. 1851)
from Santa Barbara, California—P. P. CARPENTER
invertebrate, from Arkansas, Wyoming, Colorado, Utah—C. A. White. (P. 1880)
invertebrate, of North America, check-list of-T. A. Conrad 200
invertebrate, of North America, check-list of—F. B. MERK 177, 183
invertebrate, of North America, cretaceous and jurassic-F. B. MEEK. 177
invertebrate, of North America, eocene and oligocene-T. A. Conrad 200
invertebrate, of North America, miocene—F. B. MEEK183
postpliocene, in coast range of California-W. H. Dall. (P. 1878) 312
tertiary, distribution of Californian-W. H. Dall. (P. 1878) 322
tertiary, presented by Imperial Geological Institute, Vienna. (R. 1863)

OF SMITHSONIAN PUBLICATIONS.	193
FOSTER, J. W. Ancient relics in Missouri. (R. 1863)	187
FOSTER, J. W. Catalogue of rocks, minerals, etc. (R. 1854)	75
Fourier, J., eulogy on, by F. ARAGO. (R. 1871)	2 4 9
FOURIER, J. Memoir of Delambre. (R. 1864)	188
FOWLER, J. Shell-heaps of New Brunswick. (R. 1870)	244
Fox, (yacht,) meteorological observations in Arctic Seas made on—F. L. McCLINTOCK	146
Fox, (yacht,) record of voyage of, in Arctic regions-F. L. McCLINTOCK	146
France, Emperor of, report to, by Minister of Public Instruction, on scientific expedition to Mexico. (R. 1864)	188
France—See Paris, Prize questions.	
Franklin, Sir John, expedition in search of-See Kane.	
Free freight between Germany and the United States by North German Lloyd— B. Schleiden. (R. 1858)	109
Free freight between United States and England—E. Cunabl. (R. 1859)	110
Free freight between United States and Germany—Kunhardt & Co. (R. 1861)	149
FRÉMORT, J. C., description of plants collected by—J. Torrey	46
Frémontians, plants—J. Torrey	46
French —	
Exposition of 1867, ethnological department of the. (R. 1867)	215
half-breeds of the northwest—V. HAVARD. (R. 1879)	845
Institute, history of-M. FLOURENS. (R. 1862)	150
Society of Archeology, Archeological Congress organized by. (R. 1866)	214
vocabulary with comparative words in English, Spanish, and Latin	170
weights and measures—See Tables.	
Fresh fish and other animals, circular relative to shipping	384
Fresh-water—	•••
alge of North America—H. C. Wood	241
glacial drift of northwestern States—C. WHITTLESEY	
in the ocean—W. C. DENNIS. (R. 1866)	
shells—See Shells.	
FRIEDLÄNDER, J. Plan of a bibliography. (R. 1858)	109
Friel, J. Antiquities of Hancock county, Kentucky. (R. 1877)	828
Friends of the Museum, circular addressed to—S. F. BAIRD. (P. 1881) 467, Frigate mackerel (Auxis Rochei) on New England coast—G. B. Goode. (P.	
1880)	425
FROEBEL, J. Physical geography of North American Continent. (R. 1854.)	75
Frost, disintegrating effects of, on building stones—C. G. PAGE	829
Fruits, dates of ripening of—F. B. HOUGH	182
Fuca, Straits of, Indians of—See J. G. Swan.	
Fund, Smithson, memorial of Regents to Congress relative to. (R. 1850)	28
Fund, Smithson, statement of, 1846-1877	829

Fundy, Bay of, synopsis of marine invertebrata of—W. STIMPSON	
Funeral of J. Henry	
Future of geology—J. Prestwich. (R. 1875)	
Future of the human race—A. DE CANDOLLE. (R. 1875)	
G.	
Gadus cimbrius, identity of, with Rhinonemus caudacuta—G. B. Goode; T. H. Bean. (P. 1878)	
GAINES, A. S.; CUNNINGHAM, K. M. Shell-heaps on Mobile river, Alabama. (R. 1877)	
GALE, L. D., statement of, on telegraph. (R. 1857)	
Galeorhinus galeus—D. S. JORDAN; C. H. GILBERT. (P. 1880)	
Gales of wind and appearance of aurora, connection of—R. T. KNIGHT; J. HENRY. (R. 1871)	
GALLATIN, A. Comparative vocabulary	
GALLATIN, A. On publication of Squier and Davis' work. (R. 1847) E	
Gallery, Art-See Corcoran.	
GALT, F. L. Indians of Peru. (R. 1877)	
Galvanism, recent progress in. J. MÜLLER. (R. 1855)	
Galvanometer, use of, as a measuring instrument—J. C. POGGENDORF. (R. 1859)	
Gangrene, bibliography of—W. W. KEEN	
Gardening, natural history as applied to farming and—J. G. Morris. (R. 1855)	
GARDINER, R. H.— Barometer, rain, and snow gauges. (R. 1858)	
Disappearance of ice. (R. 1860)	

GARDNER, W. H. Indians of valley of Red river of the North. (R. 1870) -- 244 GARFIELD, J. A. Biographical notice of S. P. Chase and L. Agassiz. (R.

GARMAN, S. American Rhinobatidæ, synopsis and description of. (P. 1880).- 425 Gas, hydrogen, as metal and—J. E. REYNOLDS. (R. 1870) _____ 244 Gasteropod, large fossil from Puebla, Mexico—C. A. WHITE. (P. 1880) 425 Gasterosteus, description of new species of, from Schoodic Lakes, Maine-

Gatschet, A. S.—	
Klamath lake Indian conjurer's practice. (E. 1879-80)	476
"The relapse" in Klamath lake dialect. (E. 1879-80)	476
Gauge, snow—W. E. Guest. (R. 1858)	109
Gauges, rain and snow-R. H. GARDINER. (R. 1868)	109
GAUTIER, Prof. Researches relative to nebulæ. (R. 1868)	187
General catalogue system for libraries—C. C. JEWETT. (R. 1850)	28
Generation, alternate and parthenogenesis in the animal kingdom—G. A. Korn-	
нивкв. (R. 1871)	249
Generic names of recent and fossil animals, list of-S. H. SCUDDER	470
Geneva, Society of Physics and Natural History of, report on transactions of-	
July, 1858, to June, 1859—A. A. DE LA RIVE. (R. 1859)	110
" 1860, " 1861—P. Duby. (R. 1864)	188
" 1861, " 1862—A. DE CANDOLLE. (R. 1864)	188
" 1862, " 1868—Prof. MARCET. (R. 1868)	187
" 1868, " 1864—Dr. Сноват. (R. 1865)	209
" 1864, " 1865—E. Plantamour. (R. 1865)	209
" 1865, " 1866—Dr. Gosse. (R. 1866)	214
" 1867, " 1868—E. WARTMANN. (R. 1868)	224
" 1868, " 1869—H. C. LOMBARD. (R. 1869)	228
" 1870, " 1871—H. DE SAUSSURE. (R. 1871)	249
" 1872, " 1878—A. A. DE LA RIVE. (R. 1874)	286
" 1878, " 1874—A. DE CANDOLLE. (R. 1875)	298
" 1874, " 1875—E. PLANTAMOUR. (R. 1877)	828
" 1875, " 1876—J. MÜLLER. (R. 1877)	828
" 1876, " 1877—A. FAVRE. (R. 1877)	828
Genitalia of male eels—S. T. CATTIE. (P. 1880)	4 25
GENTH, F. A.; GIBBS, W. Ammonia-cobalt bases	88
Geographical distribution of batrachia and reptilia—E. D. COPE	292
Geographical manuscripts, catalogue of collection of-L. Berlandier. (R.	
1854)	75
Geography—See Physical Geography.	
Geography of North American Continent-J. FROEBEL. (R. 1854)	75
Geological—	
researches in China, Mongolia, and Japan—R. PUMPELLY	202
specimens, catalogue of—D. D. Owen. (R. 1854)	75
survey in Michigan, catalogue of rocks, minerals, and ores collected on-	
C. T. JACKSON. (R. 1854)	75
Geological Survey, directory of officers and employés of	466
Geological surveys—See annual reports; also Hayden, Powell, Wheeler.	
Geology—	
and history, boundary line between—E. Suess. (R. 1872)	271
economic, of Trinidad, report on survey of-G. P. WALL; J. G. SAW-	
KINS. (R. 1856)	91
of Kerguelen Island-F. M. ENDLICH	294

Geology—Continued.	
of lower Louisiana and salt deposit on Petite Anse Island—E. W. Hil-	248
past and future of-J. Prestwice. (R. 1875)	
progress in, in 1879, 1880—G. W. HAWES. (R. 1880) 442,	
surface, illustrations of—E. HITCHCOCK	90
Georgia—	
aboriginal structures in—C. C. Jones. (R. 1877) 823,	400
ancient mounds in—M. F. STEVENSON. (R. 1870)	244
Bartow county, mound in-M. F. STEVENSON. (R. 1872)	271
description of fishes of Alleghany region of—D. S. JORDAN; A. W. BRAYTON	308
habits of pouched rat, or salamander of-W. GESNER. (R. 1860)	147
McIntosh and Early counties, mounds in-W. McKinley. (R. 1872).	271
microscopical observations in-J. W. BAILEY	2
notes on collection of fishes from-T. H. BRAN. (P. 1879)	331
occurrence of Stricklandia salteri and Stricklandia Davidsoni in—C. A. White. (P. 1880)	42
remarkable forms of hail stones in—S. ABICH. (R. 1869)	
shell-heaps in—D. Brown (R. 1871)	
Spalding county, antiquities of-W. B. F. BAILEY. (R. 1877)	
Gephyrma of northeast coast of America—A. E. VERRILL. (P. 1879)	
Germany and United States, free freight between, by North German Lloyd- R. Schleiden. (R. 1858)	
Germany and United States, free freight between—Kunhardt & Co. (R. 1861)	
GESNER, W. Habits of pouched rat, or salamander of Georgia. (R. 1860)	147
GESNER, W. Mica beds in Alabama. (R. 1879)	
GIBBONS, H. Climate of San Francisco. (R. 1854)	78
GIBBES, R. W. Memoir on Mosasaurus and allied new genera, Holcodus, Comosaurus, Amphorosteus	14
Gіввя, G.—	
Archeology in the United States. (R. 1861)	
Bibliography of Chinook jargon	101
Comparative vocabulary	17V 187
Dictionary of Chinook jargon, or trade language of Oregon	ror
Ethnological instructions	TON
Ethnological map of the United States. (R. 1862)	150
Ethnological suggestions	2Uî
Indian languages. (B. 1865)	(U)
Indian vocabularies. (R. 1862)	150
Instructions for archeological investigations. (R. 1861)	149
Instructions for ethnology and philology	.60

OF SMITHSONIAN PUBLICATIONS.	197
GIBBS, G.—Continued.	
Intermixture of races. (R. 1864)	188
Language of aboriginal Indians of America. (R. 1870)	244
Makah Indians	220
Philological circular. (B. 1862)	150
Physical atlas of North America. (R. 1866)	214
Suggestions for scientific investigations in Russian America	207
Gibbs, G., memoir of, by J. A. STEVENS. (R. 1878)	275
GIRBS, G., and others. Recommendation of Shea's Indian linguistics. (R. 1861)	149
GIBBS, G., and others. Tinneh or Chepewyan Indians of British and Russian America. (R. 1866)	365
GIBBS, G. J. Stone celts in the West Indies and Africa. (R. 1877)	828
GIBBS, W.; GENTH, F. A. Researches in ammonia-cobalt bases	88
Gideon's Farm, Hennepin county, Minnesota, mounds on—F. H. NUTTER. (R. 1879)	8 4 5
GILBERT, C. H.—See D. S. Jordan; C. H. Gilbert.	
GILBERT, Sir D. Notice of Smithson. (R. 1858) 67	, 880
Gill, T.—	
Arrangement of families of fishes	247
Arrangement of families of mammals	230
Arrangement of families of mollusks	227
Bibliography of fishes	247
Bibliography of fishes of Pacific coast of United States.	468
Bibliography of mammals	280
Bibliography of mollusks	227
Catalogue of fishes of east coast of North America	283
Fishes of Kerguelen Island	294
Fishes of New York. (R. 1856)	91
Identity of the genus Leurynnis Lockington, with Lycodopsis Collett. (P. 1880)	425
Note on Antennariida. (P. 1878)	882
Note on Ceratiidæ. (P. 1878)	882
Note on the Latiloid genera. (P. 1881)	467
Note on Maltheidæ. (P. 1878)	882
Recent progress in zoölogy. (R. 1880)	431
Status of ichthyology	247
Synopsis of pediculate fishes of the eastern coast of extratropical North America. (P. 1878)	882
Synoptical tables of mammals	280
GILLISS, J. M. Solar eclipse, Peru, September 7, 1858	
GILLMAN, H. Characteristics of ancient man in Michigan. (R. 1875) 298,	
GILLMAN, H. Mound builders and platycnemism in Michigan. (R. 1878.) 275,	

GILMAN, W. S. Lightning discharges. (R. 1867)	215
Girard, C.—	
Bibliography of American natural history for 1851	48
Monograph of Cottoids	
GIRARD, C.; BAIRD, S. F. Catalogue of North American reptiles	49
Girard College observations—See A. D. Bache.	
GIVEN, W. F. Remarkable electric phenomenon. (R. 1865)	
Glacial drift (fresh water) of the northwestern States—C. WHITTLESEY	97
Glaciers, traces of, in Massachusetts and Vermont—E. HITCHCOCK	
GLAISHER, J. Account of balloon ascensions. (R. 1863)	
GLAZIER, W. C. W. On destruction of fish in Gulf of Mexico. (P. 1881)	
Gliddon mummy case—C. Pickering	
Globe, winds of the—J. H. COFFIN	268
Glyptocephalus cynoglossus on coast of North America—G. B. Goode; T. H.	
BEAN. (P. 1878)	
Goat, Rocky Mountain, habits of—J. C. MERRILL. (P. 1879)	500
(P. 1881)	467
Gobioid fish, description of a new, from San Diego, California—R. SMITH. (P. 1881)	467
Gold and silver coins, assay of, at United States Mint—J. Pollock. (R. 1968.)	
Gold and silver coins, foreign, table of. (R. 1868)	224
Gold ornament from a mound in Florida, observations on—C. RAU. (R. 1877.) 823, 440, 4	
Goode, G. B.—	
Catalogue of collection to illustrate animal resources and fisheries of the United States, exhibited at Philadelphia, 1876	26
Catalogue of fishes of the Bermudas2	
Catalogue of United States Fish Commission's exhibit at Berlin, 1880 4:	
Classification of collection of animal resources2	
Clupea tyrannus of Latrobe. (P. 1878)	32
Description of new species of amber fish, (Seriola Stearnsii,) obtained from Florida by S. Stearns. (P. 1879)	
Descriptions of new species of fishes from deep soundings on southern New England coast, with diagnoses of two undescribed geners of flounders and genus related to Merlucius. (P. 1880)	
First decade of the United States Fish Commission. (R. 1880)	42
Fishes from Bermuda mistakenly described as new by Günther. $(P. 1878)^{-33}$	32
Fishes from deep waters on south coast of New England obtained by Fish Commission in 1880. (P. 1880)	
Frigate mackerel, (Auxis Rochei,) on New England coast. (P. 1880) 42	25
List of substances derived from animal kingdom	,
Methods of capture and utilization of animals)7

Goode, G. B.—Continued.	
Notacanthus phasganorus, new species of Notacanthidæ from Grand Banks of Newfoundland. (P. 1880)	425
Occurrence of Belone latinanus in Buzzard's Bay, Mass. (P. 1878)	882
Occurrence of Canada porcupine in West Virginia. (P. 1878)	882
Occurrence of Hippocampus antiquorum on St. George's Banks. (P. 1878)	382
Organization and objects of the U.S. National Museum. (P. 1881.) 467,	
Plan of organization and regulations of the U. S. National Museum. (P. 1881)467,	
Plans for installation of collections in the U.S. National Museum	
Preliminary catalogue of fishes of St. John's river and east coast of Florida, with descriptions of new genus and three new species. (P. 1879)	888
Provisional classification of the food collection. (P. 1881) 467,	455
Revision of American species of genus Brevoortia, with description of new species from Gulf of Mexico. (P. 1878)	8 82
Scheme of classification of collections in the U. S. National Museum. (P. 1881)467,	457
Study of trunk fishes, (Ostraciontidæ,) with notes on American species. (P. 1879)	388
Synopsis of useful and injurpus animals	297
Taxonomic relations and geographical distribution of swordfish family, (Xiphiidæ.) (P. 1881)	4 67
Voices of crustaceans. (P. 1878)	882
GOODE, G. B.; BEAN, T. H	
Catalogue of Stearns's fishes of Florida and descriptions of new species. (P. 1879)	888
Catalogue of Velie's fishes in Gulf of Mexico and description of new species. (P. 1879)	838
Craig flounder of Europe (Glyptocephalus cynoglossus) on coast of North America. (P. 1878)	882
Description of Alepocephalus Bairdii, new deep-sca fish, western Atlantic, (P. 1879)	388
Description of Argentina systemsium, new deep-sea fish, Sable Island bank. (P. 1878)	882
Description of Caulolatilus microps from Gulf coast of Florida. (P. 1878)	882
Description of gadoid fishes, Phycis Chesteri and Haloporphyrus viola, from northwestern Atlantic. (P. 1878)	882
Description of Lycodes paxillus. (P. 1879)	888
Description of new genus and species of fish, Lopholatilus chamæleonticeps, from New England. (P. 1879)	888
Description of new species of fish, Apogon pandionis, from mouth of Chesapeake Bay. (P. 1881)	467

GOODE, G. B.; BEAN, T. H.—Continued.	
Description of new species of fish (Lutjanus Blackfordii and Lutjanus Stearnsii) from coast of Florida. (P. 1878)	832
Identity of Brosmius brosme Americanus, Gill, with Brosmius brosme (Müller) White. (P. 1878)	
Identity of Rhinonemus caudacuta with Gadus cimbrius. (P. 1878)	
New genus of fishes, Benthodesmus. (P. 1881)	467
New serranoid fish, Epinephelus Drummond-Hayi, from Bermudas and Florida. (P. 1878)	332
New species of Liparus (L. ranula) from Halifax. (P. 1879)	
Note on Platessa ferruginea and Platessa rostrata. (P. 1878)	
Note upon black grouper (Epinephelus nigritus) of the Southern coast. (P. 1878)	332
Occurrence of Lycodes Vahlii on La Have and Grand Banks. (P. 1879)	833
Oceanic bonito on coast of United States. (P. 1878)	832
Gopher of Illinois, habits of the-J. B. PARVIN. (B. 1854)	75
Gosse, Dr. Report on transactions of the Society of Physics and Natural History of Geneva, 1865-1866. (R. 1866)	214
GOULD, B. A.— Account of astronomical observatory at Cordoba, Argentine Republic.	
(R. 1878)	210
Discussion of Piazzi's astronomical observations. (R. 1863)	
List of scientific papers of A. D. Bache. (R. 1870) 244, 329,	
Report on history of discovery of Neptune	18
Report on history of discovery of Neptune Transatlantic longitude	18 223
Report on history of discovery of Neptune Transatlantic longitude GOULD, J. Acknowledgment of birds. (R. 1867)	18 223
Report on history of discovery of Neptune	18 223 21.:
Report on history of discovery of Neptune Transatlantic longitude GOULD, J. Acknowledgment of birds. (R. 1867) Government Museum—See Museum. Government of Bremen, exchange system. (B. 1865)	18 223 21 204
Report on history of discovery of Neptune Transatlantic longitude GOULD, J. Acknowledgment of birds. (R. 1867) Government Museum—See Museum. Government of Bremen, exchange system. (R. 1865) Graham, T., scientific work of—W. Odling. (R. 1871)	18 223 21.:
Report on history of discovery of Neptune Transatlantic longitude	18 223 21:- 209 249
Report on history of discovery of Neptune Transatlantic longitude	18 223 21:- 204 249 275
Report on history of discovery of Neptune Transatlantic longitude Grould, J. Acknowledgment of birds. (R. 1867) Government Museum—See Museum. Government of Bremen, exchange system. (R. 1865) Graham, T., scientific work of—W. Odling. (B. 1871) Grammar and dictionary of the Carib or Karif language—C. H. Berendt. (R. 1878) Grammar and dictionary of the Dakota language—S. R. Riggs	18 223 21 204 249 275 40
Report on history of discovery of Neptune Transatlantic longitude	18 223 21 204 249 275 40
Report on history of discovery of Neptune Transatlantic longitude	18 223 215 209 249 275 40 98
Report on history of discovery of Neptune Transatlantic longitude	18 223 215 209 249 275 40 98
Report on history of discovery of Neptune Transatlantic longitude	18 223 215 209 249 275 40 98 425
Report on history of discovery of Neptune Transatlantic longitude GOULD, J. Acknowledgment of birds. (R. 1867) Government Museum—See Museum. Government of Bremen, exchange system. (R. 1865) Graham, T., scientific work of—W. Odling. (R. 1871) Grammar and dictionary of the Carib or Karif language—C. H. Berendt. (R. 1878) Grammar and dictionary of the Dakota language—S. R. RIGGS Grammar and dictionary of the Yoruba language—T. J. Bowen Grammar of Yoruba language—W. W. Turner Grand Bank, new species of Notacanthidæ from—G. B. Goode. (P. 1880) Grand Bank, occurrence of Lycodes Vahlii on—G. B., Goode; T. H. Bean. (P. 1879) Grand Manan, marine invertebrata of—W. STIMPSON	18 223 215 209 249 275 40 98 425 333 50
Report on history of discovery of Neptune Transatlantic longitude GOULD, J. Acknowledgment of birds. (R. 1867) Government Museum—See Museum. Government of Bremen, exchange system. (R. 1865) Graham, T., scientific work of—W. Odling. (R. 1871) Grammar and dictionary of the Carib or Karif language—C. H. Berendt. (R. 1878) Grammar and dictionary of the Dakota language—S. R. RIGGS Grammar and dictionary of the Yoruba language—T. J. Bowen Grammar of Yoruba language—W. W. Turner Grand Bank, new species of Notacanthidæ from—G. B. Goode. (P. 1880) Grand Bank, occurrence of Lycodes Vahlii on—G. B., Goode; T. H. Bean. (P. 1879) Grand Manan, marine invertebrata of—W. Stimpson Grant, E. M. Account of discovery of stone image in Tennessee. (R. 1870.)	18 223 21:- 20:- 249 275 40 98 425 333 50 244
Report on history of discovery of Neptune Transatlantic longitude GOULD, J. Acknowledgment of birds. (R. 1867) Government Museum—See Museum. Government of Bremen, exchange system. (R. 1865) Graham, T., scientific work of—W. Odling. (R. 1871) Grammar and dictionary of the Carib or Karif language—C. H. Berendt. (R. 1878) Grammar and dictionary of the Dakota language—S. R. RIGGS Grammar and dictionary of the Yoruba language—T. J. Bowen Grammar of Yoruba language—W. W. Turner Grand Bank, new species of Notacanthidæ from—G. B. Goode. (P. 1880) Grand Bank, occurrence of Lycodes Vahlii on—G. B., Goode; T. H. Bean. (P. 1879) Grand Manan, marine invertebrata of—W. Stimpson Grand Manan, marine invertebrata of—W. Stimpson Grand F. M. Account of discovery of stone image in Tennessee. (R. 1870.) Grasshopper, means of destroying—V. Motschulsky. (R. 1858)	18 223 21: 20+ 249 275 40 98 425 333 50 244 109
Report on history of discovery of Neptune Transatlantic longitude GOULD, J. Acknowledgment of birds. (R. 1867) Government Museum—See Museum. Government of Bremen, exchange system. (R. 1865) Graham, T., scientific work of—W. Odling. (R. 1871) Grammar and dictionary of the Carib or Karif language—C. H. Berendt. (R. 1878) Grammar and dictionary of the Dakota language—S. R. Riggs Grammar and dictionary of the Yoruba language—T. J. Bowen Grammar of Yoruba language—W. W. Turner Grammar of Yoruba language—W. W. Turner Grand Bank, new species of Notacanthidæ from—G. B. Goode. (P. 1880) Grand Bank, occurrence of Lycodes Vahlii on—G. B., Goode; T. H. Bean. (P. 1879) Grand Manan, marine invertebrata of—W. Stimpson Grand Manan, marine invertebrata of—W. Stimpson Grasshopper, means of destroying—V. Motschulsky. (R. 1858) Grasshopper, wingless, of California—E. P. Vollum. (R. 1860)	18 223 21:- 209 249 275 40 98 425 833 50 244 109 147
Report on history of discovery of Neptune Transatlantic longitude GOULD, J. Acknowledgment of birds. (R. 1867) Government Museum—See Museum. Government of Bremen, exchange system. (R. 1865) Graham, T., scientific work of—W. Odling. (R. 1871) Grammar and dictionary of the Carib or Karif language—C. H. Berendt. (R. 1878) Grammar and dictionary of the Dakota language—S. R. RIGGS Grammar and dictionary of the Yoruba language—T. J. Bowen Grammar of Yoruba language—W. W. Turner Grand Bank, new species of Notacanthidæ from—G. B. Goode. (P. 1880) Grand Bank, occurrence of Lycodes Vahlii on—G. B., Goode; T. H. Bean. (P. 1879) Grand Manan, marine invertebrata of—W. Stimpson Grand Manan, marine invertebrata of—W. Stimpson Grand F. M. Account of discovery of stone image in Tennessee. (R. 1870.) Grasshopper, means of destroying—V. Motschulsky. (R. 1858)	18 223 21: 20+ 249 275 40 98 425 333 50 244 109

Graves, ancient, and shell-heaps of California—P. Schumacher. (R. 1874) _	286
Gravitation, figures of equilibrium of liquid mass withdrawn from action of— See Plateau.	
Gravitation, kinetic theories of-W. B. TAYLOB. (R. 1876) 299,	395
GRAY, A.—	
Address at memorial of J. Henry. (R. 1878) 841	, 856
Biographical memoir of J. Henry. (R. 1878) 841,	407
Biographical notice of W. H. Harvey. (R. 1867)	215
Memoir of John Torrey. (R. 1878)	275
Plantse Wrightianse Texano-Neo-Mexicanse. Part I	22
Plants: Wrightians: Texano-Neo-Mexicans:. Part II	42
Plants of Arctic America	342
Plants of Kerguelen Island	294
Report on Lindheimer's, Fendler's, and Wright's botanical explorations	
in New Mexico and California. (R. 1849)	21
GRAY, A., and others. Report on organization of Smithsonian Institution. (R. 1853)	67
GRAY, A.; COPPÉE, H. Report of Committee of Regents on Smithsonian Museum. (B. 1874)	286
GRAY, A.; SARGENT, A. A.; CLYMER, H. Report of Committee on Museum. (R. 1876)	299
Gray, J. E.; DE CANDOLLE, A. Dominant language for science. (R. 1874.)	286
Gray substance of medulla oblongata, and trapezium—J. DEAN	173
Great Britain, history of the Royal Institution of—E. MAILLY. (R. 1867)	215
Greece, Museum of National University of, exchange of specimens. (R. 1867.)	215
Greece, ornamental album from—Miss E. B. Contaxaki. (R. 1857)	107
Green, J. Account of a new barometer. (R. 1855) 77	
GREEN, J.; WÜRDEMANN, W. On filling berometer tubes. (R. 1859)	110
	148
Green's standard barometerGreenland—	140
account of cryolite of—Lewis; Quale. (R. 1866)	214
observations in—See Kane.	
report of explorations in—L. Kumlien. (R. 1878)	841
Green river country, meteorology of—Col. Collins. (B. 1871)	249
Green river valley, rock engravings on—J. G. BRUFF. (R. 1872)	271
Grenada, catalogue of Ober's collection of birds of—G. N. LAWRENCE. (P. 1878)	882
Grinnell expedition to Arctic seas—See Kane.	
Grossmann, F. E. Pima Indians of Arizona. (R. 1871)	249
Guadeloupe, catalogue of Ober's collection of birds of—G. N. LAWRENCE. (P. 1878)	882
Guanajuato, notes on Dugès' fishes from—D. S. JORDAN. (P. 1879)	888
Quatemals—	555
antiquities in—G. WILLIAMSON. (B. 1876)	299

ALPHABETICAL INDEX

Guatemala—Continued.
collection of historical documents in-C. H. Berendt. (R. 1876)
earthquakes in—A. Canudas. (R. 1858)
new humming bird (Atthis Ellioti) from-R. RIDGWAY. (P. 1878)
sculptures of Santa Lucia Cosumalwhuapa in—S. HABEL
Guatemalan minister, A. T. De Irisarri, letter from, recommending Dr. Berendt. (R. 1865)
Guest, W. E. Ancient Indian remains near Prescott, Canada West. (R. 1856)
Guest, W. E. Snow gauge. (R. 1858)
Guide to the flora of Washington and vicinity—L. F. WARD
Guild, R. A. Biographical notice of Chas. C. Jewett. (R. 1867)
Gulf of California, lecture on shells of—P. P. CARPENTER. (R. 1859)
Gulf of California, list of fishes of—D. S. JOEDAN; C. H. GILBERT. (P. 1881) Gulf of Mexico—
analysis of water destructive to fish in—F. M. Endlich. (P. 1881)
destruction of fish in-W. C. W. GLAZIER. (P. 1881)
destruction of fish in-J. Y. PORTER. (P. 1881)
fish mortality in—E. INGERSOLL. (P. 1881)
fish mortality in-M. A. MOORE. (P. 1881)
notes on mortality among fishes in—S. H. Johnson. (P. 1881)
report on water from-W. G. Fablow. (P. 1881)
Gulf of Mexico—See Mexico.
Gulf-stream, formation of clouds over-H. M. BANNISTER. (R. 1866)
Gun-cotton and gunpowder—Lieut. von Karolyi; B. F. Craig. (R. 1864.)
GUNN, D. Egging expedition to Shoal Lake, Lake Winnipeg. (R. 1867)
GUNN, D. Indian remains in Red River Settlement, Hudson's Bay Territory. (R. 1867)
Guthrie, J., authority given by, to collectors of customs, etc., to receive and transmit specimens to Smithsonian
Gutta-percha and caoutchouc. (R. 1864)
GUYOT, A.—
Directions for meteorological observations. (R. 1855) 77, 19, 1
Meteorological tables
Meteorological and physical tables1
Metric system for scientific observations. (R. 1848)
Mountain measurements. (R. 1862)
GUYOT, A.; HENRY, J. Directions for meteorological observations. (B. 1865.)
Gyroscope, problems of rotary motion presented by-J. G. BARNARD 21

H.

Haarlem, Holland, Society of Sciences of—See Harlem.	
HAAS, H. Account of lightning discharges. (R. 1867)	215
HABEL, S. Sculptures of Santa Lucia Cosumalwhuapa in Guatemala 2	69
Habits of—	
beaver—F. R. Brunot. (R. 1878)	275
black bass of the Ohio—J. EOFF. (R. 1854)	75
gopher of Illinois—J. B. PARVIN. (R. 1854)	75
pouched rat or salamander of Georgia-W. GESNER. (R. 1860)	47
Rocky Mountain goat—J. C. MERRILL. (P. 1879)	388
species of salamander—C. MANN. (R. 1854)	75
Hæmatoma, bibliography of—W. KEEN	300
HAGEN, G. Memoir of Encke. (R. 1868)	24
Hagen, H.—	
Neuropters. (R. 1861) 1	49
	294
Synopsis of North American neuroptera1	34
Haidah Indians of Queen Charlotte Islands—J. G. SWAN 20	67
HAIDINGER, W. Honorary medal to von Martius. (R. 1863) 1	87
Hail, on the electricity of induction in-F. ZANTEDESCHI. (R. 1870) 2	44
Hail-stones in Georgia, remarkable forms of-S. ABICH. (R. 1869) 2	28
Hail-storm in Texas-G. M. BACHE. (R. 1870) 2	44
	44
HAILE, J.; MCHENRY, J. W. Antiquities of Jackson Co., Tenn. (R. 1874.) 2	86
Hainault, Belgium, Society of Science, Art, and Literature. Prize questions. (R. 1878)	:75
Hake, description of a new-T. H. BEAN. (P. 1880)	25
HALDEMAN, S. S. Coleoptera of the United States	62
HALDEMAN, S. S. On a polychrome bead from Florida. (R. 1877) 823, 46)4
HALE, C. Exploration of the Nile. (R. 1865) 2	:09
HALE, E. E. Report on Jewett's general stereotype catalogue of public libraries.	47
HALE, H. Vocabulary of Chinook jargon 1	61
Half-breeds of the northwest—V. HARVARD. (R. 1879)	45
Haliclystus auricula, anatomy and physiology of-H. J. CLARK 2	42
Halifax, description of fish obtained at—See G. B. GOODE; T. H. BEAN.	
Hall in Smithsonian building, report on use of-L. Agassiz. (R. 1867) 2	15
Hall, C. F., scientific instructions to-J. Henry; J. E. HILGARD; S. NEW-	
COMB; S. F. BAIRD; F. B. MEEK; L. AGASSIZ. (R. 1871) 2	49
Hall, J, statement of, on telegraph. (R. 1857) 107, 1	15
Halliwell manuscripts, report on -C. C. JEWETT. (R. 1852)	57

Haloporphyrus viola from deep-sea fauna of northwestern Atlantic—G. B. GOODE; T. H. BEAN. (P. 1878)	32
Hamburg Zoological Gardens, exchange of specimens. (R. 1867) 2	15
Hamilton College, Clinton, N. Y., exchange of specimens. (R. 1861)	49
Hamilton College, Clinton, N. Y., examination of Spencer's telescope. (R.	77
Hamilton, J. Bequest deposited in United States Treasury. (R. 1878) 275.3	29
Hamilton, J. Bequest of \$1,000 to Smithsonian Institution. (R. 1872) 271, 3:	×
HAMLIN, H. Address on death of Joseph Henry	
HAMLIN, H. Biographical notice of S. P. Chase. (R. 1878)	75
Hampton, Virginia, Normal and Agricultural Institute, casts of heads of Indian	
boys and girls at—R. H. PRATT. (P. 1879)	33
Hann, J., reply to criticisms of, by W. FERREL. (B. 1877) 828, 33	
Hann, J.—	
Atmospheric pressure and rain-fall. (R. 1877)	98
Diminution of aqueous vapor with increasing altitude. (R. 1877) 323, 33	
Influence of rain upon the barometer. (R. 1877) \$23, 33	
Laws of variation of temperature in ascending currents of air. (R. 1877) 823, 33	
Relation between pressure of air and velocity of wind. (R. 1877) 823, 30	
Harbors, dates of opening and closing of—F. B. Hough	
Harbors, tides and tidal action in—J. E. HILGARD. (R. 1874) 286, 39	0
HARDISTY, W. L. The Loucheux Indians. (R. 1866)	
HARE, R.—	
Explosiveness of nitre	7
Letter relative to gift of apparatus. (R. 1848)	1
Method of forming small weights. (R. 1858) 10	9
On John Wise's observation of a thunder storm. (R. 1854)	
HARGER, O. Notes on New England Isopoda. (P. 1879)	3
HARKNESS, W. Observations on terrestrial magnetism and deviation of compasses on iron-clads	9
Harlem, Holland, Society of Science. Prize questions. (R. 1861, 1864, 1868, 1878)149, 188, 224, 276	
Harmonies of the solar system—S. ALEXANDER)
HARRIOTT, J. Account of the city of Washington, 1808	0
HARRIS, E. List of birds and mammalia of Missouri river. (R. 1850) 2	8
HABBISON, A. M. Colored bead from mound in Florids. (R. 1877) 32	\$
HARRISON, W. Account of storm in Butler county, Kansas. (R. 1871) 245)
HART, J. N. Dr. Mounds and osteology of the mound builders of Wisconsin. (R. 1877)	į
Harvey, W. H., biographical notice of, by A. GRAY. (R. 1867) 216	í
HARVEY, W. H.—	
Lecture on marine algse. (R. 1855)	,
List of austic alam 96	

OF SMITHSONIAN PUBLICATIONS.	201
HABVEY, W. H.—Continued.	
Marine alge. Part I. Melanospermees	32
Marine alga. Part II. Rhodospermeas	44
Marine alga. Part III. Chlorospermea	9!
Marine algae. Parts I, II, III. Complete	90
Harwood, A. A. Account of sarcophagus from Syria. (R. 1870)	24
Hassler expedition, narrative of the—L. Agassiz. (R. 1872)	27
HASTINGS, C.S.; HOLDEN, E.S. Synopsis of Herschel's writings. (R. 1880.) 442,	
HATCH, F. W. Meteorological observations at Sacramento, Cal. (R. 1854)	7
HAUGHTON, S. Relation of food to work, and its bearing on medical practice. (R. 1870)	24
Haux, René Just, memoir of, by G. CUVIER. (R. 1860)	14
HAVARD, V. French half-breeds of the northwest. (R. 1879)	84
HAVEN, S. F. Archæology of the United States	7:
HAVEN, S. F. Report on Jewett's general stereotype catalogue of public libraries	4
Hawaiian Islands, fishes of—T. H. STREETS	80
Hawaiian Islands, natural history of-T. H. STREETS	80
Hawks, G. W	
Determination of feldspar in thin sections of rocks. (P. 1881)	46
On mineralogical composition of normal mesozoic diabase upon Atlantic border. (P. 1881)	46
Progress in geology and mineralogy, 1879-80. (R. 1880) 442,	
HAWKES, F. L., and others. On publication of Spanish works on New Mexico. (R. 1855)	7
Hawkins and Wale, blow-pipe apparatus of. (R. 1872)	27
HAY, O. P. On a collection of fishes from eastern Mississippi. (P. 1880)	42
HAYDEN, F. V. Notes on Indian history. (R. 1867)	21
	172
HAYES, I. I.—	
Astronomical observations in Arctic Seas	19
Atmospheric pressure, wind, temperature, tides, in Arctic Seas	19
Lecture on Arctic explorations. (R. 1861)	14
Magnetic observations in Arctic Seas	19
Physical observations in the Arctic Seas	196
Haystack mound, Lincoln county, Dakota—A. BARRANDT. (R. 1872)	27
HEAD, J. E. Natural history of country about Fort Ripley, Minn. (R. 1854)	7
Health, architecture in relation to—D. B. REID. (R. 1856)	9
Hearing, sense of. (R. 1866)	214
Heart, strain and over-action of the. Toner lecture No. 111—J. M. DA COSTA.	
Heat—	
and light of the sun, relative intensity of-L. W. MEECH. (R. 1856.) 91	88
animal, of fishes, experiments upon-J. H. Kidder. (P. 1879)	888

Heat—Continued.
dry, comparative action of sulphurous acid and, upon putrefactive bacteria. (P. 1881)
principles of the mechanical theory of-J. MÜLLER. (R. 1868)
radiant, report on state of knowledge of-B. Powell. (R. 1859)
recent progress in relation to the theory of—A. CAZIN. (R. 1868)
specific, table of—F. W. CLABKE
tables of expansion by-F. W. CLARKE
Heavenly bodies, results of spectrum analysis applied to—W. Huggins. (R. 1866)
Heights, circular relative to—J. HENRY
Heights of mountains in Colorado—G. ENGELMANN. (R. 1862)
Heights, quantity of rain at different-O. W. Morris; J. Henry. (R. 1855.)
HEILPRIN, A. New species of eocene mollusca from southern United States. (P. 1880)
HEINEMANN, V. Lepidoptera
Helicinidæ—W. G. BINNEY
Heligoland, birds of-H. Gärke. (P. 1879)
Hellwald, F. von. American migration. (R. 1866)
HELMHOLTZ, H. Relation of physical sciences to science in general. (R. 1871.)
HELMHOLTZ, H.; MAXWELL, J. C. Later views of connection of electricity and magnetism. (R. 1878)
Hemiptera, instructions for collecting-P. R. UHLER. (R. 1858)
Hemirhamphus rosm from coast of California, description of—D. S. JORDAN; C. H. GILBERT, (P. 1880)
Henry and the telegraph—W. B. TAYLOB. (R. 1878)
Henry as a discoverer—A. M. MAYER
Henry, J.—
biographical memoir of, by A. GRAY. (R. 1878) 841,
deposition of, in the case of Morse vs. O'Rielly, taken September, 1879
examination of, by English Scientific Commission.
life and character of, by J. C. Welling
memoir of, and sketch of his scientific work, by W. B. TAYLOR 356,
memorial of, by United States Congress
monument to, act of Congress for
report of Special Committee of Board of Regents on communication of— C. C. Felton. (R. 1857)107,
statue of, proceedings of Congress relative to
statue of, report of Executive Committee on. (R. 1880)
HENRY, J., papers by:
Account of Priestley's lens. (R. 1859)
Acoustics applied to public buildings. (R. 1856)
Address on the Smithsonian Institution. (R. 1858)

OF SMIINSONIAN PUBLICATIONS.	
NRY, J., papers by:—Continued.	
Annual reports, 1865-1877 (reprints)	
Annual reports, 1847-1877—See each annual report.	
Appendix to notice of Schoenbein, the discoverer of ozone. (B. 1868).	-
Aurora directions. (B. 1855)	-
Circular of instructions to observatories relative to telegraphic announce	}-
ment of astronomical discoveries.	
Circular relating to collections in archeology and ethnology	
Circular relative to ancient mining in Lake Superior copper region. (R	
Circular relative to collections of birds from Middle and South America	۱.
Circular relative to earthquakes	
Circular relative to heights	
. Circular respecting copyrights. (R. 1854)	-
Circular respecting new report on libraries. (R. 1854)	
Circular sent with specimens presented. (R. 1872)	
Circular to entomologists	
Circular to officers of Hudson's Bay Company	-
Color-blindness. (B. 1877)	_
Communication relative to a publication by Prof. Morse. (R. 1857.) 107, 82 Currents of atmosphere and aerial navigation. (R. 1860)	
Description of Smithsonian anemometer. (R. 1860)	-
Digest of act of organization of Smithsonian Institution	C
Directions for constructing lightning-rods	-
Earthquake directions. (R. 1855)	•
Effect of the moon on the weather. (R. 1871)	
Eulogy on A. D. Bache. (R. 1870) 24	4,
Exposition of bequest of Smithson	
History of electro-magnetic telegraph. (R. 1857) 10	7,
Instructions for observations of thunder storms	
Investigation of illuminating materials. (R. 1880) 44:	
Letter to Secretary of Treasury on payment of interest in coin. (R. 1865	•
Memoir of W. W. Seaton. (R. 1866)	
Meteorological stations, cost of establishing. (R. 1858)	
Mode of testing building materials and account of the marble used in the extension of the United States Capitol. (R. 1856)	-
Notes relative to George Catlin. (R. 1872)	
Notes to accounts of lightning discharges by G. W. Dodge and others (R. 1867)	
Notes to article on American migration by F. von Hellwald. (R. 1866.)
Notes to article on horary variations of barometer by M. Vaillant. (R	
1866)	

HENRY, J., papers by :Continued.	
Notes to article on vitality by H. H. Higgins. (R. 1866)	. 214
Notes to articles on meteorology by G. Latimer and others. (R. 1871.	
Notice of Parker Cleaveland. (R. 1859)	
On a physical observatory. (R. 1870)	
On the "Moon hoax." (R. 1878)	
Organization of local scientific societies. (R. 1875)	
Programme of organization of Smithsonian	
Queries relative to tornadoes	
Registration of periodical phenomena. (R. 1855)	
Remarks appended to meteorological articles by W. C. Dennis and others (R. 1866)	i.
Reply to memorial on Lowe's aeronautic voyage. (R. 1860)	
Report of Secretary, 1847-1877. (R. 1847-1877.) H , I , 21, 28, 51, 57, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 214, 215, 224, 22 249, 271, 275, 286, 298, 299, 328	8, 244.
Researches in sound. (R. 1878)	ı, 406
Statement in relation to the history of the electro-magnetic telegraph.	
Suggestions relative to objects of scientific investigation in Russian	1 _ 207
Syllabus of course of lectures on physics. (R. 1856)	
HENRY, J., and others. Scientific instructions to Captain Hall. (R. 1871)	249
HENRY, J.; Du Pré, W. Earthquakes in North Carolina, 1874. (R. 1874).	
HENRY, J.; GUYOT, A. Directions for meteorological observations. (R. 1855. 77, 19)
HENRY, J.; KNIGHT, R. T. Connection of gales of wind and aurora. (R 1871)	249
HENRY, J.; MORRIS, O. W. Quantity of rain at different heights. (R. 1855.)	
HENRY, J.; RUSSELL, R. Lectures and notes on meteorology. (R. 1854)	. 75
HENRY, J.; WALLACH, R. Report of Committee of Regents on fire at the	ŧ
Smithsonian. (R. 1864)	8, 3 29
Henshall, J. A., notes on collection of fishes obtained by, from east Florids— D. S. JORDAN. (P. 1880)	
Herbarium captured in Tennessee—H. R. Wirz. (R. 1862)	150
Herpetology of Hawaiian and Fanning Islands—T. H. STREETS	303
Herschel, Sir J. F. W., memoir of, by N. S. Dodge. (R. 1871)	249
Therefore, Sir J. F. W. On stome (P. 1989)	150
HERSCHEL, Sir J. F. W. On atoms. (R. 1862)	244
Herschel, Sir W., sulogy on, by F. Arago(R. 1870)	
HASTINGS. (R. 1880) 442,	
Hexanchus corinus, new species of notidanoid shark, description of—D. S. JOB-DAN; C. H. GILBERT. (P. 1880)	4%
HIGGINS. H. H. On vitality, with notes by J. Henry. (R. 1866)	214
ALTOGRAMO. AL. AL. WILLYING WILL HUNDOUT U. ALGILLY. 15, 1000	

High antiquity in Europe, study of—A. Morlor. (R. 1862; R. 1864) 156	
HILDRETH, S. P. Meteorological observations, Marietta, Ohio	12
HILGARD, E. W. Geology of lower Louisiana and salt deposit on Petite Anse Island	21
HILGARD, J. E. Description of magnetic observatory at the Smithsonian Institution. (B. 1859)	1
HILGARD, J. E. Tides and tidal action in harbors. (R. 1874) 286	39
HILGARD, J. E., and others. Scientific instructions to Capt. Hall. (R. 1871.)	24
HILL, A. J. Map of antiquities of United States. (R. 1861)	14
HILL, G. W. Ancient earthworks of Ashland county, Ohio. (R. 1877) HILL, G. W. Antiquities of northern Ohio. (R. 1874)	8: 2:
HILL, T. Map of solar eclipse of March 15, 1858, and description of the occultator	10
HILLIER, S. L. Effect of lightning. (R. 1866)	2
Hincks, W. On Smithsonian exchanges. (R. 1860)	14
Hints on public architecture—R. D. OWEN	17
Hippocampus antiquorum, occurrence of, on St. George's Bank—G. B. Goode.	
(P. 1878)	88
Hippoglossoides elassodon, new species of flounder, from Puget Sound, description of—D. S. Jordan; C. H. Gilbert. (P. 1880)	42
Hippoglossoides exilis, new flounder, from coast of California, description of-	
D. S. JORDAN; C. H. GILBERT. (P. 1880)	42
Hippoglossus vulgaris, occurrence of, at Unalaska and St. Michael's, Alaska— T. H. Bran. (P. 1879)	88
Historical documents in Guatemala—C. H. BERENDT. (R. 1876)	29
Historical manuscripts, catalogue of collection of—L. Berlandier. (R. 1854)	7
Historical sketch of Academy of Sciences of Paris—M. Flourens. (R. 1862.)	18
Historical sketch of Henry's contribution to electro-magnetic telegraph—W. B. TAYLOR. (R. 1878)	
Historical sketches of vestiges of antiquity in United States—S. F. HAVEN	7
Historical Society of Minnesota, grammar and dictionary of Dakota language.	4
Historical study of jade—S. Blondel. (R. 1876)	29
History—	20
ancient, of North America—M. MUCH. (R. 1871)	24
and antiquities of Santa Rosa Island, California—S. Bowers. (R. 1877.)	82
and archeology, Mexican—B. MAYER	8
and climate of New Mexico-T. A. McParlin. (R. 1877) 823,	39
and geology, boundary line between—E. Surss. (R. 1872)	27
and origin of Royal Society of London-C. A. ALEXANDER. (R. 1868.)	18
Indian, notes on-F. V. HAYDEN. (R. 1867)	21
of Chester county, Pennsylvania-W. Darlington. (R. 1862)	16
of discovery of planet Neptune—B. A. Gould	1
of discovery relative to magnetism. (R. 1863)	18
14	

History—Continued.
of education in the United States-F. A. PACKARD. (R. 1868)
of electro-magnetic telegraph-J. HENRY. (R. 1857) 107
of fresh-water algo of North America—H. C. Wood241
of marine alge-W. H. HARVEY 32, 43, 95, 96
of my youth, an autobiography—F. Arago. (R. 1870) 244
of North American grasshoppers, circular in reference to
of petroleum or rock-oil—T. S. Hunt. (R. 1861) 149
of Royal Institution of Great Britain—E. MAILLY. (R. 1867) 215
of Smithsonian Institution—W. J. RHEES
of transactions of Anthropological Society of Paris, 1865-1867-P.
Broca. (R. 1868)
of works of Cuvier-M. FLOURENS. (B. 1868) 224
HITCHCOCK, E.—
Illustrations of surface geology 90
On erosions of earth's surface
On geology of New England 90
Traces of glaciers in Massachusetts and Vermont
Hoax, the Moon—J. HENRY. (R. 1878)
Hodge, C. Prayer at funeral of Joseph Henry
Hodgins, J. G. Meteorological stations of Upper Canada. (R. 1858) 109
Hodgins, J. G. Meteorological system of Canada. (R. 1865)
Hodgkinson, Eaton, memoir of, by R. Rawson. (R. 1868) 224
Hodgson, W. B., and others. On publication of Spanish works on New Mexico. (R. 1855)
HOEK, M. Letter in reference to meteoric shower of November 18, 1867 217
Holcodus, memoir on—R. W. GIBBES
Holden, E. S.—
Index catalogue of books relating to nebulæ, clusters, etc
Recent progress in astronomy 1879-1880. (R. 1880) 442, 427
Reports of American and foreign observatories. (R. 1879) 345
Studies in Central American picture-writing. (E. 1879-'80)
HOLDEN, E. S.; BOEHMER, G. H. Report of observatories for 1880. (R. 1880) 442, 438
HOLDEN, E. S.; HASTINGS, C. S. Synopsis of scientific writings of Sir Wil-
liam Herschel. (R. 1880)
Holland Society of Science, Harlem. Prize questions. (R. 1861, 1864, 1868, 1878)
HOLMGREN, F. Color-blindness in relation to accidents by rail and sea. (R. 1877)
Horary variations of barometer—M. VAILLANT. (R. 1866)
HORNADAY, W. T. Classification of collections to illustrate taxidermy. (P. 1881)
HOTOHKISS, T. P. Indian remains in Caddo parish, Louisiana. (R. 1872) 271

OF SMITHSONIAN PUBLICATIONS.	211
Нотен, F. В.—	
Dates of blossoming of plants	182
Dates of defoliation or fall of leaf of plants	182
Dates of first appearance of birds	182
Dates of first appearance of fish	182
Dates of first appearance of insects	182
Dates of first appearance of reptiles	182
Dates of foliation of plants	182
Dates of opening and closing of lakes	182
Dates of opening and closing of rivers	182
Dates of ripening of fruits	182
Periodical phenomena, 1851 to 1859	182
Hough, J. Mounds in Washington county, Mississippi. (R. 1879)	845
Hours, best, to find mean temperature—C. Dewey. (R. 1857)	107
Howgate Polar Expedition, natural history of Arctic America—L. Kumlien.	
Hoy, P. R. Journal of an exploration of western Missouri. (R. 1864)	188
Hoy, P. R. On Amblystoma luridum, a salamander of Wisconsin. (B. 1854.)	75
HUBBARD, J. S. Investigations of Biela's comet. (R. 1862)	150
, ,	
Hudson's Bay, notes on fishes from—T. H. BEAN. (P. 1881)	467
Hudson's Bay Company. Kennicott's explorations. (R. 1868)	187
Hudson's Bay Company, circular to officers of J. HENRY	
Hudson's Bay Company, letter to officers of—G. SIMPSON	187
Hudson's Bay Territory, observations in—B. R. Ross. (R. 1859)	110
Hudson's Bay Territory, Red river settlements, Indian remains in—D. Gunn. (R. 1867)	215
Huggins, W. Results of spectrum analysis applied to the heavenly bodies.	214
(R. 1866)	
Human race, probable future of the—A. DE CANDOLLE. (B. 1875)	298
Human remains from Patagonia—A. RIED. (R. 1862)	150
Humidity, relative, tables of—J. H. Coffin	87
Humming-bird, a new, (Atthis Ellioti), from Guatemala—R. RIDGWAY. (P. 1878)	882
Humming-birds, described, list of-D. G. Ellior	334
Humming-birds, monograph of—D. G. ELLIOT	
HUMPHREYS, A. A. Method of ascertaining amount of water in rivers. (R. 1858)	109
Hungarian National Museum, acknowledgment for birds. (R. 1863)	187
Hungary, prehistoric antiquities of—F. F. Romes. (R. 1876) 299, 440,	
HUNT, T. S. Chemistry of the earth. (R. 1869)	
Hunt, T. S. History of petroleum or rock oil. (R. 1861)	149
Huntington, G. C. Climate of Kelley's Island, Ohio. (R. 1866)	214
Hurricane at Tortola, St. Thomas, and Porto Rico—G. A. LATIMER. (R. 1867.)	215
Hurricane in Island of St. Thomas. (R. 1867)	215

•

•

HUXLEY, T. H. Principles and methods of palsontology. (R. 1869)	228
Hydrobiinse, researches upon the—W. STIMPSON	
Hydrogen as a gas and as a metal—J. E. REYNOLDS. (R. 1870)	244
Hygrometrical tables—A. GUYOT	154
Hymenoptera—	
instructions for collecting—B. CLEMENS. (R. 1858)	109
letter relative to catalogue of-H. DE SAUSSURE. (R. 1862)	150
monograph of-H. DE SAUSSURE	254
of Arctic regions—S. H. SCUDDER and others	342
Hypsometrical tables—A. Guyot	158

I.

Ice—	
disappearance of—R. H. GARDINER. (R. 1860)	
formation of, at the bottom of water—M. Engelhardt. (R. 1866)	
observations on—D. WALKER	146
Iceland, vegetable colonization of—C. MARTINS. (R. 1858)	100
Ichthyology—	
Contributions to North American-	
Part I. Review Rafinesque's memoirs—D. S. JORDAN	305
Part II. A. Notes on Cottidæ, (etc.) B. Synopsis Siluridæ—D. S. Jordan	
Part III. A. Distribution of fishes of Alleghany region, (etc.)—D. S. JORDAN; A. W. BRAYTON. B. Synopsis of family Catostomidæ—D. S. JORDAN	
of Hawaiian and Fanning Islands and California-T. H. STREETS	303
status of-T. GILL	247
Ichthyology—See Fishes.	
Icichthys Lockingtoni, new species deep-water fish from coast of California- D. S. JORDAN; C. H. GILBERT. (P. 1880)	
Icosteus senigmaticus of California, description of—W. N. Lockington. (P. 1880)	
Icterus, description of new species of, from West Indies—G. N. LAWRESCE. (P. 1880)	
Idaho, distribution of forest trees in—W. W. JOHNSON. (B. 1870)	
Identification of artisan and artist—N. WISEMAN. (R. 1870)	
Illinois—	
Adams county, mounds in, near Quincy-W. G. ANDERSON. (B. 1879.)	145
Albany, mounds near-R. J. FAROUHARSON. (R. 1874)	

Illinois—Continued.
deposits of flint implements in-J. F. SNYDER. (R. 1876) 299
habits of the gopher of-J. B. PARVIN. (R. 1854)
Lawrence county, antiquities of—A. PATTON. (R. 1878)
Madison county, stone cists in, near Highland—A. Ornier. (R. 1879.) 845
Mason county, antiquities of-J. Cochrane. (R. 1877) 828
Mercer county, ancient mounds of—T. McWHORTER. (R. 1874) 286
Pike county, mounds in-B. MITCHELL. (R. 1879) 845
Rock Bluff, description of human skull from-J. A. MEIGS. (R. 1867.) 215
Rock Island, shell-bed skull from—A. S. TIFFANY. (R. 1874) 286
Rock Island county, mounds in—T. THOMPSON. (R. 1879) 845
Rock Island county, antiquities of—A. TOELLNER. (R. 1879) 845
Rock river valley, mound builders in—J. SHAW. (R. 1877) 823
Spoon river valley, mounds in-W. H. ADAMS. (R. 1879) 845
southern, agricultural flint implements in—C. RAU. (R. 1868.) 224, 440, 370
Union county, antiquities of—T. M. PERRINE. (R. 1878) 275
Union county, mounds in, near Anna—T. M. PERRINE. (R. 1872) 271
Whitesides county, antiquities of—W. H. Pratt. (R. 1874) 286
Illinois Board of Education. Meteorological system for every State. (R. 1855.) 77
Illuminating materials, investigation of—J. HENRY. (R. 1880)
Illustrations of method of recording Indian languages—J. O. Dorsey; A. S.
GATSCHET; S. R. RIGGS. (E. 1879-80)
Illustrations of surface geology—E. HITCHCOCK 90
Image, stone, in Tennessee, account of discovery of-E. M. GRANT. (R. 1870.) 244
Imperial Academy of Sciences, Belles-lettres, and Arts, of Bordeaux. Prize
questions. (R. 1868) 224
Imperial Academy of Sciences, Vienna. Prize questions. (R. 1864; R. 1865.) 188, 209
Imperial Geological Institute, Vienna, tertiary fossils presented by. (R. 1863.) 187
Imperial Library of Vienna, books presented by. (R. 1865) 209
Imperial Society of Natural Sciences of Cherbourg. Prize questions. (R. 1864.) 188
Imperial Society of Science, Agriculture, and Arts of Lille. Prize questions.
(R. 1865)
Implement, wood, ancient, found in Connecticut—E. W. Ellsworth. (R. 1876)
Implements—
agricultural, North American stone period—C. RAU. (R. 1863) 187, 440
flint, agricultural, in southern Illinois—C. RAU. (R. 1868) 224, 440, 370
flint, deposits of, in Illinois—J. F. SNYDER. (R. 1876) 299
flint, in Holmes county, Ohio—H. B. CASE. (R. 1877)
stone, North American—C. RAU. (R. 1872) 271, 440, 382
Improvements, recent, in chemical arts—J. C. BOOTH; C. MORFIT
Income, majority and minority reports on distribution of—J. A. PEABCE; J. MEACHAM. (R. 1858)

Incorporation, act of, of Smithsonian Institution—See Congress.
Index-catalogue of books and memoirs on nebulæ and clusters-E. S. Holden. 311
Index—
of genera of birds—S. F. BAIRD
of Smithsonian publications 48
systematic and alphabetical, of publications of the Smithsonian Institu- tion to July, 1869. (R. 1868) 24
systematic, to list of foreign correspondents257
to genera of recent and fossil animals—S. H. SCUDDER 470
to names applied to subdivisions of class Brachiopoda-W. H. DALL 304
to North American botany—S. Watson 258
to papers on anthropology published by Smithsonian Institution, 1847- 1878—G. H. BOEHMER. (R. 1879) 346, 421
Indian—
boys and girls at Hampton Normal Institute, Va., catalogue of casts of
heads of—R. H. PRATT. (P. 1879)
burial in California-W. M. KING. (R. 1874)
conjurer's practice, details of—A. S. GATSCHET. (E. 1879-80)
engravings on rocks along Green river valley—J. G. Bruyy. (R. 1872)
forts and dwellings, Indian Territory—W. E. DOYLE. (R. 1876) 29
history, notes on—F. V. HAYDEN. (R. 1867) 215
languages—G. Flachenecker. (R. 1862)
languages, method of recording—J. O. Dorsey; A. S. Gatscher; S. R. Riges. (E. 1879–80)
languages—See G. Gibbs, F. L. O. Roehrig, S. R. Riggs, J. G. Shea, W. L. Hardisty.
linguistics, account of library of-J. G. SHEA. (R. 1861)
linguistics, recommendation of Shea's—G. GIBBS and others. (R. 1860.)
mode of making arrow-heads and obtaining fire—G. CROOK. (R. 1871. 249
mounds in southern Florida, explorations of—S. T. WALKER. (R. 1879) 345
mounds near Fort Wadsworth, Dakota—A. J. Comfort. (R. 1871) 249
philology-W. W. Turner. (R. 1851)
photographs, catalogue of 216
portraits and sketches of scenery, catalogue of—J. M. STANLEY 53
portraits, report of Committee of Regents on Stanley's gallery of. (R. 1857) 107
pottery—C. RAU. (R. 1866)214, 440. 368
prisoners at St. Augustine, Florida, catalogue of casts of heads of—R. H. Pratt. (P. 1878)
relics, catalogue of cabinet of, presented by J. H. Devereux. (R. 1872.)
relics from Schoharie, New York-F. D. Andrews. (R. 1879) 345
remains in Caddo parish, Louisiana—T. P. Hotunkiss. (R. 1872) 271
remains in Red River Settlement, Hudson's Bay Territory—D. Guss. (R. 1867)

Indian—Continued
remains near Prescott, Canada West-W. E. Guzet. (R. 1856)
tribes, cessions of land by, to United States-C. C. ROYCE. (E. 1879-80) 4
tribes of upper Missouri-T. A. Culbertson. (R. 1850)
village (Kushkushkee) near Newcastle, Pennsylvania.—E. M. McCon- NELL. (R. 1871)
vocabularies—G. GIBBS. (R. 1862)1
vocabularies received from the Wheeler survey, list of—G. GIBBS. (R. 1874)
Indian Ocean, cyclone in the—N. Pike. (R. 1867) 2
Indian Territory, Indian forts and dwellings in-W. E. DOYLE. (R. 1876) 2
Indiana—
Allen county, antiquities of—R. S. Robertson. (R. 1874) 2
De Kalb county, antiquities of—R. S. ROBERTSON. (R. 1874)
Franklin county, mounds in—E. R. Quick. (R. 1879)
Knox county, antiquities of—A. PATTON. (R. 1878)
La Porte county, antiquities of—R. S. ROBERTSON. (R. 1874)
Rush county, mounds in—F. Jackman. (R. 1879)
tornado in—J. Chappelsmith
Indians—
aboriginal, of America, language of—G. GIBBS. (R. 1870) 24
Cree, system of relationship of—E. A. WATKINS. (R. 1862) 10
Dakota or Sioux, language of—F. L. O. ROEHRIG. (R. 1871)
Haidah, of Queen Charlotte's Islands, British Columbia—J. G. SWAN 26
Kutchin tribes—S. Jones. (R. 1866)
Loucheux—W. L. HARDISTY. (R. 1866)
Makah—J. G. Swan
mortuary customs of-H. C. YARROW. (E. 1879-80)
mythology of-J. W. Powkll. (E. 1879-80) 4
Navajo, sketch of the-J. Letterman. (R. 1855)
of British America, account of—E. Petitot. (R. 1865) 20
of Cape Flattery, Washington Territory-J. G. Swan 22
of Peru—F. L. GALT. (R. 1877)
of valley of Red River of the North, ethnology of-W. H. GARDNER.
of western Nevada and California, Centennial mission to—S. Powers.
(R. 1876) 20
Pima, of Arizona-F. E. Grossman. (R. 1871) 24
Quillehute. Method of taking surf smelt—J. G. Swan. (P. 1880) 42
sign language of-G. Mallery. (E. 1879-80) 47
Sioux or Dakota—A. G. Brackett. (R. 1876) 29
Sioux or Dakota, language of-F. L. O. Roehrig. (R. 1871) 249, 37

Indians—Continued.
Tinneh or Chepewyan, of British and Russian America—G. GIBBS and others. (R. 1866)
uses of brain and marrow of animals by-T. R. PEALE. (R. 1870) 24
Indians—See California, Cape Flattery, Carib, Dakota, Haidah, Nevada, Kutchin, Loucheux, Makah, Mandan, Maya, Navajo, Peru, Portraits, Red River of the North, Shoshone, Sioux, Tinneh, Tonto.
Induction and deduction—J. VON LIEBIG. (R. 1870) 24
Induction, electricity of, in clouds—F. ZANTEDESCHI. (R. 1870) 244
Inflammation in arteries after ligature, acupressure, and torsion. Toner lecture No. VII—E. O. SHAKESPEARE321
Influence of aurora on the telegraph—W. D. SARGENT. (R. 1870)
Influence of rain upon the barometer—J. HANN. (R. 1877) 323. 379
Infusoria—J. W. Bailey
Infusorial earths, examination of—A. M. EDWARDS
INGERSOLL, E. Fish mortality in Gulf of Mexico. (P. 1881) 45
Inquiries relative to crawfish and other crustacea
Inquiries relative to disease known as chorea—S. W. MITCHELL. (R. 1874) 286
Inquiry relative to food-fishes of United States—S. F. BAIRD 231
Insect instincts and transformations, lecture on-J. G. Morris. (R. 1855)
Insects—
son. (P. 1878) 322
circular concerning the department of—S. F. BAIRD 448
classification of, from embryological data—L. AGASSIZ 16
dates of first appearance of—F. B. Hough
directions for collecting and preserving—A. S. PACKARD 261
instructions for collecting—S. F. BAIRD. (R. 1858)
of Arctic America—L. Kumlien
of Kerguelen Island-C. R. OSTEN SACKEN; H. A. HAGEN 294
Installation of collections in National Museum, plans for—G. B. Goode. (P. 1881)
Instincts and transformations of insects, lecture on-J. G. Morris. (B. 1855) 7
Institute of Bologna, Academy of Sciences of the. Prize questions. (R. 1862.) 150
Institute of France, historical sketch of, by M. FLOURENS. (R. 1862) 150
Institute of Rupert's Land, circular of. (R. 1861)
Institute, Royal Scientific and Literary, of Lombardy. Prize questions. (R. 1865)
Institution of Civil Engineers, London. Prize questions. (R. 1862) 150
Institutions—
domestic, in correspondence with Smithsonian Institution, list of. (R. 1858)67, 69, 238
foreign, in correspondence with Smithsonian Institution, list of. (R. 1864; R. 1865) 188, 202, 64, 154, 225, 243, 309, 469

Institutions—Continued.	
in United States and British Provinces of North America, list of—W. J. RHEES.	116
in which phonography is taught. (R. 1856)	91
scientific and literary, in United States, list of principal	335
Instructions—	
for archeological investigations-G. GIBBS. (R. 1861)	149
for collecting coleoptera—J. L. LE CONTE. (R. 1858)	109
for collecting diatomaces—A. M. Edwards	366
for collecting dipters-H. LOEW; R. OSTEN SACKEN. (R. 1858)	109
for collecting hemipters—P. R. UHLEB. (R. 1858)	109
for collecting hymenopters—B. CLEMENS. (R. 1858)	109
for collecting insects—S. F. BAIRD. (R. 1858)	109
for collecting land and fresh-water shells-J. LEWIS	363
for collecting lepidoptera-B. CLEMENS. (R. 1858)	109
for collecting myriapods, phalangidse, etc.—H. C. Wood. (R. 1866)	214
for collecting nests and eggs of North American birds. (R. 1858)	189
for collecting neuroptera-P. R. UHLER. (R. 1858)	109
for collecting orthoptera—P. R. UHLER. (R. 1858)	109
for observations of thunder storms—J. HENRY	235
for research relative to ethnology and philology of America—G. GIBBS	160
in reference to collecting nests and eggs of North American birds—T. M. Brewer	189
to Captain Hall for Arctic observation—J. HENRY and others. (R. 1871)	249
to observatories relative to telegraphic announcement of astronomical discoveries—J. HENRY	263
Instruments, meteorological,	
articles on, in Smithsonian. (R. 1874)	286
description of—L. CASELLA. (R. 1859)	110
description of—A. Guyor; J. Henry. (R. 1855) 77, 19,	
Integrals, general, of planetary motion—S. Newcomb	
Intensity of heat and light of the sun—L. W. MEECH. (R. 1856) 91	
Interest, letter to Secretary of Treasury on payment of, in coin—J. HENRY.	209
(R. 1865)	188
Intermixture of races—G. GIBBS. (R. 1864)	
Internal structure of the earth—J. G. BARNARD	
International Archeological Congress, Antwerp, 1866. (R. 1866)	214
International code of symbols of archeology—G. De Mortillet; E. Chantre. (R. 1875)	298
International exchange, report on—G. H. BOEHMER	
International Exhibition, Philadelphia—See Centennial.	
International Fisheries Exhibition, Berlin, catalogue of United States exhibit	
at—G. B. Goode	413

Interpolation, methods of, applicable to graduation of irregular series. Parts 1, 11.—E. L. DE FOREST. (R. 1871; R. 1878)	275
Introduction, letters of. (R. 1865)	
Introduction to study of Coptic language-M. KABIS. (B. 1867)	
Invertebrata, marine, from New England coast distributed by United States Fish Commission, list of—A. E. Verrill; R. Rathbun. (P. 1879)	33
Invertebrata, marine, of Grand Manan—W. STIMPSON	50
Invertebrata, marine, of northeastern coast of America, notice of recent additions to—A. E. VERBILL:	
Part I. Annelida, etc. (P. 1879)	
Part II. Mollusca, with notes on Annelida, etc. (P. 1880)	
Part III. Catalogue of Mollusca recently added to fauna of southern New England. (P. 1880)	
Invertebrate fossils— check-list of—T. A. Conrad	200
from Arkansas, Wyoming, Colorado, and Utah, descriptions of new- C. A. WHITE. (P. 1880)	
of North America, check-list of—F. B. MEEK	183
report on-F. B. MEEK; F. V. HAYDEN	172
Invertebrates, marine, distributed by U. S. National Museum, list of—R. BATE-BUN. (P. 1881)465, 467,	47]
Invertebrates-See Marine invertebrates.	
Investigation of—	
anatomy and physiology of the rattlesnake—S. W. MITCHELL	135
disturbances of horizontal component of magnetic force—A. D. BACHE	
disturbances of vertical component of magnetic force-A. D. BACHE	175
eleven year period in amplitude of solar-diurnal variation and disturbances of magnetic declination—A. D. BACHE	
illuminating materials—J. HENRY. (R. 1880) 442,	
influence of moon on magnetic declination—A. D. BACHE	
orbit of Neptune and tables of its motion—S. NEWCOMB	
orbit of Uranus, with tables of its motion—S. NEWCOMB	
Investigations— .	
anthropological, in 1879—O. T. MASON. (R. 1879) 345,	
archeological, instructions for-G. GIBBS. (R. 1861)	149
chemical and physiological, relative to certain American vertebrata—J. JONES	
of Biela's comet—J. S. Hubbard. (B. 1862)	150
scientific, in Russian America, suggestions relative to-J. HENRY	107
Iowa—	
ancient relics in—J. B. Cutts. (R. 1872)	271
Delaware county, mounds in-M. W. MOULTON. (R. 1877)	3.3
Muscatine county, mounds in—T. THOMPSON. (R. 1879)	345
IRBY, J. R. McD. Works and character of Smithson	27

, **J.**

JACKMAN, F. Mounds and earthworks of Rush county, Indiana. (R. 1879.) 34
JACKSON C. T. Catalogue of rooks minerals and area collected on molecial
survey in Michigan. (R. 1854)
JACKSON, J. B. S. Arsenic acid for protecting anatomical preparations from insects. (P. 1878)
Jade. Study of the mineral called yu by the Chinese—S. BLONDEL. (R. 1876.)
JAMES, T. P. Mosses of Kerguelen Island
Jamin, J. Photo-chemistry. (R. 1867)
Jamin, J. Vegetation and the atmosphere. (R. 1864)18
Japan—
analysis of coals from—J. A. MACDONALD
catalogue of Japanese woods from—L. F. WARD. (P. 1881) 46
examination of infusorial earth from—A. M. Edwards
geological researches in—R. PUMPELLY 20
Government of, collection of cotton fibre presented by. (P. 1881) 46
Jargon, Chinook-
bibliography of—G. GIBBS
dictionary of-G. GIBBS; H. HALE
vocabulary of-M. LIONNET; B. R. MITCHELL; W. W. TURNER 6
JEFFERSON, J. P. Mortality of fishes in Gulf of Mexico. (P. 1878) 33
JEFFERSON, J. P.; PORTER, J. Y.; MOORE, T. Destruction of fish in vicinity of the Tortugas. (P. 1878)
JEVONS, W. S. On a national library. (R. 1878) 273
JEWETT, C. C.—
Construction of catalogues of libraries and their publication by separate
stereotyped titles47
Copyright books from 1846 to 1849. (R. 1850) 2
General catalogue system for libraries. (R. 1850)
Report on catalogue system. (R. 1849)
Report on copyright system. (R. 1851)
Report on Halliwell manuscripts. (R. 1852)
Report on plan of library. (R. 1847)
Report on public libraries of the United States. (R. 1849)21,25
Reports on library. (R. 1848-1853) I, 21, 28, 51, 57, 57
Jewett, C. C., biographical notice of, by R. A. Guild. (R. 1867)
Jewett, E., diagnoses of mollusca collected by—P. P. CARPENTER 22
Johns' (Bishop) library, deposit of—E. CANBY. (R. 1862)
JOHNSON, S. H. Notes on mortality among fishes of Gulf of Mexico. (P. 1881.)
JOHNSON, S. W. Lectures on agricultural chemistry. (R. 1859)
JOHNSON W R Scientific character and researches of Smithson

OF SMITHSONIAN PUBLICATIONS.	22
JOHNSON, W. W. Distribution of forest trees in Montana, Idaho, and Washington. (R. 1870)	24
Joints, bibliography of diseases of the-W. W. KEEN	30
JONES, C. C., Jr. Aboriginal structures in Georgia. (R. 1877) 828,	400
Jones, C. C., Jr. Primitive manufacture of spear and arrow points on Savan-	
nah river. (R. 1879)	84
Jones, J.—	
Analysis of blood	8
Chemical and physiological investigations relative to vertebrata	82
Explorations of aboriginal remains of Tennessee	
Modes of burial	25
Observations on the liver, pancreas, spleen, kidney, etc	8
On burial caves	259
On earthworks.	25
On mounds	25
On relics	25
Jones, J. M. Kjækken-mædding in Nova Scotia. (R. 1868)	18
Jones, S. The Kutchin Indians. (R. 1866)	, 86
Jordan, D. S.—	
Contributions to North American ichthyology: Part I. Review of Rafinesque's memoirs	80
Part II. Notes on Cottidæ, etc.; synopsis Siluridæ	800
Part III. Distribution of fishes of Alleghany region; synopsis of family Catostomida	800
Description of new species of Caranx (Caranx Beani) from Beaufort, North Carolina. (P. 1880)	42
Description of new species of North American fishes. (P. 1879)	888
Forgotten paper of Dr. Ayres and its bearing on nomenclature of cyprinoid fishes of San Francisco. (P. 1880)	42
Notes on collection of fishes from Clackamas river, Oregon. (P. 1878.)	382
Notes on Cottidæ, Etheostomatidæ, Percidæ, Centrarchidæ, Aphododeridæ, Dorysomatidæ, and Cyprinidæ	806
Notes on Curtiss' collection of fishes from Florida. (P. 1880)	426
Notes on Duges' collection of fishes from Mexico. (P. 1879)	888
Notes on Henshall's collection of fishes from east Florids. (P. 1880)	428
Notes on Sema and Dacentrus. (P. 1880)	426
Notes on typical specimens of American fishes in British Museum and in Museum d'Histoire Naturelle, Paris. (P. 1879)	888
Review of Rafinesque's memoirs on North American fishes	308

Synopsis of Catostomidæ

Description	of new agonoid fish (Agonus vulsus) from California. (P. 1880
Description (P. 188	of new agonoid fish (Brachiopsis xyosternus) from California
fornia,	of new embiotocoid fish (Abeona aurora) from Monterey, Cali- with notes on a related species. (P. 1880)
of Culi	of new embiotocoid fish (Cymatogaster rosaceus) from consornis. (P. 1880)
ifornia.	of new embiotocoid fish (Ditrema atripes) from coast of Cal (P. 1880)
fornia.	of new flounder (Hippoglossoides exilis) from coast of Cali (P. 1880)
Description Californ	of new flounder (Platysomatichthys stomias) from coast of is. (P. 1880)
Description	of new flounder (Pleuronichthys verticalis) from California
	tes on other species. (P. 1880)
Island,	of new flounder (Xystreurys liolepis) from Santa Catalina California. (P. 1880)
	of new ray (Platyrhina triseriata) from coast of California
Description Califor	of new scorpsenoid fish (Sebastichthys maliger) from cost of is. (P. 1880)
Description rey Bay	of new scorpænoid fish (Sebastichthys proriger) from Monte , California. (P. 1880)
Description coast of	of new species of deep-water fish (<i>Icichthys Lockingtoni</i>) from California. (P. 1880)
Description coast of	of new species of Hemirhampus (Hemirhampus Rosa) from California. (P. 1880)
Description	of new species of Nemichthys (Nemichthys avocetta) from ound. (P. 1880)
Description	of new species of notidanoid shark (<i>Hexanchus corinus</i>) from coast of United States. (P. 1880)
Description	of new species of Paralepis (Paralepis coruscans) from Strait de Fuca. (P. 1880)
Description	of new species of Pomadosys from Mazatlan. (P. 1881)
	of new species of <i>Ptychochilus (Ptychochilus Harfordii</i>) from onto river. (P. 1881)
Description	of new species of ray (Raia rhina) from coast of California
Description	of new species of ray (Raia stellulata) from Monterey, Cal (P. 1880)
Description	of new species of rock cod (Sebastichthys serriceps) from compornia. (P. 1880)
Description	of new species of rockfish (Sebastichthys carnatus) from coss

OF SMITHSONIAN PUBLICATIONS.

JORDAN, D. S.; GILBERT, C. H.—Continued.
Description of new species of rockfish (Sebastichthys chrysomelas) from coast of California. (P. 1880)
Description of new species of Sebastichthys (Sebastichthys miniatus) from Monterey bay, California. (P. 1880)
Description of new species of Xenichthys (Xenichthys ocyurus) from west coast of Central America. (P. 1881)
Description of new species of Xiphister and Apodichthys from Monterey, California. (P. 1880)
Description of Sebastichthys mystinus. (P. 1881)
Description of seven new species of sebastoid fishes from coast of California. (P. 1880)
Description of two new species of fishes (Ascelicthys rhodorus and Scyta- lina cerdale) from Neah bay, Washington Territory. (P. 1880)
Description of two new species of flounders (Parophrys ischyurus and Hippoglossoides elassodon) from Puget Sound. (P. 1880)
Description of two new species of Sebastichthys (Sebastichthys entomelas and Sebastichthys rhodochloris) from Monterey bay, Cal. (P. 1880.)
Description of two new species of scopeloid fishes (Sudis ringens and Myctophum crenulare) from Santa Barbara channel, Cal. (P. 1880.)
Descriptions of five new species of fishes from Mazatlan, Mexico. (P. 1881)
Descriptions of thirty-three new species of fish from Mazatlan, Mexico. (P. 1881)
Generic relations of Belone exilis. (P. 1880)
Generic relations of Platyrhina exasperata. (P. 1880)
List of fishes collected by Lt. H. E. Nichols in Gulf of California and on west coast of Lower California, with descriptions of four new species. (P. 1881)
List of fishes of Pacific coast of United States, with table showing distri- bution of species. (P. 1880)
Note on Raia inornata. (P. 1881)
Notes on collection of fishes from San Diego, California. (P. 1880)
Notes on collection of fishes from Utah Lake. (P. 1880)
Notes on collection of fishes made by Lt. H. E. Nichols on west coast of Mexico, with descriptions of new species. (P. 1881)
Notes on fishes of Beaufort Harbor, North Carolina. (P. 1878)
Notes on fishes of Pacific coast of United States. (P. 1881)
Notes on sharks from coast of California. (P. 1880)
Occurrence of Cephaloscyllium laticeps (Duméril), Gill, in California. (P. 1880)
Oil shark of southern California (Galeorhinus galeus). (P. 1880)
JORDAN, D. S.; JOUY, P. L. Check-list of duplicate fishes of Pacific, distributed
by Smithsonian Institution in 1881. (P. 1881)
Journal of exploration of western Missouri in 1854—P. R. Hoy. (R. 1864)

Journals of Board of Regents—See Regents.	
Journey to the Yukon, Russian America-W. W. KIRBY. (R. 1864) 1	88
JOUY, P. L. Description of new species of Squalius (Squalius alicia) from Utah Lake. (P. 1881)	167
JOUY, P. L.; JORDAN, D. S. Check-list of duplicates of fishes of Pacific, distributed by Smithsonian Institution in 1881. (P. 1881)	16 7
Judiciary Committee, Senate, report of, on management of Smithsonian Insti- tution—A. P. Butler. (R. 1855)	7.
Jupiter, orbit and tables of—J. N. STOCKWELL	27:
Jupiter, small planets between Mars and—Prof. LESPIAULT. (R. 1861)	14.
Jurassic fossils, check-list of-F. B. MEEK	77
Jussieus, the, and the natural method—M. FLOURENS. (R. 1867)	Zlá

K.

KABIS, M. Introduction to study of Coptic language. (R. 1867)	215
Kämtz; Dufour, C. Scintillation of the stars. (R. 1861)	144
KANE, E. K	
Astronomical observations in the Arctic Seas	129
Magnetic observations in the Arctic Seas	97
Meteorological observations in the Arctic Seas	104
Physical observations in the Arctic Seas	196
Tidal observations in the Arctic Seas	130
Kansas—	
Butler county, account of storm in-W. HARRISON. (R. 1871)	24:
climate of—R. S. Elliott. (R. 1870)	244
coleoptera of-J. L. LE CONTE	126
new cretaceous invertebrate fossils from—C. A. WHITE. (P. 1879)	33:
Kansas City, Missouri, antiquities of—W. H. R. LYKINS. (R. 1877)	27
Karif or Carib language and people—C. H. BERENDT. (R. 1878)	7.5
KAROLYI, Lieut. von; CRAIG, B. F. Products of combustion of gun-cotton	
and gunpowder. (R. 1864)	
KEEN, W. W. Bibliography of works on diseases	
KEEN, W. W. Surgical complications and sequels of continued fevers. Toner lecture No. V.	
Keller, Dr., abstract of report of, on settlements in Switzerland—A. Morlor. (R. 1868)	
KELLEY, O. H. Ancient town in Minnesota. (R. 1868)	18.
Kelley's Island, Ohio, climate of—G. C. Huntington. (R. 1866)	214

Kennebec river, Maine, opening and closing of-R. H. GARDINER. (R. 1858.)	109
KENNEDY, J. P. Authority to naval officers to receive and transmit specimens	
to the Smithsonian Institution	84
Kennicott's explorations—Hudson's Bay Company. (R. 1863)	187
Kentucky—	
ancient mounds in—R. Peter. (R. 1872)	271
antiquities from—S. S. Lyon. (R. 1858)	109
Breckinridge county, age of stone and troglodytes of—R. S. Robertson. (R. 1874)	286
Hancock county, antiquities of-J. FRIEL. (R. 1877)	328
Lexington, ancient mound near—R. PETER. (R. 1871)	249
Union county, exploration of ancient mounds in—S. S. Lyon. (R. 1870.)	244
Kepler, his life and works, by M. BERTHRAND. (R. 1869)	228
Kerguelen Island, contributions to natural history of-	
1. Ornithology-J. H. KIDDER; E. COUES	293
II. Oölogy, botany, etc.—J. H. KIDDER and others	294
Kershner collection from Kerguelen Island	294
Kew Observatory, apparatus and processes employed in verification of barome-	
ters at—J. Welsh. (R. 1859)	110
Kew Observatory, experiments on aneroid barometers made at-B. STEWART.	
(R. 1868)	224
KIDDER, J. H. Experiments on animal heat of fishes. (P. 1879)	888
KIDDER, J. H. Mammals of Kerguelen Island	294
KIDDER, J. H., and others. Natural history of Kerguelen Island	294
KIDDER, J. H.; COUES, E. Birds of Kerguelen Island	293
Kidney, observations on—J. Jones	82
Kinetic theories of gravitation—W. B. TAYLOB. (R. 1876) 299,	
KING, C. B. Catalogue of engravings presented to Smithsonian. (R. 1861)	149
King, W. M. Account of Indian burial in California. (R. 1874)	286
KIPP, J. Accuracy of Catlin's account of Mandan ceremonies. (R. 1872)	271
KIRBY, W. W. Journey to the Yukon, Russian America. (R. 1864)	188
KIRKWOOD, D. Asteroids between Mars and Jupiter. (R. 1876)	299
Kitchen-midden in Costa Rica, note on shells from—W. H. Dall. (P. 1878.)	882
Kitchen-midden—See Kjækken-mæddings.	
Kjækken-mæddings in Nova Scotia—J. M. Jones. (R. 1868)	187
Kjækken-mæddings on northwest coast of America—P. SCHUMACHEB. (R.	
1878; R. 1874) 275	, 286
Klamath lake dialect, details of Indian conjurer's practice in—A. S. GATSCHET.	
(E. 1879–80)	476
Klamath lake dialect, "The Relapse" in—A. S. GATSCHET. (E. 1879-80)	476
KLETZINSKY, Prof. Nitrogen bodies of modern chemistry. (R. 1872)	271
KNAPP, G. Earthworks on Arkansas river. (R. 1877)	828

Киюнт, Е. H. Study of savage weapons at Centennial Exhibition. (В. 1879) 345,	415
KNIGHT, G. H. New system of weights and measures. (R. 1867)	215
KNIGHT, R. T.; HENRY, J. Connection of gales of wind and appearance of	
aurora. (R. 1871)	249
Knowledge of cryptogamous plants, present state of—W. REICHARDT. (R. 1871)	
Kohl, J. G. On collection of charts and maps of America. (R. 1856)	
KORNHUBER, G. A. Alternate generation and parthenogenesis in the animal kingdom. (R. 1871)	
KRON, F. J. Antiquities of Stanley and Montgomery counties, North Carolina. (B. 1874)	
KRON, H. J. Lightning discharges. (R. 1867)	
Kumlien, L.—	
Ethnology, mammals, and birds of Arctic America	342
Howgate Polar Expedition, natural history of Arctic America	
Report on explorations in Greenland. (R. 1878)	
KUNHARDT AND COMPANY. Free freight to Smithsonian Institution between United States and Germany. (R. 1861)	
Kushkushkee, Indian village near Newcastle, Penn., account of-E. M. Mc-	
	949
CONNELL. (R. 1871)	220
CONNELL. (R. 1871)	
CONNELL. (R. 1871)	
CONNELL. (R. 1871)	865
CONNELL. (R. 1871) Kutchin Indians—S. Jones. (R. 1866) 214, Labels for collections, etc	164
CONNELL. (R. 1871) Kutchin Indians—S. Jones. (R. 1866) 214, Labels for collections, etc Laboulaye, E. Acknowledgment for books on education. (R. 1867)	365 164 215
CONNELL. (R. 1871)	365 164 215
CONNELL. (R. 1871)	164 215 149
CONNELL. (R. 1871)	164 215 149
CONNELL. (R. 1871) Kutchin Indians—S. Jones. (R. 1866) Labels for collections, etc. Laboulaye, E. Acknowledgment for books on education. (R. 1867) Lacustrian cities of Switzerland—F. Troyon. (R. 1861) Lacustrian constructions or palafittes of the lake of Neuchâtel—E. Desor. (R. 1865) Lacustrian habitations of Switzerland—F. Troyon. (R. 1861) Lacustrian settlements in Switzerland, abstract of Dr. Keller's report on—A.	164 215 149 360 149
CONNELL. (R. 1871) Kutchin Indians—S. Jones. (R. 1866) Labels for collections, etc. Laboulaye, E. Acknowledgment for books on education. (R. 1867) Lacustrian cities of Switzerland—F. Troyon. (R. 1861) Lacustrian constructions or palafittes of the lake of Neuchâtel—E. Desor. (R. 1865) (R. 1865) Lacustrian habitations of Switzerland—F. Troyon. (R. 1861) Lacustrian settlements in Switzerland, abstract of Dr. Keller's report on—A. Morlot. (R. 1868) La Have Banks, occurrence of Lycodes vahlii on—G. B. Goode; T. H. Bean.	164 215 149 360 149
CONNELL. (R. 1871) Kutchin Indians—S. Jones. (R. 1866) Labels for collections, etc. Laboulaye, E. Acknowledgment for books on education. (R. 1867) Lacustrian cities of Switzerland—F. Troyon. (R. 1861) Lacustrian constructions or palafittes of the lake of Neuchâtel—E. Desor. (R. 1865) Lacustrian habitations of Switzerland—F. Troyon. (R. 1861) Lacustrian settlements in Switzerland, abstract of Dr. Keller's report on—A. Morlot. (R. 1863) La Have Banks, occurrence of Lycodes vahlii on—G. B. Goode; T. H. Bean. (P. 1879)	164 215 149 260 149 187
CONNELL. (R. 1871) Kutchin Indians—S. Jones. (R. 1866) Labels for collections, etc. Laboulaye, E. Acknowledgment for books on education. (R. 1867) Lacustrian cities of Switzerland—F. Troyon. (R. 1861) Lacustrian constructions or palafittes of the lake of Neuchâtel—E. Desor. (R. 1865) Lacustrian habitations of Switzerland—F. Troyon. (R. 1861) Lacustrian settlements in Switzerland, abstract of Dr. Keller's report on—A. Morlot. (R. 1868) La Have Banks, occurrence of Lycodes vahlii on—G. B. Goode; T. H. Bean. (P. 1879) Lake Neuchâtel, lacustrian constructions of—E. Desor. (R. 1865)	164 215 149 260 149 187 333 209
CONNELL. (R. 1871) Kutchin Indians—S. Jones. (R. 1866)	164 215 149 360 149 187 333 209 214
CONNELL. (R. 1871) Kutchin Indians—S. Jones. (R. 1866) Labels for collections, etc. Laboulaye, E. Acknowledgment for books on education. (R. 1867) Lacustrian cities of Switzerland—F. Troyon. (R. 1861) Lacustrian constructions or palafittes of the lake of Neuchâtel—E. Desor. (R. 1865) Lacustrian habitations of Switzerland—F. Troyon. (R. 1861) Lacustrian settlements in Switzerland, abstract of Dr. Keller's report on—A. Morlot. (R. 1868) La Have Banks, occurrence of Lycodes vahlii on—G. B. Goode; T. H. Bean. (P. 1879) Lake Neuchâtel, lacustrian constructions of—E. Desor. (R. 1865)	164 215 149 260 149 187 333 209 214 155

OF SMITHSONIAN PUBLICATIONS.	227
Lake Winnipeg, notes on egging expedition to—D. Gunn. (R. 1867)	215
Lakes, dates of opening and closing of—F. B. Hough	182
Lakes, North American, fluctuations of level in—C. WHITTLESEY	119
Lakes, Norwegian, statistics relative to—O. E. DREUTZER. (R. 1866)	214
Lalande observations, 1795—S. C. WALKER	5
LAMARCK, J. B. Distinction between tornadoes and tempests. (R. 1871)	249
LAMONT, J. The solar eclipse of July 18, 1860. (R. 1864)	188
Land and fresh-water shells—See Shells.	
Land, cessions of; by Indian tribes to United States—C. C. ROYCE. (E. 1879-80.)	476
Land, gradual approach of sea upon—S. P. MAYBERRY. (R. 1867)	215
Language—	
Carib or Karif—C. H. BERENDT. (R. 1873)	275
Coptic, introduction to the study of—M. KABIS. (R. 1867)	215
Dakota, grammar and dictionary of—S. R. Riggs	40
dominant, for science—A. DE CANDOLLE; J. E. GRAY. (R. 1874)	286
evolution of—J. W. Powell. (E. 1879-80)	476
of aboriginal Indians of America—G. GIBBS. (R. 1870)	244
of Dakota or Sioux Indians—F. L. O. Roehrig. (R. 1871) 249,	
of Navajos, said to resemble the Welsh—S. Y. McMasters. (R. 1865.)	209
phonetic, vocal sounds of L. Bridgman compared with elements of—F. LIEBER	12
sign, among North American Indians compared with that among other peoples and deaf mutes—G. MALLERY. (E. 1879–80)	476
trade, of Oregon, dictionary of-G. GIBBS	161
trade, of Oregon, vocabulary of-B. R. MITCHELL; W. W. TURMER	68
Yoruba, of Africa, grammar and dictionary of-T. J. Bowen	98
Languages—	
Indian-G. Flachenecker. (R. 1862)	150
Indian—G. Gibbs. (R. 1865)	209
Indian, method of recording—J. O. Dorsey; A. S. Gatschet; S. R. Riegs. (E. 1879-80)	4 76
LAPHAM, I. A. Antiquities of Wisconsin	70
Lapidary, aboriginal, stock-in-trade of-C. RAU. (R. 1877) 828, 440,	402
La Place, eulogy on, by F. Arago. (R. 1874)	286
La Rive—See De La Rive.	
Lartet, E., scientific labors of, by F. FISHER. (R. 1872)	271
Larynx, bibliography of diseases of the-W. W. KEEN.	800
Latiloid genera, note on-T. GILL. (P. 1881)	467
Latimer collection of antiquities from Porto Rico-O. T. Mason. (R. 1876.) 299,	397
LATIMER, G. A.—	
Earthquakes in St. Thomas. (R. 1867)	215
Hurricane at Tortola, St. Thomas, and Porto Rico. (R. 1867)	215

Latin vocabulary with comparative words in English, Spanish, and French !	70
Latitudes and longitudes, Arctic—E. K. KANE	29
LAUGEL, A. The sun, its chemical analysis. (R. 1861)	149
Lausanne, report on ethnological collections of Museum at—F. Trovon. (R. 1861)	149
LAUTENBACH, B. F. Effect of irritation of a polarized nerve. (R. 1878.) 341,4	11
LAW, A. E. Antiquities of Blount county, Tennessee. (R. 1874)	256
Law of deposit of flood-tide—C. H. Davis	33
Law of variation of temperature in ascending currents of air—J. HANN. (R. 1877)	198
Law of variation of temperature in ascending moist currents—L. Sohneke. (R. 1877)	198
LAWRENCE, G. N.—	
Birds of southwestern Mexico collected by F. E. Sumichrast	19 5
Catalogue of Ober's collection of birds of Antigua and Barbuda, with his notes. (P. 1878)	332
Catalogue of Ober's collection of birds of Dominica, with his notes and observations. (P. 1878)	322
Catalogue of Ober's collection of birds of Grenada, including others seen, but not obtained. (P. 1878)	3\$ 2
Catalogue of Ober's collection of birds of Guadeloupe. (P. 1878)	332
Catalogue of Ober's collection of birds of Lesser Antilles, with table showing distribution. (P. 1878)	<u>332</u>
Catalogue of Ober's collection of birds of Martinique. (P. 1878)	3≈
Catalogue of Ober's collection of birds of St. Vincent, with his notes. (P. 1878)	37:
Description of new species of bird of family <i>Turdidæ</i> from Dominica, W. I. (P. 1880)	
Description of new species of Icterus from West Indies. (P. 1880)	
Description of new species of parrot, of genus Chrysotis, from Dominica. (P. 1880)	
Description of new subspecies of Loxigilla from West Indies. (P. 1881.)	
Laws of atmospheric circulation over the earth—J. H. COFFIN	
LEA, I. Check-list of the shells of North America	
LE CONTE, JOHN. Constants of nature. (R. 1878)	3 4 i
LE CONTE, J. L.—	
Classification of coleoptera of North America. Parts I, II 136,	
Coleopters of Kansas and eastern New Mexico	,26
Instructions for collecting coleoptera. (R. 1858)	[(])
List of coleopters of North America. Part I	res TAN
New species of North American coleoptera. Parts 1, 11 167, 2	62
Revision of Melsheimer's catalogue of coleopters.	10°
LE CONTE, JOSEPH. Lecture on coal. (R. 1857)	75
Lecture on American fire-alarm telegraph—W. F. CHANNING. (R. 1854)	

OF SMITHSONIAN PUBLICATIONS.	229
Lectures on—	
agricultural chemistry—S. W. Johnson. (R. 1859)	110
Arctic explorations—I. I. HAYES. (R. 1861)	149
astronomy—A. Caswell. (R. 1858)	109
bridges—F. Rogers. (R. 1861)	149
bridges and roads—F. Rogers. (R. 1860)	147
the camel—G. P. Marsh. (R. 1854)	78
cancerous tumors. Toner lecture No. I-J. J. WOODWARD	266
coal—J. Le Conte. (R. 1857)	107
dual character of the brain. Toner lecture No. 11-C. E. Brown-Séquard.	291
electro-physiology—C. MATTEUCCI. (R. 1865)	209
insect instincts and transformations—J. G. Morris. (R. 1855)	77
marine algae—W. H. HARVEY. (R. 1855)	77
meteoric stones—J. L. SMITH. (R. 1855)	77
meteorology-R. Russell, with notes by J. Henry. (R. 1854)	78
mollusca, or shell-fish and their allies-P. P. CARPENTER. (R. 1860.) 147,	152
natural history as applied to farming and gardening—J. G. Morrais. (R. 1855)	77
nature and cure of bites of serpents and wounds of poisoned arrows— D. Brainard. (R. 1854)	78
nature of reparatory inflammation in arteries after ligature, etc. Toner	
lecture No. VII—E. O. SHAKESPEARE	321
oxygen and its combinations—G. I. CHACE. (R. 1855)	77
physical ethnology—D. Wilson. (R. 1862)	150
planetary disturbances—E. S. SNELL. (R. 1855)	77
principles of linguistic science—W. D. WHITNEY. (R. 1863) 187,	352
relations of time and space—S. ALEXANDER. (R. 1861)	149
sanitary drainage of Washington. Toner lecture No. VIII—G. E. WARING, Jr.	34 9
shells of Gulf of California-P. P. CARPENTER. (R. 1859)	110
strain and over-action of the heart. Toner lecture No. III—J. M. DA	279
study of high antiquity in Europe-A. Morlot. (R. 1862; R. 1864.) 150	
study of nature and mechanism of fever. Toner lecture No. IV—H. C. Wood	282
surgical complications and sequels of fevers. Toner lecture No. v—W. W. Keen	
Switzerland-A. D. BACHE. (R. 1870)	244
syllabus of a course of, on physics—J. HENRY. (R. 1856)	91
undulatory theory of light—F. A. P. BARNARD. (R. 1862)	150

the Union-H. REED. (R. 1854)

zone of small planets between Mars and Jupiter-E. Loomis. (R. 1854.)

vastness of visible creation—S. ALEXANDER. (R. 1857)

75

Lectures, Toner—	
No. 1. Cancerous tumors—Woodward	
II. Dual character of brain—Brown-Sfquard	291
III. Over-action of heart—Da Costa	279
IV. Study of fever—Wood	282
v. Continued fevers—KEEN	300
VI. Sub-cutaneous surgery—ADAMS	302
VII. Reparatory inflammation—SHAKESPEARE.	321
VIII. Sanitary drainage of Washington-WARING	349
LEE, Dr. Progress of astronomical photography. (R. 1861)	143
LEE, J. C. Y. Ancient ruin in Arizona. (R. 1872)	9-1
Legacy—See Bequest.	
Legendre, memoir of, by E. DE BEAUMONT. (R. 1867)	213
LEIDY, J.—	
Ancient fauna of Nebraska	58
Cretaceous reptiles of the United States	
Extinct sloth tribe of North America	72
Extinct species of American ox	41
Fauna and flora within living animals	. #
On parasites	4
Report on fossils from Nebraska. (R. 1851)	5.
Review of cretaceous reptiles of United States. (R. 1864)	
Leidy, J.; Tryon, G. W. Report on shells presented to Academy of Natural Sciences. (R. 1865)	T.
Leipsic Museum of Ethnology—A. Schott; O. T. Mason. (R. 1873)	2,
LEMSTRÖM, S.; DE LA RIVE, A. A. Electricity of the atmosphere and the aurora borealis. (R. 1874)	
Lens, account of Priestley's-J. HENRY. (R. 1859)	110
Lepidopsetta isolepis, new flat flah, found in markets of San Francisco, note on—W. N. Lockington. (P. 1880)	
Lepidoptera—	
appendix to Morris' synopsis of—B. CLEMENS	134
described, of North America, catalogue of-J. G. Morris	118
described, of North America, synopsis of. Part 1-J. G. Morris	133
diurnal, of Arctic America—W. H. EDWARDS	342
instructions for collecting—B. CLEMENS. (R. 1858)	Ich
introduction to Morris' synopsis of—V. Heinemann	13.
method of preserving-T. R. PEALE. (R. 1868)	15
nocturnal, of Arctic America—S. H. SCUDDER	91-
notes to Morris' synopsis of—W. H. EDWARDS	12
LEPSIUS, R. Presentation of books on Egypt. (R. 1860)	141
Leptocardii, arrangement of families of—T. GILL	**
I per per I P On the classification of books (R. 1862)	150

LESPIAULT, Prof. Small planets between Mars and Jupiter. (R. 1861) 1. Lesser Antilles, catalogue of Ober's collection of birds of—G. N. LAWBENCE.	
· · · · · · · · · · · · · · · · · · ·	82
Letter—	
by M. Hoek in reference to meteoric shower of November 18, 1867 21	17
from Robert Hare relative to gift of apparatus I, 8	29
from Joseph Henry to Secretary Treasury on payment of interest in coin. (R. 1865)	:09
from Richard Rush relative to Smithson. (R. 1853) 67, 8	28
LETTERMAN, J. Sketch of the Navajo Indians. (R. 1855)	77
Letters—	
of introduction. (R. 1865) 2	209
on work of the Museum—B. PHILLIPS. (P. 1881) 467, 48	54
relative to programme of organization of Smithsonian Institution 8	28
relative to Smithson's bequest 8	28
Leurynnis, Lockington, identity of, with Lycodopsis, Collet—T. GILL. (P. 1880) 4	25
	19
LEWIS, J. Instructions for collecting land and fresh-water shells. (R. 1866.) 214, 30	
	258
	14
	49
	244
	49
Libraries—	
circular respecting new report on-J. HENRY. (R. 1854)	75
construction of catalogues of, and their publication by means of separate	
· . · · · · · · · · · · · · · ·	47
general catalogue system for—C. C. JEWETT. (R. 1850)	28
general catalogue system for, report of Commission on. (R. 1850)	28
public, in United States and British Provinces, list of-W. J. RHEES. 116, 2:	38
public of United States, notices of-C. C. JEWETT. (R. 1849) 21,	25
report of Commission on stereotype catalogue of	47
rules for cataloguing—C. C. JEWETT	47
Library—	
2021000, 10F (50
Bishop Johns', deposit of—E. CANBY. (R. 1862)	150
Imperial, of Vienna, books presented by. (R. 1865) 2	209
of Bureau of Ethnology, catalogue of linguistic manuscripts in—J. C. PILLING. (E. 1879-80)4	1 76
of Indian linguistics, account of-J. G. SHEA. (R. 1861) 1	49
of National Museum, circular asking contributions to. (P. 1881) 467, 4	58
	275

Library of Congress—
act of Congress to transfer Smithsonian library to. (R. 1865) 209, 328
catalogue of publications of societies and of periodical works belonging to the Smithsonian Institution deposited in
meteorological articles received by the Smithsonian Institution and deposited in. (R. 1871; R. 1873) 249, 275
Library of Smithsonian Institution—
act of Congress to transfer, to Library of Congress. (B. 1865) 209, 328
additions to—See each annual report.
and copyright system—C. C. JEWETT. (R. 1851)
and Halliwell manuscripts, report on—C. C. JEWETT. (R. 1852)
catalogue of publications of societies and other periodical works in, 117, 179
donations to, from foreign institutions. (R. 1864) 188
publications of learned societies and periodicals in. Parts 1, 11 73, 85
report on—C. C. JEWETT. (R. 1848, 1849, 1858)
report on plan of—C. C. JEWETT. (R. 1847)
statistics of, 1846–1877 \$29
Lichens of Arctic America—E. Tuckerman 342
Lichens of Kerguelen Island—E. TUCKERMAN 294
LIEBER, F. Vocal sounds of Laura Bridgman compared with elements of phonetic language
LIEBIG, J. VON. Induction and deduction. (R. 1870) 2H
Life—
and character of Joseph Henry—J. C. WELLING 356, 338
and labors of Henry Gustavus Magnus. (R. 1870) 24
and scientific labors of Stefano Marianini—C. MATTEUCCI. (R. 1869) 228
and works of Michael Faraday—A. A. DE LA RIVE. (R. 1867) 215
and works of Kepler-M. BERTHRAND. (R. 1869) 23
of George Catlin—J. HENRY. (R. 1872) 271
of Prof. Chester Dewey-M. B. Anderson. (R. 1870)
Life—See Biography, Eulogy, Memoirs.
Light—
and heat of sun, relative intensity of—L. W. MERCH. (R. 1856) 91,83
polar, or aurora borealis, its phenomena and laws—E. Loomis. (R. 1865.) 209
undulatory theory of, lectures on-F. A. P. BARNARD. (R. 1862) 150
velocity of, essay on—C. Delaunay. (R. 1864) 188, 354
Light-House Board—
investigations by, relative to illuminating materials—J. HERRY. (B. 1880)442, 389
proceedings of, on death of J. Henry 356
researches by, in sound—J. HENRY. (R. 1878)341, 406
Lighting, architecture in relation to-D. B. REID. (R. 1856)

OF SMITHSONIAN PUBLICATIONS.	288
Lightning—	
and thunder, observations on—S. MASTERMAN. (R. 1855)	77
discharges, accounts of-G. W. Dodge and others. (R. 1867)	215
effect of—S. L. HILLIER. (R. 1866)	214
Lightning-rods, directions for constructing—J. HENRY	237
Lille, Imperial Society of Science, Agriculture, and Arts of. Prize questions. (R. 1865)	209
LILLJEBORG, W. Outline of systematic review of the classification of birds. (R. 1865) 209,	364
Limitations to use of some anthropological data—J. W. Powell. (E. 1879-80.)	476
Limpets from deep waters off eastern coast of U.S.—W. H. Dall. (P. 1881.)	467
Limpets of Alaska and Arctic region, report on—W. H. Dall. (P. 1878)	882
Lindheimer's botanical exploration in New Mexico and California, account of— A. Gray. (R. 1849)	21
Linguistic manuscripts in library of Bureau of Ethnology, catalogue of—J. C. PILLING. (E. 1879-80)	476
Linguistic science, lectures on principles of—W. D. WHITNEY. (R. 1863.) 187,	
Linguistics, Indian, account of library of—J. G. Shea. (R. 1861)	149
Linguistics, Indian, recommendation of Shea's—G. GIBBS and others. (R. 1861.)	149
LIONNET, M. Vocabulary of the Chinook jargon	68
Liparis ranula, new species of, obtained by U. S. Fish Commission off Halifax,	
Nova Scotia-G. B. Goode; T. H. Bean. (P. 1879)	888
Liquids, expansion of, by heat—F. W. CLARKE	289
Liquids, specific heats of—F. W. CLARKE	276
Lisboa, M. M., books on Brazil, presented by. (R. 1865)	209
List of—	
additions to Museum—See each annual report.	
addresses of foreign institutions, 1862-1865. (R. 1865)	209
algæ of Rhode Island—S. T. OLNEY	241
American correspondents	69
American libraries and public institutions—W. J. RHEES.	238
anthropological publications of Charles Rau. (P. 1881)	467
apparatus available for scientific research. (R. 1878)	841
articles deposited by Smithsonian Institution in Corcoran Gallery of Art. (R. 1874)	286
birds and mammalia of Missouri river—E. HARRIS. (R. 1850)	28
birds of central California, partial—L. Belding. (P. 1878)	882
birds of District of Columbia—E. Cours; S. S. Prentiss. (R. 1861)	149
casts of heads of Indian boys and girls at Hampton, Va.—R. H. PRATT. (P. 1879)	888
coleopters of North America. Part I-J. L. LE CONTE	140
collections presented by foreign Centennial commissioners. (R. 1876)	299

•

. . ____

٠

.

List of—Continued.
described birds of Mexico, Central America, and West Indies not in Smithsonian Institution
described species of humming-birds—D. G. Elliot 334
desiderata among North American birds-R. RIDGWAY. (P. 1881) 467
domestic institutions in correspondence with Smithsonian Institution. 69, 238
duplicates of fishes of Pacific coast distributed by Smithsonian Institution in 1881—D. S. JORDAN; P. L. JOUY. (P. 1881)
European fishes in National Museum—T. H. BEAN. (P. 1879) 333
expeditions from which specimens in Museum have been derived—S. F. BAIRD. (R. 1867)
fishes collected by Lieut. H. E. Nichols in Gulf of California—D. S. JORDAN; C. H. GILBERT. (P. 1881)
fishes of Pacific coast of United States—D. S. JORDAN; C. H. GILBERT. (P. 1880)
foreign agents of Smithsonian Institution in 1876. (See also annual reports)
foreign correspondents of Smithsonian Institution. 64, 154, 225, 243, 309, 469
foreign correspondents of Smithsonian Institution, systematic index to 257
generic names of animals—S. H. SCUDDER 470
Indian vocabularies received from the Wheeler Expedition. (R. 1874., 25.
institutions, libraries, colleges, and other establishments in correspondence with the Smithsonian Institution—W. J. RHEES
invertebrate fossils of North America— Cretaceous and jurassic—F. B. Meek
Eocene and oligocene—T. A. CONBAD 200
Miocene-F. B. Merk183
marine invertebrata of New England distributed by United States Fish Commission—A. E. VERRILL; R. RATHBUN. (P. 1879)
marine invertebrates, mainly from New England, distributed by U. S. National Museum—R. RATHBUN. (P. 1881) 467, 465. 471
meteorological material contributed to Smithsonian Institution. (R. 1860- 1865, 1867-1871) 147, 149, 150, 187, 188, 209, 215, 224, 228, 244. 24
meteorological stations and observers. (R. 1849; R. 1853–1873) == 21, 67, 73, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 214, 215, 224, 228, 244, 249, 271, 275.
minerals in United States National Museum—F. M. ENDLICH. (R. 1873; P. 1880)
North American batrachia and reptilia—E. D. Cope 292
North American species of myriapods of family Lysiopetalide—J. A. RYDBB. (P. 1880)
official publications of the United States Government between 1868 and 1881—G. H. BOEHMER 45
papers presented to Royal Society by James Smithson. (R. 1853) 67. 330
periodicals received by the Institution. (R. 1880) Q, 442, 436

List of—Continued.
photographic portraits of North American Indians in Smithsonian Insti- tution 216
plants of the upper Missouri-T. C. PORTER. (R. 1850) 26
plants of Washington and vicinity-L. F. WARD 460
principal literary and scientific institutions in United States, 1879 335
public libraries, institutions, and societies in United States and British Provinces of North America—W. J. Rhees
publications of the Smithsonian Institution. (R. 1868) 74, 203, 224, 226 245, 278, 290, 301, 344, 487, 478
publications of the United States National Museum. (P. 1881) 467, 474
Regents, officers, and assistants of the Smithsonian Institution and U.S.
National Museum. (See also each annual report)466, 290
scientific papers of A. D. Bache—B. A. Gould. (R. 1870) 244, 329, 375
shells of exploring expedition
shells of North America—I. LEA; P. P. CARPENTER; W. STIMPSON;
W. G. Binney; T. Prime
species of Middle and South American birds not in United States National
Museum—R. Ridgway. (P. 1881)
substances derived from animal kingdom—G. B. Goode
Literary exchanges—See Exchanges.
Literary study of jade—S. Blondel. (R. 1876)
Littoral marine fauna of Provincetown, Mass.—R. RATHBUN. (P. 1880) 426
Liver, observations on the—J. Jones82
LIVERMORE, G. Report on Jewett's general stereotype catalogue of public libraries47
Living animals, flora and fauna within-J. LEIDY 44
Lloyd, North German, free freight between Germany and the United States. (R. 1858)
LLOYD, W. A. Exchange of specimens. (R. 1867) 215
LLOYD, W. A. Sparrows sent to the United States. (R. 1867) 215
LOCKE, J. Catalogue of rocks, minerals, ores, and fossils. (R. 1854) 75
Locke, J. Observations on terrestrial magnetism 35
LOCKETT, S. H. Mounds in Louisiana. (R. 1872) 271
Lockington, W. N.—
Description of new chiroid fish (Myriolepis zonifer) from Monterey Bay, California. (P. 1880)
Description of new fish from Alaska (Uranidea microstoma). (P. 1880.) 425
Description of new genus and some new species of California fishes (Icosteus ænigmaticus and Osmerus attenuatus). (P. 1880)
Description of new genus and species of Cottidæ. (P. 1881) 467
Description of new sparoid fish (Sparus brachysomus) from Lower Cali-
fornia. (P. 1880)

Lockington, W. N.—Continued.	
Description of new species of Agonidæ (Brachyopsis verrucosus) from coast of California. (P. 1880)	425
Description of new species of Prionotus (Prionotus stephanophyrs) from coast of California. (P. 1880)	
Descriptions of new genera and species of fishes from coast of California. (P. 1879)	833
Note on new flat fish (Lepidopsetta isolepis) found in markets of San Francisco. (P. 1880)	425
Remarks on species of genus Chirus found in San Francisco market. (P. 1880)	42 5
Review of the Pleuronectidae of San Francisco. (P. 1879)	
Locusts and grasshoppers of America—A. S. TAYLOR. (R. 1858)	
LOEW, H.—	
Monographs of the diptera of North America-	
Part 1. Edited by R. OSTEN SACKEN	
Part II. Edited by R. OSTEN SACKEN	
Part III	
Review of North American Trypetina	
LOEW, H.; OSTEN SACKEN, R. Instructions for collecting diptera. (R. 1858.)	109
Logan, T. M.— . Climate of California. (R. 1855)	77
· Meteorological observations at Sacramento, California. (R. 1854)	75
Meteorology of Sacramento, California. (R. 1857)	107
LOGAN, W. E. Request for duplicate shells. (R. 1859)	
LOMBARD, H. C. Report on the transactions of Society of Physics and Natural History of Geneva, July, 1868, to June, 1869. (R. 1869)	
Lombardy, Royal Scientific and Literary Institute of. Prize questions. (R. 1865.)	
London Institution of Civil Engineers. Prize questions. (R. 1862)	
London Royal Horticultural Society. Exchange of publications. (R. 1861)	
London, Royal Society of, origin and history of—C. A. ALEXANDER. (R. 1868.)	187
LONGFELLOW, H. W., and others. Report of American Academy of Arts and Sciences on organization of Smithsonian Institution. (R. 1858)	l
Long Island, fishes on coast of New Jersey and—S. F. BAIRD. (R. 1854.) 75	
Longitude, transatlantic—B. A. Gould	
Longitudes, Arctic—E. K. KANE	
LOOMIS, E.—	
Aurora borealis, or polar light, its phenomena and laws. (R. 1865)	209
Lecture on zone of small planets between Mars and Jupiter. (R. 1854.)	75
On certain storms in Europe and America, December, 1836	127
Report on meteorology of the United States. (R. 1847)	
Lopholatilus chamæleonticeps, new genus and species of fish from New England, description of—G. B. Goode; T. H. Bean. (P. 1879)	
Tanahara Taliana W. I. Wannyame (D. 1966)	

•	
LOUD, F. H. Discussion of Prof. Snell's barometric observations. (R. 1880.) 442, 4	35
Louisiana, Caddo parish, Indian remains in—T. P. HOTCHKISS. (R. 1872) 2	271
Louisiana, lower, geology of, and salt deposit of Petite Anse Island—E. W.	40
	48 271
	856
	100
	147
Lowe, T. S. C., memorial of citizens of Philadelphia relative to aeronautic voyage of, across the Atlantic. (R. 1860)	147
Loxigilla, description of new subspecies of, from West Indies-G. N. LAW-	1 67
• •	150
·	228
	42
LUDEWIG, H. E., and others. On publication of Spanish works on New Mexico.	
(R. 1855)	77
LUGGER, O. Occurrence of Canada porcupine in Maryland. (P. 1881)	167
Lunar effect on magnetic force—A. D. BACHE	195
LUPTON, N. T. Breeding habits of sea catfish (Ariopsis milberti). (P. 1878.)	882
Luray cavern, blind form of Lysiopetalides from-J. A. RYDER. (P. 1880) 4	125
Luray cavern, report of visit to-O. T. MASON and others. (R. 1880) 442, 4	33
Lussac, Gay, eulogy on, by F. ARAGO. (R. 1876)	299
Lutjanus Blackfordii, from coast of Florida, description of—G. B. Goode; T. H. Bean. (P. 1878)	382
Lutjanus Stearnsii, from coast of Florida, description of-G. B. GOODE; T. H.	382
Lycodes paxillus, description of species of-G. B. GOODE; T. H. BEAN. (P. 1879)	383
	38
Lycodes Vahlii, occurrence of, on La Have and Grand Banks—G. B. Goode; T. H. Bean. (P. 1879)	388
Lycodopsis, Collett, identity of, with Leurynnis, Lockington-T. GILL. (P.	125
_	294
	328
_	150
	109
	244
	249
	25

M.

McCLINTOCK, F. L. Meteorological observations in the Arctic Seas	146
McConnell, E. M. Account of old Indian village, Kushkushkee, near New-	
castle, Pennsylvania. (R. 1871)	
McCosh, J. Prayer at memorial service of Prof. Joseph Henry	356
MACDONALD, J. A. Analyses of Chinese and Japanese coals	202
McHenry, J. W.; Haile, J. Antiquities of Jackson Co., Tenn. (R. 1874.)	280
McKAY, C. L. Review of Centrarchida. (P. 1881)	467
McKinley, W. Mounds in McIntosh and Early counties, Georgia. (R. 1872.)	271
McMasters, S. Y. Language of Navajos said to resemble the Welsh. (R. 1865.)	209
McParlin, T. A. Notes on history and climate of New Mexico. (R. 1877.) 323	, 396
McWhorter, T. Ancient mounds of Mercer county, Illinois. (R. 1874)	286
Mackerel, the frigate, (Auxis Rochei,) on New England coast-G. B. Goode.	
(P. 1880)	
Madrid Royal Academy of Science. Exchanges. (R. 1861)	
MAEDLER, J. H. Movement of the stars around a central point. (R. 1859)	
Magendie, F., memoir of, by M. FLOURENS. (R. 1866)	214
Magnetic—	
observations at Girard College, discussion of—A. D. BACHE: Part I. Eleven year period in amplitude of solar diurnal variation	113
Part II. Solar diurnal variation in magnetic declination	
Part III. Influence of moon on magnetic declination	
Parts IV, v, vI. Horizontal force	
Parts vii, viii, ix. Vertical force	
Parts x, x1, x11. Dip and total force	
_	
observations in the Arctic Seas—E. K. KANE	
observations, on continuance of—E. Sabine. (R. 1858)	
observations on the iron-clad "Monadnock"—W. HARKNESS	
observatory at the Smithsonian Institution, description of—J. E. Hil- GABD. (R. 1859)	110
storms—E. Sabine. (R. 1860)	
survey of Pennsylvania, (etc.)—A. D. BACHE	166
telegraph—See Telegraph.	
Magnetism—	
history of discovery relative to. (R. 1863)	
later views of connection of electricity and—H. Helmholtz; J. C. Maxwell. (R. 1878)	275
terrestrial, observations on-W. HARKNESS	
terrestrial, observations on-J. LOCKE	35
terrestrial observations on in Maxico Beron von Müller: A. Sonntag.	

,	
•	
	•
OF SMITHSONIAN PUBLICATIONS.	239
Magnus, Henry Gustavus, life and labors of. (R. 1870)	244
MAGRINI, L. Continuous vibratory movement of all matter. (R. 1868)	224
Mailly, E.—	
Estimate of population of the world. (R. 1878)	275
Eulogy on Quetelet. (R. 1874)	286
History of Royal Institution of Great Britain. (R. 1867)	215
Maine—	
Brunswick, results of meteorological observations at—P. CLEAVELAND.	204
Schoodic lakes, description of apparently new species of Gasterosteus from—T. H. BEAN. (P. 1879)	888
Kennebec river, opening and closing of—R. H. GARDINER. (R. 1858.)	109
Portland Society of Natural History, account of—E. C. Bolles. (R. 1867)	215
remains of walrus (?) in—C. H. Boyn. (P. 1881)	467
Makah Indians, account of the—G. GIBBS	220
Makah Indians, memoir on the—J. G. SWAN	220
Makah vocabulary—J. G. Swan	220
MALLERY, G. Sign language among North American Indians compared with that among other peoples and deaf-mutes. (E. 1879-80)	476
MALLET, R. On observations of earthquake phenomena. (R. 1859)	110
MALLINKROOT, C. Changes of wind. (R. 1866)	214
Maltheids, note on the—T. GILL. (P. 1878)	882
Mammals—	JU2
and birds of Missouri river, list of—E. HARRIS. (R. 1850)	28
arrangement of families of—T. GILL	230
bibliography of—T. GILL	280
extinct, of Nebraska—J. LEIDY	58
North American, catalogue of—S. F. BAIRD	105
of Arctic America—L. Kumlien	342
of Kerguelen Island—J. H. Kidder	294
synoptical tables of—T. GILL	280
Man-	
as the contemporary of the mammoth and reindeer in middle Europe. (R. 1867)	215
ancient, in Michigan, characteristics pertaining to—H. GILLMAN. (R. 1875)	393
later prehistoric, remains of, from caves in Alaska-W. H. Dall	
on traces of the early mental condition of—E. B. TAYLOR. (R. 1867)	215
social and religious condition of the lower races of—J. LUBBOCK. (R. 1869.)	228
Management of the Smithsonian Institution, report of Senate Judiciary Com-	
mittee on—A. P. Butler. (R. 1855)	77
Mandan ceremonies, accuracy of Catlin's account of-J. KIPP. (R. 1872)	271
MANN, C. Habits of a species of salamander. (R. 1854)	75
Manners of Shoshone Indians—A. G. BRACKETT. (R. 1879)	845

(R. 1879)
Manuscripts, Halliwell, report on-C. C. JEWETT. (R. 1852)
Manuscripts, etc., historical and geographical, catalogue of collection of—L. Beblandier. (R. 1854)
Manuscripts, linguistic, in library of Bureau of Ethnology, catalogue of— J. C. Pilling. (E. 1879-80)
Map of—
antiquities of United States, proposed—A. J. HILL. (R. 1861)
solar eclipse of March 15, 1858—T. HILL
stars for observations on meteorites
stars near North Pole for observations on auroras
United States, ethnological, proposed—G. GIBBS. (R. 1862)
Мара—
and charts of America, on a collection of-J. G. Kohl. (R. 1856)
catalogue of collection of—L. Berlandier. (R. 1854)
ethnological, of North America, suggestions relative to—L. H. Morgan. (R. 1861)
of winds of the globe—S. J. COFFIN
Marble used in extension of U.S. Capitol, account of—J. HENRY. (R. 1856.)
MARCET, Prof. Report on transactions of Society of Physics and Natural History of Geneva, July, 1862, to June, 1868. (R. 1868)
MAREY, E. J. Natural history of organized bodies. (R. 1867)
MARRY, E. J. Phenomena of flight in the animal kingdom. (R. 1869)
Marianini, Stefano, life and scientific labors of, by C. MATTEUCCI. (R. 1869.)
Marietta, Ohio, meteorological observations at—S. P. HILDRETH; J. WOOD.
Marine algse, lecture on-W. H. HARVEY. (R. 1855)
Marine algae of North America—W. H. HARVEY:
Part I. Melanospermea
Part II. Rhodospermea
Part III. Chlorospermeæ
Parts 1, 11, 111, complete
Marine invertebrates—
distributed by U. S. National Museum—R. RATHBUN. (P. 1881.) 465,
of Grand Manan-W. STIMPSON
of New England coast distributed by United States Fish Commission— A. E. VERRILL; R. RATHBUN. (P. 1879)
of northeastern coast of America, notice of recent additions to, with descriptions of new genera and species—A. E. VERRILL:
Part I. Annelida, etc. (P. 1879)
Part II. Mollusca, with notes on Annelida, etc. (P. 1880)
Part III. Catalogue of Mollusca recently added to fauna of southern
New England. (P. 1880)
on collection and preservation of-W. STIMPSON

OF SMITHSONIAN PUBLICATIONS.	241
Maritime digasters of the Antilles. (R. 1867)	215
Marrow of animals, uses of, among Indians—T. R. PEALE. (R. 1870)	244
Mars, secular variations of elements of orbit of-J. N. STOCKWELL	282
Mars, small planets between Jupiter and-Prof. LESPIAULT. (R. 1861)	149
Marschall, A., list of generic names supplemental to those catalogued by—S. H. SCUDDER	470
MARSH, G. P. Lecture on the camel. (R. 1854)	75
MARSH, G. P., and others. On publication of Squier and Davis' ancient mon- uments. (R. 1847)	
Marsipobranchii, arrangement of families of—T. GILL	247
MARTIN, S. D. Account of lightning discharges. (R. 1867)	215
Martinique, catalogue of Ober's collection of birds of—G. N. LAWRENCE. (P. 1878)	882
MARTINS, C. Vegetable colonization of the British Isles of Shetland, Faroe, and Iceland. (R. 1858)	109
Martius, C. F. P. von, honorary medal to-W. HAIDINGEB. (R. 1868)	187
Martius, C. F. P. von, memoir of, by C. RAU. (R. 1869) 440,	251
Maryland, Charles county, antiquities of-O. N. BRYAN. (R. 1874)	286
Maryland, occurrence of Canada porcupine in-O. LUGGER. (P. 1881)	467
MASON, C. Statement relative to the telegraph. (R. 1857)	107
MASON, O. T.—	
Abstracts of anthropological correspondence, 1880. (R. 1880)	442
Anthropological investigations in 1879. (R. 1879) 845,	420
Anthropological investigations in 1880. (R. 1880) 442,	
Bibliography of anthropology. (R. 1879; R. 1880) 845,	
Latimer collection of antiquities from Porto Rico. (R. 1876) 299,	
Leipsic Museum of Ethnology. (R. 1878)	
Record of progress in anthropology, 1879, 1880. (R. 1880) 442,	
Report on Luray Cavern. (R. 1880) 442,	
Summary of anthropological correspondence, 1879. (R. 1879) 845,	
Massachusetts—	
Buzzards' Bay, occurrence of Belone latimanus in—G. B. Goode. (P. 1878)	882
Provincetown, Cape Cod, littoral marine fauna of—R. RATHBUN. (P. 1880)	425
Vineyard Sound, occurrence of oceanic bonito in—V. N. EDWARDS. (P. 1878)	882
MASTERMAN, S. Observations on natural phenomena, shooting stars, aurora,	
etc. (R. 1857)	107
MASTERMAN, S. Observations on thunder and lightning. (R. 1855)	77
Materia medica collection, classification and arrangement of-J. M. FLINT.	
(P. 1881) 467,	450
Materia medica section of the U.S. National Museum, memoranda for collectors	
of drugs for the—J. M. FLINT. (P. 1881)467,	452

Matter, continuous vibratory movement of—L. Magrini. (R. 1868) 224
Matteucci, C.—
Electrical currents of the earth. (R. 1867; R. 1869) 215,228
Lectures on electro-physiology. (B. 1865) 299
Life and scientific labors of Stefano Marianini. (R. 1869)
Mauvaises Terres, expedition to the upper Missouri and the—T. A. CULBERT- SON. (R. 1850)
Mauvaises Terres of Nebraska, extinct mammalia and chelonia from the—J. LEIDY
MAXWELL, J. C.; HELMHOLTZ, H. Later views of connection of electricity and magnetism. (R. 1878)
Maya sculpture, remarks on ancient relic of—A. Schott. (R. 1871), 249
MAYBERRY, S. P. Gradual approach of sea upon land. (R. 1867) 215
MAYBERRY, S. P. Shell heaps at mouth of St. John's river, Florida. (R. 1877)
MAYER, A. M. Henry as a discoverer. Memorial address
MAYER, A. M., translation by, of Delaunay's essay on velocity of light. (R. 1864)
MAYER, B. Observations on Mexican history and archeology
Mazztlan, descriptions of new fishes from—D. S. JORDAN; C. H. GILBERT. (P. 1881)
MEACHAM, J. Minority report of Committee of Regents on distribution of Smithson income. (R. 1858)
MEADS, O. Memorial of Joseph Henry
Means available for correcting measures of sun's distance—G. B. AIRY. (R. 1859)110
Means of destroying the grasshopper-V. Motschulsky. (R. 1858) 109
Measurements, anthropological, table of—Scherzer; Schwarz. (R. 1866) 214
Measurements, mountain—A. Guyot. (R. 1862)
and weights, English and French, tables of. (R. 1863; R. 1864) 187, 183
and weights, metric system of, with tables—H. A. NEWTON. (R. 1865.) 209,371
and weights, new system of, with 8 as the metrical number—G. H. KNIGHT. (R. 1867)
of sun's distance—G. B. AIRY. (R. 1859)
Measuring instrument, use of galvanometer as—J. C. POGGENDORFF. (R. 1859.) 110
Mechanics and artisans, scientific education of—A. P. PEABODY (R. 1872.) 271,380
Mechanism of fever, study of. Toner lecture No. IV-H. C. Wood 282
Medal to Von Martius-W. Haldinger. (R. 1863)
Medical practice, relation of food to work and its bearing on—S. HAUGHTOF. (R. 1870)
Medicines, classification of forms in which drugs and, appear and are administered.—J. M. FLINT. (P. 1881)
Medulls oblongate and transzium, grav substance of the-J. DRAN

MERCH, L. W. Relative intensity of heat and light of the sun. (R. 1856.) 91,.83
MREK, F. B. Check-list of invertebrate fossils of North America. Oretaceous
and jurassic
MEEK, F. B. Check-list of invertebrate fossils of North America. Miocene. 183
MEEK, F. B, and others. Scientific instructions to Capt. Hall. (R. 1871) 249
MEEK, F. B.; HAYDEN, F. V. Paleontology of upper Missouri. Part 1 172
MEEK, J. B. Account of tornado in Spruce Creek valley, Centre county, Penn.
(R. 1871)
MEIGS, J. A. Description of human skull from Rock Bluff, Illinois. (R. 1867.) 215
Melanians of North America—G. W. TRYON, Jr 258
Melanospermeze of North America, descriptions of—W. H. HARVEY 32
Melbourne Mining Department. Exchange system. (R. 1865) 209
MELSHEIMER, F. E. Catalogue of described coleopters of the United States 62
Melting points, bibliography of-F. W. CLARKE 255
Melting points, tables of. Part I-F. W. CLARKE 255
Melting points, tables of. First supplement—F. W. CLARKE 288
Memoir-See Biography, Life, Eulogy.
Memoir of—
Agassiz, Louis, by R. P. STEBBINS. (R. 1878) 275
Babbage, Charles, by N. S. Dodge. (R. 1878) 275
Beautemps-Beaupré, C. F, by E. DE BEAUMONT. (R. 1868) 187
Blainville, Ducrotay de, by M. FLOURENS. (R. 1865) 209
Bravais, Auguste, by E. De Beaumont. (R. 1869) 228
Buch, Leopold von, by M. FLOURENS. (R. 1862) 150
Candolle, Pyramus de, by M. FLOURENS. (R. 1859)
Clark, H. J., by A. S. PACKARD
Cuvier, G., by M. FLOURENS. (R. 1868) 224
Delambre, J. B., by J. FOURIER. (R. 1864) 188
Encke, J. F., by G. Hagen. (R. 1868)
Gibbs, George, by J. A. STEVENS. (R. 1878)
Hauy, René Just, by G. CUVIER. (R. 1860) 147
Henry, Joseph, by A. Gray. (R. 1878) 841, 856, 407
Henry, Joseph, by J. LOVEBING
Henry, Joseph, by S. Newcomb
Henry, Joseph, by W. B. TAYLOR 856, 339
Herschel, Sir John Frederick William, by N. S. Dodge. (R. 1871) 249
Hodgkinson, Eaton, by R. RAWSON. (R. 1868) 224
Legendre, A. M., by E. DE BEAUMONT. (R. 1867) 215
Magendie, F., by M. FLOURENS. (R. 1868) 214
Martius, C. F. P. von, by C. RAU. (R. 1869) 228, 440, 251
Oersted, H. C., by E. DE BEAUMONT. (R. 1868) 224
Peltier, J. C. A., by F. A. Peltier. (R. 1867) 216

Memoir of—Continued.
Priestley, J., by G. CUVIER. (R. 1858)
Saint-Hilaire, Isidore Geoffroy, by J. L. QUATREFAGES. (R. 1862) 150
Saint-Hilaire, Geoffroy, by M. FLOURENS. (R. 1861) 149
Seaton, W. W., by J. HENRY. (R. 1866)
Thénard, Louis Jacques, by M. FLOURENS. (R. 1862) 150
Torrey, John, by A. GRAY. (R. 1878)
Memoir on—
anatomy and physiology of lucernarise—H. J. CLARK 242
explosiveness of nitre—R. HARE
extinct sloth tribe of North America—J. LEIDY
extinct species of American ox—J. LEIDY
meteorological subjects, by J. Hann and others. Translated by C. ABBL. (R. 1877)
methods of interpolation—E. L. DE FOREST. (R. 1871, 1878) 249.275
mosasaurus—R. W. Gibbes14
Memoranda for collectors of drugs for materia medica section of the U. S. National Museum—J. M. FLINT. (P. 1881) 467, 452
Memoranda of inquiry relative to food fishes of United States 231
Memorial—
address on Joseph Henry—A. M. MAYER 356, 417
of citizens of Philadelphia relative to Lowe's aeronautic voyage across the Atlantic. (R. 1860)
of Joseph Henry. (Published by Congress)
of Regents to Congress asking appropriations for Museum. (B. 1867.) 215, 329
of Regents to Congress relative to Smithson fund. (R. 1850; R. 1896.) 28, 214, 329
to Congress in behalf of Regents relative to new museum building—G. BANCROFT. (R. 1876)
Menhaden, gulf, (Brevoortia patronus), note on—S. STEARNS. (P. 1878) 332
Mental condition of man, early traces of—E. B. TYLOR. (R. 1867) 215
Mcrcury, secular variations of elements of orbit of, with tables—J. N. STOCK-
MERINO, St. M. Figure of the earth. (R. 1868)
Merlucius, diagnosis of genus of fish related to—G. B. Goode. (P. 1880) 425
MERRILL, J. C. Habits of Rocky Mountain goat. (P. 1879)
MERRILL, J. C. List of birds observed near Fort Brown, Texas, 1876-1878. (P. 1878)
Mesozoic rocks of Arkansas, Wyoming, Colorado, and Utah, descriptions of new invertebrate fossils from—C. A. White. (P. 1880) 45
Metal, hydrogen as gas and—J. E. REYNOLDS. (R. 1870)
Metallic castings of delicate natural objects. (P. 1881)
Metamorphism and formation of crystalline rocks—G. A. DAUBRÉE. (R. 1861.) 149
Metaorie fine hall subit and phonomena of T. H. Courry

OF SMITHSONIAN PUBLICATIONS.	245
Meteoric shower of November 18, 1867, letter of M. Hoek, relative to	_ 217
Meteoric stones, lectures on—J. L. SMITH. (R. 1855)	. 77
Meteorite—	
in Mexico—A. Woodworth. (B. 1867)	. 215
in Mexico, account of—R. Simson. (R. 1867)	. 215
in Mexico, discovery of large—W. M. PIERSON. (R. 1878)	
Tucson, account of—S. AINSA. (R. 1868)	. 187
Tucson, account of-B. J. D. IRWIN. (R. 1863)	. 187
Meteorites—	
extract from discourse on—F. Brenndecke. (R. 1869)	. 228
in Mexico-J. H. CARLETON. (R. 1865)	. 209
in mineralogical collection of Yale College, catalogue of—G. J. Brush (R. 1868)	
Synthetic experiments relative to-G. A. DAUBRÉE. (R. 1868)	. 224
Meteorograph, universal, for detached observatories—E. H. VON BAUMHAUER (R. 1879)	
Meteorological—	
articles in periodicals received by Smithsonian Institution and deposited in Library of Congress. (R. 1871; R. 1878)	
correspondence of the Institution, extracts from. (R. 1866)	
discovery-F. L. CAPEN. (R. 1866)	
instruments, description of—L. CASELLA. (R. 1859)	
material contributed to the Smithsonian Institution. (R. 1860-1866, 1868 1871, 1874) 147, 149, 150, 187, 188, 209, 214, 224, 228, 244, 24	,
memoirs—J. Hann; L. Sohncke; T. Reye; W. Ferrel; A. Cold ing; M. Peslin; C. Abbe. (R. 1877)	
publications deposited in Library of Congress, classified list of. (R. 1878.)	
reports preserved in Smithsonian Institution, classified list of. (R. 1878.)	
stations, cost of establishment of-J. HENRY. (R. 1858)	. 109
stations and observers, list of. (R. 1849, 1853–1872.) 21, 67, 75, 77, 91, 10' 110, 147, 149, 150, 157, 187, 188, 209, 214, 215, 224, 228, 244, 249, 271	
subjects, short memoirs on, by J. Hann and others. Translated by C ABBE. (R. 1877)828	
suggestions for Russian America-J. HENRY	207
system for every State—Illinois State Board of Education. (R. 1855.)	77
system of Canada—J. G. Hodgins. (R. 1865)	
system of Smithsonian Institution, report on—E. FOREMAN. (R. 1851) R. 1852)	;
tables—A. Guyor31	
telegrams by Western Union Telegraph Company—H. Sibler. (R. 1862)	
Keteorological observations—	, 200
at Brunswick, Maine—P. CLEAVELAND	204
at Marietta, Ohio—S. P. HILDRETH; J. WOOD	

•

ALPHABETICAL INDEX

Meteorological observations—Continued.
at Philadelphia, Girard College, discussion of-A. D. BACHE. 113, 121,
132, 162, 175, 196, 19
at Providence, Rhode Island, 1881-1860—A. CASWELL
at Providence, Rhode Island, 1881-1876-A. CASWELL
at Sacramento, California-F. W. HATCH. (R. 1854)
at Sacramento, California—T. M. LOGAN. (R. 1854)
at San Francisco, California—H. GIBBONS. (R. 1854)
at Washington, Arkansas-N. D. SMITH
combined, report of American Association for Advancement of Science on system of. (R. 1851)
directions for-A. GUYOT; J. HENRY. (R. 1855) 77, 19, 19
in the Arctic Seas—I. I. HAYES
in the Arctic Seas—E. K. KANE 10
in the Arctic Seas—F. L. McCLINTOCK14
results of, from 1854 to 1859. Vols. I, II157,18
Smithsonian, for the year 1855
Meteorology-
articles on. (R. 1867; R. 1870) 215, 24
articles on, with notes by J. Henry. (R. 1871)
communication relative to—J. P. Espy. (R. 1847)
lectures and notes on—R. RUSSELL; J. HENRY. (R. 1854)
of Caracas, South America—G. A. Ernst. (R. 1867)
of Colonia Tovar, Venezuela—A. FENDLEB. (R. 1857) 10
of District of Columbia—J. WISSNER. (R. 1857) 10
of Green river country, Utah—Col. Collins. (R. 1871)
of Porto Rico-G. Latimer. (R. 1871)
of Russia—A. Workoff. (R. 1872)
of Sacramento, California—T. M. LOGAN. (R. 1857) 10
of the United States, report on—E. Loomis. (R. 1847)
Meteorology-See Atmospheric pressure, Temperature, Rain, Winds, etc.
Meteors—
articles on, preserved in Smithsonian. (R. 1874)
observation of—L. Andrews. (R. 1866)
of November 18, 1867
planisphere of the heavens for observations of
Method of—
ascertaining amount of water in rivers—A. A. HUMPHREYS. (R. 1858.)
forming small weights—R. HARE. (R. 1858)
preserving lepidoptera—T. R. PEALE. (R. 1863)
Methods of—
capture and utilization of animals—G. B. Goods
installation in the National Museum-G. B. Goods. (P. 1881) 467, 472

Methods of—Continued.	
interpolation applicable to graduation of irregular series-E. L. Dz	
FOREST. (R. 1871)	249
interpolation, additions to memoir on—E. L. DE FOREST. (R. 1878)	275
making and preserving plaster casts—A. PIRZ. (P. 1881)	467
Metric system for scientific observation—A. GUYOT. (R. 1848)	I
Metric system of weights and measures with tables—H. A. NEWTON. (R. 1865.) 209,	371
Metric system with 8 as base—G. H. KNIGHT. (R. 1867)	215
METTENIUS, G. Acknowledgment for ferns. (R. 1862)	150
Mexican axolotl, change of, to an amblystoma—A. Weismann. (R. 1877.) 828,	401
Mexican boundary, forests and trees of Florida and the-J. G. Cooper. (R. 1860.)	147
Mexican history and archeology, observations on—B. MAYER	86
Mexican Society of Geography and Statistics. Exchanges. (R. 1861; R. 1865.) 149,	209
Mexico—	
description of large fossil gasteropod from-C. A. WHITE. (P. 1880)	425
description of new fishes from-D. S. JORDAN; U. H. GILBERT. (P. 1881.)	467
description of new species of genus Brevoortia from—G. B. Goode. (P. 1878)	832
eastern, earthquake in, January, 1866-C. Sartorius. (R. 1866)	214
explorations of John Xantus in-M. Romeno. (R. 1862)	150
Guanajuato and Chapala lake, notes on Dugès' collection of fishes from— D. S. JORDAN. (P. 1879)	883
	185
magnetic observations in—Baron von MÜLLER; A. SONNTAG	114
meteorite in—R. Simson. (R. 1867)	215
meteorite in-A. WOODWORTH. (R. 1867)	215
meteorite in, discovery of-W. M. PIERSON. (R. 1878)	275
meteorites in-J. H. CARLETON. (R. 1865)	209
notes on fishes from, collected by Lieut. H. E. Nichols—D. S. JORDAN; C. H. GILBERT. (P. 1881)	467
scientific expedition to, report addressed to Emperor of France by Minister of Public Instruction. (R. 1864)	188
southwestern, birds of, collected by F. E. Sumichrast—G. N. LAWRENCE.	295
Vera Cruz, account of antiquities in—H. FINCK. (R. 1870)	244
Mexico, Gulf of, catalogue of fishes collected in, by J. W. Velie-G. B. Goode; T. H. Bean. (P. 1879)	888
Mexico, Gulf of, mortality of fishes in—J. P. JEFFERSON. (P. 1878)	382
Mica beds in Alabama—W. GESNER. (R. 1879)	845
Mica mines, ancient, in North Carolina—C. D. SMITH. (R. 1876)	299
Michigan-	
ancient mining in—C. WHITTLESEY	155
Beaver Island, natural history of—J. J. STRANG. (R. 1854)	75

Michigan—Continued.
catalogue of rocks, minerals, and ores collected on geological survey in— C. T. Jackson. (R. 1854)
characteristics pertaining to ancient man in-H. GILLMAN. (R. 1875.) 298, 393
Isle Royale, Lake Superior, antiquities of-A. C. DAVIS. (R. 1874) 28
mound-builders and platycnemism in—H. GILLMAN. (R. 1873) 275, 393
Microscope, the. (R. 1860)
Microscopic organisms—
directions for collecting6
directions for collecting, preserving, and transporting-A. M. Edwards. 366
notes on new species and localities of-J. W. BAILEY
Microscopical examination of Japanese infusorial earths -A. M. Edwards 33
Microscopical examination of soundings made by U. S. Coast Survey off At-
lantic coast of United States-J. W. BAILEY. 20
Microscopical observations in South Carolina, Georgia, and Florida—J. W. BAILEY
Middle America—
birds of—S. F. BAIRD
birds of, circular for collecting 168
birds of, not in National Museum—R. RIDGWAY. (P. 1881) 45
(explanation of term) 18
Migration, the American—F. von Hellwald. (R. 1866) 28
Migrations and nesting habits of west coast birds—J. G. Cooper. (P. 1879) 334
Milan Agricultural Association. Exchange of publications. (R. 1863) 185
Milky way, bibliography of works relating to—E. S. Holden
MILLER, F. Mound in Trumbull county, Ohio. (R. 1877)
Mills, Clark, casts of heads of Indian boys and girls at Hampton, Virginia,
taken by. (P. 1879)
Mills, Clark, casts of heads of Indian prisoners in Florida taken by. (P. 1878.) 332
Mineral, jade, study of—S. BLONDEL. (R. 1876)
Mineralogical collection, arrangement of—C. U. SHEPARD. (R. 1861) 149
Mineralogical collection of Yale College, catalogue of meteorites in the—G. J. Brush. (R. 1868)
Mineralogical composition of the normal mesozoic diabase upon Atlantic border—G. W. HAWES. (P. 1881)
Mineralogy, progress in, in 1879 and 1880—G. W. HAWES. (R. 1880) 442, 428
Minerals—
and ores collected on geological survey in Michigan, catalogue of—C. T. JACKSON. (R. 1854)
and rocks, catalogue of-J. W. FOSTER. (R. 1854)
and rocks, catalogue of—J. D. WHITNEY. (R. 1854)
catalogue of, with their formulas, etc.—T. EGLESTON156
in U. S. National Museum in 1878, list of-F. M. ENDLICH. (R. 1873.) 275

ALPHARETICAL INDEX

Missouri—Continued.
prehistoric evidences in-G. C. BROADHEAD. (R. 1879) 34
St. Louis, ancient mound at—T. R. PEALE. (R. 1861)
St. Louis, temperature of—A. FENDLER. (R. 1860)
upper, expedition to Mauvaises Terres and—A. Culbertson. (R. 1850.)
upper, list of plants of—T. C. PORTER. (B. 1850)
upper, palsontology of—F. B. MEEK; F. V. HAYDEN
western, exploration of—P. R. Hoy. (R. 1864)
Missouri river, ancient earthworks on—A. Barrandt. (R. 1870)
Missouri river, list of birds and mammalia of—E. HARRIS. (R. 1850)
MITCHELL, A. Antiquities of Florida. (R. 1874)
MITCHELL, B. Mounds in Pike county, Illinois. (R. 1879)
MITCHELL, B. R.; TURNER, W. W. Vocabulary of the jargon of Oregon 6
MITCHELL, S. S. Funeral sermon of Joseph Henry
MITCHELL, S. W. Inquiries relative to disease known as chorea. (R. 1874) 28
MITCHELL, S. W. Researches upon the venom of the rattlesnake
MITCHELL, S. W.; Morehouse, G. R. Researches upon anatomy and physi-
ology of respiration in chelonia
Mitla, drawings of—J. G. SAWKINS
Mixed races in Liberia—E. D. BLYDEN. (R. 1870)24
Mobile river, shell-heaps on—A. S. GAINES; K. M. CUNNINGHAM. (R. 1877.)
Modern theory of chemical types—C. M. WETHERILL. (R. 1863)
Modes of flight in relation to aeronautics—J. B. Pettigrew. (R. 1867) 21
Moigno, F. N. Accidental or subjective colors. (R. 1866)
MOLINA, L., Costa Rican minister, recommending Dr. Berendt. (R. 1865)
Mollusca of Northeastern coast of America. Parts I, II, III—A. E. VERBILL.
(P. 1879; P. 1880)
Mollusca, or shellfish, and their allies, lectures on—P. P. CARPENTER. (R. 1860)
Molluscoids of Arctic America—A. E. VERRILL
Mollusks-
arrangement of families of-T. GILL 27
bibliography of—T. GILL
eocene, some new species of, from southern United States—A. Heilpris. (P. 1880)
new forms of, from Alaska-W. H. Dall. (P. 1878)
of Arctic America—W. H. Dall
of Kerguelen Island—W. H. Dall
of western North America—P. P. CARPENTER252
Monadrock, ironclad, deviation of compasses on-W. HARKNESS
Mongolia, geological researches in—R. PUMPELLY 202
Monograph of—
American corbiculadæ, recent and fossil—T. PRIME

OF SMITHSONIAN PUBLICATIONS.	25
Monograph of-Continued.	
bats of North America—H. ALLEN	16
cottoids of North America—C. GIRARD	`a
dolichopodidæ-H. Loew	1
ortalidæ-H. Lorw	2
strepomatidæ, (American melanians)-G. W. TRYON	25
tipulidæ—C. R. Osten Sacken	2
wasps—H. DE SAUSSURE	
Monographs of dipters of North America—	
Part I-H. LOEW; R. OSTEN SACKEN	14
Part II-H. LOEW; R. OSTEN SACKEN.	
Part III—H. Lorw	
Part IV-R. OSTEN SACKEN	
Montana, distribution of forest trees in-W. W. Johnson. (R. 1870)	
Montana, prehistoric remains in—P. W. Norris. (R. 1879)	
Monterey Bay, California, new fishes from-D. S. JORDAN; C. H. GILBER	T.
(P. 1880)	
Monument to Joseph Henry, proceedings of Congress relative to	8
Monuments—	
aboriginal, of State of New York—E. G. SQUIER	:
ancient, of the Mississippi valley—E. G. SQUIER; E. H. DAVIS	
and antiquities of Denmark, preservation of J. A. Worsaar. (R. 1878)	9.) 8
Moon—	
effect of, on the magnetic force—A. D. BACHE	
effect of, on the weather—J. HENRY. (R. 1871)	
influence of, on magnetic declination—A. D. BACHE	1
occultations of planets by—See Downes, J.	
Moon-hoax, on the-J. HENRY. (R. 1873)	2
MOORE, M. A. Fish mortality in the Gulf of Mexico. (P. 1881)	4
MOORE, T., and others. Destruction of fish near the Tortugas. (P. 1878)	8
Morehouse, G. W.; MITCHELL, S. W. Anatomy and physiology of respir	
Мовятт, С; Воотн, J. С. Recent improvements in chemical arts	
Morgan, L. H.—	
Circular in reference to degrees of relationship among different nations	1
Proposed ethnological map of North America. (R. 1861)	
Systems of consanguinity and affinity of the human family	
Morgan, L. H., circular to diplomatic agents by State Department relative	
	1
MORIN, A. Warming and ventilating buildings. Parts 1, 11. (R. 1878; 1874) 275, 2	
Morlot, A.—	. • -
Abstract of Dr. Keller's report on lacustrian settlements in Switzerlan	d.
(R. 1868)	

.

.

.

MORLOT, A.—Continued.
Archæology. (R. 1861)149
General views on archeology. (R. 1860)
Lecture on study of high antiquity in Europe. (R. 1862; R. 1864.) 150, 188
Morris, J. G.—
Catalogue of described lepidopters of North America 118
Lecture on insect instincts and transformations. (R. 1855)
Lecture on natural history as applied to farming and gardening. (R. 1855.)
Synopsis of described lepidopters
MORRIS, O. W.; HENRY, J. Quantity of rain at different heights. (R. 1855.)
Morse, S. F. B., account of origin and development of invention of—W. B. TAYLOR. (R. 1878)
Morse, S. F. B., communication relative to a publication of—J. HENRY. (R.
1857) 107, 115. 829
Mortality—
of fishes in Gulf of Mexico—E. INGERSOLL. (P. 1881)457
of fishes in Gulf of Mexico—J. P. JEFFERSON. (P. 1878)
of fishes in Gulf of Mexico—S. H. Johnson. (P. 1881)467
of fishes in Gulf of Mexico—M. A. Moore. (P. 1881) 46
of fishes in vicinity of Tortugas—J. P. JEFFERSON; T. MOORE; J. Y. PORTER. (P. 1878)
tables of, methods of interpolation applied to graduation of—E. L. Dz. Forest. (R. 1871; R. 1873) :249, 275
MORTILLET, G. DE; CHANTER, E. International code of symbols for charts of prehistoric archeology. (R. 1875)
MOBTON, S. G., and others. On publication of Squier and Davis' ancient monuments. (R. 1847)
Mortuary customs of North American Indians—H. C. YARROW. (E. 1879-80.) 476
Mosasaurus and the three allied genera, Holcodus, Conosaurus, and Amphoros- teus, memoir on—R. W. GIBBES
Motion, planetary, general integrals of—S. Newcomb281
Motion, rotary, problems of, presented by gyroscope, precession of equinoze, and pendulum—J. G. BARNARD
MOTSCHULSKY, V. On means of destroying the grasshopper. (R. 1858) 109
Moulton, M. W. Mounds in Delaware county, Iowa. (R. 1877)
Mound in—
Dakota, Lincoln county—A. BARRANDT. (R. 1872)
Florida, colored bead from—A. M. HARRISON. (R. 1877) 323
Florida, gold ornament from—C. RAU. (R. 1877) 323, 440, 403
Indiana—W. Pidgeon. (R. 1867) 215
Kentucky, near Lexington—R. PETER. (R. 1871) 249
Missouri, St. Louis-T. R. Peale. (R. 1861)
Ohio, Trumbull county—F. MILLER. (R. 1877) 323
Pennsylvania, western—W. M. TAYLOR. (R. 1877)

Mound in-Continued.
Tennessee, near Chattanooga—M. C. READ. (B. 1867) 215
Tennessee, east—A. F. Danilsen. (R. 1868)
Wisconsin—C. K. DEAN. (R. 1872)
Wisconsin, Grant county-J. WARNER. (R. 1872) 271
Mound-builders—
and platycnemism in Michigan—H. GILLMAN. (R. 1878) 275, 393
in Rock river valley, Illinois—J. SHAW. (R. 1877) 828
of Wisconsin, mounds and osteology of—J. N. DE HART. (R. 1877) 828
Mounds and osteology of the mound-builders of Wisconsin-J. N. DE HART.
(R. 1877)
Dakota, Fort Wadsworth-A. J. Comport. (R. 1871) 249
Florida, (southern,) explorations among—S. T. WALKER. (R. 1879) 845
Georgia-M. F. Stephenson. (R. 1870) 244
Georgia, Bartow county-M. F. STEPHENSON. (R. 1872) 271
Georgia, McIntosh and Early counties—W. McKinley. (R. 1872) 271
Illinois, Mercer county—T. McWhorter. (R. 1874) 286
Illinois, near Albany, skull and bones from-R. J. FARQUHARSON.
(R. 1874) 286
Illinois, Pike county—B. MITCHELL. (R. 1879) 845
Illinois, Rock Island county—T. THOMPSON. (R. 1879)
Illinois, Spoon river valley—W. H. ADAMS. (R. 1879) 845
Illinois, Union county, near Anna-T. M. PERRINE. (R. 1872) 271
Indiana, Franklin county—E. R. Quick. (R. 1879) 846
Indiana, Rush county—F. JACKMAN. (R. 1879) 845
Iowa, Delaware county—M. W. Moulton. (B. 1877)
Iowa, Des Moines valley, notes on -S. B. Evans. (R. 1879)
Iowa, Muscatine county—T. THOMPSON. (R. 1879) 346
Kentucky—R. Peter. (R. 1872) 271
Kentucky, exploration of—S. S. Lyon. (R. 1870) 244
Kentucky, Union county, exploration of—S. S. LYON. (R. 1870) 244
Louisiana—S. H. Lockett. (R. 1872) 271
Louisiana, Moorehouse parish—B. H. BRODNAX. (R. 1879) 846
Minnesota, Hennepin county, on Gideon's farm—F. H. NUTTER. (R. 1879) 846
Mississippi—S. A. Agnew. (R. 1867)
Mississippi, Washington county—J. Hough. (R. 1879) 846
New Brunswick and New England, notes on—S. F. BAIRD. (P. 1881.) 467
Ohio, Trumbull county—F. MILLER. (R. 1877) 828
Pennsylvania, (western)—W. M. TAYLOB. (R. 1877) 828
Tennessee-J. Jones 256
Wisconsin, Grant county—M. STRONG. (R. 1879) 296
Wisconsin Winnehogo county T ARMSTRONG (R. 1879) 84

Mountain, Black, North Carolina, topography of—T. L. CLINGMAN. (R. 1856.)	77
Mountain measurements—A. Guyor. (R. 1862)	150
Mountains, electric resonance of—H. DE SAUSSURE. (R. 1868)	24
Mountains in Colorado, heights of-G. Engelmann. (B. 1862)	150
Mountains, lakes, and the snow-line, Norwegian—O. E. DEEUTZER. (R. 1866.)	214
Movement of the stars around a central point—J. H. MAEDLER. (R. 1859)	110
MUCH, M. Ancient history of North America. (R. 1871)	249
MUDGE, B. F. Account of lightning discharges. (R. 1867)	215
MÜLLER, Baron von. Observations on terrestrial magnetism in Mexico	114
MÜLLER, F. Smithsonian exchanges. (R. 1860)	147
MÜLLER, J.—	
Principles of the mechanical theory of heat. (R. 1868)	224
Report on recent progress in physics—electricity. (R. 1856; R. 1857.) 91,	107
Report on recent progress in physics—galvanism. (R. 1855; R. 1857; R. 1858) 77, 107.	109
Report on transactions of Geneva Society of Physics and Natural His-	
tory, June, 1875, to July, 1876. (R. 1877)	
Mummy and skulls from Patagonia—A. REID. (R. 1862)	
Mummy case, Gliddon—C. PICKERING	108
MURPHEY, H. C., and others. On publication of Spanish works on New Mexico. (R. 1855)	;;
Musci of Kerguelen Island—T. P. JAMES.	294
Museum—	
at Lausanne, ethnological collections of-F. TROYON. (R. 1861)	
Blackmore, Salisbury, England, notice of. (R. 1868)	
British, statistics of—S. F. BAIRD. (R. 1850)	25
d'Histoire Naturelle, Paris, notes on typical American fishes in—D. S. JORDAN. (P. 1879)	333
Hungarian National. Acknowledgment for birds. (R. 1863)	
miscellanea—S. F. BAIRD	
of ethnology, Leipsic-A. SCHOTT; O. T. MASON. (R. 1878)	
of National University of Greece. Exchange of specimens. (R. 1867.)	
of Norway, Ethnological-L. K. DAA. (R. 1862)	
of Toronto University, objects of. (R. 1865)	209
on formation of a-L. Agassiz. (R. 1849)	21
Smithsonian—See Museum, U. S. National.	
Museum, U. S. National—	
acts and resolutions of Congress relative to—See Congress.	
additions to, list of—See each annual report.	
birds of Middle and South America not in—R. RIDGWAY. (P. 1881)	467
Bulletins of—See Bulletins.	
catalogue and index of the Proceedings and Bulletins of	4.8
catalogue of collection of Japanese cotton fibre presented to. (P. 1881.)	467

Museum, U. S. National—Continued.
catalogue of collection of Japanese woods presented to—L. F. WARD.
(P. 1881) 467
catalogue of Old World birds in-R. RIDGWAY. (P. 1881) 467, 462
circular asking contributions to library of. (P. 1881) 467, 458
circulars of—See Circulars.
classification of collections of-G. B. GOODE. (P. 1881) 467, 457
classification of collections of, to illustrate taxidermy—W. T. HORNADAY. (P. 1881)467, 456
collections furnished to, by explorations, 1888 to 1877—S. F. BAIRD. (R. 1877) 823
contributions to, and their acknowledgment. (P. 1881) 467, 473
directory of officers, collaborators, and employés of 466
distribution by, of marine invertebrates—R. RATHBUN. (P. 1881.) 467, 465, 471
establishment and officers of Smithsanian Institution and. (P. 1881.) 467, 449
letters on work of—B. PHILLIPS. (P. 1881)
list of publications of. (P. 1881) 467, 478, 474
memorial of Regents of Smithsonian Institution to Congress asking
appropriations for. (B. 1867)
memorial to Congress in behalf of Regents of Smithsonian Institution
for new building for—G. BANCROFT. (R. 1876) 299, 829
organization and objects of—G. B. Goode. (P. 1881) 467, 459
outline of scheme of classification for collections in—G. B. Goode. (P 1881)
plan of organization and regulations of—G. B. Goode. (P. 1881)_467, 445
plans for installation of collections in—G. B. Goode. (P. 1881) 467, 472
proceedings in Congress relative to 328
Proceedings of. Vols. I-IV, 1878-1881 332, 333, 425, 467
report of Committee of Regents on-A. GRAY; H. COPPÉE. (R. 1874.) 286
report of Committee of Regents on—A. GRAY; A. A. SARGENT; H. CLYMER. (R. 1876)
report on, and statistics of British Museum—S. F. BAIRD. (R. 1850) 28
reports of Building Commission of, for 1879, 1880. (R. 1879; R. 1880.)
845, 442, 409, 434
reports on, 1858-1867, 1878-1877—S. F. BAIRD. (R. 1858-1867, 1878-1877) 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187,
188, 209, 214, 215, 275, 286, 298, 299, 328
Mysdestes obscurus, description of two new races of—L. STEJNEGER. (P. 1881.) 467
Myctophum crenulare, description of—T. H. BEAN. (P. 1881) 467
Myctophum crenulare from Santa Barbara channel, California, description of—
D. S. JOBDAN; C. H. GILBERT. (P. 1880)
Myological specimens, rapid preparation of—F. Plateau. (P. 1881) 467

Myriapods, list of North American species of, belonging to family of Lyriope- talidæ—J. A. RYDER. (P. 1880)
Myriapods, phalangidse, etc., instructions for collecting—H. C. Wood. (R. 1866)
Myriolepis zonifer, a new chiroid fish from Monterey Bay, Calfornia, descrip-
tion of—W. N. LOCKINGTON. (P. 1880)—————————425
Mythology of North American IndiansJ. W. Powell. (E. 1879-80) 476
Myths, Indian. (E. 1879-80) 476
·
N.
•
NAILL, D. W. Dispersion of a cloud by electrical discharge. (B. 1858) 100
Naples, Italy, Royal Institute for Encouragement of Natural, Economical, and
Technological Sciences. Prize questions. (R. 1878) 275
Narrative of the Hassler expedition—L. AGASSIZ. (R. 1872)
National Academy of Sciences, bequest to—A. D. BACHE. (R. 1872) 271,829
National library, on a-W. S. JEVONS. (R. 1878) 275
National Museum—See Museum, U. S. National.
Natural history—
applied to farming and gardening, lecture on—J. G. Morris. (R. 1855.)
bibliography of American—C. GIRARD 48
directions for collecting, preserving, and transporting specimens of— S. F. Baird. (R. 1856)
explorations in, in the United States, 1851—S. F. BAIRD. (R. 1851) 51
of Arctic America—L. Kumlien
of Beaver Island, Michigan—J. J. Strang. (R. 1854) 75
of country about Fort Ripley, Minnesota—J. E. HEAD. (R. 1854) 75
of Fanning Islands—T. H. STREETS
of fresh-water fishes of North America—C. GIRARD
of Hawaiian Islands, etc.—T. H. STREETS
of Kerguelen Island—J. H. KIDDER and others 294
of lower California—T. H. Streets
of organized bodies-E. J. MARRY. (R. 1867) 215
of Pacific Islands—W. H. PEASE. (R. 1862)
suggestions relative to investigations in, in Russian America—8. F. Baird
Natural History Society—See Geneva.
Natural method, the Jussieus and the—M. FLOURENS. (R. 1867)
Naturalists, generic names of animals employed by-S. H. SOUDDER470

Nature—	
and art, on tables of the constants of-C. BABBAGE. (R. 1856)	91
and mechanism of fever. Toner lecture No. IV-H. C. Wood	282
and origin of force-W. B. TAYLOR. (R. 1870) 244	, 875
of reparatory inflammation in arterice after ligature. Toner lecture No.	321
Nature—See Constants of Nature.	
Navajo Indians, language of, said to resemble Welsh—S. Y. McMasters. (R. 1865)	209
Navajo Indians, sketch of the-J. LETTERMAN. (R. 1865)	77
Naval officers, authority to, to receive and transmit specimens to Smithsonian	84
Navigation of Ohio and other rivers, improvement of-C. ELLET	13
Navy Department, authority given by, to naval officers to receive and transmit	
specimens to the Smithsonian Institution	84
Neah Bay, Washington Territory, description of two new species of fishes from—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Nebraska-	
ancient fauna of—J. LEIDY	58
palsontology of-F. B. MERK; F. V. HAYDEN	172
report on fossils from—J. LEIDY. (R. 1851)	51
Nebulæ, index-catalogue of books relative to—E. S. Holden	311
Nebulæ, researches relative to the-Prof. GAUTIEE. (R. 1863)	187
Nebular hypothesis, bibliography of the-E. S. HOLDEN	811
Nematoda of northeast coast of North America - A. E. VERRILL. (P. 1879)	888
Nemertina of northeast coast of America—A. E. VERRILL. (P. 1879)	888
Nemichthys avocetta, new species of Nemichthys, description of—D. S. Jordan; C. H. Gilbert. (P. 1880)	425
Neptune—	
ephemerides of, 1848, 1846-49, 1850, 1851, 1852-S. C. WALKER. 4, 5, 6, 7	7, 24
history of discovery of—B. A. GOULD	18
investigation of orbit of, with tables of its motion—S. NEWCOMB	199
researches relative to—S. C. WALKER	3
secular variations of elements of orbit of-J. N. STOCKWELL	232
Nereis-Boreali-Americana—W. H. HARVEY: Part I. Melanospermeæ	32
Part II. Rhodospermeæ	43
Part III. Chlorospermes	95
Parts I, II, complete	96
Neritide, fresh-water—W. G. BINNEY	144
Nerve, polarized, effect of irritation of—B. F. LAUTENBACH. (B. 1878)_ 841,	
Nervous system of Rana pipiens, anatomy of—J. WYMAN	45
Nests—	
and eggs of American birds, instructions for collecting—S. F. BAIRD. (R. 1858)	139

Nests—Continued.
and eggs of the eight North American species of empidonaces—T. M. Brewer. (P. 1879)
instructions for collecting and preserving—T. M. BREWER139
of west coast birds-J. G. Cooper. (P. 1879)
Netherlands, the, Royal Academy of. Prize questions. (R. 1861) 149
Neuchâtel, lake of, palafittes or lacustrian constructions of—E. Desos. (R. 1865)
Neuroptera-
instructions for collecting-P. R. UHLER. (R. 1858) 109
North American, synopsis of—H. HAGEN. (R. 1861) 149, 134
of Arctic America—S. H. SCUDDER 342
of North America, synopsis of—H. HAGEN134
of North America, synopsis of—P. R. Uuler
pseudo-, of Kerguelen Island—H. A. HAGEN
Nevada, western, Centennial mission to Indians of—S. Powers. (R. 1876) 299 New-
classification of clouds—A. Poër. (R. 1870)
species and localities of microscopical organisms, notes on—J. W. BAILEY.
species of North American coleoptera. Parts I, II-J. L. LE CONTE. 167, 264
system of weights and measures, with 8 as metrical number—G. H. KNIGHT. (R. 1867)
tables for determining values of coefficients in perturbative function of planetary motion—J. D. RUNKLE
New Brunswick— Bay of Fundy, marine invertebrata of region about mouth of—W. Stimpson 50
shell heaps in—J. Fowler. (R. 1870)
shell mounds of—S. F. BAIRD. (P. 1881)467
New England—crustacea dredged off south coast of—S. I. SMITH. (P. 1880) 425
fishes from south coast of, descriptions of new genera and species of—6. B. Goode. (P. 1880)
frigate mackerel (Auxis Rochei) on coast of-G. B. Goode. (P. 1890) 425
geology of—E. Hitchcock
marine invertebrata from, distributed by U. S. Fish Commission—A. E. Verrill; R. Rathbun. (P. 1879)
marine invertebrates from, distributed by U. S. National Museum-R. 'RATHBUN. (P. 1881)467, 465, 471
mollusca added to fauna of—A. E. VERRILL. (P. 1880)
shell mounds on coast of—S. F. BAIRD. (P. 1881)67
New Harmony, Indiana, account of tornado near, April 80, 1852—J. CHAPPEL-8MITH
NEW HAVEN JOURNAL. Account of lightning discharges. (R. 1867) 215

New Jersey—	
artificial shell deposits in—C. RAU. (R. 1864) 188, 440,	
fishes on the coast of Long Island and—S. F. BAIRD. (R. 1854) 75,	348
stone age in—C. C. Abbott. (R. 1875)	394
New Jersey Historical Society, address of Prof. Henry before the	B
New Madrid, Missouri, earthquake at—T. DUDLEY. (R. 1858)	109
New Mexico—	
account of Lindheimer's, Fendler's, and Wright's botanical explorations	
in—A. Gray. (R. 1849)	21
antiquities in-W. B. Lyon. (R. 1871)	249
communications relative to publication of Spanish works on—E. Robinson; H. E. Ludewic; E. G. Squier; H. C. Murphry; W. B.	
HODGSON; W. IRVING; W. H. PRESCOTT; J. SPARKS; G. BAN- CROFT; F. L. HAWKS. (R. 1855)	7 7
description of new cretaceous Pinna from—C. A. WHITE. (P 1880)	
•	425 75
diary of excursion in—J. H. Carleton. (R. 1854)eastern, coleoptera of Knnsas and—J. L. Le Conte	
notes on the history and climate of—T. A. McParlin. (R. 1877). 823,	
plants of, collected by C. Wright. Parts 1, 11—A. Gray 2	
report of explorations in Arizona and—J. STEVENSON. (R. 1880)	
New York—	442
aboriginal monuments of—E. G. SQUIER	15
Clinton, Humilton College. Exchange of specimens. (R. 1861)	149
fishes of—T. Gill. (R. 1858)	91
Fishkill, deposit of arrow-heads near—E. M. Shepard. (R. 1877)	828
New York, explosion of nitre in, July, 1845-R. HARE	17
Orleans county, antiquities of—F. H. Cushing. (R. 1874)	286
Schoharie, Indian relies from—F. D. Andrews. (R. 1879)	845
Tompkins county, ancient fort and burial ground in—D. TROWBRIDGE.	010
(R. 1863)	187
University of State of, Regents of. Acknowledgment of specimens. (R. 1865)	209
NEWBERRY, J. S. Description of fossil plants from Chinese coal-bearing rocks.	202
Newcastle, Pennsylvania, account of old Indian village near—E. M. McCon-	202
NELL. (R. 1871)	249
NEWCOMB, S.—	
General integrals of planetary motion	281
Investigation of orbit of Neptune, with tables of its motion	199
Investigation of orbit of Uranus, with tables of its motion	262
Memorial address on Joseph Henry	856
NEWCOMB, S., and others. Scientific instructions to Capt. Hall. (R. 1871)	249
Newfoundland, new species of fishes from Grand Banks of—G. B. Goods. (P. 1880)	425
NEWTON. A. Suggestions for forming collections of birds' eggs	189

NEWTON, A. Suggestions for saving parts of skeletons of birds. (R. 1860)	147
NEWTON, H. A. Metric system of weights and measures, with tables. (R. 1865)	
Nicaragua—	
antiquities of—E. G. SQUIER. (R. 1850)	28
archeological researches in—J. F. Bransford	383
eruption of volcano in—A. B. DICKINSON. (R. 1867)	215
Nichols, Lieut. H. E.—	
list of fishes collected by, in Gulf of California and on west coast of lower	
California-D. S. JORDAN; C. H. GILBERT. (P. 1881)	467
notes on fishes collected by, in British Columbia and southern Alaska— T. H. BEAN. (P. 1881)	
notes on fishes collected by, on west coast of MexicoD. S. JORDAN; C.	
H. GILBERT. (P. 1881)	467
NICKLÉS, F. J. Scientific Congress of Carlsruhe, 1858. (R. 1860)	147
Nile, explorations of the—C. HALE. (R. 1865)	209
Nitre, memoir on explosiveness of—R. HARE	17
Nitrification—B. F. CRAIG. (R. 1861)	149
Nitrogen bodies of modern chemistry—Prof. KLETZINSKY. (R. 1872)	271
Nomenclator zoologicus. Alphabetical list of generic names employed by naturalists. Part I—S. H. SCUDDER	470
Nomenclature of—	
certain North American birds, revisions of—R. RIDGWAY. (P. 1880)	
clouds	
North American birds—R. Ridgway	
NORRIS, P. W. Prehistoric remains in Montana. (R. 1879)	849
North America—	
abbreviations of names of States and Territories of	
alge of, fresh-water, history of—H. C. Wood, Jr	341
algæ of, marine—See W. H. Harvey.	
ancient aboriginal trade in—C. RAU. (R. 1872) 271, 440,	
ancient history of—M. Much. (R. 1871)	249
atlas of, physical, proposed—G. GIBBS. (R. 1866)	714
batrachia and reptiles of, check-list of—E. D. COPE	292
bats of, monograph of—H. ALLEN	165
birds added by L. Belding to fauna of-R. RIDGWAY. (P. 1881)	467
birds of, catalogue of—S. F. BAIRD	106
birds of, instructions for collecting nests and eggs of	L39
birds of, nomenclature and catalogue of-R. RIDGWAY	22
birds of, review of. Part I—S. F. BAIRD	81
botany of, bibliographical index to—S. WATSON	58
coleoptera of, classification of. Parts I, II—J. L. LE CONTE 136, 2	65
coleopters of, list of—J. L. LE CONTE.	40

North America—Continued.	
coleoptera of, new species of. Parts I, II—J. L. LE CONTE 167,	264
combined meteorological system for, report of Committee of American	
Association for Advancement of Science on. (R. 1851)	51
conchology of, bibliography of. Parts I, II-W. G. BINNEY 142,	
contributions to natural history of fresh-water fishes of. Part 1. Mono-	
graph of cottoids-U. GIRARD	30
diptera of, catalogue of-C. R. OSTEN SACKEN 102,	27 0
diptera of, monographs of-H. LOEW; R. OSTEN SACKEN. 141, 171, 219,	256
duck new to fauna of—R. RIDGWAY. (P. 1881)	467
explorations on western coast of-W. H. Dall. (R. 1873)	275
extinct sloth tribe of—J. LEIDY	72
fishes of east coast of, catalogue of—T. GILL	283
fishes of Pacific coast of, duplicates of, distributed in 1881—D. S. Jor- DAN; P. L. JOUY. (P. 1881)	467
forests and trees of, distribution of-J. G. Cooper. (R. 1858) 109,	351
grasshoppers of, circular relative to history of	163
Indians of, uses of brain and marrow of animals among-T. R. Peale.	244
lakes of, fluctuations of level of—C. WHITTLESEY	
lepidoptera of, catalogue of-J. G. Morris	
lepidopters of, synopsis of—J. G. Morris	
libraries, public, in, list of-W. J. RHEES	
mammals of, catalogue of—S. F. BAIRD	
map of, ethnological, suggestions relative to-L. H. MORGAN. (R. 1861.)	
meteorological stations and observers in. (R. 1868) 224,	
mollusks of western—P. P. CARPENTER.	
myriapods of, belonging to family Lysiopetalidæ—J. A. RYDER. (P. 1880)	
neuroptera of, synopsis of—H. Hagen	
orthopters of, catalogue of—S. H SCUDDER	
rain and snow fall of, tables of—C. A. Schott 222, 3	
reptiles of, catalogue of—S. F. BAIRD; C. GIRARD	
shells of, catalogue of—I. Lea; P. P. Carpenter; W. Stimpson; W. G. Binney; T. Prime	
shells of, circular in reference to collecting	
shells of, land and fresh-water. Parts I-IV-W. G. BINNEY; T. BLAND.	
143, 144, 194, 2	253
silurids of, synopsis of—D. S. Jordan	806
stone implements of—C. RAU. (R. 1872)	382
North American—	
archeology-J. Lubbock. (R. 1862)	150
birds, catalogue of S. F. BAIRD	LOS

North American—Continued.	
birds, catalogue of-R. RIDGWAY. (P. 1880)	122
birds, desiderata among—R. RIDGWAY. (P. 1881)	
Continent, physical geography of-J. FROEBEL. (R. 1854)	
fishes, duplicates of, distributed by Smithsonian Institution—T. H. Bear. (P. 1880)	
Indians, catalogue of portraits of—J. M. STANLEY	
Indians, list of photographic portraits of	
stone implements—C. RAU. (R. 1872)	
stone period, agricultural implements of—C. RAU. (R. 1863) 187,	
tortoises, land, of genus Xerobates—F. W. TRUE. (P. 1881)	
North American Ichthyology, contributions to—	•••
Part 1. Review Rafinesque's memoirs—D. S. JORDAN.	305
Part II. A. Notes on Cottidæ, (etc.) B. Synopsis Siluridæ—D. S. JORDAN	
Part III. A. Distribution of fishes of Alleghany region, (etc.)—D. S. JORDAN; A. W. BRAYTON. B. Synopsis of family Cutostomidæ—D. S. JORDAN	
North American Oölogy. Part I. Raptores and fissirostres-T. M. Brewer.	
North Carolina—	
ancient mica mines in—C. D. SMITH. (R. 1876)	299
Beaufort, description of new species of fish (Caranz Beani) from—D. S. JORDAN. (P. 1880)	
Beaufort, notes on fishes of—D. S. JORDAN; C. H. GILBERT. (P. 1878.)	
Black Mountain, topography of-T. L. CLINGMAN. (R 1855)	
earthquakes in, in 1874—W. Du Pré; J. Henry. (R. 1874)	
Lenoir county, antiquities in—J. M. SPAINHOUR. (R. 1871)	
Montgomery county, antiquities of—F. J. Kron. (R. 1874)	
occurrence of Physis regius in—T. H. BEAN. (P. 1880)	425
Stanley county, antiquities of—F. J. KRON. (R. 1874)	286
North German Lloyd, free freight between Germany and United States by— R. Schleiden. (R. 1858)	
North Pacific Surveying Expedition, contributions to natural history made in connection with—T. H. STREETS	
North Pole, expedition toward the, scientific instructions to—J. Henry; J. B. Hilgard; S. Newcomb; S. F. Baird; F. B. Meek; L. Agassil.	
(R. 1871)	249
North Pole, map of stars near the, for observations on the aurora	50
Northern Hemisphere, winds of the—J. H. COFFIN.	52
Northern latitudes, record of auroral phenomena observed in—P. Force	84
Northern seas, the—J. Babiner. (R. 1869)	228
Northumberland, Duke of, account of the	; 30
Northumberland, Duke of, presentation of books by. (R. 1859)	110

Northwest coast of America, kjökken-möddings on the—P. SCHUMACHER. (R. 1873)	275
Norton, E. On wasps	254
Norway, Ethnological Museum of-L. K. DAA. (R. 1862)	150
Norway, University of Christiania, ethnological specimens from. (R. 1868)	187
Norwegian mountains, lakes, and the snow-line—O. E. DREUTZER. (R. 1866.)	214
Notacanthus phasganorus from Newfoundland-G. B. GOODE. (P. 1880)	425
Note on occurrence of Phycis regius in South Carolina—T. H. BEAN. (P. 1880.)	425
Note on the latiloid genera—T. GILL. (P. 1881)	467
Notes—	
of egging expedition to Shoal Lake, Lake Winnipeg—D. Gunn. (R. 1867)	215
on aboriginal shell mounds of New Brunswick and New England—S. F. BAIRD. (P. 1881)	467
on American migration—F. von Hellwald. (R. 1866)	214
on fishes—See D. S. Jordan; C. H. Gilbert; also, Proceedings National Museum.	
on fishes of Pacific coast of United States—D. S. JORDAN; C. H. GIL- BERT. (P. 1881)	467
on history and climate of New Mexico—T. A. McParlin. (R. 1877.) 328,	396
on horary variations of barometer-M. VAILLANT; J. HENRY. (R. 1866.)	214
on Indian history—F. V. HAYDEN. (R. 1867)	215
on life and character of Joseph Henry-J. C. WELLING 856,	338
on mortality among fishes of Gulf of Mexico—S. H. Johnson. (P. 1881.)	467
on new species and localities of microscopical organisms—J. W. BAILEY.	63
on physical geography of North America—J. G. Cooper. (R. 1858.) 109,	351
on Salmonidæ of upper Columbia—C. BENDIRE. (P. 1881)	467
on some Costa Rican birds—R. RIDGWAY. (P. 1881)	467
on Tinneh or Chepewyan Indians of British and Russian America-G.	
GIBBS and others. (R. 1866) 214,	365
on Tonto Apaches—C. SMART. (R. 1867)	215
relative to George Catlin-J. HENRY. (R. 1872)	271
to accounts of lightning discharges, by G. W. Dodge and others—J. HENRY. (R. 1867)	215
to article on vitality, by H. H. Higgins-J. HENRY. (R. 1866)	214
to articles on meteorology, by G. Latimer and others—J. HENRY. (R. 1871)	249
to lectures on meteorology, by R. Russell-J. HENRY. (R. 1854)	75
Notice, biographical, of—	
Agassiz, L., by J. A. GARFIELD. (R. 1878)	275
Agassiz, L., by P. PARKER. (R. 1878)	275
Blackmore Museum, Salisbury, England. (R. 1868)	224
Chase, S. P., by J. A. GARFIELD. (R. 1873)	275

Notice, biographical, of-Continued.
Chase, S. P., by H. Hamlin. (R. 1878)
Cleaveland, Parker, by J. HENRY. (R. 1859)
Espy, James P., by A. D. BACHE. (R. 1859)
Harvey, W. H., by A. GRAY. (R. 1867) 215
Irving, Washington, by C. C. FELTON. (R. 1859)
Jewett, C. C., by R. A. Guild. (R. 1867)
Rush, Richard, by J. A. PEARCE. (R. 1859) 110
Schoenbein, Christian Frederic, the discoverer of ozone. (R. 1868) 224
Smithson, James, by D. GILBERT, President of the Royal Society. (R. 1853) 67,330
Smithson, James, by W. R. Johnson; J. R. McD. Irby 327
Smithson, James, by W. J. RHEES. (R. 1879) 349, 330
Turner, W. W., by C. C. FELTON. (R. 1859) 110
Würdemann, G., by A. D. BACHE. (R. 1859)110
Notice of recent additions to marine invertebrates of northeast coast of America— A. E. VERRILL. (P. 1880)
Notices of public libraries in the United States-C. C. JEWETT 25
Nova Scotia-
birds of-Blackiston; T. Bland; J. R. Willis. (R. 1858) 109
fishes obtained at—See G. B. Goode; T. H. Bean.
kjökken-möddings in—J. M. Jones. (R. 1868)
Numeration, report on improved system of—W. B. TAYLOR. (B. 1867) 215
Nuovi Lincei, Pontifical Academy of the. Prize questions. (R. 1865; E. 1867)
Nutation, phenomena of—J. G. BARNARD
NUTTER, F. H. Mounds on Gideon's farm, near Excelsior, Hennepin county, Minnesota. (R. 1879)

O.

Ober, F. A.—	
birds of Antigua and Barbuda collected by, catalogue of—G. N. LAW- RENCE. (P. 1878)	882
birds of Dominica collected by, catalogue of—G. N. LAWRENCE. (P. 1878.).	
birds of Grenada collected by, catalogue of-G. N. LAWRENCE. (P. 1878.)	882
birds of Guadeloupe collected by, catalogue of—G. N. LAWRENCE. (P. 1878)	882
birds of Lesser Antilles collected by, catalogue of—G. N. LAWRENCE. (P. 1878)	882
birds of Martinique collected by, catalogue of—G. N. LAWRENCE. (P. 1878)	882
birds of St. Vincent collected by, catalogue of—G. N. LAWRENCE. (P. 1878)	882
OBER, F. A. Ornithology of the Caribbee Islands. (R. 1878)	841
Obsequies of Joseph Henry	856
Observations-	
astronomical—W. HARKNESS	289
astronomical, discussion of Piazzi's—B. A. Gould. (R. 1868)	187
astronomical, in Arctic Seas-E. K. KANE	129
in Hudson's Bay Territory—B. R. Ross. (R. 1859)	110
magnetic, in the Arctic Seas—E. K. KANE	97
physical, in Arctic Seas—I. I. HAYES	196
physical, in Arctic Seas. Parts I-IV—E. K. KANE	198
scientific, metric system for—A. Guyor. (R. 1848)	I
Observations-See Magnetic, Meteorological, Physical, Tidal.	
Observations, meteorological—	
at Brunswick, Maine-P. CLEAVELAND	204
at Providence, Rhode Island—A. CASWELL 103,	443
at Sacramento, California-F. W. HATCH. (R. 1854)	75
at Sacramento, California—T. M. Logan. (R. 1854)	75
at Washington, Arkansas-N. D. Smith	131
combined, report of committee of American Association for Advance- ment of Science on system of. (R. 1851)	51
directions for-A. GUYOT	19
for year 1855	93
in the Arctic Seas—E. K. KANE	104
Observations of—	
earthquake phenomena—R. MALLET. (R. 1859)	110
Siredon lichenoides-W. E. CARLIN. (P. 1881)	467
temperature, best hours for-C. DEWEY. (R. 1860)	147
thunder storm, John Wise's-R. HARE. (R. 1854)	75
thunder storms, instructions for—J. HENRY	235

Observations on—
auroras, map of stars near North Pole for
Batis maritima of Linnsus—J. Torrey
electric resonance of mountains—H. DE SAUSSURE. (R. 1868) 22
gold ornament from mound in Florida—C. RAU. (R. 1877) 323, 440, 400
ice—D. WALKER
Mexican history and archeology—B. MAYER
natural phenomena, shooting stars, aurora, etc.—S. Masterman. (R. 1857)
prehistoric mounds of Grant county, Wisconsin-M. STRONG. (R. 1876.) 29
terrestrial magnetism—J. Locke 3
terrestrial magnetism and deviation of compasses of iron-clad "Monad-nock"—W. HARKNESS
terrestrial magnetism in Mexico-Baron von Müllen; A. Sonntag 11
thunder and lightning-S. MASTERMAN. (R. 1855)
Observatories—
at Dorpat and Poulkova, description of—C. ABBE. (R. 1867) 216, 365
detached, universal meteorograph for—E. H. von BAUMHAUER. (R. 1879.) 34
instructions to, relative to telegraphic announcements of astronomical discoveries—J. Henry
reports of, 1879—E. S. Holden. (R. 1879)
reports of, 1880—E. S. HOLDEN; G. H. BOEHMER. (R. 1880) 442, 438
Observatory—
at Cordova, Argentine Republic, account of-B. A. Gould. (R. 1873.) 27
at St. Martin, Isle Jesus, Canada East, description of—C. SMALLWOOD. (R. 1856)
Girard College-See Bache, A. D.
Kew, verification of barometers at-J. WELSH. (R. 1859) 110
magnetic, at Smithsonian Institution, description of-J. E HILGARD
(R. 1859) 110
physical, on a—J. Henry. (R. 1870) 24
Toronto, instructions by, for aurora observations 148
Observers, meteorological, of Smithsonian Institution, list of. (R. 1868) 224,373
Observers—See Meteorological stations and observers.
Occultations—
in 1852—C. H. DAVIS
of planets and stars by the moon, 1858—J. Downes
visible in the United States during the years 1848-1852-J. Downs.
8, 9, 10, 11, 29
Occultator, account of—T. HILL
Ocean, fresh water in the—W. C. DENNIS. (R. 1866)
Ocean, Indian, cyclone in the—N. PIKE (R. 1867)
Oceanic bonito on coast of United States-G. B. GOODE; T. H. BEAN. (P. 1878)

EDWARDS. (P 1878)
• • • • • • • • • • • • • • • • • • • •
Oersted, H. C., memoir of, by E. De Beaumont. (R. 1868) 224
Officers— and establishment of the Smithsonian Institution and National Museum. 449
collaborators, employé-, etc., of Smithsonian Institution, National Mu-
seum, Geological Survey, Bureau of Ethnology, and Fish Commission, directory of
and Regents of Smithsonian Institution, 1846, list of N
Ohio—
ancient earthworks in, sketch of—I. DILLE. (R. 1866)
ancient works in, descriptions of—C. WHITTLESEY
Ashland county, earthworks in—G. W. HILL. (R. 1877) 823
Ashtabula county, double-walled earthwork in—S. D. PEET. (R. 1876.) 299
Holmes county, flint instruments in—H. B. CASE. (R. 1877) 328
Kelley's Island, climate of—G. C. Huntington. (R. 1866)
Marietta, meteorological observations at—S. P. HILDRETH; J. Wood 120
northern, antiquities of—G. W. Hill. (R. 1874)
Perry county, antiquities of—W. Anderson. (R. 1874) 286
Trumbull county, mound in—F. MILLER. (R. 1877)
Ohio river, habits of black bass of the—J. EOFF. (R. 1854)
Ohio river, suggestions for improvement of navigation of—C. ELLET, Jr 18
Oil—
coal, explosibility of—Z. Allen. (R. 1861)
porpoise, manufacture of—U. Cook. (P. 1878)
rock, or petroleum, history of—T. S. HUNT. (R. 1861) 149
Oil-shark of southern California—D. S. JORDAN; C. H. GILBERT. (P. 1880.) 425
Old-World birds in National Museum, catalogue of—R. RIDGWAY. (P. 1881.) 467
Oligocene foseils, check-list of—T. A. Conrad
Olmos, Peru, account of solar eclipse at—J. M. GILLISS
OLMSTED, D. Aurora borealis, recent secular period of
OLNEY, S. T. List of Rhode Island algee 241
Omaha myth—J. O. Dorsey. (E. 1879-80)
Oblogy of Kerguelen Island—J. H. KIDDER; E. COUES
Opening and closing of Kennebec river, Maine—R. H. GARDINER. (R. 1858.) 109
Opheosaurus ventralis, remarks on osteology of—R. W. SHUFELDT. (P. 1881.) 467
Orbit—
and phenomena of meteoric fire-ball—J. H. COFFIN
of Neptune, investigation of, with tables of its motion—S. Newcomb 262

Orbits, planetary, secular variations of-J. N. STOCKWELL. (R. 1871) 249, 2	32
Orcynus pelamys (oceanic bonito), occurrence of, in Vineyard Sound, Massa-	
chusetts-V. N. EDWARDS. (P. 1878)	332
Oregon—	
fishes of, notes on collection of—D. S. JORDAN. (P. 1878)	832
jargon of-B. R. MITCHELL; W. W. TURNER	
trade-language of-G. GIBBS	161
Ores collected on geological survey in Michigan, catalogue of—C. T. JACKSON. (R. 1854)	75
Ores, rocks, minerals, and fossils, catalogue of—J. Locke. (R. 1854)	
Organic bases—A. BAUER. (R. 1872)	
Organisms—See Microscopic.	
Organization and objects of the United States National Museum—G. B. Goods. (P. 1881)	59
Organization, plan of, and regulations of the U.S. National Museum-G.B.	
Goode. (P. 1881)	145
Organization of—	
· ·	298
Smithsonian Institution, programme ofP, J,	32
Smithsonian Institution, report of American Academy of Arts and Sciences on—E. EVERETT; J. SPARKS; B. PEIRCE; H. W. LONG-FELLOW; A. GRAY. (R. 1858)	
Smithsonian Institution, report of committee of Regents on B, L.	
Organized bodies, natural history of—E. J. MARRY. (R. 1867)	
O'Rielly, Henry, deposition of J. Henry in case of S. F. B. Morse versus	115
Origin—	
and history of the Smithsonian Institution-W. J. RHEES 3	
and propagation of disease—J. C. Dalton. (R. 1878)	
of force—W. B. TAYLOR. (R. 1870)	75
Ornament, gold, from a mound in Florida, observations on—C. RAU. (R. 1877.) 323, 440. 4	03
Ornithology—	24!
of Caribbee Islands—F. A. OBER. (R. 1878)	
of Hawaiian and Fanning Islands and California—T. H. STREETS	
of Kerguelen Islands—J. H. KIDDER; E. COURS	, 336
Ortalidæ, monograph of—H. Lorw	200
Orthoptera—	
bibliography of—S. H. SCUDDER	109
instructions for collecting—P. R. UHLER. (R. 1858)	89
of North America, catalogue of—S. H. SCUDDER	215
OSBORNE, J. A. Account of lightning discharges. (R. 1867)	
Osmerus attenuatus, new species of California fish, description of—W. N. LOCK-	125
INULUA. I C. 10001	

OF SMITHSONIAN PUBLICATIONS.

OSTEN SACKEN, R.—
Bibliography of diptera 270
Catalogue of described diptera of North America 102, 270
Diptera of Kerguelen Island 294
Directions for collecting and preserving dipters 102
Monograph of Tipulida 219
Monographs of diptera of North America. Part IV
OSTEN SACKEN, R.; LOEW, H. Instructions for collecting dipters. (R. 1858.) 109
OSTEN SACKEN, R.; LOEW, H. Monographs of diptera of North America. Parts I, II
Osteology and mounds of the mound-builders of Wisconsin—J. N. DE HART. (R. 1877)
Osteology of Opheosaurus ventralis-R. W. SHUFELDT. (P. 1881) 467
Ostraciontide (trunk fishes), a study of, with notes on American species—G. B. GOODE. (P. 1879)
Othonops eos, a new gobioid fish from San Diego, Cal.—R. SMITH. (P. 1881.) 467
Outline of scheme of classification of collections in U. S. National Museum— G. B. Goode. (P. 1881)
Outline of systematic review of the class of birds—W. Lilljeborg 364
OWEN, D. D. Catalogue of geological specimens. (R. 1854)75
OWEN, D. D. Report on building stones. (R. 1847) H, 829
OWEN, R. D. Hints on public architecture
Owl, description of a new, from Porto Rico—R. RIDGWAY. (P. 1881) 467
Ox, American, extinct species of—J. LEIDY
Oxygen and its combinations, lecture on—G. I. CHACE. (R. 1855)
Ozone and antozone—C. M. WETHERILL. (R. 1864)

P.

fishes from, descriptions of new—D. S. JORDAN; C. H. GILBERT. (P. 1881)
fishes from, distributed in 1881—D. S. JORDAN; P. L. JOUY. (P. 1881.) 467 fishes of—See D. S. Jordan; C. H. Gilbert.
fishes of, bibliography of—T. H. BEAN. (P. 1881)
fishes of, bibliography of—T. Gill
fishes of, notes on—D. S. Jordan; C. H. Gilbert. (P. 1881) 467
Pacific Islands, natural history of—W. H. PEASE. (R. 1862)
PACKARD, A. S. Directions for collecting and preserving insects 261
PACKARD, A. S. Memoir of H. J. Clark
PACKARD, F. A. Project of outline history of education in the United States. (R. 1868)
PAGE, C. G. Report on effect of frost on building stones. (R. 1847) H. 329
Palsontology.
list of generic names employed in—S. H. SCUDDER
of the upper Missouri-F. B. MEEK; F. V. HAYDEN
principles and methods of-T. H. HUXLEY. (R. 1869)
Palafittes or lacustrian constructions of the lake of Neuchâtel—E. Desor. (R. 1865)
Palenque tablet in the U.S. National Museum—C. RAU
Palermo, evaporation observed at, in 1865, 1866-P. TACCHINI. (R. 1870) 34
PALMIERI, Prof. Electro-magnetic seismograph. (R. 1870)
PALMIERI, Prof. Presence of electricity during fall of rain. (R. 1870) 24
Panama, shells of, review of C. B. Adams' catalogue of—P. P. CARPENTER 252
Pancreas, observations on the-J. JONES
Pandoride, contributions towards a monograph of the-P. P. CARPENTER 25
Paralepis coruscans, new species of <i>Paralepis</i> , description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)425
Parasites-J. Leidy
Paris—
Academy of Sciences, historical sketch of the-M. Flourens. (R. 1862.) 150
Anthropological Society of, history of transactions of, 1865 to 1867—P. Broca. (R. 1868)
Exposition—See French Exposition.
Muséum d'Histoire Naturelle, typical specimens of American fishes in
the—D. S. JORDAN. (P. 1879)
PARKER, P. Biographical notice of L. Agassiz. (R. 1878) 275, 323
PARKER, P. Eulogy on Henry Wilson. (R. 1875) 298, 329
Parophrys ischyurus, new species of flounders from Puget Sound, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)

Parrot, new species of, of genus Chirus, from Dominica—G. N. LAWRENCE. (P. 1880)
Parthenogenesis in the animal kingdom—G. A. KORNHUBER. (R. 1871)
PARVIN, J. B. Habits of the gopher of Illinois. (R. 1854)
Past and future of geology—J. Prestwich. (R. 1875)
Patagonia, human remains from—A. Rikd. (R. 1862)
Patent Office, U. S., meteorological observations under the direction of. Vols.
ı, ıı 157,
PATTERSON, R.; SHARPLESS, T. Phonography. (R. 1856)
PATTON, A. Antiquities of Knox county, Indiana. (R. 1878)
PATTON, A. Antiquities of Lawrence county, Illinois. (R. 1873)
PKABODY, A. P. Scientific education of mechanics and artisans. (R. 1872.) 271,
PEALE, T. R.—
Ancient mound in St. Louis, Missouri. (R. 1861)
Method of preserving lepidoptera. (R. 1868)
Prehistoric remains in vicinity of city of Washington, D. C. (R. 1872.)
Uses of brain and marrow of animals among Indians of North America. (R. 1870)
Pearce, J. A., eulogy on, by A. D. BACHE. (R. 1862)
PEARCE, J. A. Notice of Richard Rush. (R. 1859)
PEARCE, J. A. Report of Committee of Regents on distribution of Smithsonian income. (R 1858)
PEASE, W. H. Acknowledgment for shells, etc. (R. 1861)
PEASE, W. H. Natural history of Pacific Islands. (R. 1862)
Pedro II, Dom, biographical sketch of, by A. FIALHO. (R. 1876)
PRET, S. D. Double-walled earthwork in Ashtabula county, Ohio. (R. 1876.)
PEIRCE, B., and others. Report of American Academy of Arts and Sciences
on organization of Smithsonian Institution. (R. 1858)
PELTIER, F. A. Memoir of J. C. A. Peltier. (R. 1867)
Peltier, J. C. A., memoir of, by F. A. Peltier. (R. 1867)
Peltier, J. C. A., scientific researches of. (R. 1867)
Pendulum, problems of rotary motion presented by the-J. G. BARNARD
Ponnsylvania—
Centre county, Spruce creek valley, account of a tornado in—J. B. MERK. (R. 1871)
Chester county, birds of—V. BARNARD. (R. 1860)
Chester county, history of—W. DARLINGTON. (R. 1862)
magnetic survey of—A. D. BACHE
Newcastle, old Indian village, Kushkushkee, near—E. M. McConnell.
(R. 1871)
western, ancient mound in-W. M. TAYLOR. (R. 1877)
Pensacola, Florida, description of new species of umber fish (Seriola Stearnsii)
Com (1 P. Coope (1970)

Pepin, Lake, antiquities on the banks of—L. C. Estes. (R. 1866) 214
· · · · · · · · · · · · · · · · · · ·
Perca flavescens, notes on—F. Steindachner. (P. 1878)
Percidæ, notes on—D. S. Jordan 306
Perennibranchiates, acknowledgment of—J. G. FISHER. (R. 1868) 187
Periodical phenomena—
directions for registry of
observations on, 1851 to 1859—F. B. HOUGH185
registration of—J. HENRY. (R. 1855)
registry of 65, 148
Periodicals—
deposited in Library of Congress, meteorological articles in. (R. 1873) 275
in Library of Smithsonian Institution. Parts I, II; foreign works; 1866. 73, 85, 117, 179
received by the Institution, list of. (R. 1880) 442, 436
received in reading-room of Smithsonian Institution in 1858, check-list of. Q
PERRINE, T. M. Antiquities of Union county, Illinois. (R. 1878) 275
PERRINE, T. M. Mounds near Anna, Union county, Illinois. (R. 1872) 271
Persistence of images—F. N. Moigno. (R. 1866)
Perspiration and respiration, apparatus for testing results of—M. Pettenkoffe. (R. 1864)
Peru-
earthquake in, August 18, 1868—J. V. CAMPBELL. (R. 1870) 344
Indians of—F. L. Galt. (R. 1877)
Olmos, total eclipse of sun, September 7, 1858, observed near—J. M. GILLISS
PESLIN, M. Relation between barometric variations and general atmospheric
currents. (B. 1877)
PETER, R. Ancient mounds in Kentucky. (R. 1872) 271
Peter, R. Ancient mound near Lexington, Kentucky. (R. 1871) 29
Petite Anse Island, salt deposit on—E. W. HILGARD 248
PETITOT, E. Account of the Indians of British America. (B. 1865)
etrel, supposed new, from Sandwich Islands, description of—R. RIDGWAY.
(P. 1881)
etroleum collections, circular in reference to—S. F. BAIRD. (P. 1881) 467, 447
etroleum or rock oil, history of—T. S. HUNT. (B. 1861)
ETTENKOFER, M. Apparatus for testing the results of perspiration and re-
piration. (R. 1864)
PETTIGEEW, J. B. Modes of flight in relation to aeronautics. (R. 1867) 215
benogamia of Kerguelen Island—A. Gray
Phalangidse, instructions for collecting—H. C. Wood. (R. 1866)
Phenomens—
accompanying the propagation of electricity in highly rarefled elastic fluids—A. De La Rive. (R. 1868)
in telegraphic lines during the aurora borealis—G. B. Donati. (R. 1872) 271

OF SMITHSONIAN PUBLICATIONS.	278
Phenomena—Continued.	
natural, observations on—S. MASTERMAN. (R. 1857)	107
of contact, studies on—T. L. Phipson. (R. 1862)	150
of flight in animal kingdom-E. J. MARRY. (R. 1869)	228
of meteoric fire-ball—J. H. COFFIN	221
of precession and nutation—J. G. BARNARD	810
periodical registration of—J. HENRY. (B. 1855)	77
Philadelphia— .	
exhibit at—See Centennial.	
magnetic and meteorological observations at—See A. D. Bache.	
memorial of citizens of, relative to Lowe's aeronautic voyages across Atlantic. (R. 1860)	147
Philadelphia Academy of Natural Sciences, report on shells presented to—J. LEIDY; G. W. TRYON. (R. 1865)	209
PHILLIPS, B. Two letters on work of the Museum. (P. 1881) 467,	454
Philological circular—G. GIBBS. (R. 1862)	150
Philology, Indian-W. W. Turner. (R. 1861)	51
Philology, instructions for research relative to—G. GIBBS	160
Philosophical Society of Washington—	
Bulletins of. Vols. I-III. March, 1871, to June, 1880	423
memoir of Joseph Henry read before the-W. B. TAYLOB 856,	339
notes on life and character of Joseph Henry read before the-J. C.	
Welling 856	338
proceedings of, on death of Joseph Henry	856
Philosophy, annals of, contributions to, by J. Smithson. (R. 1858) 67	, 880
Philosophy, Batavian Society of Experimental, of Rotterdam. Prize questions. (R. 1861)	149
PHIPSON, T. L. The catalytic force or studies on the phenomena of contact. (R. 1862)	150
Phlegmasia, disease of, bibliography of works on—W. W. KEEN	800
Phonetic language, vocal sounds of L. Bridgman compared with elements of—	
F. Lieber	12
Phonography—T. SHARPLESS; R. PATTERSON. (R. 1856)	91
Phonography, institutions in which, is taught. (R. 1856)	91
Photo-chemistry—J. Jamin. (B. 1867)	215
Photographs of Indians, list of	216
Photography, astronomical, progress in—Dr. LEE. (R. 1861)	149
Photography, celestial, use of silvered-glass telescope in—H. DRAPER	180

Phycis Chesteri from deep-sea fauna of northwestern Atlantic, description of— G. B. GOODE; T. H. BEAN. (P. 1878)

Phycis Earlii, new hake, from South Carolina, description of—T. H. BEAN.
(P. 1880)

Phycis regius from North Carolina, note on occurrence of—T. H. BEAN. (P. 1880)

882

425

Physical—
atlas of North America, proposed—G. GIBBS. (B. 1866)
ethnology, lectures on—D. Wilson. (B. 1862)150
geography of Mississippi valley—C. ELLET, Jr13
geography of North America—J. G. Cooper. (R. 1858) 109, 351
geography of North American Continent-J. FROZBEL. (R. 1854) 75
observations in the Arctic Seas—I. I. HAYES 196
observations in the Arctic Seas—E. K. KANE 196
observations in the Arctic Seas—F. L. McCLINTOCK 146
observatory, on a-J. HENRY. (R. 1870)
sciences, on the relation of, to science in general—H. Helmholtz. (R. 1871) 249
tables—A. Guyor153
Physics—
present fundamental principles of—F. J. Pisko. (R. 1879) 345
recent progress in, 1879 and 1880—G. F. BARKER. (R. 1880) 442, 429
report on recent progress in, electricity, galvanism—J. MÜLLER. (R. 1855-1858)77, 91, 107, 109
syllabus of a course of lectures on-J. Henry. (R. 1856) 91
terrestrial, articles on—J. V. Campbell; Prof. Palmieri; W. W. Johnson; W. D. Sargent. (R. 1870)24
Physics and Natural History, Geneva Society of-See Geneva.
Physiological investigations of vertebrata—J. Jones
Physiology—
electro-, lectures on—C. MATTEUCCI. (R. 1865)
of fever—H. C. Wood 357
of lucernarians—H. J. CLARK
of rattlesnake—S. W. MITCHELL 185
of respiration in chelonia—S. W. MITCHELL; G. R. MOREHOUSE 159
Piazzi's astronomical observations, discussion of—B. A. GOULD. (B. 1863) 187
PICKERING, C. Gliddon mummy case in Museum of Smithsonian Institution. 208
Picture writing, Central American, studies in-E. S. Holden. (E. 1879-80.) 476
PIDGEON, W. Ancient burial mound in Indiana. (R. 1867) 216
PIERSON, W. M. Discovery of large meteorite in Mexico. (R. 1978) 275
PIKE, N. Cyclone in Indian Ocean. (R. 1867)
PILAR, G. Revolutions of crust of the earth. (R. 1876)
Pile work antiquities of Olmutz. (R. 1866)
PILLING, J. C. Catalogue of linguistic manuscripts in library of Bureau of Ethnology. (E. 1879-80)
Pima Indians of Arizona—F. E. Grossman. (R. 1871) 249
Pinna, cretaceous, from New Mexico, description of new—C. A. WHITE. (P. 1880)
PIRZ, A. Methods of making and preserving plaster casts. (P. 1881) 457
The state of the s

OF SMITHSONIAN PUBLICATIONS. 2	7 5
Pisko, F. J. Present fundamental principles of physics. (R. 1879)	845
	61
	328
Pittsburg, on Des Moines river, composition of ancient pottery found at—R.	8 4 5
Plan of—	
a bibliography—J. Friedländer. (R. 1858)	109
organization and regulations of U. S. National Museum—G. B. Goode. (P. 1881) 467, 4	45
research upon the atmosphere—C. M. WETHERILL. (R. 1866) 2	214
Smithsonian exhibit at Centennial—S. F. BAIRD. (R. 1875)	298
Planet—See Neptune, Uranus.	
Planetary—	
disturbances, lecture on—E. S. SNELL. (R. 1855)	77
motion, general integrals of—S. Newcomb 2	81
motion, perturbative function of, tables for determining values of coeffi- cients in—J. D. RUNKLE	94
orbits, secular variations of elements of—J. N. STOCKWELL. (R. 1871.)	
Planetoids—See Asteroids.	
Planets-	
between Mars and Jupiter—Prof. LESPIAULT. (R. 1861)	149
between Mars and Jupiter, lecture on—E. Loomis. (R. 1854)	75
occultations of—See Downes, J.	
Planisphere of the visible heavens for observations of meteoric displays 3	59
Plans for the installation of collections in the U.S. National Museum—G. B.	
GOODE, (P. 1881) 467, 4	
Plans of the Smithsonian building—R. D. Owen	P
	46
Plants Wrightians Texano-Neo-Mexicans. Parts I, II—A. GRAY 22,	42
PLANTAMOUR, E. Report on transactions of the Society of Physics and Natural History of Geneva, July, 1864, to June, 1865; July, 1874, to June, 1875. (R. 1865; R. 1877)	898
Plants—	
collected by J. C. Frémont in California—J. TORREY	46
·	249
	182
	182
	182
	160
	842
	22
of upper Missouri, list of—T. C. PORTER. (R. 1850)	28
••	61

Plants-See Torrey, J., Gray, A.
Plaster casts—
methods of making and preserving—A. PTRZ. (P. 1881)
of antique and modern statues—W. J. STONE. (R. 1855) 7
of heads of Indian boys and girls at Hampton, Virginia—R. H. Pratt. (P. 1879) 88
of heads of Indian prisoners at St. Augustine, Florida—R. H. PRATT. (P. 1878)
PLATEAU, F. Rapid preparation of myological specimens. (P. 1881) 467
PLATEAU, J. Experimental and theoretical researches on the figures of equilibrium of a liquid mass withdrawn from the action of gravity. Parts 1-v1. (R. 1868-1866)
Platessa ferruginea, note on—G. B. GOODE; T. H. BEAN. (P. 1878)
Platessa rostrata, note on-G. B. Goode; T. H. BEAN. (P. 1878)
Platycnemism in Michigan—H. GILLMAN. (R. 1878) 275, 393
Platyrhina exasperata, generic relations of—D. S. JORDAN; C. H. GILBERT. (P. 1880)
Platyrhina triseriata, new ray from coast of California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)423
Platysomatichthys stomias, new flounder from coast of California, description of—D. S. JOBDAN; C. H. GILBERT. (P. 1880)
Pleuronectes glaber, identity of, with Euchalarodus Putnami—T. H. BEAN. (P. 1878)
Pleuronectids of San Francisco, review of-W. N. LOCKINGTON. (P. 1879) \$33
Pleuronichthys verticalis, new flounder from coast of California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880) 425
POE, O. M. Account of lightning discharges. (R. 1867) 215
POET, A. New classification of clouds. (R. 1870) 244
POET, F. Notes on American species of genus Cybium. (P. 1878) 322
POGGENDORFF, J. C. Use of galvanometer as measuring instrument. (R. 1859.) 110
Points, boiling and melting-F. W. CLARKE 255, 288
Points, spear and arrow, primitive manufacture of, along line of Savannah
river—C. C. Jones, Jr. (R. 1879)
Poison—See Venom.
Poisoned arrows, lecture on nature and cure of wounds of—D. BRAINARD. (R. 1854)
Poisonous water in Gulf of Mexico, destruction of fish by—J. Y. PORTER. (P. 1881)
Polar expedition—See Hall, Capt.
Polar expedition, the Howgate, contributions to natural history of Arctic America made in connection with—L. Kumlien
Polar light, or aurora borealis, its phenomena and laws—E. Looms. (R. 1865.) 309
Polaris expedition, scientific instructions for-J. HENRY; J. E. HILGARD; S.
NEWCOMB: S. E. BAIRD: F. B. MEEK: L. AGASSIZ. (R. 1871) 249

•	
Polarized nerve, effect of irritation of-B. F. LAUTENBACH. (R. 1878) 841	411
Pollock, J. Assay of coins at the Mint of the United States. (R. 1868)	224
Polychrome bead from Florida-S. S. HALDEMAN. (R. 1877) 828,	404
Polypetals, bibliographical index of-S. Warson	258
Polypi, relations of lucernarians to—H. J. CLARK	242
Polyzon of northeastern coast of America, notice of recent additions to—A. E.	
VERRILL. (P. 1879)	888
Pomadasys, description of new species of, from Mazatlan, Mexico—D. S. Jor- DAN; C. H. GILBERT. (P. 1881)	467
Pontifical Academy of the Nuovi Lincei, Rome. Prize questions. (R. 1865; R. 1867)	, 215
Poole, H. Cone-in-cone. (R. 1868)	187
Popocatepetl, examination of—Baron von Müller; A. Sonntag	114
Population of the world, estimate of the—E. MAILLY. (R. 1873)	275
Porcupine, Canada, occurrence of, in Maryland—O. LUGGER. (P. 1881)	467
Porcupine, Canada, occurrence of, in West Virginia—G. B. Goode. (P. 1878.)	832
Porifers of northeast coast of America—A. E. VERRILL. (P. 1879)	888
Porpoise oil, manufacture of—C. Cook. (P. 1878)	882
PORTER, Commodore. Account of hail storm on the Bosphorus. (R. 1870)	244
PORTER, J. Y. Destruction of fish by poisonous water in Gulf of Mexico. (P. 1881)	467
PORTER, J. Y.; JEFFERSON, J. P.; MOORE, T. Destruction of fish near the	-0.
Tortugas. (P. 1878)	882
PORTER, T. C. List of plants of upper Missouri. (R. 1850)	28
Portland Society of Natural History, account of—E. C. Bolles. (R. 1867)	215
Porto Bico—	
description of new owl from-R. RIDGWAY. (P. 1881)	467
great hurricane at—G. A. LATIMER. (R. 1867)	215
Latimer collection of antiquities from-O. T. MASON. (R. 1876)	397
meteorology of-G. LATIMER. (R. 1871)	249
Portraits-	
of North American Indians, catalogue of-J. M. STANLEY	53
of North American Indians, Stanley, report of Committee of Regents on. (R. 1857)	107
photographic, of North American Indians, in gallery of Smithsonian Institution, list of	216
Potamocottus Bendirei, description of—T. H. BEAN. (P. 1881)	467.
Pottery—	
ancient, from Phillips county, Arkansas—J. H. DEVEREUX. (R. 1872.)	271
ancient, from Pittsburg, on Des Moines river—R. N. DAHLBERG; C. DAHLBERG. (R. 1879)	345
· Indian—C. RAU. (R. 1866)	368
Pouched rat, or salamander of Georgia, habits of-W. GESNER. (R. 1860)	
Poulkova and Dorpat, description of observatories at—C. ABBE. (R. 1867.) 215,	

Powell, B. Reports on state of knowledge of radiant heat. (R. 1869) 110
Powell, J. W.—
Evolution of language. (E. 1879–80) 476
Limitations to use of some anthropological data. (E. 1879-80) 476
Mythology of North American Indians. (E. 1879-80) 476
Report of Director of Bureau of Ethnology. (E. 1879-80) 476
Wyandot government. (E. 1879-80) 476
Powers, S. Centennial mission to Indians of western Nevada and California. (R. 1876)
PRATT, R. H. Catalogue of casts of heads of Indian boys and girls at Hampton Institute, Virginia. (P. 1879)
PRATT, R. H. Catalogue of casts of heads of Indian prisoners at St. Augustine, Florida. (P. 1878)
PRATT, W. H. Antiquities of Whiteside county, Illinois. (R. 1874) 28
Prayer at funeral of Joseph Henry-C. Hodge
Prayer at memorial of Joseph Henry—J. McCosh
Prayer at memorial of Joseph Henry—B. SUNDERLAND
Precession of the equinoxes—J. N. STOCKWELL 22
Precession of the equinoxes, problems presented by-J. G. BARNARD 240, 310
Precious stones—J. BABINET. (R. 1870)
Precipitation—
influence of rain upon the formation of—J. HANN. (R. 1877) 823, 386
of rain and snow in the United States, charts of-C. A. Schott 374
of rain and snow in the United States, tables of-C. A. SCHOTT 222, 350
Prehistoric—
antiquities of Hungary-F. F. Romes. (B. 1876) 299, 440, 392
archeology, international code of symbols for charts of—G. DE MORTIL- LET; E. CHANTRE. (R. 1875)
evidences in Missouri—G. C. BROADHEAD. (R. 1879) 345
man, remains of, from caves in Alaska—W. H. DALL
mounds of Grant county, Wisconsin-M. STRONG. (R. 1876) 299
remains in vicinity of city of Washington, D. C.—T. R. PEALE. (R. 1872)
PRENTISS, S. S.; COUES, E. List of birds of the District of Columbia. (R. 1861)
Preparation of large myological specimens—F. J. PLATEAU. (P. 1881) 467
Preparations, dry, on Semper's method of making-J. A. RYDER. (P. 1881.) 467
Prescott, Canada West, ancient Indian remains near—W. E. Guest. (R. 1856)
PRESCOTT, W. H., and others. On publication of Spanish works on New Mexico. (B. 1855)
Present state of ethnology in relation to form of human skull—A. Rezzios. (R 1859)

OF SMITHSONIAN PUBLICATIONS.	279
Presentation of—	
books—Duke of Northumberland. (R. 1859)	110
books—Imperial Library of Vienna. (R. 1865)	209
books on Brazil—M. M. Lisboa. (R. 1865)	209
books on Egypt—R. Lepsius. (R. 1860)	147
collections by foreign governments. (R. 1876)	299
engravings—C. B. King. (R. 1861)	149
specimens—University of Christiania, Norway. (R. 1868)	187
Preservation—	
and collection of marine invertebrates-W. STIMPSON	84
of antiquities and national monuments in Denmark—J. J. A. Worsaar. (R. 1879)	845
of copper and iron in salt water—A. E. BECQUEREL. (R. 1864)	188
of wood. (R. 1864)	188
Preserving-	
dipters, directions for-R. OSTEN SACKEN	102
fish, directions for—T. H. BEAN. (P. 1881) 467,	464
insects, directions for—A. S. PACKARD	
lepidopters, method of-T. R. PEALE. (R. 1868)	
nests and eggs, instructions for-T. M. BREWER	
plants, directions for—L. F. WARD	
specimens of distomaces—A. M. EDWARDS	
specimens of natural historyS. F. BAIRD. (R. 1856)	l, 34
Pressure and velocity of the wind—J. HANN. (R. 1877) 828	, 398
Pressure of the air—J. HANN and others. (R. 1877) 828	, 898
PRESTWICH, J. Past and future of geology. (R. 1875)	.298
Priestley, J., memoir of, by G. CUVIER. (R. 1858)	109
Priestley's lens, account of—J. Henry. (R. 1859)	110
PRIME, T. Monograph of American corbiculads	145
PRIME, T., and others. Check-list of shells of North America	128
Primitive manufacture of spear and arrow points along line of Savannah river— C. C. Jones, Jr. (R. 1879)	845
Prince Regent's Inlet, meteorological observations in—F. L. McCLINTOCK	146
Princeton, New Jersey, address at, by J. HENRY	E
Principles—	
and methods of palmontology—T. H. HUXLEY. (R. 1869)	228
of crystallography and crystallophysics, explanation of—A. Brezina. (B. 1872)	386
of mechanical theory of heat—J. MÜLLER. (R. 1868)	224
of physics, present fundamental—F. J. Pisko. (R. 1879)	845
of subcutaneous surgery—W. ADAMS	302
Prionotus stephanophrys, new species of Prionotus from coast of California, description of—W. N. Lockington. (P. 1880)	425

•

.

Prize questions—	
	160
Batavian Society of Experimental Philosophy of Rotterdam. (R, 1861.)	149
Dunkirk Society for Encouragement of Sciences, Letters, and Arts. (R. 1865)	209
Holland Society of Science, Harlem. (R. 1861, 1864, 1867, 1878) 149, 188, 224,	275
Imperial Academy of Sciences of Vienna. (R. 1864; R. 1865) 188,	209
Imperial Society of Natural Sciences of Cherbourg. (R. 1864)	
Imperial Society of Science, Agriculture, and Arts of Lille. (R. 1865.)	
London Institution of Civil Engineers. (R. 1862)	
Pontifical Academy of Nuovi Lincei, Rome. (R. 1865; R. 1867) 209,	
Royal Academy of Netherlands. (R. 1861)	
Royal Academy of Science, Literature, and Fine Arts, Brussels. (R. 1878.)	
Royal Danish Society of Sciences. (R. 1862, 1865, 1867) 150, 209,	
Royal Institute for Encouragement of Natural, Economical, and Tech-	
nological Science, Naples. (R. 1878)	275
Royal Prussian Academy of Sciences. (R. 1864)	
Royal Scientific and Literary Institute of Lombardy. (R. 1865)	209
Society for Encouragement of Science, Literature, and Art, Duakirk, France. (R. 1878)	975
Society of Arts and Sciences of Utrecht. (R. 1861; R. 1862) 149,	
Society of Science, Art, and Literature of Hainaut, Mons, Belgium. (R. 1873)	
Problems of rotary motion, presented by gyroscope, precession of equinoxes, and pendulum—J. G. BABNABD.	
Proceedings of-	
Board of Regents—See Regents.	
Establishment, 1858–1854. (R. 1858) 67,	329
U. S. National Museum. Vols. I-IV, 1878-1881 332, 333, 425,	467
U. S. National Museum, catalogue and index of	478
Products of combustion of gun-cotton and gunpowder—Lieut. von Karolyi; B. F. Craig. (R. 1864)	188
Productus giganteus, note on occurrence of, in California—C. A. WHITE. (P. 1880)	
Programme of organization of Smithsonian Institution	
Progress—	
in physics—See J. Müller.	
in science, record of—See Record.	
of astronomical photography—Dr. Lune. (R. 1861)	149
Project of outline history of education in United States—F. A. PACKARD. (R. 1863)	187
Promotion of science in the United States, Tyndall trust for the. (R. 1872)	271
Propagation of disease—J. C. Dalton. (R. 1878)	
Prospectus of a Bibliographia Americana—H. STEVENS. (R. 1848)	_

OF SMITHSONIAN PUBLICATIONS.	281
Providence, Rhode Island, meteorological observations at—A. CASWELL: 1881 to 1860 :	103
1881 to 1876	443
Provincetown, Massachusetts, experiments made at, upon animal heat of fishes— J. H. Kidder. (P. 1879)	888
Provincetown, Massachusetts, littoral marine fauna of—R. RATHBUN. (P. 1880.)	425
Provincial Society of Arts and Sciences, Utrecht. Prize questions. (R. 1862.)	150
Provisional classification of the food collections-G. B. Goode. (P. 1881.) 467,	455
Prussian Royal Academy of Sciences. Prize questions. (R. 1864)	188
Psychrometric observations	157
Psychrometrical tables—J. H. Coffin	87
Pterophryne, on proper specific name of—T. Gill. (P. 1878)	882
Ptychochilus Harfordi, new species of <i>Ptychochilus</i> , description of—D. S. Jor- DAN; C. H. GILBERT. (P. 1881)	467
Public-	
architecture, hints on-R. D. Owen	P
buildings, acoustics applied to-J. HENRY. (R. 1856)	91
instruction, minister of, report to Emperor of France by, on scientific expedition to Mexico. (R. 1864)	188
libraries of the United States, notices of—C. C. JEWETT. (R. 1849) 21 libraries—See Libraries.	, 25
Publications—	
exchange of—Agricultural Association of Milan. (R. 1868)	187
exchange of-Chamber of Commerce of Bordeaux. (R. 1868)	187
of learned societies and periodicals in Smithsonian library. 73, 85, 117, of Smithsonian Institution, list of. (R. 1866.) 74, 203, 226, 245, 278, 290,	179
801, 844, 487	, 4 78
of U.S. Government, 1868-1881-G. H. BOEHMER	477
of U.S. National Museum, list of. (P. 1881)467,	474
periodical received in reading-room of Smithsonian Institution	Q
report on, 1858-1866—S. F. BAIRD. (R. 1858-1866.) 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209	, 214
Puebla, Mexico, description of large fossil gasteropod from—C. A. WHITE. (P. 1880)	425
Puget Sound, description of two new species of flounders (Parophrys ischyurus and Hippoglossoides elassodon) from—D. S. JORDAN; C. H. GILBERT.	405
(P. 1880)	425
Pulmonata geophila, monograph of—W. G. BINNEY; T. BLAND	194
Pulmonata limnophila, monograph of-W. G. BINNEY	148

Q.

QUALE; LEWIS. Account of the cryolite of Greenland. (R. 1866)	214
Qualitative determinations by the blow-pipe—T. EGLESTON. (R. 1872)	271
Quartermasters, authority to, to receive and transmit specimens to Smithsonian.	34
QUATREFAGES, J. L. Memoir of Isidore Geoffroy St. Hilaire. (B. 1862)	150
Queen Charlotte's Islands, Haidah Indians of-J. G. SWAN	267
Queries about expression for anthropological inquiry—C. DARWIN, (R. 1867.)	215
Queries relative to tornadoes—J. HENRY	190
Questions relative to the food fishes of the United States—S. F. BAIRD	234
Questions—Ses Prize questions.	
Quetelet, L. A. J., eulogy on, by E. MAILLY. (R. 1874)	286
QUICK, E. R. Mounds in Franklin county, Indiana. (R. 1879)	845
Quillehute Indians of west coast of Washington Territory, method of taking	
surf smelt by-J. G. Swan. (P. 1880)	425
Quincy, Illinois, mounds near—W. G. Armstrong. (R. 1879)	345

R.

Raddit catching the sun in a trap (Omana myth)—J. O. Dorsey. (E. 1879-80.)	2/0
Race, the human, probable future of—A. DE CANDOLLE. (B. 1875)	298
Races—	
intermixture of—G. GIBBS. (R. 1864)	188
lower, social and religious condition of the-J. LUBBOCK. (R. 1869)	228
mixed, in Liberia-E. D. BLYDEN. (R. 1870)	244
Radiant heat, reports on state of knowledge of—B. Powell. (R. 1859)	110
Radiates of Arctic America—A. E. VERRILL	342
Radiation, lecture on-J. TYNDALL. (R. 1868)	224
Rafinesque's memoirs on North American fishes, review of—D. S. Jordan	
Raia inornata, note on—D. S. JORDAN; C. H. GILBERT. (P. 1881)	457
Rais rhina, new species of ray from coast of California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	
Raia stellulata, new species of ray from Monterey, California, description of— D. S. JORDAN; C. H. GILBERT. (P. 1880)	
Rain—	
and atmospheric pressure—J. HANN. (R. 1877) 323	, 398
and barometric minima—T. REYE. (R. 1877) 323	, 398
and snow, tables of precipitation in—C. A. Schott 222,	353

Rain—Continued.
articles on, preserved in Smithsonian. (R. 1874) 286, 353
gauges-R. H. GARDINER. (R. 1858)
influence of, upon the barometer—J. HANN. (R. 1877) 828, 896
observations 15
presence of electricity during fall of-Prof. PALMIERI. (R. 1870) 24
quantity of, at different heights-O. W. Morris; J. Henry. (R. 1855.)
snow, or hail, electricity of induction in strata of atmosphere surrounding cloud resolving into—F. ZANTEDESCHI. (R. 1870) 24-
Rain-charts of United States, 1, 2, 8: summer, winter, and year—C. A. Schott. 374
Rana pipiens, anatomy of nervous system of—J. WYMAN 45
Raptores, oölogy of—T. M. Brewer.
Rat, pouched, of Georgia, habits of-W. GESNER. (R. 1860) 14
RATHBUN, R. Littoral marine fauna of Provincetown, Cape Cod, Massachusetts. (P. 1880)426
RATHBUN, R. Marine invertebrates from New England, distributed by U. S. National Museum, list of. (P. 1881) 467, 465, 471
RATHBUN, R.; VERRILL, A. E. Marine invertebrata of New England, distributed by Fish Commission. (P. 1879)
Ratio between diameter and circumference of a circle—W. FERREL 233
Rattlesnake, researches upon venom of—S. W. MITCHELL
Rattlesnakes, bibliography of—S. W. MITCHELL
Rattlesnakes, genera and species of—E. D. Copp. 186
RAU, C.—
Agricultural flint implements in southern Illinois. (R. 1868.) 224, 440, 370
Agricultural implements of North American stone period. (R. 1868.) 187, 440
Ancient aboriginal trade in North America. (R. 1872) 271, 440, 385
Anthropological publications, list of. (P. 1881) 467
Archeological collection of the U.S. National Museum 287
Artificial shell-deposits in New Jersey. (R. 1864) 188, 440, 362
Drilling in stone without metal. (R. 1868) 224, 440, 372
Gold ornament from mound in Florida, observations on. (R. 1877.) 323, 440, 403
Indian pottery. (R. 1866) 214, 440, 368
Memoir of C. F. P. Von Martius. (R. 1869) 228, 440, 251
North American stone implements. (B. 1872) 271, 440, 382
Palenque tablet in the U.S. National Museum
Stock-in-trade of an aboriginal lapidary. (R. 1877) 828, 440, 402
Translation of Baegert's account of the Indians of southern California.
(R. 1868; R. 1864)
Translation of Romer's antiquities of Hungary. (R. 1876) 299, 440, 392
RAWSON, R. Memoir of Eaton Hodgkinson. (R. 1868) 224
KWAN ME C: A DOLON'S MANING NAMP C'INQUIGNOAGGA (K. 1867) YE

Reading-room of Smithsonian Institution, periodicals received in, 1868 Q
Recalculation of atomic weights. Constants of Nature Part v—F. W. CLARKE. 441
Receipts of specimens—See annual reports.
Recent—
improvements in the chemical arts—J. C. BOOTH; C. MOBFIT 27
progress in physics-J. MÜLLER. (R. 1855-1858) 77, 91, 107, 109
progress in relation to the theory of heat—A. Cazin. (R. 1868) 224
progress in science, reports on—See Record of recent progress.
researches on secular variations of planetary orbits—J. N. STOCKWELL. (R. 1871)249
Recommendation of Shea's Indian linguistics—G. GIBBS and others. (R. 1861.) 149
Record of—
auroras in higher northern latitudes—P. Force 84
meteorological material preserved in Smithsonian. (R. 1874) 286
monthly meteorological reports preserved in Smithsonian Institution. (R. 1878)275
Record of recent progress in science, 1879 and 1880—
Anthropology-O. T. MASON. (R. 1880) 442, 432
Astronomy—E. S. Holden. (R. 1880)———————————————————————————————————
Botany-W. G. Farlow. (R. 1880) 442, 430
Chemistry—G. F. BARKER. (R. 1880)
Geology—G. W. HAWES. (R. 1880) 442, 428
Mineralogy—G. W. HAWES. (R. 1880) 442, 428
Physics—G. F. Barker. (R. 1880)
Zoology—T. Gill. (R. 1880)
Recording Indian languages, method of—J. O. DORSEY; A. S. GATSCHET; S. R. RIGGS. (E. 1879-80)
Records and results of magnetic survey of Pennsylvania—A. D. BACHE 166
Red river of the North, ethnology of Indians of valley of—W. H. GARDNER. (R. 1870)
Red river settlement, Hudson's Bay Territory, Indian remains in—D. GUNN. (R. 1867)
REED, H. Lecture on "the Union." (R. 1854)
Refraction of sound—W. B. TAYLOB. (R. 1875)
Regents of Smithsonian Institution— list of, 1846
list of, 1846-1879. (See also each annual report) 329
memorial of Board of, relative to Smithson fund. (R. 1850; R. 1866.) 28, 214, 329
memorial of Board of, to Congress asking appropriations for museum.
(R. 1867)215, 829
memorial to Congress in behalf of, relative to new museum building— G. Bangroff. (R. 1876)
proceedings of Board of. (See also each annual report) A, G, H, I, M, 829
nucceedings of Doord of an dooth of Tosonh Honny

Regents of Smithsonian Institution—Continued.
proceedings of Board of, relative to the electro-magnetic telegraph 115, 829
report of Committee of, on Corcoran Art Gallery. (R. 1872) 271, 829
report of Committee of, on fire at Smithsonian—R. Wallach; J. Henry. (R. 1864) 188, 829
report of Committee of, on Museum — A. GRAY; H. COPPÉE. (R. 1874) 286, 829
report of Committee of, on Museum—A. Gray; A. A. Sargent; H. Clymer. (R. 1876) 299, 829
report of Committee of, on Stanley's gallery of Indian portraits. (R. 1857) 107, 829
report of Committee of, on communication of J. Henry relative to electro- magnetic telegraph—C. C. Felton. (R. 1867) 107, 829
reports of Committee of, on distribution of Smithsonian income. J. A. Pearce; J. Meacham. (R. 1858) 67, 329
Regents of University of State of New York. Acknowledgment for specimens.
(R. 1865) 209, 829
Registration of periodical phenomena—J. HENRY. (R. 1855)
Registry of periodical phenomena
Regulations of the Smithsonian Institution, 1872 280
Regulations of the U.S. National Museum-G.B. Goode. (P. 1881) 467, 445
REICHARDT, W. Present state of knowledge of cryptogamous plants. (R. 1871.) 249
REID, D. B. Architecture in relation to ventilation, warming, lighting, fire- proofing, acoustics, and health. (R. 1856)
Reigen Mazatlan collection of mollusks—P. P. CARPENTER 252
Reindeer, man as the contemporary of, in middle Europe. (R. 1867) 215
Relation—
between barometric variations and atmospheric currents—M. PESLIN. (R. 1877) 828, 898
of food to work, and its bearing on medical practice—S. HAUGHTON.
(R. 1870) 244
of physical sciences to science in general—H. Helmholtz. (R. 1871) 249
of time and space, lecture on—S. ALEXANDER. (R. 1861) 149
Relationship—
circular respecting degrees of—L. H. Mobgan
system of, of Cree Indians—E. A. WATKINS. (R. 1862)
systems of—L. H. Morgan 218
Relative intensity of heat and light of sun-L. W. MEECH. (R. 1856) 91, 83
Relic, ancient, of Maya sculpture, remarks on—A. Schott. (R. 1871) 249
Relies—
ancient, in Missouri—J. W. Foster. (R. 1868) 187
ancient Indian-J. Jones 259
ancient, of northwestern Iowa—J. B. Outrs. (R. 1872) 271
presented by J. H. Devereux. (R. 1872) 271

Religion of Shoshone Indians—A. G. BRACKETT. (R. 1879)	5
Religious condition of lower races of man—J. LUBBOCK. (R. 1869)	
Remains—	
aboriginal, of Tennessee-J. JONES 2	9
ancient, in Colorado—E. L. BERTHOUD. (R. 1867)21	15
human, from Patagonia—A. RIED. (R. 1862)	
Indian, in Caddo parish, Louisiana—T. P. HOTCHKISS. (R. 1872) 2	
of later prehistoric man from caves in Alaska-W. H. Dall 31	
of walrus (?) in Maine—C. H. Boyd. (P. 1881)	
prehistoric, in Montana—P. W. Norris. (R. 1879)	
prehistoric, in vicinity of city of Washington, D. C.—T. R. PEALE. (R. 1872)	
Zapotec, account of—J. G. SAWKINS; B. MAYER.	
Remains—See Ancient remains.	
Remarkable forms of hail-stones in Georgia—S. ABICH. (R. 1869)2	28
Remarks—	
concerning nature of currents of air—A. Colding. (R. 1877) 323, 33	96
on ancient relic of Maya sculpture—A. Schott. (R. 1871)2	
on Cara gigantesea of Yzamal in Yucatan—A. Schott. (B. 1869) 2	
on kjökken-möddings on northwest coast of America—P. Schumacher. (R. 1878)	
on osteology of Opheosaurus ventralis—R. W. SHUFELDT. (P. 1881) 4	
on physical geography of North America—J. FROEBEL. (R. 1854)	
on species of genus Chiurus—W. N. Lockington. (P. 1880)	
Reminiscences of Joseph Henry by S. B. Dop	
Reparatory inflammation in arteries after ligature, etc., nature of. Toner lec-	,,,
ture No. VII—E. O. SHAKESPEARE 32	
Reply to criticisms of J. Hann by W. FERREL. (R. 1877) 822, 32	98
Report—See Henry, J., Baird, S. F., Regents.	
Report of—	
American Academy of Arts and Sciences on organization of Smithsonian—E. Everett; J. Sparks; B. Pierce; H. W. Longfellow; A. Grav. (R. 1858)	67
architect—B. S. Alexander. (R. 1854)	
architects, 1879, 1880—Cluss and Schulze. (R. 1879; R. 1880) 345,	
Board of Regents of Smithsonian Institution, 1847-1889 G, H, I, 21,	
28, 51, 57, 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 214, 215, 224, 228, 244, 249, 271, 275, 286, 298, 299, 828, 841, 345, 4	42
Bureau of Ethnology, firstJ. W. Powell. (E. 1879-80)	
chemist—F. W. TAYLOR. (R. 1880)	
Commission on general stereotype catalogue of public libraries. (R. 1850.) 28,	
Committee of American Association on system of combined meteoro-	
	51

Report of—Continued.
Executive Committee (See also each annual report)
Executive Committee on Henry statue. (R. 1880)
experiments on animal heat of fishes—J. H. Kidden. (P. 1879) 888
explorations in New Mexico and Arizona-J. STEVENSON. (R. 1880) 442
Government explorations and surveys. (R. 1878) 84
National Museum Building Commission for 1879. (R. 1879) 845, 409
National Museum Building Commission for 1880. (R. 1880) 442, 434
Secretary of Smithsonian Institution, 1847-1877—J. Henry. (R. 1847-1877) F, H, I, 21, 28, 51, 57, 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 214, 215, 224, 228, 244, 249, 271, 275, 286, 298, 299, 824
Secretary of Smithsonian Institution, 1878-1880—S. F. BAIRD. (R. 1878-1880)
Senate Judiciary Committee on management of Smithsonian—A. P. But- LER. (R. 1855)
Smithsonian Institution—See Report of Board of Regents.
visit to Luray Cavern-O. T. MASON. (R. 1880) 442, 433
Report of Committee of Regents of Smithsonian Institution on-
best use of new hall of the Institution—L. AGASSIZ. (R. 1867) 215, 824
communication from S. F. B. Morse—C. C. Felton. (R. 1857) 107, 826
Corcoran Art Gallery. (R. 1872) 271, 826
electro-magnetic telegraph—C. C. Felton. (R. 1857) 107, 825
fire of January, 1865-R. Wallach; J. Henry. (R. 1864) 188, 826
Greek album82
Museum—A. Gray; H. Coppée. (R. 1874)
Museum—A. Gray; A. A. SARGENT; H. CLYMER. (R. 1876) 296
organization of InstitutionB, L, 826
Stanley's gallery of Indian portraits. (R. 1857) 107, 826
system of accounts. (R. 1866) 214
Washington canal—R. DELAFIELD. (R. 1868) 224, 826
Report on—
building stones—D. D. Owen 826
catalogue system—C. C. JEWETT. (R. 1849)2
Centennial Exhibition—S. F. BAIRD. (R. 1876) 299, 307
contents of bottles of water from Gulf of Mexico—W. G. Farlow. (P. 1881)
copyright system—C. C. JEWETT. (R. 1851) 51
effect of frost on stones—C. G. PAGE 829
ethnological collections of Museum of Lausanne—F. Troyon. (R. 1861) 149
exploration of ancient mounds in Union county, Kentucky—S. S. Lyon. (R. 1870)244
fishes of New Jersey coast, as observed in 1854—S. F. BAIRD. (R. 1854)
fossils from Nebrasku-J. LEIDY. (R. 1851)

Report on—Continued.
Halliwell manuscripts—C. C. Jewerr. (R. 1852)
history of discovery of Neptune—B. A. Gould 18
improved system of numeration—W. B. TAYLOB. (R. 1867)
international exchanges—G. H. BOEHMER 477
lacustrian settlements—A. Morlot. (R. 1868) 187
library—C. C. JEWETT. (R. 1848-1858) I, 21, 28, 51, 57, 67
limpets and chitons of Alaskan and Arctic regions—W. H. DALL. (P. 1878)
magnetic telegraph 115
meteorological system—E. FOREMAN. (R. 1852)
meteorology of the United States—E. LOOMIS. (R. 1847) H
mollusca of west coast of North America—P. P. CAPENTER 252
Museum and statistics of British Museum—S. F. BAIRD. (R. 1850) 23
paleontological collections made by Lieut. G. K. Warren—F. B. MEEK; F. V. HAYDEN
plan of library—C. C. JEWETT. (R. 1847)
proposed exhibit by Smithsonian Institution at the Centennial Exhibition—S. F. BAIRD. (R. 1875) 298
public libraries of United States—C. C. JEWETT. (R. 1849) 21
publications, exchanges, explorations, and Museum, 1851–1877—S. F. BAIRD. (R. 1851–1877) 51, 57, 67, 75, 77, 91, 107, 109, 110, 147, 149, 150, 187, 188, 209, 214, 215, 224, 228, 244, 249, 271, 275, 286, 298, 299, 323
recent progress in physics: electricity, galvanism—J. MÜLLER. (R. 1855–1857) 77, 91, 107
shell heaps of Tampa Bay, Florida—S. T. WALKER. (R. 1879) 345
shells presented to Academy of Natural Sciences—J. LEIDY; G. W. TRYON. (R. 1865)
state of knowledge of radiant heat—B. Powell. (R. 1859) 110
survey of economic geology of Trinidad—G. P. Wall; J. G. Sawkins. (R. 1856)
system of accounts. (R. 1866) 214
use of new hall in Smithsonian building—L. Agassiz. (R. 1867) 215
Report to Emperor of France by Minister of Public Instruction on scientific expedition to Mexico. (R. 1864)
Reports, meteorological, preserved in Smithsonian Institution, list of. (R. 1873.) 275
Reports of—
astronomical observatories, 1879—E. S. HOLDEN. (R. 1879) 845, 410
astronomical observatories, 1880—E. S. Holden; G. H. Boehmer. (R. 1880)
Committees of Regents, 1846-1877
Secretary of Smithsonian Institution, J. Henry, 1865-1877 343
Smitheonien Institution cetalogue and index of

2	٤	3	(9

OF SMITHSONIAN PUBLICATIONS.

Reports on transactions of Geneva Society of Physics—See Geneva.	
Reptiles—	
circular relating to collections of living	320
	192
cretaceous, review of-J. Leidy. (R. 1864)	188
dates of first appearance of—F. B. Hough	182
extinct—J. Leidy	192
North American, catalogue of. Part I. Serpents—S. F. BAIRD; C. GIRARD	4 9
Reptilia, check-list of North American—E. D. COPE.	292
Request for bison by Bern Museum. (R. 1865)	209
Request for shells-W. E. LOGAN. (R. 1859)	110
Research—	
ethnological—E. H. Davis. (R. 1866)	214
relative to ethnology and philology of America, instructions for-G.	•
GIBBS	160
upon the atmosphere, plan of a-C. M. WETHERILL. (R. 1866)	214
Researches—	
archeological, at Concise—F. TROYON. (R. 1861)	
archeological, in Nicaragua—J. F. Bransford	383
by officers of Hudson's Bay Company, letter of Sir G. Simpson	187
in sound—J. Henry. (R. 1878)	4 06
on ammonia-cobalt bases—W. GIBBS; F. A. GENTH	88
on electrical rheometry—A. Secchi	36
on figure of equilibrium of a liquid mass withdrawn from action of gravity—See J. Plateau.	
on secular variations of planetary orbits—J. N. STOCKWELL. (R. 1871.)	249
relative to nebulæ—Prof. GAUTIER. (R. 1868)	187
relative to planet Neptune—S. C. WALKER	3
scientific, list of apparatus available for. (B. 1878)	841
scientific, of Peltier—F. A. Peltier. (R. 1867)	215
upon anatomy and physiology of Chelonia—S. W. MITCHELL; G. R. MOBEHOUSE	159
upon fever—H. C. Wood	857
upon Hydrobiina and allied forms-W. STIMPSON	201
upon venom of the rattlesnake—S. W. MITCHELL	135
Residuary legacy—See Smithson.	
Resolutions of Congress—See Congress.	
Resolutions of Regents—See each annual report.	
Resonance, electric, of mountains, observations on the—H. DE SAUSSURE. (R. 1868)	224
Respiration, apparatus for testing results of—M. Pettenkofer. (R. 1864) 19	188

ALPHABETICAL INDEX

Respiration in the Chelonia, researches on—S. W. MITCHELL; G. R. MORE-HOUSE
Results of—
magnetic survey of Pennsylvania—A. D. Bache 166
meteorological observations at Brunswick, Maine—P. CLEAVELAND 204
meteorological observations at Marietta, Ohio—S. P. HILDRETH; J. WOOD120
meteorological observations at Providence, Rhode Island, 1831-1860, 1881-1876—A. CASWELL
meteorological observations in the United States, 1854–1859 157, 185
precipitation in rain and snow in United States—C. A. Schott 222, 353
spectrum analysis applied to heavenly bodies—W. Huggins. (R. 1866.) 214
RETZIUS, A. Present state of ethnology in relation to form of the skull. (B. 1859)
Review of—
American birds. Part I. North and Middle America-S. F. BAIRD 181
American species of genus Scops—R. RIDGWAY. (P. 1878) 33
classification of birds-W. LILLJEBORG. (R. 1865) 209, 364
cretaceous reptiles of the United States-J. LEIDY. (R. 1864)
genera and species of family Centrarchide-C. L. McKAY. (P. 1881) 46
genus Centurus—R. RIDGWAY. (P. 1881) 46
pleuronectids of San Francisco-W. N. Lockington. (P. 1879) 33
Rafinesque's memoirs on North American fishes—D. S. JORDAN
Revisions of nomenclature of North American birds—R. RIDGWAY. (P. 1880.) 42
Revolutions of crust of the earth—G. PILAR. (R. 1876)
REYE, T. Rain-fall and barometric minima. (R. 1877) 822, 395
REYNOLDS, J. E. Hydrogen as gas and metal. (B. 1870) 24
Rhees, W. J.—
Oatalogue and index of publications of the Smithsonian Institution, U. S. National Museum, etc., 1846–1881478
James Smithson and his bequest. (R. 1879) 345, 330
Journals of Board of Regents, reports, statistics, etc., of Smithsonian In-
atitution
List of public libraries, institutions, and societies in United States and British Provinces116
List of public libraries, institutions, and societies in United States in cor-
respondence with Smithsonian Institution 238
Origin and history of Smithsonian Institution 328
Scientific writings of James Smithson, (edited by)
Rheometry, electrical, researches on—A. SECCHI
Rhinobatidæ, American, synopsis and descriptions of—S. Garman. (P. 1890.) 425
Rhinonemus caudacuta, identity of with Gadus cimbrius—G. B. Goode; T. H. BEAN. (P. 1878)

. OF SMITHSONIAN PUBLICATIONS.	291
Rhode Island, list of alge of—S. T. OLNEY	241
Rhode Island, Providence, meteorological observations at, 1881-1860, 1881-1876— A. CASWELL	443
Rhodospermene-W. H. HARVEY	48
RIDGWAY, R.—	
Catalogue of birds of North America. (P. 1880)	425
Catalogue of Old World birds in U. S. National Museum. (P. 1881)	467
Catalogue of Trochilidæ in U. S. National Museum. (P. 1880)	425
Description of new fly-catcher and new petrel from Sandwich Islands. (P. 1881)	467
Description of new owl from Porto Rico. (P. 1881)	467
Description of new species of birds from Costa Rica and notes on rare	
species. (P. 1878)	882
Descriptions of new species and races of American birds and synopsis of genus Tyrannus. (P. 1878)	882
Descriptions of new species and races of birds in U.S. National Museum. (P. 1878)	882
Descriptions of two new thrushes from the United States. (P. 1881)	467
List of special desiderata among North American birds. (P. 1881)	467
List of species of Middle and South American birds not in National Museum. (P. 1881)	467
Nomenclature of North American birds chiefly in National Museum	422
Notes on some Costa Rican birds. (P. 1881)	467
On Amazilia Yucatanensis and Amazilia cerviniventris. (P. 1881)	467
On duck new to American fauna. (P. 1881)	467
On new humming bird, Atthis Elliotti, from Guatemala. (P. 1878)	882
On two recent additions to North American bird fauna by L. Belding. (P. 1881)	467
Review of American species of genus Scops. (P. 1878)	882
Review of genus Centurus. (P. 1881.)	467
RIED, A. Human remains from Patagonia. (R. 1862)	150
RIGGS, S. R. A dog's revenge (Dakota fable.) (E. 1879-80)	476
Rices, S. R. Grammar and dictionary of Dakota language	40
Ripley, Fort, Minnesota, natural history of the country about—J. E. HEAD. (R. 1854)	75
Rissoidæ, fresh-water—W. G. BINNEY	144
River, Kennebec, opening and closing of the—R. H. GARDINER. (R. 1858)	109
Rivers—	
dates of opening and closing of—F. B. HOUGH.	182
diminution of water in-H. G. WEX. (R. 1875)	298
improvement of navigation of—C. ELLET, Jr.	18
method of ascertaining amount of water in—A. A. HUMPHREYS. (R. 1858)	109
Roads and bridges, lectures on—F. Rogers. (R. 1860; R. 1861) 147	, 1:19

•

Robertson, R. S.—
Age of stone and troglodytes of Breckenridge county, Kentucky. (B. 1874)
Antiquities of Allen and De Kalb counties, Indiana. (B. 1874)
Antiquities of La Porte county, Indiana. (R. 1874)
Antiquities of Nashville, Tennessee. (R. 1877)
ROBINSON, E., and others. On publication of Spanish works on New Mexico.
(R. 1855)
ROBINSON, E., and others. On publication of Squier and Davis' ancient monuments
Rock Bluff, Illinois, description of human skull from-J. A. MRIGS. (R. 1867.)
Rock-oil, or petroleum, history of—T. S. HUNT. (R. 1861)
Rock river valley, Illinois, mound builders in-J. SHAW. (R. 1877)
Rocks-
catalogue of—J. W. FOSTER. (R. 1854)
catalogue of—C. T. Jackson. (R. 1854)
catalogue of—J. Locke. (R. 1854)
catalogue of-J. D. WHITNEY. (R. 1854)
crystalline, metamorphism, and formation of—G. A. DAUBRÉE. (R. 1861.)
in Green river valley, Indian engravings on face of—J. G. BRUFF. (R. 1872)
Rocky Mountain goat, habits of the-J. C. MEBRILL. (P. 1879) 3
Rodent, rure, (Cricetodipus parvus)-F. W. TRUE. (P. 1881)
ROEHRIG, F. L. O. Language of Dakota or Sioux Indians. (R. 1871) 249, 3
ROGERS, F. Lectures on roads and bridges. (R. 1860; R. 1861)
ROGERS, W. B. Memorial address on Joseph Henry
Rome, Pontifical Academy—See Prize questions.
ROMER, F. F. Prehistoric antiquities of Hungary. (B. 1876) 299, 440, 35
ROMEBO, M. Explorations of John Xantus in Mexico. (R. 1862)
Rosing, J. Exchange system. (R. 1865)
Ross, B. R. Eastern Tinneh Indians. (R. 1866) 214, 3
Ross, B. R. Observations in Hudson's Bay Territory. (R. 1859)
Rotary motion, problems of—J. G. BARNARD
ROTHBOCK, J. T. Sketch of flora of Alaska. (R. 1867)215,3
Rotterdam, Batavian Society of Experimental Philosophy of. Prize questions. (R. 1861)
Royal—
Academy of Netherlands. Prize questions. (R. 1861)
Academy of Science, Madrid. On exchanges. (R. 1861)1
Danish Society of Sciences. Prize questions. (R. 1862, 1865, 1867.) 150, 209, 21
Horticultural Society, London. Exchange of publications. (R. 1861) 14
Institution of Great Britain, history of—E. MAILLY. (R. 1867) 21
Prussian Academy of Sciences, Prize questions. (R. 1864)

Royal—Continued. Scientific and Literary Institute of Lombardy. Prize questions. (R. 1865.) 209
Society of London, list of Smithson's papers presented to. (R. 1858.) 67, 880
Society of London, notice of Smithson, by D. Gilbert, president of. (B. 1868)
Society of London, origin and history of—C. A. ALEXANDER. (R. 1868.) 187
Society of Victoria, address of president of—R. L. J. ELLERY. (R. 1868.) 224
ROYCE, C. C. Cessions of land by Indian tribes to United States. (E. 1879-80.) 476
Ruin, ancient, in Arizona—J. C. Y. Lee. (R. 1872) 271
Ruins-
at Savannah, Tennessee—J. P. Stelle. (R. 1870)
in Hardin county, Tennessee—J. P. Stelle. (R. 1870) 244
in White river canon, Pima county, Arizona—R. T. Burn. (R. 1879.) 845
Rules—
for cataloguing libraries—C. C. JEWETT 47
for examination of specimens. (R. 1880) 442
of distribution of Smithsonian publications 290
RUNKLE, J. D. New tables for determining perturbation of planets 79
RUNKLE, J. D. Supplement for asteroid perturbations 94
Rupert's Land, Institute of, circular of the. (R. 1861) 149
Rush, R.—
correspondence of, relative to Smithson828
letter from, relative to James Smithson. (R. 1853) 67, 328
notice of, by J. A. Pearce. (R. 1859) 110
RUSSELL, R. Lectures on meteorology, with notes by J. Henry. (R. 1854) 75
Russia, meteorology in—A. Woeikoff. (R. 1872) 271
Russian America—
ethnological investigations in, suggestions for-G. GIBBS 207
suggestions relative to objects of scientific investigation in—J. Henry 207
Tinneh Indians of—G. GIBBS; B. Ross; W. L. HARDISTY; S. JONES. (R. 1866)
Yukon, journey to the—W. W. KIBBY. (R. 1864) 188
RUTIMEYER, L. Fauna of middle Europe during the stone age. (R. 1861) 149 RYDER, J. A.—
Camaraphysema, a new type of sponge, (P. 1880)425
List of North American species of myriapods belonging to family of Lysiopetalidæ, with description of blind form from Luray Cave, Virginia. (P. 1880).
ginia. (P. 1880)

S.

SABINE, E. Magnetic storms. (R. 1860)	147
SABINE, E. On continuance of magnetic observations. (R. 1858)	109
Sable Island Bank, description of Argentina systemsium, new deep-sea fish from-	
G. B. GOODE; T. H. BEAN. (P. 1878)	882
Sacken, Osten—See Osten Sacken.	
Sacramento, California—	
meteorological observations at—F. W. HATCH. (R. 1854)	75
meteorological observations at—T. M. Logan. (R. 1854)	75
meteorology of—T. M. Logan. (R. 1857)	107
Sacramento river, description of new species of Ptychochilus (Ptychochilus Herfordi) from—D. S. JORDAN; C. H. GILBERT. (P. 1881)	467
Saint Augustine, Florida, catalogue of casts of heads of Indian prisoners at— R. H. Pratt. (P. 1878)	382
Saint Christopher, West Indies, description of new subspecies of saxigilla from—G. N. LAWRENCE. (P. 1881)	457
Saint Croix Island, flora of—H. F. A. Eggers	813
Saint George's Banks, occurrence of Hippocampus antiquorum on—G. B. Goods. (P. 1878)	832
Saint Hilaire, Etienne Geoffroy, memoir of, by M. FLOURENS. (R. 1861)	
Saint Hilaire, Isidore Geoffroy, memoir of-J. L. QUATREFAGES. (B. 1862)	
Saint John's river, Florida, catalogue of fishes of-G. B. Goode. (P. 1879)	
Saint John's river, Florida, shell-heaps at mouth of—S. P. MAYBERRY. (R. 1877)	
Saint Louis, Missouri, ancient mound in—T. R. PEALE. (R. 1861)	
Saint Louis, Missouri, temperature of—A. FENDLER. (R. 1860)	147
Saint Martin, Isle Jesus, Canada East, description of observatory at—C. SMALL-wood. (R. 1856)	91
Saint Michael's, Alaska, occurrence of Hippoglossus vulgaris at—T. H. Bran. (P. 1879)	833
Saint Michael's, Alaska, occurrence of Stichous punctatus at—T. H. Bran.	
	332
Saint Petersburg Academy of Sciences. Exchange system. (B. 1867)	215
Saint Thomas—	
earthquakes in—G. A. LATIMER. (R. 1867)	
Bullicate in the island of—G. M. Dalland. (16. 1007)	215
whirlwind at, August 21, 1871—A. Colding. (B. 1877) 323,	39 8
Saint Vincent, catalogue of Ober's collection of birds of-G. N. LAWRENCE.	832
Salamander—	
habits of a species of—C. MANN. (B. 1854)	75
inhabiting Wisconsin-P. R. Hov. (R. 1854)	75
or noushed not of Goorgie hebits of W Greyne (P 1980)	147

alisbury, England, notice of Blackmore Museum at. (B. 1868)	•	
alisbury, England, notice of Blackmore Museum at. (B. 1868)		•
almon in Columbia river, Chinook names of—S. B. SMITH. (P. 1881)	OF SMITHSONIAN PUBLICATIONS.	295
almon in Columbia river, Chinook names of—S. B. SMITH. (P. 1881)	isbury, England, notice of Blackmore Museum at. (B. 1868)	224
almonides of upper Columbia, notes on the—C. Bendies. (P. 1881)		467
alt deposit on Petite Anse Island—E. W. HILGARD	•	467
alt water, preservation of copper and iron in—A. E. BECQUEREL. (R. 1884.) amoan Islands, fishes from the—T. H. STERETS	••	248
amoan Islands, fishes from the—T. H. STERETS	•	188
description of new gobioid fish (Othonops cos) from—R. SMITH. (P. 1881.) description of new species of Gobiesox (Gobiesox rhessodon) from—R. SMITH. (P. 1881)		808
description of new gobioid fish (Othonops cos) from—R. SMITH. (P. 1881.) description of new species of Gobiesox (Gobiesox rhessodon) from—R. SMITH. (P. 1881)		
description of new species of Gobiesox (Gobiesox rhessodon) from—R. SMITH. (P. 1881)		467
SMITH. (P. 1881)	• • • • • • • • • • • • • • • • • • • •	
occurrence of species of cremnobates at—R. SMITH. (P. 1880)		467
occurrence of species of cremnobates at—R. SMITH. (P. 1880)		
an Francisco, California— climate of—H. Gibbons. (R. 1854) fishes of—See Jordan, D. S., Gilbert, C. H., Leckington, W. N. review of Pleuronectidæ of—W. N. Lockington. (P. 1879) andwich Islands, description of new fly-catcher and new petrel from—R. RIDGWAY. (P. 1881) G. E. WARING Anitary drainage of Washington, suggestions for the. Toner lecture No. VIII— G. E. WARING Anta Barbara, California, pleistocene fossils collected at, by E. Jewett—P. P. CARPENTER Anta Barbara channel, California, description of two new species of scopeloid fishes (Sudis ringens and Myctophum crenulare) from—D. S. Jordan; C. H. Gilbert. (P. 1880) Anta Catalina Island, California, description of new flounder (Xystreurys liolepis) from—D. S. Jordan; C. H. Gilbert. (P. 1880) Anta Lucia Cosumalwhuapa, Guatemala, sculptures of—S. Habel Anta Rosa Island, history and antiquities of—S. Bowers. (R. 1877) ARGENT, A. A., and others. Report on Museum. (R. 1876) ARGENT, W. D. Influence of aurora on the telegraph. (R. 1870) ARGENT, W. D. Influence of aurora on the telegraph. (R. 1870) ARGENT, W. D. Influence of aurora on the telegraph. (R. 1870) ARTORIUS, C. Earthquakes in eastern Mexico, January, 1866. (R. 1866) ARTORIUS, C. Eruption of volcano of Colima. (R. 1869) ALTORIUS, C. Eruption of volcano of Colima. (R. 1869) ALTORIUS, C. Gommander. Tidal record at Wolstenholm Sound, 1849, 1850 AUNDERS, Commander. Tidal record at Wolstenholm Sound, 1849, 1850 AUNDERS, Commander. Tidal record at Wolstenholm Sound, 1849, 1850 AUNDERS, Commander. Tidal record at Wolstenholm Sound, 1849, 1850 AUNDERS, Commander. Tidal record at Wolstenholm Sound, 1849, 1850 AUNDERS, Commander. Tidal record at Wolstenholm Sound, 1849, 1850		428
climate of—H. Gibbons. (R. 1854)	occurrence of species of cremnobates at—R. SMITH. (P. 1880)	426
fishes of—See Jordan, D. S., Gilbert, C. H., Leckington, W. N. review of Pleuronectidæ of—W. N. Lockington. (P. 1879)	·	
review of Pleuronectidæ of—W. N. Lockington. (P. 1879)	·	78
andwich Islands, description of new fly-catcher and new petrel from—R. RIDGWAY. (P. 1881)	• • • • • • • • • • • • • • • • • • • •	
RIDGWAY. (P. 1881)	•	888
anitary drainage of Washington, suggestions for the. Toner lecture No. VIII— G. E. WARING		AQT
G. E. Waring anta Barbara, California, pleistocene fossils collected at, by E. Jewett—P. P. Carpenter anta Barbara channel, California, description of two new species of scopeloid fishes (Sudis ringens and Myctophum crenulare) from—D. S. Jordan; C. H. Gilbert. (P. 1880) anta Catalina Island, California, description of new flounder (Xystreurys liolepis) from—D. S. Jordan; C. H. Gilbert. (P. 1880) anta Lucia Cosumalwhuapa, Guatemala, sculptures of—S. Habel anta Rosa Island, history and antiquities of—S. Bowers. (R. 1877) arcophagus from Beirut, Syria, account of—A. A. Harwood. (R. 1870) argus from Beirut, Syria, account of—A. A. Harwood. (R. 1870) argus Holbrookii, a new sparoid fish from Savannah Bank, description of— T. H. Bean. (P. 1878) artorius, C. Earthquakes in eastern Mexico, January, 1866. (R. 1866) aturn, secular variations of elements of orbit of, with tables—J. N. Stock— well aunders, Commander. Tidal record at Wolstenholm Sound, 1849, 1850 aussure—See De Saussure.	· · · · · · · · · · · · · · · · · · ·	
anta Barbara, California, pleistocene fossils collected at, by E. Jewett—P. P. CARPENTER anta Barbara channel, California, description of two new species of scopeloid fishes (Sudis ringens and Myctophum crenulare) from—D. S. Jordan; C. H. Gilbert. (P. 1880) anta Catalina Island, California, description of new flounder (Xystreurys liolepis) from—D. S. Jordan; C. H. Gilbert. (P. 1880) anta Lucia Cosumalwhuapa, Guatemala, sculptures of—S. Habel anta Rosa Island, history and antiquities of—S. Bowers. (R. 1877) arcophagus from Beirut, Syria, account of—A. A. Harwood. (R. 1870) aredard, A. A., and others. Report on Museum. (R. 1876) argus Holbrookii, a new sparoid fish from Savannah Bank, description of— T. H. Bean. (P. 1878) artorius, C. Earthquakes in eastern Mexico, January, 1866. (R. 1866) aturn, secular variations of elements of orbit of, with tables—J. N. Stock— well aunders, Commander. Tidal record at Wolstenholm Sound, 1849, 1850 aussure—See De Saussure.		
CARPENTER anta Barbara channel, California, description of two new species of scopeloid fishes (Sudis ringens and Myctophum crenulare) from—D. S. Jordan; C. H. Gilbert. (P. 1880)		
fishes (Sudis ringens and Myctophum crenulare) from—D. S. Jordan; C. H. Gilbert. (P. 1880)		252
fishes (Sudis ringens and Myctophum crenulare) from—D. S. Jordan; C. H. Gilbert. (P. 1880)	nta Barbara channel, California, description of two new species of scopeloid	
anta Catalina Island, California, description of new flounder (Xystreurys liolepis) from—D. S. Jordan; C. H. Gilbert. (P. 1880)	fishes (Sudis ringens and Myctophum crenulare) from-D. S. JORDAN	
liolepis) from—D. S. JORDAN; C. H. GILBERT. (P. 1880)		428
anta Lucia Cosumalwhuapa, Guatemala, sculptures of—S. HABEL	· · · · · · · · · · · · · · · · · · ·	
anta Rosa Island, history and antiquities of—S. Bowers. (R. 1877)		
ARGENT, A. A., and others. Report on Museum. (R. 1876)		
ABGENT, A. A., and others. Report on Museum. (R. 1876)		828
ARGENT, W. D. Influence of aurora on the telegraph. (R. 1870)		
ARTORIUS, C. Earthquakes in eastern Mexico, January, 1866. (R. 1866)		
T. H. BEAN. (P. 1878)	· · · · · · · · · · · · · · · · · · ·	
ARTORIUS, C. Earthquakes in eastern Mexico, January, 1866. (R. 1866)	, , ,	882
ABTORIUS, C. Eruption of volcano of Colima. (R. 1869)		
Aturn, secular variations of elements of orbit of, with tables—J. N. STOCK—WELL		
WELL 22 AUNDERS, Commander. Tidal record at Wolstenholm Sound, 1849, 1850 18 aussure—See De Saussure.	• • • • • • • • • • • • • • • • • • • •	0
AUNDERS, Commander. Tidal record at Wolstenholm Sound, 1849, 1850 18 sussure—See De Saussure.		232
aussure—See De Saussure.		
	rage weapons at Centennial Exhibition, study of-E. H. KNIGHT. (R.	

Savannah Bank, description of new sparoid fish (Sargus Holbrockii) from—T. H. BEAN. (P. 1878)
Savannah river, primitive manufacture of spear and arrow points along line of— C. O. Jones, Jr. (R. 1879)
Sawkins, J. G., drawings of Zapotec remains by
SAWKINS, J. G.; WALL, G. P. Economic geology of Trinidad. (R. 1856) 9
Scenery, catalogue of sketches of-J. M. STANLEY.
Scheme for qualitative determinations by the blow-pipe—T. Egleston. (R. 1872)
Scheme of classification for collections of U. S. National Museum—G. B. Goode. (P. 1881)
SCHERZER; SCHWARZ. Table of anthropological measurements. (R. 1866) 214
SCHLAGINTWEIT, H. DE. Ethnographical collections. (R. 1862)
Schlagintweit ethnographic collection, account of the—H. ZISGENBALS. (R. 1867)
SCHLEIDEN, R. Free freight between Germany and United States by North German Lloyd. (R. 1858)
Schoenbein, C. F., notice of, by J. HENRY. (B. 1868) 22
Schoharie, New York, Indian relics from-F. D Andrews. (R. 1879) 34
Schoodic Lakes, Maine, description of new species of Gasterosteus from—T. H. BEAN. (P. 1879)
SCHOTT, A. Remarks on ancient relic of Maya sculpture. (R. 1871) 249
SCHOTT, A. Remarks on the Cara gigantesca in Yucatan. (R. 1869) 22
SCHOTT, A.; MASON, O. T. Leipsic Museum of Ethnology. (R. 1878) 273
SCHOTT, C. A.—
Base-chart of the United States416
Discussion of Caswell's meteorological observations at Providence, Rhode Island
Discussion of Cleaveland's meteorological observations at Brunswick, Maine
Discussion of Hayes' physical observations in the Arctic Seas 196
Discussion of Hildreth's and Wood's meteorological observations at Marietta, Ohio
Discussion of Kane's astronomical observations in the Arctic Seas 123
Discussion of Kane's magnetic observations in the Arctic Seas91
Discussion of Kane's meteorological observations in the Arctic Seas 104
Discussion of Kane's physical observations in the Arctic Seas
Discussion of Kane's tidal observations in the Arctic Seas130
Discussion of McClintock's meteorological observations in the Arctic
Discussion of Smith's meteorological observations made near Washington, Arkansas 131
Tables, distribution, and variation of atmospheric temperature 277
Tables of rain and snow in the United States 222, 353

OF SMITHSONIAN PUBLICATIONS.	297
SCHOTT, C. A.; EVERETT, J. D. Underground temperature. (R. 1874) Schulze—See Cluss & Schulze.	286
SCHUMACHER, P. Ancient graves and shell-heaps of California. (R. 1874)SCHUMACHER, P. Kjökken-möddings on the northwest coast of North America.	286
(R. 1878)	275
Schwarz; Scherzer. Table of anthropological measurements. (R. 1866)	214
on a dominant language for—A. Dr Candolle; J. E. Gray. (R. 1874.)	286
in general, relation of the physical sciences to—H. HELMHOLTZ. (R. 1871.)	249
recent progress in—See Record.	
trust fund for promotion of, in the United States-J. TYNDALL. (R. 1872)	271
Science, Agriculture, and Arts, Society of, Lille—See Prize questions.	
Science, Art, and Literature, Society of, Hainaut—See Prize questions.	
Sciences, Harlem Society of—See Prize questions.	
Sciences, Letters, and Arts, Society of, Dunkirk—See Prize questions.	
Sciences, Natural, Society of, Cherbourg—See Prize questions.	
Sciences, Royal Danish Society of—See Prize questions.	
Scientific—	
character and researches of Smithson—W. R. Johnson	827
Congress of Carlsruhe, 1858—F. J. NICKLÉS. (R. 1860)	147
cooperation—University of Toronto. (R. 1861)	149
education of mechanics and artisans—A. P. PEABODY. (R. 1872) 271,	380
exchanges—See Exchanges.	
expedition to Mexico by French Government. (R. 1864)	188
instructions to Capt. Hall—J. Henry; J. E. Hilgard; S. Newcomb; S. F. Baird; F. B. Meek; L. Agassiz. (R. 1871)	249
investigation in Russian America, suggestions relative to-J. HENRY 2	207
labors of Edward Lartet-F. FISCHER. (R. 1872.)	271
observations, on metric system for-A. Guyor. (R. 1848)	1
progress, reports of. (R. 1880) 442, 427, 428, 429, 430, 431,	1 32
researches of J. C. A. Peltier-F. A. Peltier. (R. 1867)	215
societies, local, organization of—J. HENRY. (R. 1875)	298
writings of Sir William Herschel—E. S. Holden; C. S. Hastings. (R. 1880)	1 26
•	827
Scientific and Literary Institute of Lombardy—See Prize questions.	
Scintillation of the stars—C. Dufour; Kämtz. (R. 1861)	149
	332
Scudder, S. H.—	
	189
• • •	189

SCUDDER, S. H., and others, Insects of Arctic America	842
Sculpture, Maya, remarks on ancient relic of-A. Schott. (R. 1871)	249
Sculptures of Santa Lucia Cosumalwhuapa in Guatemala—S. HABEL	269
Scytalina cerdale, new species of fish from Neah Bay, Washington Territory, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Sea, gradual approach of, upon land—S. P. MAYBERRY. (R. 1867)	215
Seas, the northern-J. BABINET. (R. 1869)	228
Seaton, W. W., memoir of, by J. HENRY. (R. 1866)	214
Sebastichthys carnatus, new species of rockfish from coast of California, description of—D. S. Jordan; C. H. Gilbert. (P. 1880)	425
Sebastichthys chrysomelas, new species of rockfish from coast of California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Sebastichthys entomelas, new species of Sebastichthys from Monterey Bay, California, description of—D. S. Johdan; C. H. Gilbert. (P. 1880)	425
Sebastichthys maliger, new scorpsenoid fish from coast of California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Sebastichthys miniatus, new species of Sebastichthys from Monterey Bay, Cali-	
fornia, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Sebastichthys mystimus, description of—D. S. JORDAN; C. H. GILBERT. (P. 1881)	467
Sebastichthys proriger, new scorpsenoid fish from Monterey Bay, California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Sebastichthys rhodochloris, new species of Sebastichthys from Monterey Bay, California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Sebastichthys serriceps, new species of rock cod from coast of California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Sebastoid fishes, description of seven new species of—D. S. JORDAN; C. H. GIL- BERT. (P. 1880)	425
SECCHI, A. Researches on electrical rheometry	36
Secretary of Smithsonian Institution, report of-See Henry, J., Baird, S. F.	
Secretary of Treasury, letter to, on payment of interest in coin—J. HENEY. (R.	900
1865)	209 81
Secular period of aurora borealis—D. OLMSTED Secular variations of elements of planetary orbits—J. N. STOCKWELL. (R. 1871)	
Seismograph, electro-magnetic—Prof. PALMIERI. (B. 1870)	
Sema, note on—D. S. Jordan. (P. 1880)	425
Semper's method of making dry preparations—J. A. Byder. (P. 1881)	467
Senate Judiciary Committee, report of, on management of Smithsonian—A. P. BUTLER. (R. 1855)	77
Senses, the. Sense of feeling, sense of smell. (R. 1865)	209
Senses, the. Sense of taste, sense of hearing, sense of sight. (R. 1866)	
Séquard—See Brown-Séquard.	
Series, converging, expressing ratio of diameter and circumference of a circle—	233

Seriola Stearnsii, new species of amber fish from Pensacola, Florida, descrip-
tion of—G. B. Goode. (P. 1879) 888
Sermon at funeral of Joseph Henry—S. S. MITCHELL
Serpent venom, bibliography of—S. W. MITCHELL 185
Serpents, catalogue of—S. F. BAIRD; C. GIRARD
Serpents, lecture on nature and cure of bites of—D. Brainard. (R. 1854) 75
Seven cities of Cibola, Coronado's march in search of the—J. H. Simpson.
(R. 1869) 228
Sexual characters of cels—S. T. CATTIE. (P. 1880)425
Shad, western gizzard, Dorosoma cepedianum heterurum, notes on—S. WILMOT. (P. 1878)
SHAKESPEARE, E. O. The nature of reparatory inflammation in arteries after ligature, etc. Toner lecture No. VII
Shark, the oil, of southern California, (Galeorhinus galeus)—D. S. JORDAN; C. H. GILBERT. (P. 1880)
SHARPLESS, T.; PATTERSON, R. Phonography. (R. 1856)
SHAW, J. Mound-builders in Rock river valley, Illinois. (R. 1877) 828
SHEA, J. G. Account of library of Indian linguistics. (R. 1861) 149
Shea's Indian linguistics, recommendation of-G. GIBBS and others. (R. 1861.) 149
Shell-bed skull—A. S. TIFFANY. (B. 1874)
Shell-deposits, artificial, of the United States-D. G. Brinton. (R. 1866.) 214, 440
Shell-deposits, artificial, in New Jersey-C. RAU. (R. 1864)
Shell engravings granted by British Museum. (R. 1868) 187
Shellfish and their allies, lectures on-P. P. CARPENTER. (R. 1860) 147, 152
Shell-heap in Georgia—D. Brown. (R. 1871) 249
Shell-heaps—
and ancient graves of California—P. SCHUMACHER. (R. 1874) 286
at mouth of St. John's river, Florida-S. P. MAYBERRY. (R. 1877) 828
in New Brunswick-J. Fowler. (R. 1870) 244
of Tampa Bay, Florida-S. T. WALKER. (R. 1879) 845
on Mobile river, Alabama—A. S. GAINES; K. M. CUNNINGHAM. (R. 1877)828
Shell-mounds—See Mounds.
Shells-
acknowledgment for—W. H. PEASE. (R. 1861)149
bibliography of. Parts I, II-W. G. BINNEY 142, 174
duplicate, collected by United States exploring expedition-C. WILKES. 193
from California, description of new species of, in U. S. National Museum— W. H. DALL. (P. 1878)832
from Costa Rica kitchen-midden-W. H. Dall. (P. 1878) 882
land and fresh-water, instructions for collecting-J. Lewis. (R. 1866) 214, 363
land and fresh-water, of North America:
Part I. Pulmonata geophila-W. G. BINNEY; T. BLAND 194

Shells—Continued.
land and fresh-water, of North America:
Part II. Pulmonata limnophila and Thallasophila—W. G. BINNEY. 16
Part III. Ampullariidæ, Valvatidæ, (etc.)-W. G. BINNEY 19
Part IV. Strepomatidæ (American melanians)—G. W. TEYON, Jr 25
letter on—J. Lewis 2
North American, check-list of-I. LEA; P. P. CARPENTER; W. STIMP
son; W. G. Binney; T. Prime
North American, circular in reference to collecting 17
of Gulf of California, lectures on the-P. P. CARPENTER. (R. 1859) 11
of Panama, collected by C. B. Adams, descriptions of new species and
varieties of Chitonida and Acmaida in-P. P. CARPENTER 2
presented to Academy of Natural Sciences, report on—J. LEIDY; G. W. TRYON. (R. 1865)
request for duplicate—W. E. LOGAN. (R. 1859)
Shells—Ses Mollusca.
SHEPARD, C. U. Arrangement of mineralogical collection. (B. 1861)
SHEPARD, E. M. Deposit of arrow-heads near Fishkill, New York. (R. 1877.) 35
SHERMAN, W. T. Memorial address on J. Henry
SHERMAN, W. T. Report of National Museum Building Commission, 1879,
1880. (R. 1879; R. 1880)
Shetland, vegetable colonization of—C. MARTINS. (R. 1858)
Shipping fresh fish and other animals, circular relative to—S. F. Barko 36
Shoal Lake, Lake Winnipeg, egging expedition to—D. Gunn. (R. 1867) 2
Shooting stars, observations on—S. MASTERMAN. (R. 1857)
Short memoirs on meteorological subjects, translated by C. Abbe. (R. 1877.) 323, 35
Shoshone Indians, religion, superstitions, and manners of—A. G. Brackett. (R. 1879)
SHUFFLDT, R. W. Remarks upon osteology of Opheosaurus ventralis. (P. 1881.)
Siberia, descriptions of new fishes from-T. H. BEAN. (P. 1881)
Sibley, H. Meteorological telegrams by Western Union Telegraph Company. (R. 1862)
Sight, sense of. (R. 1866)
Sign language among North American Indians-G. MALLERY. (E. 1879-80.) 47
Sign language of deaf mutes-G. MALLERY. (E. 1879-80)
Silurids of fresh waters of North America, synopsis of—D. S. JORDAN 30
Silver coins, assay of, at United States Mint-J. Pollock. (R. 1868) 22
Silver coins, foreign, table of. (R. 1868)
Silvered-glass telescope, on construction of—H. DRAPER
SIMPSON, G. Letter to officers of Hudson's Bay Company relative to scientific
researches 13

SIMPSON, J. H. Coronado's march in search of the seven cities of Cibola.

of smithsonian publications. 80	1
Simson, R. Meteorite in Mexico. (R. 1867) 21	5
Sioux Indians—A. G. BRACKETT. (R. 1876) 29	9
Sioux Indians, language of the—F. L. O. ROEHRIG. (R. 1871) 249, 87	8
Sioux nation of the upper Missouri—T. A. CULBERTSON. (R. 1850) 2	_
Sioux—See Dakota.	
Siredon lichenoides, observations on—W. E. CABLIN. (P. 1881) 46	7
Six Nations, wampum belts of the—W: M. BEAUCHAMP. (B. 1879) 84	5
Skeletons of birds, suggestions for saving parts of—A. NEWTON. (R. 1860) 14	7
Sketch of—	
Academy of Science of Paris-M. FLOURENS. (R. 1862) 15	j()
ancient earthworks of Ohio-I. DILLE. (R. 1866) 21	4
ancient earthworks on upper Missouri—A. BARRANDT. (R. 1870) 24	4
flora of Alaska—J. T. ROTHROCK. (R. 1867)	7
Navajo Indians-J. Letterman. (R. 1855)	77
scientific work of Joseph Henry-W. B. TAYLOR 856, 33	9
Sketches, historical and bibliographical, of archeology of United States—S. F.	
	71
	8
Skull— and long bones from mounds near Albany, Illinois—R. J. FARQUHAR-	
son. (R. 1874) 28	36
human, from Rock Bluff, Illinois, description of-J. A. MEIGS. (R. 1867.) 21	5
human, present state of ethnology in relation to form of the—A. RETZIUS. (R. 1859)	10
shell-bed, from Rock Island, Illinois—A. S. TIFFANY. (R. 1874) 28	36
Skulls and mummy from Patagonia—A. RIED. (R. 1862) 15	0
	2
SMALLWOOD, C. Description of observatory at St. Martin, Isle Jesus, Canada)1
SMART, C. Netes on the Tonto Apaches. (R. 1867)21	
Smell, sense of. (R. 1865)	
Smelt, surf, of northwest coast, and method of taking by Indians—J. G. Swan. (P. 1880)	
	7
SMITH, C. D. Ancient mica mines in North Carolina. (R. 1876) 29	9
·	7
SMITH, J. W. C. Antiquities of Yazoo county, Mississippi. (R. 1874) 28	
SMITH, N. D. Meteorological observations made near Washington, Arkansas. 13	1
Sмітн, R.—	
Description of new gobioid fish (Othonops cos) from San Diego, California. (P. 1881)	7
Description of new species of gobiesox (Gobiesox rhessodon) from San Diego, California. (P. 1881)	. 7
Occurrence of species of cremnobates at San Diego, California. (P. 1880.) 42	

SMITH, S. B. On the Chinook names of salmon in Columbia river. (P. 1881.) 46
Sмітн, S. I.—
Crustaceans of Kerguelen Island
Notice of crustacea dredged off south coast of New England in 1880. (P. 1880) 42
Notice of new species of Willemæsia group of crustacea (recent Eryon-tidæ.) (P. 1879)
Occurrence of Chelura terebrans (crustacean destructive to timber) on coast of United States. (P. 1879)
Smithson fund, memorial of Regents to Congress relative to the. (R. 1850) 2
Smithson, James—
act of Congress to receive residuary legacy of. (R. 1866) 214, 82
catalogue of library of
character and researches of—W. R. Johnson
contributions to Annals of Philosophy by. (R. 1853) 67, 33
exposition of bequest of—J. HENRY
letter from Richard Rush relative to. (R. 1853)
letters from Fladgate, Clark, and Finch, relative to. (R. 1861) 149, 32
life of, by W. J. RHEES. (R. 1879) 845, 330
list of papers presented to Royal Society by. (R. 1858) 67, 33
notice of, by D. Gilbert, president of the Royal Society. (R. 1858) 67, 33
personal effects of 828, 33
scientific writings of 82
will of. (R. 1858) 67, 328, 83
will of, fac-simile of83
works and character of—J. R. McD. IRBY
Smithson, James, and his bequest-W. J. RHEES. (R. 1879)
Smithsonian-
building, report on use of new hall in—L. Agassiz. (R. 1867) 215
Bulletin No. 1. Vocabulary of the jargon68
Contributions to Knowledge-See Contributions.
Miscellaneous Collections—See Miscellaneous Collections.
publications—See Publications.
publications, catalogue and index of—W. J. RHEES. 478
Reports—See Reports.
Smithsonian Institution—
act to establish—See Congress.
address on the—J. HENRY. (R. 1858) E, 67
business arrangements of the 325
directory of officers, collaborators, and employés of the
establishment and officers of the
Journals of Regents, reports of committees, statistics, etc W. J. RHEES. 329
magnetic observatory at, description of—J. E. HILGARD. (R. 1859) 110

Smithsonian Institution—Continued.
origin and history of-W. J. RHEES
report of American Academy of Arts and Sciences, on the. (R. 1858) 67
report of Senate Judiciary Committee on management of-A. P. BUTLER.
(R. 1855) 77
views and plans of the—R. D. OWEN
Snake Indians, religion, superstition, and manners of—A. G. BRACKETT. (R.
1879)846
SNELL, E. S. Lecture on planetary disturbances. (R. 1855)
Snell's barometric observations, discussion of—F. H. Loud. (R. 1880) 442, 435
Snow—
charts—See Rain.
electricity during fall of-F. Zantedeschi. (R. 1870) 244
gauge—W. E. Guest. (R. 1858)
gauges-R. H. GARDINER. (R. 1858)
line, Norwegian-O. E. DREUTZER. (R. 1866)
observations of 157
tables of precipitation in—C. A. SCHOTT 222, 353
SNYDER, J. F. Deposit of stone implements in Illinois. (R. 1876)
Social and religious condition of the lower races of man-J. LUBBOCK. (R. 1869.) 228
Societies—
list of, in United States and British Provinces-W. J. RHEES 116
local scientific, organization of-J. HENRY. (R. 1875) 298
proceedings of, relative to Joseph Henry 856
publications of, in Smithsonian library, 1854, 1856, 1858, 1866. 73, 85, 117, 179
Societies—See Holland, Belgium, Publications.
Society-
Batavian, of Experimental Philosophy, Rotterdam. Prize questions. (R. 1861)
Bath and West of England, for Encouragement of Agricultural Arts, etc. Exchange system. (R. 1867)
for Encouragement of Science, Dunkirk—See Prize questions.
Society of—
Arts and Sciences, Utrecht. Prize questions. (R. 1861)
Natural History, Portland, Maine, account of-E. C. Bolles. (R. 1867.) 215
Natural Sciences, Cherbourg-See Prize questions.
Physics and Natural History of Geneva-See Geneva.
Science, Agriculture, and Arts of Lille—See Prize questions.
Science, Art, and Literature, Hainaut-See Prize questions.
Sciences, Harlem—See Prize questions.
Sciences, Letters, and Arts, Dunkirk—See Prize questions.
Sciences, Royal Danish—See Prize questions.
Victoria, address of president—R. L. J. ELLERY. (R. 1868) 224
•

SOHNCKE, L. Law of variation of temperature in ascending moist current of air. (R. 1877)
Solar-
diurnal variation in magnetic declination—A. D. BACHE
diurnal variation in vertical component of magnetic force—A. D. BACHE. 175
eclipse of April 25, 1865—Baron DE PRADOS. (R. 1864) 188
eclipse of July 18, 1860—J. LAMONT. (R. 1864) 188
system, harmonies of the—S. ALEXANDER280
Solids, specific heat of—F. W. CLARKE
Solids, tables of expansion of, by heat—F. W. CLARKE
Solitary wasps, synopsis of—H. DE SAUSSURE
SONNTAG, A. Observations on terrestrial magnetism in Mexico
Sonora, plants collected in, by C. Wright—A. GRAY
Sound, refraction of—W. B. TAYLOR. (R. 1875)
Sound, researches in—J. HENRY. (R. 1878) \$41,406
Soundings, microscopical examination of—J. W. BAILEY 20, 23
Sounds, vocal, of Laura Bridgman—F. LIEBER 12
South America—
birds of, not in U. S. National Museum—R. RIDGWAY. (P. 1881) 467
Caracas, meteorology of—G. A. Ernst. (R. 1867) 215
circular relative to collections of birds from168
rain-fall in—C. A. SCHOTT 222, 533
travels in—S. Habel 29
South Carolina—
description of new hake from—T. H. BEAN. (P. 1880)
fishes of—D. S. Jordan; A. W. Brayton
microscopical observations in—J. W. BAILEY
Southern Baptist Missionary Society. Yoruba grammar and dictionary
Southern States, antiquities in the—H. C. WILLIAMS. (R. 1880)
bouting in brains, increased that one will the the transfer of the bouting in
blace and time, results on the relations of D. Missandas. (M. 1991)
SPAINHOUR, J. Antiquities in Lenoir county, North Carolina. (R. 1871) 269
Spanish vocabulary, with comparative words in English, French, and Latin 170
Spanish works on New Mexico, by Buckingham Smith, communications relative to publication of. (R. 1855)
SPARKS, J., and others. On publication of Spanish works on New Mexico. (R. 1855)
SPARES, J., and others. Report of American Academy of Art and Sciences on organization of Smithsonian Institution. (R. 1858)
Sparoid fish, new, (Sparus brachysomus,) from lower California—W. N. Lock- INGTON. (P. 1880)
Sparrows sent to United States—W. A. LLOYD. (R. 1867)215

OF SMITHSONIAN PUBLICATIONS.

Sparus brachysomus, new sparoid fish, from lower California, description of— W. N. Lockington. (P. 1880)	425
Specific gravity—	055
bibliography of—F. W. CLARKE	255
of sea water—D. Walker	146
tables. Constants of Nature, Part I—F. W. CLARKE	255
tables. First supplement to Constants of Nature, Part I—F. W. CLARKE.	288
Specific heat, bibliography of—F. W. CLARKE	276
Specific heat tables. Constants of Nature, Part II—F. W. CLARKE	276
Specimens—	
acknowledged by British Museum. (R. 1865)	209
acknowledged by Chicago Academy of Sciences. (R. 1867)	215
authority to collectors of customs to receive and transmit to Smithsonian	
Institution	84
authority to naval officers to receive and transmit to Smithsonian Insti-	0.4
tution	84
authority to quartermasters of War Department to receive and transmit	
to Smithsonian Institution	84
derived from expeditions—S. F. BAIRD. (R. 1867)	215
exchange of—Hamburg Zoölogical Gardens. (R. 1867)	215
exchange of—W. A. LLOYD. (R. 1867)	215
exchange of—University of Costa Rica. (R. 1867)	215
exchange of—University of Greece. (R. 1867)	215
from Brazil, presented—M. M. LISBOA. (R. 1865)	209
large myological, rapid preparation of—F. PLATEAU. (P. 1881)	467
of flamingo from south Florida, presented—G. WURDEMAN. (R. 1860.)	147
of natural history, directions for collecting, preserving, and transport- ing-S. F. Baird. (B. 1856)91	24
presented, circular sent with—J. HENRY. (R. 1872)	271
	442
receipts and distribution of. (R. 1880)	
rules for examination of. (R. 1880)	442
Spectrum analysis applied to the heavenly bodies—W. Huggins. (R. 1866)	214
Spencer's telescope, examination of, for Hamilton College. (R. 1855)	77
Spheres, tangencies of—B. ALVORD	80
Spleen, observations on—J. JONES	82
Sponge, a new type of, (Camaraphysema)—J. A. RYDEB. (P. 1880)	425
Spruce Creek valley, Pennsylvania, account of tornado in—J. B. MEEK. (R. 1871)	249
Squalius alicise from Utah Lake—P. L. Joux. (P. 1881)	467
SQUIER, E. G. Aboriginal monuments of State of New York	15
SQUIER, E. G. Antiquities of Nicaragua. (R. 1850)	28
SQUIER, E. G., and others. On publication of Spanish works on New Mexico. (R. 1855)	77

SQUIER, E. G.; DAVIS, C. H. Ancient monuments of the Mississippi valley.
Squier, E. G.; Davis, C. H., correspondence relative to memoir by. (R. 1847.) H, K
Stanley's gallery of Indian portraits, report of Committee of Regents on the. (R. 1857)
STANLEY, J. M. Catalogue of portraits of North American Indians 53
STANTON, E. M. Deposit of Beaufort Library. (R. 1862) 150
Star clusters, bibliography of—E. S. HOLDEN
Stars—
index-catalogue of—E. S. Holden
movement of, around a central point—J. H. MAEDLER. (B. 1859) 110
near the North Pole, map of, for observations on auroras 350
scintillation of the—C. DUFOUE; KÄMTZ. (R. 1861)149
State Department. Circular to diplomatic agents relative to Morgan's research. 138
Statement and exposition of harmonies of the solar system—S. ALEXANDER 280
Stations, meteorological, of Smithsonian Institution (See, also, List) 373
Statistics of—
British Museum—S. F. BAIRD. (R. 1850)
Norwegian mountains, lakes, and snow line—O. E. DREUTZER. (R. 1866.) 214
Smithsonian Institution, finances, exchanges, etc., 1846-1877 329
Statue of Joseph Henry, proceedings of Congress relative to 356
Statues, on plaster casts of—W. J. STONE. (R. 1855)
STEARNS, J. Translation of Babinet on diamonds and precious stones. (B. 1870)
Stearns, S., catalogue of fishes from Pensacola, collected by—G. B. Goode; T. H. BEAN. (P. 1879)
Stearns, S., description of new species of amber fish (Seriola Stearnsii) obtained
near Pensacola, by-G. B. Goode; T. H. Bran. (P. 1879) 333
STEARNS, S. Note on Gulf menhaden, Brevoortia patronus. (P. 1878) 332
STEBBINS, R. P. Memoir of Louis Agassiz. (R. 1878) 275
STEINDACHNER, F. Note on Perca flavescens. (P. 1878)
STEJNEGER, L. Description of two new races of Myadestes obscurus. (P. 1881.) 467
STELLE, J. P. Account of aboriginal ruins in Hardin Co., Tennessee. (R. 1870.) 244
STELLE, J. P. Account of aboriginal ruins at Savannah, Tennessee. (R. 1870.) 244
STEPHENSON, M. F. Account of ancient mounds in Georgia. (R. 1870) 244
STEPHENSON, M. F. Mounds in Bartow county, Georgia. (R. 1872) 271
Stereotype catalogue of public libraries, report of Commissioners on. (R. 1850.) 28,47
Stereotyped titles for library catalogues—C. C. Jewerr 47
STEVENS, H. Prospectus of a bibliographia Americana. (R. 1848) I
STEVENS, J. A. Memoir of George Gibbs. (R. 1873)
STEVENSON, J. Report of explorations in New Mexico and Arizona. (R. 1880.) 442
STEWART, B. Experiments on aneroid barometers at Kew Observatory. (R. 1868.) 224
Stichseus punctatus, occurrence of, at St. Michael's, Alaska—T. H. BEAN. (P. 1878)

Stimpson, W.—	
Check-list of the shells of North America	128
On collection and preservation of marine invertebrates	84
Researches upon the Hydrobiinæ and allied forms	201
Synopsis of the marine invertebrata of Grand Manan	50
Stock-in-trade of an aboriginal lapidary-C. RAU. (R. 1877) 823, 440,	402
Stockholm Academy of Sciences. Acknowledgment of birds. (R. 1867)	215
STOCKWELL, J. N. Secular variations of elements of orbits of the eight principal planets. (R. 1871)	232
STONE, W. J. On plaster casts of statues. (R. 1855)	77
Stone—	
celts in the West Indies and Africa—G. J. GIBBS. (R. 1877)	828
cists near Highland, Madison county, Illinois—A. Orhler. (R. 1879.)	845
drilling in, without metal-C. RAU. (R. 1868) 224, 440,	372
image in Tennessee, account of discovery of—E. M. GRANT. (R. 1870.)	244
implements, North American—C. RAU. (R. 1872) 271, 440,	382
Stone age—	
and the troglodytes of Breckinridge county, Kentucky—R. S. Robert- son. (R. 1874)	286
fauna of middle Europe during the-L. RÜTIMEYER. (R. 1861)	149
in New Jersey—C. C. Abbott. (R. 1875) 298,	394
Stone period, North American, agricultural implements of the—C. RAU. (R. 1868) 187.	, 44 0
Stones, precious—J. Babinet. (R. 1870) 244	, 877
Storm in Butler county, Kansas, June 28, 1871—W. HARRISON. (R. 1871)	249
Storms—	
in Europe and America, December, 1836—E. Loomis	127
magnetic—E. Sabine. (R. 1860)	147
of 1859—J. H. Coffin	182
thunder, instructions for observations of—J. HENRY	285
Strain and over-action of the heart. Toner lecture No. 111-J. M. Da Costa.	279
Strait of Juan de Fuca, Indians of-J. G. SWAN	220
Strait of Juan de Fuca, description of new species of Paralepis (Paralepis coruscans) from—D. S. Jordan; C. H. Gilbert. (P. 1880)	425
STRANG, J. J. Natural History of Beaver Island, Michigan. (R. 1854)	75
STREETS, T. N. Contributions to natural history of Hawaiian Islands, Fanning	
Islands, and lower California	808
Strepomatidæ (American melanians)—G. W. TRYON	253
Stricklandia Davidsoni, note on the occurrence of, in Georgia—C. A. WHITE. (P. 1880)	425
Stricklandia Salteri, note on the occurrence of, in Georgia—C. A. WHITE. (P. 1880)	425
STRONG, M. Antiquities in Wisconsin. (R. 1877)	828

STRONG, M. Observations on the prehistoric mounds of Grant county, Wisconsin. (R. 1876)
Structure, internal, of the earth—J. G. BARNARD 310
Structure of cancerous tumors. Toner lecture No. 1—J. J. WOODWARD 266
Structures, aboriginal, in Georgia—C. C. Jones, Jr. (R. 1877) 323, 400
Study of—
chionis minor—J. H. KIDDER; E. COUES
high antiquity in Europe, lecture on-A. Morlot. (R. 1862; R. 1864.) 150, 188
jade—S. Blondel. (R. 1876) 299
mortuary customs of North American Indians—H. C. YARROW. (E. 1879-80)
nature and mechanism of fever. Toner lecture No. IV-H. C. WOOD 282
phenomena of contact—T. L. Phirson. (R. 1862) 150
physiology of fever—H. C. Wood
savage weapons at the Centennial—E. H. KNIGHT. (R. 1879) 345, 415
skulls and long bones from mounds in Illinois—R. J. FARQUHARSON. (R. 1874) 286
trunk fishes, Ostraciontida-G. B. Goode. (P. 1879) 333
Studies in Central American picture writing—E. S. HOLDEN. (E. 1879-80) 476
Subcutaneous surgery. Toner lecture No. VI-W. ADAMS
Sudis ringens, new species of scopeloid fish from Santa Barbara channel, Cali-
fornia, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880) 425
Suess, E. Boundary line between geology and history. (R. 1872) 271
Suggestions—
for ethnological researches in Russian America—G. GIBBS:207
for forming collections of birds' eggs—A. NEWTON139
for meteorological observations in Russian America—J. Henry
for sanitary drainage of Washington city—G. E. WARING 349
for saving parts of the skeletons of birds—A. Newton. (R. 1860) 147
relative to ethnological map of North America—L. H. Morgan. (B. 1861.) 149
relative to objects of scientific investigations in Russian America—J. HENRY: S. F. BAIRD
to beginners in botany—L. F. WARD
Sulphurous acid, action of, upon putrefactive bacteria. (P. 1881)
Sumichrast, F., birds of Mexico, collected by—G. N. LAWRENCE
(R. 1879)
Sun—
chemical analysis of the—A. LAUGEL. (R. 1861)
disk of the, external appearance of. (R. 1866)
eclipse of the, September 7, 1858—J. M. GILLISS
eclipse of the, March 15, 1868—T. HILL
eclipse of the, April 25, 1865—M. DE PRADOS. (R. 1864)
relative intensity of heat and light of the-L. W. MERCH. (R. 1856.) 91, 83

OF SMITHSONIAN PUBLICATIONS.	309
SUNDERLAND, B. Prayer at memorial of Joseph Henry	. 356
Sun's distance, means available for correcting measures of the—G. B. AIRY (R. 1859)	
Superstitions of Shoshone Indians—A. G. Brackett (R. 1879)	
Supplement, asteroid, to new tables of planetary motion—J. D. RUNKLE	
Surgery, subcutaneous. Toner lecture No. vi—W. Adams	
Surgical complications and sequels of continued fevers. Toner lecture No. v—	
W. W. KEEN	
Survey—	
in Michigan, catalogue of rocks, minerals, and ores collected in the—O. T. JACKSON. (R. 1854)	75
magnetic, of Pennsylvania and parts of adjacent States-A. D. BACHE.	166
of economic geology of Trinidad—G. P. Wall; J. G. Sawkins. (R. 1856)	
of State of New York—E. G. Squies.	
Surveying expedition, North Pacific, natural history of—T. H. STREETS	
Surveys, Government—See Explorations.	
SWAN, J. G.—	
Eulachon, or candlefish of northwest coast. (P. 1880)	425
Haidah Indians of Queen Charlotte's Islands, British Columbia	
Makah Indians of Cape Flattery, Strait of Fuca	
Makah vocabulary	
Surf smelt of the northwest coast, and mode of taking them by the Quillehute Indians of west coast of Washington Territory. (P. 1880.)	1
Sweden, Stockholm, Academy of Sciences. Acknowledgment of birds. (R. 1867.)	
Switzerland—	
Concise, archæological researches made at—F. TROYON. (R. 1861)	149
Crania helvetica—F. TROYON. (R. 1864)	188
lacustrian cities of-F. Troyon. (R. 1861)	
lacustrian settlements in, abstract of Dr. Keller's report on—A. Morlot. (R. 1868)	187
Lake of Neuchâtel, palafittes or lacustrian constructions of-E. DESOR.	
(R. 1865)	360
Lausanne, Cantonal Museum at, antiquarian and ethnological collections	
of the—F. Troyon. (R. 1861)	149
lecture on—A. D. BACHE. (R. 1870)	244
Syllabus of a course of lectures on physics—J. Henry. (R. 1856)	91
Symbols for charts of prehistoric archaeology, international code of—G. DE	
MORTILLET; E. CHANTRE. (R. 1875)	
Symbols used by James Smithson	827
Synopsis of— American Rhinobatida—S. GARMAN. (P. 1880)	425
AMOITOM ANDROVONION—D. GARMAN. (1. 1000)	
American wasps, solitary wasps—H. DE SAUSSURE	254

•

ALPHABETICAL INDEX

Synopsis of—Continued.
marine invertebrata of Grand Manan—W. STIMPSON 50
North American lepidoptera—J. G. Morris 133
North American neuroptera—H. HAGEN; P. R. UHLER 124
pediculate fishes of east coast of extra tropical North America—T. GILL. (P. 1878)
scientific writings of Sir William Herschel—E. S. Holden; C. S. Has- Tings. (R. 1880)42,426
siluride of fresh waters of North America—D. S. JORDAN 200
trochilids-D. G. Elliott
useful and injurious animals—G. B. GOODE
Synoptical tables of characters of mammals—T. GILL 230
Synthetic experiments relative to meteorites—G. A. DAUBRÉE. (R. 1868) 24
Syria, Beirut, sarcophagus from-A. A. HARWOOD. (R. 1870) 244
System—
meteorological, of Smithsonian Institution—E. FOREMAN. (R. 1851; R. 1852)
of accounts adopted by Board of Regents. (R. 1866)
of combined meteorological observations, report on, by committee of American Association for Advancement of Science. (R. 1851)
of numeration, improved, report on—W. B. TAYLOR. (R. 1867) 215
of weights and measures, new—G. H. KNIGHT. (R. 1867) 215
solar, harmonies of the—S. ALEXANDER 280
Systematic—
index of Smithsonian publications. (B. 1868) 224, 478
index to list of foreign correspondents 257
list of batrachia and reptilia—E. D. COPE 292
review of classification of birds-W. LILLJEBORG. (B. 1865) 209, 864
Systems of consanguinity and affinity of the human family—L. H. MORGAN 218
Systems of relationship of Cree Indians—E. A. WATKINS. (R. 1862)

T.

Table of—
anthropological measurements—SCHERZER; SCHWARZ. (R. 1866) 214
chemical equivalents of sixty-three elements. (R. 1864) 188
distribution of birds—G. N. LAWRENCE. (P. 1878) 332
foreign gold and silver coins. (R. 1868) 224
Tables—
asteroid—J. D. Runkle
barometrical—A. Guyor 158
for conversion of centigrade degrees to Fahrenheit's scale. (R. 1863) 187
for determining values of coefficients in perturbative function of planetary motion which depend on ratio of mean distances—J. D. Runkle 79
hygrometrical—A. Guyor 158
hypsometrical—A. Guyor 158
meteorological—A. Guyor
metric—H. A. NEWTON. (R. 1865) 209, 371
miscellaneous—A. Guyot158
mortality, methods of interpolation applied to. Parts I, II—E. L. DE FOREST. (R. 1871; R. 1878)249, 275
of atmospheric temperature—C. A. Schott 277
of constants of nature and art-C. BABBAGE. (R. 1856)
of expansion by heat. Constants of Nature, Part III-F. W. CLARKE. 289
of meteorological corrections—A. GUYOT
of motion of Neptune—8. Newcomb
of motion of Uranus—S. Newcomb
of precipitation in rain and snow—C. A. Schott 222, 353
of variations of elements of orbits of the eight principal planets—J. N. STOCKWELL
of weights and measures, English and French. (R. 1863-1865.) 187, 188, 209
of winds of the globe—J. H. COPPIN; S. J. COPPIN 268
physical—A. Guvor
psychrometrical—J. H. Coffin
specific gravity. Constants of Nature, Part I, and supplement—F. W.
specific heat. Constants of Nature, Part II-F. W. CLARKE 276
thermometrical—A. Guyor 153
Tablet, Palenque—C. RAU
TACCHINI, P. Evaporation observed at Palermo in 1865 and 1866. (R. 1870.) 244
Tampa Bay, Florida, shell-heaps of—S. T. WALKER. (R. 1879) 845
Tangencies of circles and of spheres—B. ALVORD
Taste, sense of. (R. 1866)

Taxidermy, classification of collections of U.S. National Museum to illustrate
art of-W. T. HORNADAY. (P. 1881) 467, 456
TAYLOR, A. S. Grasshoppers and locusts of America. (R. 1858) 109
TAYLOR, F. W. Report of chemist of Smithsonian Institution. (R. 1880) 442
TAYLOR, W. B.—
Henry and the telegraph. (R. 1878) 841, 405
Kinetic theories of gravitation. (R. 1876) 299, 395
Memoir on the scientific work of Joseph Henry 856, 339
Refraction of sound. (R. 1875) 298
Report on improved system of numeration. (R. 1867) 215
Thoughts on the nature and origin of force. (R. 1870) 244, 375
TAYLOR, W. M. Ancient mound in western Pennsylvania. (R. 1877) 323
Telegrams, astronomical, circular relative to-J. Henry
Telegrams, meteorological, by Western Union Telegraph Company—H. Sibler. (R. 1862)
Telegrams read at memorial of Joseph Henry 356
Telegraph—
American fire-alarm, lecture on the-W. F. CHANNING. (R. 1854) 75
electro-magnetic, deposition of J. Henry in relation to. (R. 1857) 107, 115
electro-magnetic, Henry's contribution to-W. B. TAYLOR. (R. 1878.) 341, 405
electro-magnetic, Morse's invention-W. B. TAYLOR. (R. 1878) 341, 405
electro-magnetic, proceedings of Board of Regents in relation to. (R. 1857)
Henry and the-W. B. TAYLOR. (R. 1878) 405
history of the—J. Henry. (R. 1857)
influence of aurors on the-W. D. SARGENT. (R. 1870) 244
lines, phenomena in, during auroras—G. B. Donati. (R. 1872) 271
report of Committee of Regents on the. (R. 1857) 107, 115, \$29
statement of S. P. Chase relative to the. (R. 1857) 107, 115, 329
statement of L. D. Gale relative to the. (R. 1857) 107, 115, 329
statement of James Hall relative to the. (R. 1857) 107, 115, 329
statement of J. Henry relative to history of. (R. 1857) 107, 115, 329
statement of Charles Mason relative to the. (R. 1857) 107, 115, 329
Telegraphic announcements of astronomical discoveries—J. Henry 263
Telescope—
Henry Draper's, account of—T. W. WEBB. (R. 1864)
silvered-glass, construction of, and its use in celestial photography—H.
DRAPER180
Spencer's, examination of, for Hamilton College, Clinton, N.Y. (R. 1855.)
Temperature—
chart of United States for the year-O. A. SCHOTT 381, 388
charts of United States, summer, winter, and year—C. A. Schott 387

Temperature—Continued.	
in ascending currents of air, laws of variation of—J. HANN. (R. 1877.) 828,	898
in ascending moist currents of air, laws of variation of-L. Sohneke.	
(B. 1877)	, 898
mean, best hours to find—C. DEWEY. (R. 1857; R. 1860) 107;	, 147
observations in the Arctic Seas—I. I. HAYES	196
observations in the Arctic Seas-E. K. KANE	104
observations in the Arctic Seas—F. L. McCLINTOCK	146
of St. Louis, Missouri-A. Fendler. (R. 1860)	147
tables for United States and adjacent parts of America—C. A. SCHOTT	277
underground-C. A. Schott; J. D. Everett. (R. 1874)	286
Temperature—See Kane, Hayes, Meteorology, etc.	
Tempests and tornadoes, distinction between-J. B. LAMARCK. (R. 1871)	249
Tennessee-	
aboriginal remains of—J. Jones	259
Alleghany region of, fishes of—D. S. JORDAN; A. W. BRAYTON.	808
antiquities in—I. DILLE. (R. 1862)	150
antiquities in—E. O. DUNNING. (R. 1870)	244
antiquities of—W. M. CLARK. (R. 1877)	828
antiquities of—J. Jones	259
antiquities of—D. F. WRIGHT. (R. 1874)	286
Blount county, antiquities of—A. E. LAW. (R. 1874)	286
Chattanooga, ancient mound near-M. C. READ. (R. 1867)	215
East, mound in—A. F. DANILSEN. (R. 1868)	187
explorations in—E. A. DAYTON. (R. 1870)	244
Hardin county, aboriginal ruins in—J. P. STELLE. (R. 1870)	244
herbarium captured in-H. R. WIRTZ. (R. 1862)	150
Jackson county, antiquities of—J. HAILE; J. W. McHENRY. (R. 1874.)	286
Nashville, antiquities of—R. S. Robertson. (R. 1877)	828
Savannah, aboriginal ruins at-J. P. STELLE. (R. 1870)	244
stone image in, discovery of—E. M. GRANT. (R. 1870)	244
Terrestrial magnetism— observations on—W. HARKNESS	289
observations on-J. Locke	35
observations on—Baron von Müllek; A. Sonntag	114
Terrestrial magnetism—See Bache, A. D.	
Terrestrial physics, articles on. (R. 1870)	244
Tertiary fossils, presented—Imperial Geological Institute, Vienna. (R.	-22
1863)	187
Testing building materials, on mode of—J. HENRY. (R. 1856)	91
Testing results of perspiration and respiration, apparatus for—M. Pettenkofer.	
(R 1864)	188

Texas—	
hail storm in—G. M. BACHE. (R. 1870)	244
new cretaceous invertebrate fossils from, descriptions of—C. A. WHITE. (P. 1879)	833
ornithology of, notes on-J. C. MERRILL. (P. 1878)	832
plants of, collected by C. Wright. Parts I, II-A. GRAY 2	
zoölogical position of—E. D. Cope	
Thalassophila—W. G. BINNEY	
Thénard, L. J., memoir of, by M. FLOURENS. (B. 1862)	150
Theoretical researches—See Plateau, J.	
Theories, kinetic, of gravitation—W. B. TAYLOB. (B. 1878)	3 95
Theory—	
mechanical, of heat, principles of-J. MÜLLER. (R. 1868)	224
modern, of chemical types—C. M. WETHERILL. (R. 1863)	187
of heat, recent progress in relation to—A. CAZIN. (R. 1868)	224
Thermometer scales, table for conversion of centigrade to Fahrenheit. (R. 1868.)	187
Thermometer, wet and dry bulb, tables for—J. H. COFFIN	87
Thermometric observations	157
Thermometrical tables—A. Guvor	153
THOMPSON, T. Mounds in Iowa and Illinois. (R. 1879)	345
Thoughts on the nature and origin of force—W. B. TAYLOR. (R. 1870) 244,	375
Thrushes, description of two new, from United States—R. RIDGWAY. (P. 1881.)	467
Thunder—	
and lightning, observations on—S. MASTERMAN. (R. 1855)	77
storm, on John Wise's observation of a-R. HARE. (R. 1854)	75
storms, instructions for observations of-J. HENRY	23 5
Tidal observations—	
at Wolstenholm Sound-Com. SAUNDERS	130
in the Arctic Seas—I. I. HAYES	196
in the Arctic Seas—E. K. KANE	130
Tide, flood, law of deposit of the-C. H. DAVIS	33
Tides and tidal action in harbors-J. E. HILGARD. (R. 1874) 286,	39 0
TIFFANY, A. S. Shell-bed skull from Illinois. (R. 1874)	236
Timber of submarine structures, occurrence of Chelura terebrans, a crustacean	
destructive to—S. I. SMITH. (P. 1879)	333
Time and space, lecture on relations of—S. ALEXANDER. (R. 1861)	149
Tinneh or Chepewyan Indians of British America—G. GIBBS; W. L. HARD-	
ISTY; S. JONES; B. R. ROSS. (R. 1866)	
Tipulidæ, monograph of—R. Osten Sacken	
TOELLNER, A. Antiquities of Rock Island county, Illinois. (R. 1879)	345
Tokio, University of, catalogue of Japanese woods presented by—L. F. WARD. (P. 1881)	467

Tompkins county, New York, ancient fort and burial ground in—D. Thow- BRIDGE. (R. 1868)	187
TONER, J. M. Deed of foundation of Toner lectures. (R. 1872) 271	
Toner lectures—	•
No. I. On the structure of cancerous tumors and the mode in which adjacent parts are invaded—J. J. WOODWARD	266
II. Dual character of the brain—C. E. BROWN-SÉQUARD	291
III. On strain and over-action of the heart-J. M. DA COSTA	279
IV. A study of the nature and mechanism of fever-H. C. Wood	282
w. On the surgical complications and sequels of the continued fevers, with a bibliography of works on diseases of the joints, bones, larynx, the eye, gangrene, hæmatoma, phlegmasia, etc.—W. W. Keen	300
VI. Subcutaneous surgery, its principles, and its recent extension in practice—W. ADAMS	302
VII. The nature of reparatory inflammation in arteries after ligature, acupressure, and torsion—E. O. Shakespeare	321
WIII. Suggestions for the sanitary drainage of Washington city—G. E. Washing, Jr.	349
Tonto Apaches, notes on the—C. SMART. (R. 1867)	215
Topography of Black Mountain, North Carolina—T. L. CLINGMAN. (R. 1855.)	77
Tornado in Spruce creek valley, Centre county, Pennsylvania—J. B. MEEK. (B. 1871)	249
Tornado near New Harmony, Indiana, April 80, 1852—J. CHAPPELSMITH	59
Tornadoes and tempests, distinction between—J. B. LAMARCK. (R. 1871)	249
Tornadoes, queries relative to—J. HENRY	190
Toronto—	
Observatory, instructions by, for observations of aurora	148
Observatory, map of stars near North Pole used by	850
University. Objects of the museum. (R. 1865)	209
University. Scientific cooperation. (R. 1861)	149
Torrey, John, memoir of, by A. GRAY. (R. 1878)	275
Tobrey, J.—	
Observations on the Batis maritima	60
On the Darlingtonia Californica, a new pitcher plant from northern California	61
Plants Frémontians, or descriptions of plants collected by J. C. Frémont in California	46
Fortoises, North American land, of genus Xerobates—F. W. TRUE. (P. 1881.)	467
Fortola, great hurricane at—G. A. LATIMER. (R. 1867)	215
Fortugas, destruction of fish in vicinity of the-J. P. JEFFERSON; J. Y. Por-	
TER; T. Moore. (P. 1878)	882
Totten, Gen. J. G., eulogy on, by J. G. BARNARD. (R. 1865)	209
Traces of early mental condition of man—E. B. TYLOB. (R. 1867)	215

Trade-

ancient aboriginal—C. RAU. (R. 1872) 271, 440	, 38 5
language of Oregon, dictionary of the-G. GIBBS	161
language of Oregon, vocabulary of the—B. R. MITCHELL; W. W. TURNER	
Transactions—See Geneva Society, Paris Anthropological Society.	
Transatlantic longitude—B. A. Gould	223
Transfer of Smithsonian library to Library of Congress, act of Congress to authorize. (R. 1865) 209, 32	
Transformations, lecture on insect instincts and—J. G. Morris. (R. 1855)	π
Transit of Venus expedition to Kerguelen Island.—J. H. KIDDER and others. 29	3, 294
Transporting specimens of Diatomaceae, directions for-A. M. EDWARDS	
Transporting specimens of natural history, directions for—S. F. BAIRD. (R. 1856)	
Trapezium, gray substance of the—J. DEAN	178
Travels in Central and South America—S. HABEL	269
Treasury Department, authority given by, to collectors of customs and others to receive and transmit specimens to the Smithsonian	
Treasury, Secretary of, letter to, on payment of interest in coin—J. HEBEY. (R. 1865)	, 329
Treasury of United States, account of, with Smithson fund 326	, 329
Trees—	
forest, distribution of, in Montana, Idaho, and Washington—W. W. Johnson. (R. 1870)	244
of Florida and the Mexican boundary—J. G. Cooper. (R. 1860)	147
of North America, distribution of-J. G. Cooper. (R. 1858) 109,	351
Trinidad, economic geology of—G. P. Wall; J. G. Sawkins. (R. 1856)	91
Trochilida—	-18
classification and synopsis of—D. G. Elliot	317
in collection of U. S. National Museum, catalogue of—R. RIDGWAY. (P. 1880)	425
list of—D. G. Elliot	
Troglodytes of Breckinridge county, Kentucky—R. S. Robertson. (R. 1874.)	
Troglodytes or cave dwellers of the valley of the Vézère-P. Broca. (R. 1872)	
TROWBRIDGE, D. Ancient fort and burial ground in Tompkins county, New York. (R. 1868)	
Troyon, F.—	
Archeological researches made at Concise. (R. 1861)	
Lacustrian cities of Switzerland. (R. 1861)	
On the Crania helvetica. (R. 1864)	186
Report on ethnological collection of Museum at Lausanne, Switzerland. (R. 1861)	149
TRUE, F. W. On North American land tortoises of genus Xerobates. (P. 1881.)	467
TRUE, F. W. On the rare rodent, Cricetodipus parvus. (P. 1881)	4 6î

OF SMITHSONIAN PUBLICATIONS.	317
Truncatellidæ—W. G. BINNEY	144
Trust fund for the promotion of science in the United States-J. TYNDALL.	
(R. 1872)	
Tryon, G. W., letter to, on shells—J. LEWIS	
Tryon, G. W. Report on shells presented to Philadelphia Academy of Natural Sciences. (R. 1865)	209
TRYON, G. W. Strepomatide or American melanians	253
Trypetina, review of-H. Lorw	256
Tuckerman, E. Lichens of Arctic America	842
TUCKERMAN, E. Lichens of Kerguelen Island	294
Tuckerr, F. F. On barometer tables. (R. 1867)	218
Tucson meteorite, account of -S. AINSA. (R. 1868)	187
Tucson meteorite, account of—B. J. D. IRWIN. (R. 1868)	187
Tumors, cancerous, structure of—J. J. WOODWARD	266
Tunicata of northeast coast of America—A. E. VERRILL. (P. 1879)	838
Turdidæ, description of new species of, from Dominica—G. N. LAWRENCE. (P. 1880)	
Turner, W. W., notice of, by C. C. Felton. (R. 1859)	110
TURNER, W. W. Indian philology. (R. 1851)	51
TURNER, W. W., and others. On publication of Squier and Davis' Ancient	
Monuments of the Mississippi valley. (R. 1847)	
TURNER, W. W.; BOWEN, T. J. Yoruba grammar and dictionary	98
TURNER, W. W.; MITCHELL, B. R. Vocabulary of the jargon of Oregon	68
TYLOR, E. B. On traces of the early mental condition of man. (R. 1867)	215
TYNDALL, J. Radiation. (R. 1868)	
TYNDALL, J. Trust fund for the promotion of science in the United States. (R. 1872) 27	
Tyrannus, synopsis of genus—R. Ridgway. (P. 1878)	

•

.

,

•

U.

Uhler, P. R.—	
. Circular in reference to the history of North American grasshoppers	168
Instructions for collecting hemipters. (R. 1858)	109
Instructions for collecting neuroptera. (B. 1858)	109
Instructions for collecting orthopters. (R. 1858)	109
Synopsis of neuropters of North America	134
Underground temperature—C. A. Schott; J. D. EVERETT. (R. 1874)	286
Union, lectures on the-H. REED. (R. 1854)	75
United States—	
animal resources and fisheries of the, catalogue of collection to illustrate the—G. B. Goode	326
animal resources of the, catalogue of collection to illustrate the—G. B. Goode	297
antiquities of the, proposed map of—A. J. Hill. (R. 1861)archeology of the—S. F. HAVEN	
base chart of the—C. A. SCHOTT	
chalk found in the—T. A. CONRAD. (R. 1865)	
coleoptera of the, catalogue of the—F. E. MELSHEIMER	
collections presented to the, by foreign commissioners, list of. (R. 1876.)	
education in the, project of outline history of—F. A. PACKARD. (R. 1863.)	187
ethnological map of the, proposed—G. GIBBS. (R. 1862)	150
exhibit of fisheries and fish culture of the, at Berlin, 1880—G. B. Goods. explorations in the—See Baird, S. F.	413
fishes, food, of the—See Baird, S. F.	
fishes of Pacific coast of the-D. S. JORDAN; C. H. GILBERT. (P. 1880.)	425
fishes of Pacific coast of the, bibliography of-T. H. BEAN. (P. 1881.)	467
fishes of Pacific coast of the, bibliography of-T. GILL	463
fishes of Pacific coast of the, notes on—D. S. JORDAN; C. H. GILBERT. (P. 1881)	
free freight between England and the—E. CUNARD. (R. 1859)	110
free freight between Germany and the-Kunhardt & Co. (R. 1861)	149
free freight between Germany and the, by North German Lloyd-R. Schleiden. (R. 1858)	109
fund for promotion of science in the-J. TYNDALL. (R. 1872)	271
institutions, scientific and literary, in the, list of the principal	335
libraries, institutions, and societies in the, list of—W. J. RHEES 116,	238
libraries, public, in the, notices of—C. C. JEWETT	25
libraries, public, of the, report on—C. C. JEWETT. (R. 1849)	21
limpets and chitons from deep waters off eastern coast of the—W. H.	467

United States—Continued.
meteorology of the, report on—E. LOOMIS. (R. 1847)
microscopic examinations of soundings off Atlantic coast of the—J. W. BAILEY
mollusca of the southern, some new species of eocene—A. Heilpein. (P. 1880)425
occultations visible in the—See Downes, J.
physical geography of the—C. Ellet, Jr18
rain and snow fall in the, tables of—C. A. Schott 222, 353
rain charts of the, 1, 2, 8: summer, winter, year—C. A. Schott 374
reptiles, cretaceous, of the-J. LEIDY 192
reptiles, cretaceous, of the, review of—J. LEIDY. (R. 1864)
shell-deposits, artificial, of the-D. G. BRINTON. (R. 1866) 214
sparrows sent to the—W. A. LLOYD. (R. 1867)
temperature chart of the, for the year—C. A. SCHOTT 381, 388
temperature charts of the, 1, 2, 8: summer, winter, year—C. A. Schott. 387
temperature tables of the—C. A. SCHOTT 277
thrushes from the, description of two new-R. RIDGWAY. (P. 1881) 467
trees of the, catalogue of—J. G. Cooper. (R. 1858) 109, 851
United States Exploring Expedition, duplicate shells collected by the 193
United States Fish Commission—See Fish Commission.
United States Government, list of publications of, 1868-1881—G. H. BOEHMER. 477
United States National Museum—See Museum, U. S. National.
United States Patent Office, results of meteorological observations under the
direction of the. Vols. I, II
United States Treasury—See Treasury.
Universal meteorograph for detached observatories—E. H. Von BAUMHAUER. (R. 1879) 345
University—
National, of Greece, Museum of. Exchange of specimens. (R. 1867.) 215
of Christiania, Norway, ethnological specimens presented by. (R. 1868.) 187
of Costa Rica. Exchange of specimens. (R. 1867) 215
of State of New York, Regents of the. Acknowledgment for specimens. (R. 1865) 209
of Tokio, catalogue of Japanese woods presented by—L. F. WARD. (P. 1881)467
of Toronto, objects of museum of. (R. 1865) 209
of Toronto, scientific cooperation. (R. 1861) 149
Uranidea marginata, new species of fish, description of-T. H. BEAN. (P. 1881.) 467
Uranidea microstoma, new fish from Alaska, description of—W. N. Locking- TON. (P. 1880)
Uranus, investigation of orbit of, with tables of its motion—S. NEWCOMB 262
Uranus, secular variation of elements of orbit of-J. N. STOCKWELL 282

Urine, observations on the—J. JONES	82
Use of galvanometer as a measuring instrument—J. C. POGGENDORFF. (R. 1859.)	110
Uses of brain and marrow of animals among Indians—T. R. Peale. (R. 1870.)	244
Utah, descriptions of new invertebrate fossils from-C. A. WHITE. (P. 1880.)	425
Utah lake, description of new species of Squalius (Squalius aliciæ) from—P. L. Joux. (P. 1881)	467
Utah lake, notes on collection of fishes from—D. S. JORDAN; C. H. GILBERT. (P. 1880)	425
Utrecht, Provincial Society of Arts and Sciences. Prize questions. (B. 1861; R. 1862)	, 150

∇_{\cdot}

VAILLANT, M.; HENRY, J. Horary variations of the barometer. (R. 1866) 214
Values of $b_s^{(i)}$ and its derivatives—J. D. Runkle
Values of coefficients in perturbative function of planetary motion depending on ratio of mean distances—J. D. Runkle
Valvatide-W. G. BINNEY 14
Van Rensselaer harbor, Arctic Seas, observations at—See Kane.
Vapor, aqueous, on the diminution of, with increasing altitude in the atmosphere—J. Hann. (R. 1877)
Vapor, aqueous, table for determining elastic force of-J. H. Corrin 87
Vastness of the visible creation, lectures on—S. ALEXANDER. (R. 1857) 107
Vegetable colonization of Shetland Islands, Faroe Islands, and Iceland—C. MARTINS. (R. 1858)
Vegetable species, causes which limit, towards the north—A. DE CANDOLLE. (R. 1858)
Vegetation and the atmosphere—J. Jamin. (R. 1864)188
Velie, J. W., catalogue and description of fishes collected by, in Gulf of Mexico-G. B. GOODE; T. H. BEAN. (P. 1879)
Velocity of light, essay on the—C. DELAUNAY. (R. 1864)
Velocity of the wind—J. HANN and others. (R. 1877)
Venezuela, Colonia Tovar, meteorology and ethnology of—A. FENDLEE. (R. 1857; R. 1866)
Venom of rattlesnake, researches on the—S. W. MITCHELL
Venom of serpents, bibliography of—S. W. MITCHELL
Ventilating and warming buildings. Parts 1, 11—A. Morin. (R. 1873; R. 1874)
Ventilation, architecture in relation to-D. B. REID. (R. 1856)
Venus, secular variations of elements of orbit of—J. N. STOCKWELL 22

o	റ	4
n	Z	

OF SMITHSONIAN PUBLICATIONS.

·
Venus—See Transit of Venus.
Vera Cruz, Mexico, antiquities in-H. Finck. (R. 1870) 244
Verification of barometers at the Kew Observatory—J. WELSH. (R. 1859) 110
Verrill, A. E.—
Annelids, echinoderms, and anthozoa of Kerguelen Island 294
Annelids, molluscoids, and radiates of Arctic America 842
Marine invertebrata of New England. (P. 1879) 888
Mollusca recently added to fauna of southern New England, catalogue of. (P. 1880)
Mollusca, with notes on annelida, echinodermata, etc. (P. 1880) 425
Recent additions to marine invertebrata of northeast coast of America,
with new genera and species. Parts 1, 11, 111. (P. 1879; P. 1880.) 888, 425
VERRILL, A. E.; RATHBUN, R. Marine invertebrata of northeast coast of America distributed by Fish Commission. (P. 1879)833
Vertebrata, chemical and physiological investigations relative to-J. Jones 82
Vespidæ, synopsis of—H. Dr Saussure
Vestiges of antiquity in the United States—S. F. HAVEN 71
Vézère, valley of the, cave-dwellers of the—P. Broca. (R. 1872) 271
Vibratory movement of matter-L. Magrini. (R. 1868) 224
Victoria Royal Society, address of the president—R. L. J. ELLERY. (R. 1868.) 224
Vienna—
Imperial Academy of Sciences of. Prize questions. (R. 1864; R. 1865.) 188, 209
Imperial Geological Institute, tertiary fossils presented by. (R. 1863) 187
Imperial Library of, books presented. (R. 1865) 209
Views and plans of Smithsonian Institution—R. D. Owen
Vineyard Sound, Massachusetts, occurrence of oceanic bonito in—V. N. EDWARDS. (P. 1878)
Virgin Islands, flora of the—H. F. A. EGGERS
Virginia—
Hampton, catalogue of casts of heads of Indian boys and girls at—R. H. PRATT. (P. 1879)
Luray Cave, blind myriapod from-J. A. RYDER. (P. 1880) 425
Luray Cave, report of visit to-O. T. Mason and others. (R. 1880.) 442, 433
Vitality-H. H. HIGGINS; J. HENRY. (R. 1866) 214
Viviparidso-W. G. BINNEY 144
Vocabularies, Indian—G. Gibbs. (R. 1862)
Vocabularies, Indian, received from the Wheeler expedition, list of. (R. 1874.) 286
Vocabulary—
comparative—A. GALLATIN
comparative, English, Spanish, French, Latin 170
Makah-J. G. SWAN 220
of Chinook jargon, or trade language of Oregon-G. GIBBS 161
of Chinook jargon, or trade language of Oregon—B. R. MITCHELL; W. W. TURNER; G. GIBBS

Vocal sounds of Laura Bridgman—F. LIEBER	12
Voices of crustaceans—G. B. Goode. (P. 1878)	832
Volcano-	
in Nicaragua, eruption of—A. B. Dickinson. (R. 1867)	215
of Colima, eruption of—C. SARTORIUS. (R. 1869)	228
of Popocatepetl, examination of—Baron von MÜLLER; A. SONNIG	114
VOLLUM, E. P. On the wingless grasshopper of California. (R. 1860)	147
Volta, Alexander, eulogy on, by F. Arago. (R. 1875)	298
VON BAUMHAUER, E. H. Universal meteorograph for detached observatories. (R. 1879)	
Von Buch, Leopold, memoir of, by M. Flourens. (R. 1862)	
VON HELLWALD, F. The American migration. (R. 1866)	214
VON KAROLYI, Lieut.; CRAIG, B. F. Products of combustion of gun-cotton	
and gunpowder. (R. 1864)	
VON LIEBIG, J. Induction and deduction. (R. 1870)	244
Von Martius, C. F. P., honorary medal to—W. HAIDINGER. (R. 1868)	187
Von Martius, C. F. P., memoir of, by C. RAU. (R. 1869) 228, 440	, 2 51
VON MÜLLER, Baron. Observations on terrestrial magnetism in Mexico	114

∇

WAITE, M. R. Memorial of Joseph Henry	\$56
WALE; HAWKINS. Blow-pipe apparatus. (R. 1872)	
WALKER, D. Observations on Arctic auroras and specific gravity of sea water.	146
WALKER, D. Observations on ice	146
WALKER, S. C. Ephemerides of planet Neptune, 1848, 1846-1849, 1850, 1851, 1852	, 24
	3
(16, 1010)	345
WALKER, S. T. Shell-heaps of Tampa Bay, Florida. (R. 1879)	345
Wall, G. P.; Sawkins, J. G. Report on survey of economic geology of Trinidad. (R. 1856)	91
Wallach, R.; Henry, J. Report of Committee of Regents on the fire at Smitheonian. (R. 1864)	188
Walrus, remains of, in Maine—C. H. Boyd. (P. 1881)	467
Wampum belts of the six nations—W. M. BEAUCHAMP. (R. 1879)	345
War Department, authority given by, to officers of Quartermaster's Department to receive and transmit specimens to Smithsonian	

OF SMITHSONIAN PUBLICATION3.	323
WARD, L. F.—	
Barometer tube breaking suddenly. (R. 1866)	214
Catalogue of collection of Japanese woods presented by University of Tokio, Japan. (P. 1881)	467
Check-list of plants of Washington and vicinity	
Directions for collecting and preserving plants	460
Guide to flora of Washington and vicinity	444
WARING, G. E., Jr. Suggestions for sanitary drainage of Washington city. Toner lecture No. VIII.	849
Warming and ventilating buildings. Parts I, II—A. MORIN. (R. 1878; R.	010
1874) 275, 286,	439
Warming, architecture in relation to—D. B. REID. (R. 1856)	91
WARNER, J. Big elephant mound in Grant county, Wisconsin. (R. 1872)	271
Warren, G. K., report on collections made by—F. B. MEEK; F. V. HAYDEN.	172
WARTMANN, E. Report on transactions of the Society of Physics and Natural	112
History of Geneva, July, 1867, to June, 1868. (R. 1868)	224
Washington, Arkansas, meteorological observations near—N. D. Smith	131
Washington, D. C.—	
canal of, report of Committee of Regentson—R. Delafield. (R. 1868.)	994
flora of, and vicinity, guide to the—L. F. WARD	
notice of, in Harriot's travels	880
notice of, in Isaac Weld's travels	880
prehistoric remains in vicinity of—T. R. PEALE. (R. 1872)	271
• • • • • • • • • • • • • • • • • • • •	
sanitary drainage of. Toner lecture No. VIII—G. E. WARING, Jr	349
Washington Philosophical Society, bulletin of. Vols. I-III; March, 1871, to June, 1880	428
Washington Territory—	220
fishes of—T. H. BEAN. (P. 1881)	467
forest trees in, distribution of—W. W. Johnson. (R. 1870)	244
Indians of—J. G. Swan	220
method of taking surf smelt by Indians of—J. G. SWAN. (P. 1880)	425
Neah Bay, description of two new species of fishes (Ascelicthys rhodorus and Scytalina cerdale) from—D. S. Jordan; C. H. Gilbert. (P.	120
. 1880)	425
Wasps, American solitary, synopsis of—H. DE SAUSSURE	
Wasps, translation of De Saussure's monograph of—E. Norton	254
Water—	
formation of ice at bottom of—M. ENGELHARDT. (R. 1866)	214
from Gulf of Mexico, report on-W. G. FARLOW. (P. 1881)	467
in rivers, diminution of—H. G. WEX. (R. 1875)	298
in rivers, method of ascertaining the amount of-A. A. HUMPHREYS.	
(R. 1858)	109
polluted, in Gulf of Mexico, destruction of fish by—W. C. W. GLAZIER. (P. 1881)	467

Water—Continued.
salt, preservation of copper and iron in—A. R. BECQUEBEL. (R. 1864.)
sea, specific gravity of—D. Walker 146
WATKINS, E. A. System of relationship of Cree Indians. (R. 1862)
petalæ
Weapons, savage, at Centennial Exhibition, study of—E. H. KNIGHT. (R. 1879)
Weather, effect of moon on the-J. Henry. (R. 1871) 249
WEBB, T. W. Account of Henry Draper's telescope. (R. 1864)
Weights, atomic, recalculation of. Constants of Nature, Part v-F. W. CLARKE. 41
Weights, small, method of forming—R. HARE. (R. 1858)
Weights and measures—
English and French, tables of. (R. 1863; R. 1864) 187, 188
metric system of, with tables—H. A. NEWTON. (R. 1865) 209, 371
new system of, with 8 as the metrical number—G. H. KNIGHT. (B. 1867.) 215
WEISMANN, A. Change of Mexican axolotl to amblystoma. (R. 1877) 323, 401
WELD, I. Account of city of Washington, 1807
Welling, J. C. Notes on the life and character of Joseph Henry 856, 338
Welsh, language of Navajos said to resemble—S. Y. McMasters. (R. 1865.) 209
Welsh, J. Construction and verification of standard barometers. (R. 1859.) 110
West Indies—
birds of the—See Lawrence, G. N., Ober, F. A.
list of described birds of the, not in Smithsonian collection185
loxigilla from, description of new subspecies of—G. N. LAWBENCE. (P. 1881)
stone celts in the—G. J. GIBBS. (R. 1877)
West Virginia, occurrence of Canada porcupine in—G. B. Goode. (P. 1878.) 332
Wet and dry bulb thermometer, tables for—J. H. COFFIN 87
WETHERILL, C. M.—
Modern theory of chemical types. (R. 1863)187
Ozone and antozone. (R. 1864) 188, 355
Plan of a research upon the atmosphere. (R. 1866)214
WEX, H. G. Diminution of the water of rivers and streams. (R. 1875) 238
Wheeler expedition, list of Indian vocabularies received from the. (R. 1874.) 286
Wheeler expedition - See Explorations.
Whirlwind at St. Thomas, August 21, 1871—A. Colding. (R. 1877) 823, 398
Wпіте, С. А.—
Description of new cretaceous pinna from New Mexico. (P. 1880) 425
Description of very large fossil gasteropod from State of Puebla, Mexico. (P. 1880)
Descriptions of new cretaceous invertebrate fossils from Kansas and Texas. (P. 1879)

WHITE, C. A.—Continued.	
Descriptions of new invertebrate fossils from the mesozoic and cenozoic	425
Descriptions of new species of carboniferous invertebrate fossils. (P. 1879.)	888
Note on acrothele. (P. 1880)	425
	888
	383
·	425
Note on occurrence of Stricklandia salteri and Stricklandia Davidsoni	425
	4 67
White river canon, Pima county, Arizona, ruins in-R. T. Burn. (R. 1879.)	845
WHITNEY, J. D. Catalogue of rocks, minerals, etc. (R. 1854)	75
	215
WHITNEY, W. D. Lectures on principles of linguistic science. (R. 1863.) 187, 3	152
WHITTLESEY, C	
Ancient mining on shores of Lake Superior1	L 55
Descriptions of ancient works in Ohio	37
On fluctuations of level in North American lakes 1	19
On fresh-water glacial drift of the northwestern States	.97
Wilkes, C., duplicate shells of exploring expedition under1	193
Will of Smithson. (R. 1853)	880
Willemæsia group of crustacea, (recent Eryontidæ,) notice of new species of the—S. I. SMITH. (P. 1879)	88 8
WILLIAMS, H. C. On antiquities in some of the southern States. (R. 1870.)	244
WILLIAMSON, G. Antiquities in Guatemala. (R. 1876)	299
WILLIS, J. R.; BLACKISTON; BLAND. Birds of Nova Scotis. (R. 1858) I	109
	109
WILMOT, S. Notes on western gizzard shad, Dorosoma cepedianum heterurum. (P. 1878)	3 32
Wilson, D. Lectures on physical ethnology. (R. 1862)	150
Wilson, Henry, eulogy on, by P. PARKER. (R. 1875)	298
Wind-	
· ,	214
, , , , , , , , , , , , , , , , , , ,	286
()	214
connection of gales of, and appearance of aurora—R. T. KNIGHT; J. HENRY. (R. 1871)	249
observations of, in the Arctic Seas-I. I. HAYES	196
observations of, in the Arctic Seas—E. K. KANE	104
observations of, in the Arctic Seas-F. L. McCLINTOCK 1	146
•	157
relation between difference of pressure and velocity of -J. HANN. (R.	398

Winds—
in Florida-J. Baltzell. (R. 1866)
of the globe—J. H. Coffin2
of the northern hemisphere—J. H. COFFIN
upper, importance of attention to the—C. DEWEY. (R. 1866)
Winnipeg Lake, egging expedition to-D. Gunn. (R. 1867)
Wirtz, H. R. Herbarium captured in Tennessee. (R. 1862)
Wisconsin—
antiquities in-M. Strong. (R. 1877)
antiquities of—I. LAPHAM
earthworks in—E. E. Breed. (R. 1872)
Grant county, elephant mound in—J. WARNER. (B. 1872)
Grant county, observations on prehistoric mounds of—M. STRONG. (R. 1876)
mound in—C. K. DEAN. (R. 1872)
mounds and osteology of mound-builders of—J. N. DE HART. (R. 1877.)
mounds in—W. G. Andreson. (R. 1879)
salamander of—P. R. Hov. (R. 1854)
Winnebago county, mounds in—T. Armstrong. (R. 1879)
Wise, J., on observation by, of a thunder storm—R. HARE. (R. 1854)
WISEMAN, N. Identification of the artisan and artist. (R. 1870) 2
WISSNER, J. Meteorology of the District of Columbia. (R. 1857)
WITHERS, R. E. Memorial address on Joseph Henry
WOELKOFF, A. Discussion of winds of the globe
,
Wood, ancient implement of, from Connecticut—E. W. ELISWORTH. (R. 1876) 29 Wood, preservation of. (R. 1864)
Wood, H. C.—
Bibliography of alge 2
Contribution to the history of the fresh-water algo of North America 24
Fever. A study in morbid and normal physiology. (R. 1878) 341, 35
Instructions for collecting myriapods, phalangids, etc. (R. 1866) 2
Study of nature and mechanism of fever. Toner lecture No. IV 2
WOOD, J. Meteorological observations at Marietta, Ohio1
Woods of Japan, collection of, presented to U. S. National Museum, catalogue of—L. F. WARD. (P. 1881)
WOODWAED, J. J. Structure of cancerous tumors and mode in which adjacent parts are invaded
WOODWORTH, A. Mcteorite in Mexico. (R. 1867)
WOOLSEY, T. D. Eulogy on C. C. Felton. (R. 1861) 14
Work, relation of food to, and its bearing on medical practice—S. HAUGHTON.

OF SMITHSONIAN PUBLICATIONS.	327
Works published by Smithsonian Institution to January, 1866	208 275
Worsaar, J. J. A. Preservation of antiquities and national monuments in Denmark. (R. 1879)	845
Wounds of poisoned arrows, lecture on nature and cure of—D. BRAINARD. (R. 1854)	75
Wright, C., account of botanical explorations of, in New Mexico and California—A. Grav. (R. 1849)	21
Wright, C., account of plants collected by, in Texas and New Mexico. Parts 1, 11—A. GRAY	2. 42
WRIGHT, D. F. Antiquities of Tennessee. (R. 1874)	286
•	
WRIGHT, G. Account of lightning discharges. (R. 1867)	215
S. Hastings. (R. 1880) 442	426
Writings, scientific, of James Smithson	827
Würdemann, G., notice of, by A. D. BACHE. (R. 1859)	110
WÜRDEMANN, G. Specimens of flamingo, etc., from south Florida. (R. 1860.)	147
WÜRDEMANN, W.; GREEN, J. On filling barometer tubes. (R. 1859)	110
Wyandot government—J. W. Powell. (E. 1879-80)	476
WYMAN, J. Anatomy of nervous system of Rana pipiens	45
Wyoming, new invertebrate fossils from—C. A. WHITE. (P. 1880)	425
X.	
Xantus, John, explorations of, in Mexico—M. Romero. (R. 1862)	150
America, description of—D. S. JORDAN; C. H. GILBERT. (P. 1881)	467
Xerobates, land tortoises of genus—F. W. TRUE. (P. 1881)	467
Xiphister, description of new species of, from Monterey, California—D. S. Jor- DAN; C. H. GILBERT. (P. 1880)	
Xystreurys liolepis, new flounder from Santa Catalina Island, California, description of—D. S. JORDAN; C. H. GILBERT. (P. 1880)	

Y.

Yacht "Fox," meteorological observations made on-F. L. McCLINTOCK	146
Yale College, catalogue of meteorites in—G. J. Brusn. (R. 1868)	224
YARROW, H. C. Study of mortuary customs of the North American Indians. (E. 1879-80)	476
Yellowstone river, prehistoric remains near the—P. W. Norris. (R. 1879)	34 5
Yoruba, Africa—	
country and people of, description of-T. J. Bowen	98
language of, bibliography of	98
language of, grammar and dictionary of-T. J. Bowkn; W. W. TURNER.	98
language of, grammar and dictionary of-Southern Baptist Mission-	
ARY SOCIETY	98
YOUNG, C. B. Translation of Morin on warming and ventilating buildings. (R. 1878; R. 1874)	439
YOUNG, C. B. Translation of Von Baumhauer on a universal meteorograph. (R. 1879)	34 5
Young, Thomas, eulogy on, by F. Arago. (R. 1869)	
Young, W. J. Cloud bursts. (R. 1867)	215
Yu, (Chinese name of jade,) study of—S. BLONDEL. (R. 1876)	
Yucatan, remarks on Cara gigantesca of Yzamal in-A. Schott. (R. 1869)	228
Yukon, Russian America, journey to the-W. W. KIRBY. (R. 1864)	188
Yzamal, Yucatan, remarks on Cara gigantesca of-A. Schott. (R. 1869)	228
	•

Z.

ZANTEDESCHI, F. R. Electricity of induction in aerial strata of atmosphere	
surrounding cloud resolving into rain, etc. (R. 1870)	244
Zapotec remains, notice of—B. MAYER; J. G. SAWKINS	86
ZISGENBALS, H. Schlagintweit ethnographic collection. (R. 1867)	215
Zone of small planets between Mars and Jupiter, lecture on the—E. Loomis. (R. 1854)	75
Zoölogical—	
Gardens, Hamburg. Exchange of specimens. (R. 1867)	215
position of Texas—E. D. Cope	412
Record, list of generic names supplemental to those indexed in—S. H. SCUDDER	470
Zoölogicus, Nomenclator. A list of generic names of recent and fossil animals—S. H. Scudder	470
Zoölogy, list of generic names employed in—S. H. Scudder	470
Zoölogy, record of recent progress in, 1879, 1880-T. Gill. (R. 1880) 442,	431

·			

	•	
·	·	

	·		
	·		

